



# Timeless in animal circadian clocks and beyond

Yao D. Cai  and Joanna C. Chiu 

Department of Entomology and Nematology, College of Agricultural and Environmental Sciences, University of California Davis, CA, USA

## Keywords

cell cycle; circadian clock; DNA replication; *Drosophila timeless*; mammalian *timeless*; seasonal biology; *timeout*

## Correspondence

J. C. Chiu, Department of Entomology and Nematology, College of Agricultural and Environmental Sciences, University of California Davis, One Shields Ave, Davis, CA 95616, USA  
Tel: +1-530-752-1839  
E-mail: jochiu@ucdavis.edu

(Received 1 August 2021, revised 9 October 2021, accepted 26 October 2021)

doi:10.1111/febs.16253

TIMELESS (TIM) was first identified as a molecular cog in the *Drosophila* circadian clock. Almost three decades of investigations have resulted in an insightful model describing the critical role of *Drosophila* TIM (dTIM) in circadian timekeeping in insects, including its function in mediating light entrainment and temperature compensation of the molecular clock. Furthermore, exciting discoveries on its sequence polymorphism and thermosensitive alternative RNA splicing have also established its role in regulating seasonal biology. Although mammalian TIM (mTIM), its mammalian paralog, was first identified as a potential circadian clock component in 1990s due to sequence similarity to dTIM, its role in clock regulation has been more controversial. Mammalian TIM has now been characterized as a DNA replication fork component and has been shown to promote fork progression and participate in cell cycle checkpoint signaling in response to DNA damage. Despite defective circadian rhythms displayed by *mtim* mutants, it remains controversial whether the regulation of circadian clocks by mTIM is direct, especially given the interconnection between the cell cycle and circadian clocks. In this review, we provide a historical perspective on the identification of animal *tim* genes, summarize the roles of TIM proteins in biological timing and genomic stability, and draw parallels between dTIM and mTIM despite apparent functional divergence.

## Introduction

Circadian rhythms are common features in all domains of life and are driven by molecular clockworks [1–6]. Molecular clocks incorporate a range of environmental time cues, such as light–dark and temperature signals, and metabolic signals to orchestrate daily rhythms in physiology and behavior [4,6,7]. This allows organisms to synchronize their biology to their external environment, thereby promoting organismal health and fitness [8–11]. The animal circadian clock is powered by cell-autonomous interlocked transcription–translation feedback loops (TTFLs) [6]. In the primary TTFL in *Drosophila*, which relies heavily on *Drosophila* TIM (dTIM) function, transcription factors CLOCK (CLK)

(ortholog of mammalian CLOCK) and CYCLE (CYC) (ortholog of mammalian BMAL1) are positive elements that heterodimerize and activate the expression of negative elements, PERIOD (PER) (ortholog of mammalian PER1, PER2, and PER3) and dTIM (functionally replaced by CRYPTOCHROMES (CRYs) in mammalian clockworks). In addition to core clock components, CLK-CYC also activates the transcription of other clock-controlled output genes [12–14], often in tissue-specific manner [15–17]. To complete the TTFL, PER, and dTIM form a repressor complex that enters the nucleus in a time-of-day-dependent manner [18–22] to repress CLK-CYC

## Abbreviations

CLK, CLOCK; CRY, CRYPTOCHROME; CYC, CYCLE; dTIM, *Drosophila* TIM; mTIM, mammalian TIM; PER, PERIOD; PTM, posttranslational modification; TIM, TIMELESS; TTFL, transcription–translation feedback loop.

transcription activity [23–25]. This repression is relieved when both PER and dTIM are degraded in a proteasome-dependent manner [26–30]. In addition to its role within the molecular clock, thermosensitive alternative splicing of *dtim* RNA [31–34] and light sensitivity [35–38] of TIM protein are key features that allow dTIM to function at the interface between circadian and seasonal timing.

In the mammalian clock, CRYs replace TIM to partner with PERs to maintain circadian rhythms [39–44]. Whether mammalian TIM (mTIM) is a key component of the mammalian clock has been heavily debated since it was first characterized [45–47]. On the other hand, evidence supporting the role of mTIM in DNA replication and DNA damage response is strong. We will discuss the controversial role of mTIM in timekeeping below.

This review summarizes the various roles played by dTIM in *Drosophila* circadian clocks, in the regulation of seasonal biology, and other non-circadian processes. We will then discuss the circadian and non-circadian functions of mTIM, highlighting data that either support its role in circadian timekeeping or are in conflict with the notion. Finally, we conclude the review by summarizing recent findings on the potential functional parallel between dTIM and mTIM.

## ***Drosophila* TIM plays critical roles in circadian timekeeping**


### ***Drosophila* TIM in the molecular clock**

Circadian timekeeping relies on cycling genes and proteins that maintain a free-running period of approximately 24 h. Investigations to elucidate the inner workings of the molecular clockwork started around 50 years ago, when Konopka and Benzer [48] isolated the first three clock mutants in *Drosophila melanogaster* via genetic screening. The mutations were all located in the same loci, which were later confirmed as the key clock gene, *period* (*per*) [49–53]. Hardin *et al.* [54] suggested that PER may feedback to repress its own mRNA expression to establish molecular oscillations that manifest into behavioral and physiology rhythms. In the next few years, taking advantage of high throughput genetic screening in *Drosophila*, Sehgal *et al.* [55] identified *dtim* as the second clock gene. This gene encodes a protein with novel structure at the time and the only recognizable sequence feature the authors highlighted was a stretch of acidic residues [56]. The arrhythmic PER nuclear localization as well as locomotor activity in *dtim* null mutants has led to the model illustrating how the coordination of *per* and

*dtim* may generate 24-h free-running period via negative feedback: (a) transcriptional activation of *per* and *dtim* in midday due to the absence of nuclear PER; (b) PER and dTIM heterodimerize and enter the nucleus at dusk; (c) increasing amount of nuclear PER blocks *per* and *dtim* mRNA transcription and accumulation at night; (d) nuclear PER and dTIM decline because of inhibited mRNA production and subsequent protein turnover in late night to early morning (Fig. 1) [57]. This model was eventually expanded to incorporate CLK [58,59] and CYC [60] after their characterization, thereby establishing the TTFL model of the *Drosophila* clock.

As a negative component in the molecular clockwork, dTIM does not have intrinsic repression activity. Instead, it is essential in maintaining rhythmic PER expression and activity (Fig. 1). This is strongly supported by observations that PER rhythmic expression and behavioral rhythmicity are abolished in *dtim* null mutant [18] and mutants that are defective in TIM nuclear entry [61,62]. Early studies suggest that dTIM binds to and blocks the cytoplasmic localization domain (CLD) of PER and thus reduces PER cytoplasmic retention [63]. Another study described a mechanism by which dTIM antagonizes the activity of DOUBLETIME (DBT, homolog of mammalian casein kinase 1 delta/epsilon) in inhibiting PER nuclear entry [22]. dTIM also acts as the major cargo recognized by the Importin- $\alpha$ 1 (IMP $\alpha$ 1) nuclear entry machinery, thus transporting PER into the nucleus [64]. Saez *et al.* [61] identified a functional nuclear localization signal (NLS) that is potentially recognized by IMP $\alpha$ 1 (Fig. 2). Once in the nucleus, dTIM appears to be bound to PER constitutively and facilitates PER repression [25,65]. Sun *et al.* [66] suggested that dTIM may act as a scaffold to promote PER-CLK interaction. Alternatively, dTIM may facilitate yet-to-be-characterized CLK kinase(s) [23,24,67] in the PER-dTIM repressor complex to phosphorylate CLK and inactivate transcriptional activity.

dTIM function is extensively regulated by posttranslational modifications (PTMs). Notably, phosphorylation is the best-studied protein modification to achieve dTIM time-of-day specific functions. Casein kinase 2 (CK2) and SHAGGY [SGG, homolog of mammalian glycogen synthase kinase-3 $\beta$  (GSK3 $\beta$ )] have been shown to phosphorylate both PER and dTIM and promote nuclear entry [68–72] (Fig. 1). Interestingly, once in the nucleus, PER-dTIM complexes are subjected to phosphorylation-dependent nuclear export, providing an additional means to control nuclear accumulation [21,67]. Protein phosphatases also participate in regulating PER-dTIM nuclear accumulation [73–




**Fig. 1.** *Drosophila* TIM (dTIM) is a core component of the molecular oscillator. During the day, CLK-CYC heterodimers activate the transcription of rhythmic genes, including *per* and *tim* in the nucleus [6]. In the cytoplasm, dTIM undergoes proteasomal degradation mediated by CRYPTOCHROME (CRY) [35–38] and JETLAG (JET) [28,88] upon light exposure. CULLIN-3 (CUL-3) has also been observed to mediate dTIM degradation in a light-independent manner [27]. Early in the night, SHAGGY (SGG) [68], casein kinase 1 $\alpha$  (CK1 $\alpha$ ) [187], casein kinase 2 (CK2) [69,70], Importin- $\alpha$ 1 (Imp $\alpha$ 1) [64] and phosphatase of regenerating liver-1 (PRL-1) [75] promote nuclear accumulation of PER-dTIM complex. This is antagonized by DOUBLETIME (DBT) [25], protein phosphatase 1 (PP1) [74] and protein phosphatase 2A (PP2A) [73]. Once PER-dTIM complex is in the nucleus, CK2-dependent phosphorylation of dTIM (S1404) inhibits PER-dTIM nuclear export by exportin 1 (XPO1) complex, retaining PER-dTIM complex in the nucleus [67]. At midnight, nuclear PER-dTIM complex interacts with CLK-CYC and represses their transcriptional activity [23,25]. From late night to early morning, CRY and JET mediate light-dependent TIM degradation [28,88], whereas DBT and SUPERNUMERARY LIMBS (SLIMB) mediate PER degradation [26,29]. There have also been reports suggesting the involvement of SLIMB in TIM degradation [27].

75]. Over the past 10 years, site-specific functions of dTIM phosphorylation have been characterized in a few studies (Fig. 2). *In vivo* functional analysis leveraging mutagenesis of dTIM protein revealed that T113 is critical for rhythmic dTIM expression [62]. Mutating T113 to non-phosphorylatable alanine (A) abolishes dTIM nuclear entry, whereas mutations at a nearby proline (P115) produce similar defects. Combining genetic and biochemical studies, Top *et al.* [72] showed that SGG and CK2 phosphorylate five residues at ST region (S297, T301, T305, S309, and S313) to promote dTIM nuclear accumulation. Interestingly, SGG and CK2 appear to regulate PER-dTIM only in a subset of clock neurons, which may contribute to the divergent functions of specific neuronal groups within the circadian neuronal circuitry. This could potentially explain how alteration in TIM phosphorylation in flies

carrying *tim<sup>blind</sup>* allele (A1128V, L1131M) results in lengthened locomotor activity rhythms but normal eclosion rhythms [76]. Activity and eclosion rhythms are two well characterized output of the *Drosophila* clock and are normally altered to the same extent in most fly mutants, including the three *per* mutants Konopka and Benzer identified in 1971 [48]. The mechanisms by which kinases phosphorylate PER-dTIM in specific neurons remain unclear. Since alternative pre-mRNA splicing patterns were observed in different clock neurons including for *sgg* mRNAs [77], we speculate that this may result in cell-type-specific posttranslational modification programs for key clock proteins, including dTIM.

Recently, two studies harnessed mass spectrometry proteomics to identify dTIM phosphorylation sites [67,75] (Fig. 2). Kula-Eversole *et al.* [75] identified five dTIM



**Fig. 2.** Schematic illustrating domain structure of TIM isoforms generated from alternative splicing. All amino acid numbering is based on the TIM-L<sub>1421</sub> isoform. ‘TIM-S start’ denotes alternative translation start site for TIM-S. Previously described domains of TIM: 32 amino acid region (amino acid [aa] 260–291) [188], also known as serine-rich domain (SRD) (aa 260–292) [71]; serine/threonine (ST)-rich region (aa 293–312) [72]; a stretch of acidic amino acid residues (acidic aa) (aa 383–412) [56]; PER binding domain 1 (PER BD1) (aa 536–610) [61]; nuclear localization sequence (NLS) (aa 558–593) [61]; C-terminal tail-like sequence (CTL) (aa 640–649) [87]; PER binding domain 2 (PER BD2) (aa 747–946) [61]; and cytoplasmic localization domain (CLD) (aa 1261–1421) [61]. P = phosphorylation sites [62,67,72,75]. Phosphorylation sites in black = identified via mass spectrometry; blue = identified via *in vivo* functional analysis but have not been validated by mass spectrometry or phospho-specific antibodies. TIM-cold, TIM-SC, TIM-M isoforms are based on Shakhmantsir *et al.*, Foley *et al.*, Martin Anduaga *et al.* [31–33].

phosphorylation sites in *Drosophila* S2R<sup>+</sup> cells coexpressing dTIM and relevant kinases (SGG and CK2). S586 and T991 are shown to be dephosphorylated by Phosphatase of Regenerating Liver-1 (PRL-1), which in turn promotes dTIM nuclear accumulation. In Cai *et al.* [67], we identified 12 phosphorylation sites in PER-bound dTIM from *Drosophila* tissues. In particular, we showed that S1404 phosphorylation inhibits the interaction between dTIM and the nuclear export complex, thereby promoting dTIM nuclear accumulation. S1404 phosphorylation status in fly tissues was confirmed using phospho-specific antibody.

In addition to nuclear accumulation, phosphorylation also regulates dTIM protein turnover. CULLIN-3 (CUL-3) and SKP1-CUL1-F-box-protein/SUPERNUMERARY LIMB complex (SCF/SLIMB) differentially facilitates dTIM degradation depending on its phosphorylation status [27,78], thus fine-tuning dTIM phase-specific functions (Fig. 1). Besides phosphorylation, O-GlcNAcylation at multiple residues on dTIM was also identified [67]. Since O-GlcNAcylation modifies serine/threonine residues and regulates the function of many proteins including PER and CLK [79–82], it will be interesting to determine how the two types of PTMs coordinate to regulate dTIM phase-specific functions. Given O-GlcNAcylation is nutrient-

sensitive, this could be a mechanism by which metabolic signals can integrate with time-of-day environmental signals to promote robust circadian rhythms.

Finally, besides PTMs, *dtim* expression is regulated by posttranscriptional mechanisms. Carbon catabolite repression-negative on TATA-less deadenylation complex (CCR4-NOT) has been shown to regulate *dtim* mRNA stability to support phase-specific dTIM function [83]. *Drosophila tim* also exhibits alternative splicing pattern in response to environmental conditions, which will be described later.

### ***Drosophila* TIM and light entrainment of circadian rhythms**

To confer fitness, a circadian clock must be synchronized to local time. Environmental time cues such as daily light–dark or temperature cycles entrain the circadian clock [84]. Identification of clock genes paved the way to investigations on molecular components that mediate clock entrainment. Two years after the identification of *dtim* in 1994, four exciting papers showed that dTIM displays light sensitivity, thus coupling the molecular clockwork to photic input from the environment [35–38] (Fig. 1). CRY is the major photoreceptor that mediates TIM light-dependent

degradation [85–87]. Light induces CRY conformational change, thus enabling CRY to bind to dTIM. Thereafter, E3 ubiquitin ligase JETLAG (JET) along with CRY promotes rapid TIM proteasomal degradation [28,87,88] upon yet uncharacterized TIM tyrosine phosphorylation [89]. QUASIMODO (QSM), a light-responsive protein expressing predominantly in CRY-negative clock neurons, also trigger dTIM degradation upon light exposure [90]. dTIM degradation promotes PER turnover, thus resetting the circadian clock [37].

### ***Drosophila* TIM and temperature compensation of the circadian clock**

Whereas rates of chemical reactions are often temperature-dependent on a molecular level, a clock is only meaningful if its period length stays constant over a wide range of temperatures. The circadian clock has the property of temperature compensation; its pace is stable over a wide range of temperatures [84]. PER was first identified to participate in this process. A repetitive threonine-glycine (Thr-Gly) tract in PER exhibits more flexible conformation in higher temperature [91], which correlates with the observation that flies expressing PER with a deletion in the Thr-Gly tract display impaired temperature compensation of the circadian clock [92]. In wild *D. melanogaster* populations, the Thr-Gly tract is polymorphic in length; this is adaptive and enables flies to maintain the pace of the clock in environments with different range of temperatures [93].


*dtim* has also been demonstrated to contribute to temperature compensation of the clock. At the posttranscriptional level, manipulating *dtim* thermosensitive splicing results in defective temperature compensation [32,33]. Elucidating the function of each *dtim* isoform under different temperatures could help understand how they regulate temperature compensation in future studies. At the posttranslational level, mutant lines bearing a number of amino acid substitutions, *tim<sup>rit</sup>* (P1116A) and *tim<sup>blind</sup>*, exhibit impaired temperature compensation [94,95]. The mechanism by which dTIM regulates temperature compensation remains unclear. One possibility is that temperature directly modulates PER-dTIM interaction. Another possibility is that temperature may indirectly modulate site-specific phosphorylation to regulate phase-specific functions of PER-dTIM and achieve temperature compensation. In mammalian systems, temperature has been shown to determine the priority of competing phosphorylation sites to regulate PER2 turnover rate [96,97]. Therefore, mass spectrometry-based phosphorylation site mapping in combination with molecular genetics may further expand our understanding of how dTIM phosphorylation confers temperature compensation in flies.

### **Sequence polymorphism and alternative splicing of *Drosophila tim* regulates seasonal biology**

To prepare for seasonal changes, plants and animals rely on internal photoperiodic timers, allowing them to undergo physiological and behavioral changes to survive unfavorable times [98–100]. Genetic analysis of wild *D. melanogaster* populations as well as molecular studies revealed that polymorphism at the *dtim* locus facilitates seasonal adaptation (Fig. 3A). *ls-tim* is a derived *dtim* allele that evolved 300–3000 years ago in Europe [101] and has a G nucleotide insertion upstream of the original ATG translational start site [102,103]. This generates an extra ATG 23 amino acids upstream of the TIM-S start codon. *ls-tim* allele thus generates two protein isoforms: TIM-S and a 23-aa longer TIM-L (Fig. 2) (TIM-S and TIM-L were originally named S-TIM and L-TIM but we are renaming them to follow the convention used in more recent publications describing other TIM protein isoforms resulting from alternative pre-mRNA splicing). TIM-L displays reduced light sensitivity, largely due to its reduced binding affinity to CRY [88]. Since light-dependent degradation of dTIM is critical to the resetting of the clock, reduced light sensitivity is thought to keep the molecular clockwork rhythmic in long summer days [104]. Furthermore, in anticipation of the onset of winter, flies carrying *ls-tim* alleles enter reproductive dormancy earlier in autumn as compared with flies carrying only *s-tim* alleles [103]. This is expected to be adaptive for flies inhabiting higher latitudes where harsh conditions are common in winter. For this reason, it was surprising that Tauber *et al.* [103] initially found the highest *ls-tim* allele frequency in southeastern Italy and decrease of *ls-tim* as the sampling distance increases both northward and southward. Subsequent analysis now suggests that this derived allele is only 300–3000 years old; it is still under selection and has not yet achieved fixation [101]. In fact, more extensive sampling in Spain [101] and in North America [105] reported a strong latitudinal cline where *ls-tim* allele increases in frequency as latitude increases.

In addition to sequence polymorphism at the *dtim* locus, *dtim* displays thermosensitive alternative splicing. This has been proposed to be a temperature sensing mechanism to regulate *D. melanogaster* seasonal biology. In response to temperature changes, *dtim* produces four splice variants: *tim-cold*, *tim-short* and *cold* (*tim-sc*), *tim-M* (also called *tim-tiny*), and *tim-L* (full-length isoform) (Fig. 2). At moderate temperature (25 °C), constitutively spliced *tim-L* is the major





**Fig. 3.** Role of *Drosophila* TIM in regulating seasonal biology. (A) Flies carrying *s-tim* allele express TIM-S, whereas flies carrying *Is-tim* allele express both TIM-L and TIM-S. Sampling of flies in North America [105] and on the eastern side of the Iberian Peninsula [101] showed that *Is-tim* allele frequency exhibits a latitudinal cline and increases with latitude. Since TIM-L is less susceptible to light-activated CRY-dependent degradation, flies carrying *Is-tim* allele interpret light signal differently and have higher inducibility of reproductive dormancy at the onset of winter to survive harsh conditions [103]. (B) High temperature promotes accumulation of TIM-M isoform [31]. TIM-L is the major isoform at warm temperature [32]. Cold temperature promotes the accumulation of TIM-SC and TIM-cold isoforms [32,33]. TIM-SC can potentially stabilize EYES ABSENT (EYA) to promote reproductive dormancy [34].

isoform and produces full-length TIM [32] (Fig. 3B). *tim-cold* and *tim-sc* are major isoforms in colder temperatures (10–18 °C) [32–34,106], whereas *tim-tiny* intron is retained in higher temperatures, resulting in high levels of *tim-M* isoform (29–35 °C) [31,32,107]. Thermosensitive alternative splicing is also observed in three other *Drosophila* species, indicating this could be a conserved mechanism across the genus [32]. Less is known regarding the functional divergence of each *dtim* splice variant and how the pattern of splicing modulates the circadian clock in different seasonal conditions. Since some splicing events generate truncated TIM proteins, they could differentially affect TIM function in the circadian clock. For example, the TIM-SC protein lacks the C-terminal CLD and part of PER-binding domain, which may compromise nuclear accumulation of the PER-dTIM complex. Further functional studies on TIM isoforms are required to test this hypothesis.

There has been a substantial amount of evidence to support the role of *dtim* in regulating seasonal biology in addition to the studies mentioned above. They include the observed correlation between *tim* alleles

and photoperiodic diapause in *D. triauraria* [108], changes in *tim* expression levels in response to photoperiod in several insect species [109,110], and differential photosensitive alternative splicing of *tim* observed in cold-adapted *D. montana* populations collected in a wide latitudinal range [111]. We recently provided evidence supporting the role of dTIM in seasonal physiology in *D. melanogaster* [34] (Fig. 3B). We showed that *dtim* null mutants fail to enter reproductive dormancy in simulated winter condition, while flies overexpressing *dtim* exhibit higher incidence of reproductive dormancy. We report evidence indicating that the cold-induced and light-insensitive isoform TIM-SC facilitates the accumulation of EYES ABSENT (EYA) protein in winter condition, an event that is sufficient to promote reproductive dormancy. It remains unclear why TIM-SC is not subjected to light-dependent degradation and how it interacts with EYA. One possibility is that the truncated protein reduces the binding affinity to CRY and/or JET, and somehow stabilizes EYA via yet unknown mechanisms. A temperature-dependent alternative splicing event is also observed in *frequency* (*frq*), a key repressor in the

*Neurospora* clockwork [112–114]. It is possible that this temperature-regulated event also contributes to *Neurospora* seasonal adaptation.

What is the mechanism by which temperature regulates *dtim* alternative splicing? So far, splicing regulator P-element somatic inhibitor (PSI) [33] and triple small nuclear ribonucleoprotein (tri-snRNP) spliceosome [31] have been shown to regulate *dtim* splicing. Temperature is known to modulate alternative splicing at multiple levels, including the expression of splicing-related genes [115,116], PTMs [117], spliceosome assembly [118], and spliceosome localization [119,120].

### Non-circadian roles of *Drosophila* TIM

The fact that dTIM is expressed and differentially regulated in non-clock cells has led to the investigation of non-circadian roles of dTIM. A few studies revealed unexpected results regarding dTIM circadian expression pattern and light sensitivity in non-clock cells. dTIM and its binding partner PER remain constitutively cytoplasmic in the fly ovary, which is known to lack intracellular molecular clocks [121–123]. This is unlike the subcellular shuttling of PER-dTIM observed consistently in clock neurons. Furthermore, dTIM in the follicle cells is not susceptible to light-induced degradation [123,124]. It is noteworthy that egg-laying rhythms persist under constant light, in contrast to the arrhythmic eclosion and locomotor activity rhythms in the same condition [125]. Whether the peculiar PER-dTIM behavior in ovaries relates to rhythmic egg laying under constant light remains unclear. Although *dtim* null mutants display reduced fitness in terms of female fertility and fecundity [123], it has been proposed that this is likely due to the overall loss of the circadian clock [11]. To examine non-circadian roles of *dtim*, it is necessary to manipulate *dtim* specifically in target cells/tissues. One possibility is that *dtim* expressed in non-clock cells has a residual role in maintaining chromosome integrity inferred from its ancestral paralog dTIMEOUT, the homolog of mTIM [126] (Fig. 4A). The non-circadian function of mTIM will be discussed below.

### Debate on mammalian TIM function in circadian timekeeping

#### Evidence supporting the role of mammalian TIM in the circadian clock

Whether mTIM is a core component in the mammalian clock has been controversial. Due to their sequence similarity, mTIM was first identified as the


homolog of dTIM in late 1990s [127–130]. Because of its rhythmic mRNA expression in the mammalian brain [127,131] and physical interaction to core clock proteins mPER1/2/3 [130,132] and CRY1/2 [133–136], mTIM was implicated as a clock protein. In addition, short-term mTIM knockdown causes phase resetting, whereas long-term knockdown of mTIM disrupts circadian neuronal activity rhythms [132]. Recently, Kurien *et al.* [137] reported a mutation in human TIM (hTIM) that causes familial advanced sleep phase syndrome (FASPS), reviving the discussion of the potential role of mTIM in mammalian clockworks. This mutation inhibits TIM nuclear accumulation and destabilizes PER/CRY2 repressor complex at the molecular level.

#### Evidence contradicting a direct role of mammalian TIM in regulating circadian rhythms

Multiple lines of evidence argue against a direct role of mTIM in the molecular clock. Homozygous *mTim* mutant mice are lethal in embryonic stage, whereas other homozygous clock mutants remain viable, suggesting a critical non-circadian role of mTIM [45]. The binding of mTIM to CRY1/2 does not necessarily support a circadian role of mTIM given that CRY1/2 also participates in non-circadian processes. CRY1 and CRY2 are known to modulate DNA damage response [138] and cell proliferation [139], and the interaction of mTIM-CRY1 and mTIM-CRY2 are critical for checkpoint activation [140,141]. Furthermore, phylogenetic analysis revealed that mTIM is an ortholog of dTIMEOUT [142]. *Drosophila* TIMEOUT is the widely conserved ancestral paralog of dTIM among eukaryotes that originated from gene duplication at the time of Cambrian Explosion [45,46,143]. Unlike dTIM, dTIMEOUT is an essential gene in *Drosophila* development and maintenance of chromosome integrity [126].

#### Non-circadian roles of mammalian TIM

There have been extensive investigations focusing on non-circadian roles of mTIM (Fig. 4B,C). Similar to its yeast homolog topoisomerase 1-associated factor 1 (*tof1*) [144], mTIM and its evolutionally conserved partner Tim-interacting protein (TIPIN) maintain replisome stability [145,146] and promote fork progression through hard-to-replicate regions [147–151]. In response to DNA damage, mTIM collaborates with cardinal signaling kinases ataxia telangiectasia-mutated checkpoint kinase 1 (ATR-CHK1) [140,152], ataxia telangiectasia and Rad3-related checkpoint kinase 2 (ATM-CHK2) [153], and poly [ADP-ribose]




**Fig. 4.** *Drosophila* TIMEOUT and mammalian TIMELESS in genome maintenance. (A) *Drosophila* TIMEOUT interacts with Ataxia telangiectasia and Rad3-related (ATR) (genetically) to maintain genomic stability [126]. (B) mTIM and Tim-interacting protein (TIPIN) couple replicative DNA helicase CMG (CDC45, MCM2-7, GINS) and DNA polymerase (Pol) [145,146] in progressing replication fork. (C) In response to DNA damage, mTIM physically interacts with and recruits poly [ADP-ribose] polymerase 1 (PARP1) to damaged sites [153,154]. ATR and ataxia telangiectasia mutated (ATM) can both sense DNA damage and phosphorylate checkpoint kinase 1/2 (CHK1/2) [189]. This is dependent on a number of partner proteins including mTIM [140,152,153].

polymerase 1 (PARP1) [154,155] to facilitate proper checkpoint control and DNA repair [156–158]. Because of its role in genome maintenance, it is not surprising that mTIM dysregulation is commonly found in many cancer types [153,159,160]. Specifically, mTIM promotes cancer development by protecting cancer cells from replication stress and cell cycle arrest [153,161,162]. Thus, mTIM appears to be a promising target for anticancer treatment. However, given its ability to influence the circadian clock, the side effect of clock disruption needs to be considered, as clock disruption has been linked to increased risks of many diseases including metabolic disorders and cancers [163,164].

Considering the role of mTIM discussed in this section, it is noteworthy that the period shortening phenotype on the molecular clock resulting from the

mTIM(R1081X) mutation is limited to proliferative cells [137]. Since the circadian clock ticks regardless of cell proliferation status, why was the period shortening phenotype only observed in proliferating cells? We speculate that mTIM modulates the circadian clock through its role in other cellular processes occurring only in proliferating cells. Specifically, its elevated expression in proliferative tissues such as spleen and thymus are consistent with its cell cycle-related function [137,165]. DNA damage has been shown to induce a circadian phase shift [166–168], with mTIM downregulation attenuating this effect [165]. Interestingly, the FASPS mutation found in hTIM lacks the C-terminal domain critical for mTIM-mediated DNA repair and checkpoint activation through replication stress response regulator SDE2 and PARP1 binding, respectively [153,154,162]. Taken together, it is





**Fig. 5.** Functional parallel between *Drosophila* and mammalian TIM. *Drosophila* TIM(S1404A) elevates PER-dTIM nuclear export [67]. The reduced abundance of nuclear PER-dTIM repressor complex leads to altered phosphorylation status of CLK and transcriptional activity of CLK-CYC, resulting in advanced behavioral rhythms. Mammalian TIM (R1081X) results in reduced nuclear mTIM [137], similar to the phenotype observed in dTIM(S1404A). This promotes destabilization of PER2-CRY2 repressor complex, thus altering transcriptional activity of CLOCK-BMAL1 and resulting in advanced sleep phase syndrome. Phosphorylation status of CLOCK or BMAL1 was not examined in [137].

plausible that the period shortening effect in proliferating cells can be attributed to a non-circadian role of mTIM.

Despite functional divergence of mTIM and dTIM, there are still some parallels. *Drosophila* TIMEOUT is expressed in the optic lobe of adult *Drosophila* and contributes to light entrainment, analogous to light sensitivity of dTIM [126]. Decreased dTIM and mTIM nuclear accumulation in *Drosophila* and mammals respectively both lead to similar outcome in circadian rhythms at the molecular and behavioral levels [67,137] (Fig. 5). This highlights an unexpected functional parallel between mTIM and dTIM in circadian regulation.

## Conclusion and perspectives

The very name of the *timeless* gene hints at its critical function in biological timing. Since its discovery, almost three decades ago in *D. melanogaster*, a large body of work have uncovered the role of dTIM as a cardinal clock protein necessary to maintain circadian timekeeping, mediate light entrainment, and modulate temperature compensation. Thermosensitive splicing of *tim* mRNA in combination with the light sensitivity of dTIM protein enables its role in regulating seasonal physiology. Its ancestral paralog *timeout* (mTIM in mammals) surprisingly plays a distinct role in the maintenance of genomic stability. An important unanswered question regarding the role of dTIM in biological rhythms is how splice variants affect dTIM protein function in response to thermal and photic

cues. The answer would clarify how the circadian clock interplays with seasonal timing. Another area of interest is to elucidate how mTIM regulates the molecular clockwork and potentially sits at the intersection between circadian clocks and cell cycle regulation. This would further shed light on the functional similarity and divergence of the two TIM paralogs. More importantly, this would extend our understanding of the interconnection between the circadian clock and the cell cycle. Circadian regulation of the cell cycle has been found in all domains of life [169–178], and the cell cycle also influences the phase and amplitude of circadian rhythms [166,179,180]. Given the accumulating evidence on circadian regulation of the cell cycle in the context of cancer and tissue regeneration upon injury [181–186], understanding the interaction of the circadian clock and the cell cycle could pave the way for innovative therapeutics for cancer and improved recovery of patients who suffered injuries.

## Acknowledgements

We thank Gary K. Chow for critical reading of the manuscript. This work was funded by NIH R01 DK124068 to JCC. We apologize to any colleagues whose works were not included in this review owing to space limitations.

## Conflict of interest

The authors declare no conflict of interest.

## Author contributions

YDC wrote the initial draft of the manuscript with input from JCC. JCC edited the manuscript for submission.

## References

- Johnson CH, Zhao C, Xu Y & Mori T (2017) Timing the day: what makes bacteria clocks tick? *Nat Rev Microbiol* **15**, 232–242.
- Swan JA, Golden SS, LiWang A & Partch CL (2018) Structure, function, and mechanism of the core circadian clock in cyanobacteria. *J Biol Chem* **293**, 5026–5034.
- Cox KH & Takahashi JS (2019) Circadian clock genes and the transcriptional architecture of the clock mechanism. *J Mol Endocrinol* **63**, R93–R102.
- Creux N & Harmer S (2019) Circadian rhythms in plants. *Cold Spring Harb Perspect Biol* **11**, a034611.
- Dunlap JC & Loros JJ (2017) Making time: conservation of biological clocks from fungi to animals. *Microbiol Spectr* **5**, 10.
- Patke A, Young MW & Axelrod S (2020) Molecular mechanisms and physiological importance of circadian rhythms. *Nat Rev Mol Cell Biol* **21**, 67–84.
- Steed G, Ramirez DC, Hannah MA & Webb AAR (2021) Chronoculture, harnessing the circadian clock to improve crop yield and sustainability. *Science* **372**, eabc9141.
- Ouyang Y, Andersson CR, Kondo T, Golden SS & Johnson CH (1998) Resonating circadian clocks enhance fitness in cyanobacteria. *Proc Natl Acad Sci USA* **95**, 8660–8664.
- Woelfle MA, Ouyang Y, Phanvijhitsiri K & Johnson CH (2004) The adaptive value of circadian clocks: an experimental assessment in cyanobacteria. *Curr Biol* **14**, 1481–1486.
- Dodd AN, Salathia N, Hall A, Kévei E, Tóth R, Nagy F, Hibberd JM, Millar AJ & Webb AAR (2005) Plant circadian clocks increase photosynthesis, growth, survival and competitive advantage. *Science* **309**, 630–633.
- Horn M, Mitesser O, Hovestadt T, Yoshi T, Rieger D & Helfrich-Föster C (2019) The circadian clock improves fitness in the fruit fly, *Drosophila melanogaster*. *Front Physiol* **10**, 1374.
- Claridge-Chang A, Wijnen H, Naef F, Boothroyd C, Rajewsky N & Young MW (2001) Circadian regulation of gene expression systems in the *Drosophila* head. *Neuron* **32**, 657–671.
- McDonald MJ & Rosbash M (2001) Microarray analysis and organization of circadian gene expression in *Drosophila*. *Cell* **107**, 567–578.
- Abruzzi KC, Rodriguez J, Mente JS, Desrochers J, Zadina A, Luo W, Tkachev S & Rosbash M (2011) *Drosophila* CLOCK target gene characterization: implications for circadian tissue-specific gene expression. *Genes Dev* **25**, 2374–2386.
- Zhang R, Lahens NF, Ballance HI, Hughes ME & Hogenesch JB (2014) A circadian gene expression atlas in mammals: implications for biology and medicine. *Proc Natl Acad Sci USA* **111**, 16219–16224.
- Mure LS, Le HD, Benegiamo G, Chang MW, Rios L, Jillani N, Ngotho M, Kariuki T, Dkhisssi-Benyahya O, Cooper HM *et al.* (2018) Diurnal transcriptome atlas of a primate across major neural and peripheral tissues. *Science* **359**, eaao0318.
- Beytebiere JR, Trott AJ, Greenwell BJ, Osborne CA, Vitet H, Spence J, Yoo SH, Chen Z, Takahashi JS, Ghaffari N *et al.* (2019) Tissue-specific BMAL1 cistromes reveal that rhythmic transcription is associated with rhythmic enhancer-enhancer interactions. *Genes Dev* **33**, 294–309.
- Vosshall LB, Price JL, Sehgal A, Saez L & Young MW (1994) Block in nuclear localization of *period* protein by a second clock mutation, *timeless*. *Science* **263**, 1606–1609.
- Price JL, Dembinska ME, Young MW & Rosbash M (1995) Suppression of PERIOD protein abundance and circadian cycling by the *Drosophila* clock mutation *timeless*. *EMBO J* **14**, 4044–4049.
- Price JL, Blau J, Rothenfluh A, Abodeely M, Kloss B & Young MW (1998) *double-time* is a novel *Drosophila* clock gene that regulates PERIOD protein accumulation. *Cell* **94**, 83–95.
- Ashmore L, Sathyanarayanan S, Silvestre DW, Emerson MM, Schotland P & Sehgal A (2003) Novel insights into the regulation of the *timeless* protein. *J Neurosci* **23**, 7810–7819.
- Cyran SA, Yiannoulos G, Buchsbaum AM, Saez L & Young MW (2005) The *double-time* protein kinase regulates the subcellular localization of the *Drosophila* clock protein *period*. *J Neurosci* **25**, 5430–5437.
- Yu W, Zheng H, Houl JH, Dauwalder B & Hardin PE (2006) PER-dependent rhythms in CLK phosphorylation and E-box binding regulate circadian transcription. *Genes Dev* **20**, 723–733.
- Yu W, Zheng H, Price JL & Hardin PE (2009) DOUBLETIME plays a noncatalytic role to mediate CLOCK phosphorylation and repress CLOCK-dependent transcription within the *Drosophila* circadian clock. *Mol Cell Biol* **29**, 1452–1458.
- Menet JS, Abruzzi KC, Desrochers J, Rodriguez J & Rosbash M (2010) Dynamic PER repression mechanisms in the *Drosophila* circadian clock: from on-DNA to off-DNA. *Genes Dev* **24**, 358–367.
- Grima B, Lamouroux A, Chélot E, Papin C, Limbourg-Bouchon B & Rouyer F (2002) The F-box protein Slimb controls the levels of clock proteins Period and Timeless. *Nature* **420**, 178–182.

- 27 Grima B, Dognon A, Lamouroux A, Chélot E & Rouyer F (2012) CULLIN-3 controls TIMELESS oscillations in the *Drosophila* circadian clock. *PLoS Biol* **10**, e1001367.
- 28 Koh K, Zheng X & Sehgal A (2006) JETLAG resets the *Drosophila* circadian clock by promoting light-induced degradation of TIMELESS. *Science* **312**, 1809–1812.
- 29 Chiu JC, Vanselow JT, Kramer A & Edery I (2008) The phospho-occupancy of an atypical SLIMB-binding site on PERIOD that is phosphorylated by DOUBLETIME controls the pace of the clock. *Genes Dev* **22**, 1758–1772.
- 30 Chiu JC, Ko KW & Edery I (2011) NEMO/NLK phosphorylates PERIOD to initiate a time-delay phosphorylation circuit that sets circadian clock speed. *Cell* **145**, 357–370.
- 31 Shakhmantsir I, Nayak S, Grant GR & Sehgal A (2018) Spliceosome factors target *timeless (tim)* mRNA to control clock protein accumulation and circadian behavior in *Drosophila*. *eLife* **7**, e39821.
- 32 Martin Anduaga A, Evantal N, Patop IL, Bartok O, Weiss R & Kadener S (2019) Thermosensitive alternative splicing senses and mediates temperature adaptation in *Drosophila*. *eLife* **8**, e44642.
- 33 Foley LE, Ling J, Joshi R, Evantal N, Kadener S & Emery P (2019) *Drosophila* PSI controls circadian period and the phase of circadian behavior under temperature cycle via *tim* splicing. *eLife* **8**, e50063.
- 34 Abrieux A, Xue Y, Cai Y, Lewald KM, Nguyen HN, Zhang Y & Chiu JC (2020) EYES ABSENT and TIMELESS integrate photoperiodic and temperature cues to regulate seasonal physiology in *Drosophila*. *Proc Natl Acad Sci USA* **117**, 15293–15304.
- 35 Lee C, Parikh V, Itsukaichi T, Bae K & Edery I (1996) Resetting the *Drosophila* clock by photic regulation of PER and a PER-TIM complex. *Science* **271**, 1740–1744.
- 36 Hunter-Ensor M, Ousley A & Sehgal A (1996) Regulation of the *Drosophila* protein *timeless* suggests a mechanism for resetting the circadian clock by light. *Cell* **84**, 677–685.
- 37 Zeng H, Qian Z, Myers MP & Rosbash M (1996) A light-entrainment mechanism for the *Drosophila* circadian clock. *Nature* **380**, 129–135.
- 38 Myers MP, Wager-Smith K, Rothenfluh-Hilfiker A & Young MW (1996) Light-induced degradation of TIMELESS and entrainment of the *Drosophila* circadian clock. *Science* **271**, 1736–1740.
- 39 Miyazaki K, Mesaki M & Ishida N (2001) Nuclear entry mechanism of rat PER2 (rPER2): role of rPER2 in nuclear localization of CRY protein. *Mol Cell Biol* **21**, 6651–6659.
- 40 Yagita K, Tamanini F, Yasuda M, Hoeijmakers JHJ, van der Horst GTJ & Okamura H (2002) Nuclearcytoplasmic shuttling and mCRY-dependent inhibition of ubiquitylation of the mPER2 clock protein. *EMBO J* **21**, 1301–1314.
- 41 Busino L, Bassermann F, Maiolica A, Lee C, Nolan PM, Godinho SIH, Draetta GF & Pagano M (2007) SCF<sup>Fbx13</sup> controls the oscillation of the circadian clock by directing the degradation of Cryptochrome proteins. *Science* **316**, 900–904.
- 42 Godinho SIH, Maywood ES, Shaw L, Tucci V, Barnard AR, Busino L, Pagano M, Kendall R, Quwailid MM, Romero MR *et al.* (2007) The *after-hours* mutant reveals a role for Fbx13 in determining mammalian circadian period. *Science* **316**, 897–900.
- 43 Siepka SM, Yoo S-H, Park J, Song W, Kumar V, Hu Y, Lee C & Takahashi JS (2007) Circadian mutant *Overtime* reveals F-box protein FBXL3 regulation of *Cryptochrome* and *Period* gene expression. *Cell* **129**, 1011–1023.
- 44 Brenna A, Olejniczak I, Chavan R, Ripperger JA, Langmesser S, Cameroni E, Hu Z, Virgilio CD, Dengele J & Albrecht U (2019) Cyclin-dependent kinase 5 (CDK5) regulates the circadian clock. *eLife* **8**, e50925.
- 45 Gotter AL, Manganaro T, Weaver DR, Kolakowski LF, Possidente B, Sriram S, MacLaughlin DT & Reppert SM (2000) A time-less function for mouse *timeless*. *Nat Neurosci* **3**, 755–756.
- 46 Gotter AL (2006) A *Timeless* debate: resolving TIM's noncircadian roles with possible clock function. *NeuroReport* **17**, 1229–1233.
- 47 Mazzoccoli G, Laukkanen MO, Vinciguerra M, Colangelo T & Colantuoni V (2016) A *Timeless* link between circadian patterns and disease. *Trends Mol Med* **22**, 68–81.
- 48 Konopka RJ & Benzer S (1971) Clock mutants of *Drosophila melanogaster*. *Proc Natl Acad Sci USA* **68**, 2112–2116.
- 49 Bargiello TA & Young MW (1984) Molecular genetics of a biological clock in *Drosophila*. *Proc Natl Acad Sci USA* **81**, 2142–2146.
- 50 Hamblen M, Zehring WA, Kyriacou CP, Reddy P, Yu Q, Wheeler DA, Zwiebel LJ, Konopka RJ, Rosbash M & Hall JC (1986) Germ-line transformation involving DNA from the *period* locus in *Drosophila melanogaster*: overlapping genomic fragments that restore circadian and ultradian rhythmicity to *per<sup>0</sup>* and *per<sup>-</sup>* mutants. *J Neurogenet* **3**, 249–291.
- 51 Baylies MK, Bargiello TA, Jackson FR & Young MW (1987) Changes in abundance or structure of the *per* gene product can alter periodicity of the *Drosophila* clock. *Nature* **326**, 390–392.
- 52 Yu Q, Colot HV, Kyriacou CP, Hall JC & Rosbash M (1987) Behaviour modification by *in vitro* mutagenesis of a variable region within the period gene of *Drosophila*. *Nature* **326**, 765–769.

- 53 Lorenz LJ, Hall JC & Rosbash M (1989) Expression of a *Drosophila* mRNA is under circadian control during pupation. *Development* **107**, 869–880.
- 54 Hardin PE, Hall JC & Rosbash M (1990) Feedback of the *Drosophila period* gene product on circadian cycling of its messenger RNA levels. *Nature* **343**, 536–540.
- 55 Sehgal A, Price JL, Man B & Young MW (1994) Loss of circadian behavioral rhythms and *per* RNA oscillations in the *Drosophila* mutant *timeless*. *Science* **263**, 1603–1606.
- 56 Myers MP, Wager-Smith K, Wesley CS, Young MW & Sehgal A (1995) Positional cloning and sequence analysis of the *Drosophila* clock gene, *timeless*. *Science* **270**, 805–808.
- 57 Sehgal A, Rothenfluh-Hilfiker A, Hunter-Ensor M, Chen Y, Myers MP & Young MW (1995) Rhythmic expression of *timeless*: a basis for promoting circadian cycles in *period* gene autoregulation. *Science* **270**, 808–810.
- 58 Allada R, White NE, So WV, Hall JC & Rosbash M (1998) A mutant *Drosophila* homolog of mammalian *Clock* disrupts circadian rhythms and transcription of *period* and *timeless*. *Cell* **93**, 791–804.
- 59 Darlington TK, Wager-Smith K, Ceriani MF, Staknis D, Gekakis N, Steeves TD, Weitz CJ, Takahashi JS & Kay SA (1998) Closing the circadian loop: CLOCK-induced transcription of its own inhibitors *per* and *tim*. *Science* **280**, 1599–1603.
- 60 Rutila JE, Suri V, Le M, So WV, Rosbash M & Hall JC (1998) CYCLE is a second bHLH-PAS clock protein essential for circadian rhythmicity and transcription of *Drosophila period* and *timeless*. *Cell* **93**, 805–814.
- 61 Saez L, Derasmo M, Meyer P, Stieglitz J & Young MW (2011) A key temporal delay in the circadian cycle of *Drosophila* is mediated by a nuclear localization signal in the *timeless* protein. *Genetics* **188**, 591–600.
- 62 Hara T, Koh K, Combs DJ & Sehgal A (2011) Post-translational regulation and nuclear entry of TIMELESS and PERIOD are affected in new *timeless* mutant. *J Neurosci* **31**, 9882–9890.
- 63 Saez L & Young MW (1996) Regulation of nuclear entry of the *Drosophila* clock proteins *period* and *timeless*. *Neuron* **17**, 911–920.
- 64 Jang AR, Moravcevic K, Saez L, Young MW & Sehgal A (2015) *Drosophila* TIM binds importin  $\alpha 1$ , and acts as an adapter to transport PER to the nucleus. *PLoS Genet* **11**, e1004974.
- 65 Meyer P, Saez L & Young MW (2006) PER-TIM interactions in living *Drosophila* cells: an interval timer for the circadian clock. *Science* **311**, 226–229.
- 66 Sun WC, Jeong EH, Jeong HJ, Ko HW, Edery I & Kim EY (2010) Two distinct modes of PERIOD recruitment onto dCLOCK reveal a novel role for TIMELESS in circadian transcription. *J Neurosci* **30**, 14458–14469.
- 67 Cai YD, Xue Y, Truong CC, Carmen-Li JD, Ochoa C, Vanselow JT, Murphy KA, Li YH, Liu X, Kunimoto BL *et al.* (2021) CK2 inhibits TIMELESS nuclear export and modulates CLOCK transcriptional activity to regulate circadian rhythms. *Curr Biol* **31**, 502–514.
- 68 Martinek S, Inonog S, Manoukian AS & Young MW (2001) A role for the segment polarity gene *shaggy*/GSK-3 in the *Drosophila* circadian clock. *Cell* **105**, 769–779.
- 69 Lin JM, Kilman VL, Keegan K, Paddock B, Emery-Le M, Rosbash M & Allada R (2002) A role for casein kinase 2 $\alpha$  in the *Drosophila* circadian clock. *Nature* **420**, 816–820.
- 70 Akten B, Jauch E, Genova GK, Kim EY, Edery I, Raabe T & Jackson FR (2003) A role for CK2 in the *Drosophila* circadian oscillator. *Nat Neurosci* **6**, 251–257.
- 71 Meissner RA, Kilman VL, Lin JM & Allada R (2008) TIMELESS is an important mediator of CK2 effects on circadian clock function *in vivo*. *J Neurosci* **28**, 9732–9740.
- 72 Top D, Harms E, Syed Y, Adams EL & Saez L (2016) GSK-3 and CK2 kinases converge on Timeless to regulate the master clock. *Cell Rep* **16**, 357–367.
- 73 Sathyanarayanan S, Zheng X, Xiao R & Sehgal A (2004) Posttranslational regulation of *Drosophila* PERIOD protein by protein phosphatase 2A. *Cell* **116**, 603–615.
- 74 Fang Y, Sathyanarayanan S & Sehgal A (2007) Post-translational regulation of the *Drosophila* circadian clock requires protein phosphatase 1 (PP1). *Genes Dev* **21**, 1506–1518.
- 75 Kula-Eversole E, Lee DH, Samba I, Yildirim E, Levine DC, Hong HK, Lear BC, Bass J, Rosbash M & Allada R (2021) Phosphatase of Regenerating Liver-1 selectively times circadian behavior in darkness via function in PDF neurons and dephosphorylation of TIMELESS. *Curr Biol* **31**, 138–149.
- 76 Wülbeck C, Szabo G, Shafer OT, Helfrich-Förster C & Stanewsky R (2005) The novel *Drosophila tim*<sup>(blind)</sup> mutation affects behavioral rhythms but not periodic eclosion. *Genetics* **169**, 751–766.
- 77 Wang Q, Abruzzi KC, Rosbash M & Rio DC (2018) Striking circadian neuron diversity and cycling of *Drosophila* alternative splicing. *eLife* **7**, e35618.
- 78 Szabó Á, Papin C, Cornu D, Chélot E, Lipinszki Z, Udvardy A, Redeker V, Mayor U & Rouyer F (2018) Ubiquitylation dynamics of the clock cell proteome and TIMELESS during a circadian cycle. *Cell Rep* **23**, 2273–2282.
- 79 Kim EY, Jeong EH, Park S, Jeong H-J, Edery I & Cho JW (2012) A role for O-GlcNAcylation in setting circadian clock speed. *Genes Dev* **26**, 490–502.
- 80 Kaasik K, Kivimäe S, Allen JJ, Chalkley RJ, Huang Y, Baer K, Kissel H, Burlingame AL, Shokat KM,

- Ptáček LJ *et al.* (2013) Glucose sensor O-GlcNAcylation coordinates with phosphorylation to regulate circadian clock. *Cell Metab* **17**, 291–302.
- 81 Li YH, Liu XL, Vanselow JT, Zheng H, Schlosser A & Chiu JC (2019) O-GlcNAcylation of PERIOD regulates its interaction with CLOCK and timing of circadian transcriptional repression. *PLoS Genet* **15**, e1007953.
- 82 Liu X, Blaženović I, Contreras AJ, Phan TM, Tabuloc CA, Li YH, Ji J, Fiehn O & Chiu JC (2021) Hexosamine biosynthetic pathway and O-GlcNAc-processing enzymes regulate daily rhythms in protein O-GlcNAcylation. *Nat Commun* **12**, 4173.
- 83 Grima B, Papin C, Martin B, Chélot E, Ponien P, Jacquet E & Rouyer F (2019) PERIOD-controlled deadenylation of the *timeless* transcript in the *Drosophila* circadian clock. *Proc Natl Acad Sci USA* **116**, 5721–5726.
- 84 Pittendrigh CS (1974) Circadian oscillations in cells and the circadian organization of multicellular systems. In *The Neurosciences Third Study Program* (Schmitt FO & Worden FG, eds), pp. 437–458. MIT Press, Cambridge, MA.
- 85 Ceriani MF, Darlington TK, Más DS, Petti AA, Weitz J & Kay SA (1999) Light-dependent sequestration of TIMELESS by CRYPTOCHROME. *Science* **285**, 553–556.
- 86 Ozturk N, Selby CP, Annayev Y, Zhong D & Sancar A (2011) Reaction mechanism of *Drosophila* cryptochrome. *Proc Natl Acad Sci USA* **108**, 516–521.
- 87 Vaidya AT, Top D, Manahan CC, Tokuda JM, Zhang S, Pollack L, Young MW & Crane BR (2013) Flavin reduction activates *Drosophila* cryptochrome. *Proc Natl Acad Sci USA* **110**, 20455–20460.
- 88 Peschel N, Chen KF, Szabo G & Stanewsky R (2009) Light-dependent interactions between the *Drosophila* circadian clock factors *cryptochrome*, *jetlag*, and *timeless*. *Curr Biol* **19**, 241–247.
- 89 Naidoo N, Song W, Hunter-Ensor M & Sehgal A (1999) A role for the proteasome in the light response of the *timeless* clock protein. *Science* **285**, 1737–1741.
- 90 Chen KF, Peschel N, Zavadska R, Sehadova H & Stanewsky R (2011) QUASIMODO, a novel GPI-anchored zone pellucida protein involved in light input to the *Drosophila* circadian clock. *Curr Biol* **21**, 719–729.
- 91 Castiglione-Morelli MA, Guantieri V, Villani V, Kyriacou CP, Costa R & Tamburro AM (1995) Conformational study of the Thr-Gly repeat in the *Drosophila* clock protein, PERIOD. *Proc Natl Acad Sci USA* **260**, 155–163.
- 92 Ewer J, Hamblen-Coyle M, Rosbash M & Hall JC (1990) Requirement for *period* gene expression in the adult and not during development for locomotor activity rhythms of imaginal *Drosophila melanogaster*. *J Neurogenet* **7**, 31–73.
- 93 Sawyer LA, Hennessy JM, Peixoto AA, Rosato E, Parkinson H, Costa R & Kyriacou CP (1997) Natural variation in a *Drosophila* clock gene and temperature compensation. *Science* **278**, 2117–2120.
- 94 Matsumoto A, Tomioka K, Chiba Y & Tanimura T (1999) *tim<sup>rit</sup>* lengthens circadian period in a temperature-dependent manner through suppression of PERIOD protein cycling and nuclear localization. *Mol Cell Biol* **19**, 4343–4354.
- 95 Singh S, Giesecke A, Damulewicz M, Fexova S, Mazzotta GM, Stanewsky R & Dolezel D (2019) New *Drosophila* circadian clock mutants affecting temperature compensation induced by targeted mutagenesis of *Timeless*. *Front Physiol* **10**, 1442.
- 96 Kidd PB, Young MW & Sigma ED (2015) Temperature compensation and temperature sensation in the circadian clock. *Proc Natl Acad Sci USA* **112**, E6284–E6292.
- 97 Zhou M, Kim JK, Eng GWL, Forger DB & Virshup DM (2015) A Period2 phosphoswitch regulates and temperature compensates circadian period. *Mol Cell* **60**, 77–88.
- 98 Yanovsky MJ & Kay SA (2003) Living by the calendar: how plants know when to flower. *Nat Rev Mol Cell Biol* **4**, 265–275.
- 99 Merlin C, Iiams SE & Lugena AB (2020) Monarch butterfly migration moving into the genetic era. *Trends Genet* **36**, 689–701.
- 100 Reiter RJ & Sharma R (2021) Central and peripheral actions of melatonin on reproduction in seasonal and continuous breeding mammals. *Gen Comp Endocrinol* **300**, 1136020.
- 101 Zonato V, Vaniò S, Costa R, Tauber E & Kyriacou CP (2018) Inverse European latitudinal cline at the *timeless* locus of *Drosophila melanogaster* reveals selection on a clock gene: population genetics of *l<sup>s</sup>-tim*. *J Biol Rhythms* **33**, 15–23.
- 102 Sandrelli F, Tauber E, Pegoraro M, Mazzotta G, Risotto P, Landskron J, Stanewsky R, Piccin A, Rosato E, Zordan M *et al.* (2007) A molecular basis for natural selection at the *timeless* locus in *Drosophila melanogaster*. *Science* **316**, 1898–1900.
- 103 Tauber E, Zordan M, Sandrelli F, Pegoraro M, Osterwalder N, Breda C, Daga A, Selmin A, Monger K, Benna C *et al.* (2007) Natural selection favors a newly derived *timeless* allele in *Drosophila melanogaster*. *Science* **316**, 1895–1898.
- 104 Beer K & Helfrich-Föster C (2020) Model and non-model insects in chronobiology. *Front Behav Neurosci* **14**, 601676.
- 105 Pegoraro M, Zonato V, Tyler ER, Fedele G, Kyriacou CP & Tauber E (2017) Geographical analysis of diapause inducibility in European *Drosophila melanogaster* populations. *J Insect Physiol* **98**, 238–244.



- 106 Boothroyd C, Wijnen H, Naef F, Saez L & Young MW (2007) Integration of light and temperature in the regulation of circadian gene expression in *Drosophila*. *PLoS Genet* **3**, e54.
- 107 Montelli S, Mazzotta G, Vanin S, Caccin L, Corrà S, De Pittà C, Boothroyd C, Greem EW, Kyriacou CP & Costa R (2015) *period* and *timeless* mRNA splicing profiles under natural conditions in *Drosophila melanogaster*. *J Biol Rhythms* **30**, 217–227.
- 108 Yamada H & Yamamoto MT (2011) Association between circadian clock genes and diapause incidence in *Drosophila triauraria*. *PLoS One* **6**, e27493.
- 109 Stehlík J, Závodská R, Shimada K, Sauman I & Kostál V (2008) Photoperiodic induction of diapause requires regulated transcription of *timeless* in the larval brain of *Chymomyza costata*. *J Biol Rhythms* **23**, 129–139.
- 110 Huang X, Poelchau MF & Armbruster PA (2015) Global transcriptional dynamics of diapause induction in non-blood-fed and blood-fed *Aedes albopictus*. *PLoS Negl Trop Dis* **9**, e0003724.
- 111 Tapanainen R, Parker DJ & Kankare M (2018) Photosensitive alternative splicing of the circadian clock gene *timeless* is population specific in a cold-adapted fly, *Drosophila montana*. *G3: Genes - Genomes - Genetics* **8**, 1291–1297.
- 112 Liu Y, Garceau NY, Loros JJ & Dunlap JC (1997) Thermally regulated translational control of FRQ mediates aspects of temperature responses in the *Neurospora* circadian clock. *Cell* **89**, 477–486.
- 113 Colot HV, Loros JJ & Dunlap JC (2005) Temperature-modulated alternative splicing and promoter use in the Circadian clock gene *frequency*. *Mol Biol Cell* **16**, 5563–5571.
- 114 Diernfellner A, Colot HV, Dintsis O, Loros JJ, Dunlap JC & Brunner M (2007) Long and short isoforms of *Neurospora* clock protein FRQ support temperature-compensated circadian rhythms. *FEBS Lett* **581**, 5759–5764.
- 115 Lee BH, Kapoor A, Zhu J & Zhu JK (2006) STABILIZED1, a stress-upregulated nuclear protein, is required for pre-mRNA splicing, mRNA turnover, and stress tolerance in *Arabidopsis*. *Plant Cell* **18**, 1736–1749.
- 116 Kim GD, Cho YH, Lee BH & Yoo SD (2017) STABILIZED1 modulates pre-mRNA splicing for thermotolerance. *Plant Physiol* **173**, 2370–2382.
- 117 Liu GT, Jiang JF, Liu XN, Jiang JZ, Sun L, Duan W, Li RM, Wang Y, Lecourieux D, Liu CH *et al.* (2019) New insights into the heat responses of grape leaves via combined phosphoproteomic and acetylproteomic analyses. *Hortic Res* **6**, 100.
- 118 Schlaen RG, Mancini E, Sanchez SE, Perez-Santángelo S, Rugnone ML, Simpson CG, Brown JWS, Zhang X, Chernomoretz A & Yanovsky MJ (2015) The spliceosome assembly factor GEMIN2 attenuates the effects of temperature on alternative splicing and circadian rhythms. *Proc Natl Acad Sci USA* **112**, 9382–9387.
- 119 Weber C, Nover L & Fauth M (2008) Plant stress granules and mRNA processing bodies are distinct from heat stress granules. *Plant J* **56**, 517–530.
- 120 Reddy ASN, Day IS, Göhring J & Barta A (2012) Localization and dynamics of nuclear speckles in plants. *Plant Physiol* **158**, 67–77.
- 121 Saez L & Young MW (1988) *In situ* localization of the *per* clock protein during development of *Drosophila melanogaster*. *Mol Cell Biol* **8**, 5378–5385.
- 122 Hardin PE (1994) Analysis of *period* mRNA cycling in *Drosophila* head and body tissues indicates that body oscillators behave differently from head oscillators. *Mol Cell Biol* **14**, 7211–7218.
- 123 Beaver LM, Rush BL, Gvakharia BO & Giebultowicz JM (2003) Noncircadian regulation and function of clock genes *period* and *timeless* in oogenesis of *Drosophila melanogaster*. *J Biol Rhythms* **18**, 463–472.
- 124 Rush BL, Murad A, Emery P & Giebultowicz JM (2006) Ectopic CRYPTOCHROME renders TIM light sensitive in the *Drosophila* ovary. *J Biol Rhythms* **21**, 272–278.
- 125 Howlader G & Sharma VK (2006) Circadian regulation of egg-laying behavior in fruit flies *Drosophila melanogaster*. *J Insect Physiol* **52**, 779–785.
- 126 Benna C, Bonaccorsi S, Wülbeck C, Helfrich-Förster C, Gatti M, Kyriacou CP, Costa R & Sandrelli F (2010) *Drosophila timeless2* is required for chromosome stability and circadian photoreception. *Curr Biol* **20**, 346–352.
- 127 Koike N, Hida A, Numano R, Hirose M, Sakaki Y & Tei H (1998) Identification of the mammalian homologues of the *Drosophila timeless* gene, *Timeless1*. *FEBS Lett* **441**, 427–431.
- 128 Sangoram AM, Saez L, Antoch MP, Gekakis N, Staknis D, Whiteley A, Fruechte EM, Vitaterna MH, Shimomura K, King DP *et al.* (1998) Mammalian circadian autoregulatory loop: a *timeless* ortholog and *mPer1* interact and negatively regulate CLOCK-BMAL1-induced transcription. *Neuron* **21**, 1101–1113.
- 129 Zylka MJ, Shearman LP, Levine JD, Jin X, Weaver DR & Reppert SM (1998) Molecular analysis of mammalian timeless. *Neuron* **21**, 1115–1122.
- 130 Takumi T, Nagamine Y, Miyake S, Matsubara C, Taguchi K, Takekida S, Sakakida Y, Nishikawa K, Kishimoto T, Niwa S *et al.* (1999) A mammalian ortholog of *Drosophila timeless*, highly expressed in SCN and retina, forms a complex with mPER1. *Genes Cells* **4**, 67–75.
- 131 Tischkau SA, Barnes JA, Lin FJ, Myers EM, Barnes JW, Meyer-Bernstein EL, Hurst WJ, Burgoon PW, Chen D, Sehgal A *et al.* (1999) Oscillation and light

- induction of *timeless* mRNA in the mammalian circadian clock. *J Neurosci* **19**, RC15.
- 132 Barnes JW, Tischkau SA, Barnes JA, Mitchell JW, Burgoon PW, Hickok JR & Gillette MU (2003) Requirement of mammalian *Timeless* for circadian rhythmicity. *Science* **302**, 439–442.
- 133 Griffin Jr EA, Staknis D & Weitz CJ (1999) Light-independent role of CRY1 and CRY2 in the mammalian circadian clock. *Science* **286**, 768–771.
- 134 Kume K, Zylka MJ, Sriram S, Shearman LP, Weaver DR, Jin X, Maywood ES, Hastings MH & Reppert SM (1999) mCRY1 and mCRY2 are essential components of the negative limb of the circadian clock feedback loop. *Cell* **98**, 193–205.
- 135 Field MD, Maywood ES, O'Brien JA, Weaver DR, Reppert SM & Hastings MH (2000) Analysis of clock proteins in mouse SCN demonstrates phylogenetic divergence of the circadian clockwork and resetting mechanisms. *Neuron* **25**, 437–447.
- 136 Gotter AL (2003) Tipin, a novel *timeless*-interacting protein, is developmentally co-expressed with *timeless* and disrupts its self-association. *J Mol Biol* **331**, 167–176.
- 137 Kurien P, Hsu PK, Leon J, Wu D, McMahon T, Shi G, Xu Y, Lipzen A, Pennacchio LA, Jones CR *et al.* (2019) TIMELESS mutation alters phase responsiveness and causes advanced sleep phase. *Proc Natl Acad Sci USA* **116**, 12045–12053.
- 138 Shafi AA, McNair CM, McCann JJ, Alshalalfa M, Shostak A, Severson TM, Zhu Y, Bergman A, Gordon N, Mandigo AC *et al.* (2021) The circadian cryptochrome, CRY1, is a pro-tumorigenic factor that rhythmically modulates DNA repair. *Nat Commun* **12**, 401.
- 139 Huber AL, Papp SJ, Chan AB, Henriksson E, Jordan SD, Kriebs A, Nguyen M, Wallace M, Li Z, Metallo CM & Lamia KA (2016) CRY2 and FBXL3 cooperatively degrade c-MYC. *Mol Cell* **64**, 774–789.
- 140 Ünsal-Kaçmaz K, Mullen TE, Kaufmann WK & Sancar A (2005) Coupling of human circadian and cell cycles by the *timeless* protein. *Mol Cell Biol* **25**, 3109–3116.
- 141 Kang TH & Leem SH (2014) Modulation of ATR-mediated DNA damage checkpoint response by cryptochrome 1. *Nucleic Acids Res* **42**, 4427–4434.
- 142 Benna C, Scannapieco P, Piccin A, Sandrelli F, Zordan M, Rosato E, Kyriacou CP, Valle G & Costa R (2000) A second *timeless* gene in *Drosophila* shares greater sequence similarity with mammalian *tim*. *Curr Biol* **10**, R512–R513.
- 143 Rubin EB, Shemesh Y, Cohen M, Elgavish S, Elgavish S, Robertson HM & Bloch G (2006) Molecular and phylogenetic analyses reveal mammalian-like clockwork in the honey bee (*Apis mellifera*) and shed new light on the molecular evolution of the circadian clock. *Genome Res* **16**, 1352–1365.
- 144 Foss EJ (2001) Tof1p regulates DNA damage responses during S phase in *Saccharomyces cerevisiae*. *Genetics* **157**, 567–577.
- 145 Gotter AL, Suppa C & Emanuel BS (2006) Mammalian TIMELESS and Tipin are evolutionarily conserved replication fork-associated factors. *J Mol Biol* **366**, 36–52.
- 146 Cho W-H, Kang Y-H, An Y-Y, Tappin I, Hurwitz J & Lee J-K (2013) Human Tim-Tipin complex affects the biochemical properties of the replicative DNA helicase and DNA polymerases. *Proc Natl Acad Sci USA* **110**, 2523–2527.
- 147 Urtishak KA, Smith KD, Chanoux RA, Greenberg RA, Johnson FB & Brown EJ (2009) Timeless maintains genomic stability and suppresses sister chromatid exchange during unperturbed DNA replication. *J Biol Chem* **284**, 8777–8785.
- 148 Smith KD, Fu MA & Brown EJ (2009) Tim-Tipin dysfunction creates an indispensable reliance on the ATR-Chk1 pathway for continued DNA synthesis. *J Cell Biol* **187**, 15–23.
- 149 Cali F, Bharti SK, Di Perna R, Brosh RM, Pisani FM (2016) Tim/Timeless, a member of the replication fork protection complex, operates with the Warsaw breakage syndrome DNA helicase DDX11 in the same fork recovery pathway. *Nucleic Acids Res* **44**, 705–717.
- 150 Shyian M, Albert B, Zupan AM, Ivanitsa V, Charbonnet G, Dilg D & Shore D (2020) Fork pausing complex engages topoisomerases at the replisome. *Genes Dev* **34**, 87–98.
- 151 Westhorpe R, Keszthelyi A, Minchell NE, Jones D & Baxter J (2020) Separable functions of Tof1/Timeless in intra-S-checkpoint signalling, replisome stability and DNA topological stress. *Nucleic Acids Res* **48**, 12169–12187.
- 152 Kemp MG, Akan Z, Yilmaz S, Grillo M, Smith-Roe SL, Kang TH, Cordeiro-Stone M, Kaufmann WK, Abraham RT, Sancar A *et al.* (2010) Tipin-replication protein A interaction mediates Chk1 phosphorylation by ATR in response to genotoxic stress. *J Biol Chem* **285**, 16562–16571.
- 153 Yang X, Wood PA & Hrushesky WJM (2010) Mammalian TIMELESS is required for ATM-dependent CHK2 activation and G2/M checkpoint control. *J Biol Chem* **285**, 3030–3034.
- 154 Xie S, Mortusewicz O, Ma HT, Herr P, Poon RYC, Helleday T & Qian (2015) Timeless interacts with PARP-1 to promote homologous recombination repair. *Mol Cell* **60**, 163–176.
- 155 Young LM, Marzio A, Perez-Duran P, Reid DA, Meredith DN, Roberti D, Star A, Rothenberg E, Ueberheide B & Pagano M (2015) TIMELESS forms a complex with PARP1 distinct from its complex with TIPIN and plays a role in the DNA damage response. *Cell Rep* **13**, 451–459.

- 156 Chou DM & Elledge SJ (2006) Tipin and Timeless form a mutually protective complex required for genotoxic stress resistance and checkpoint function. *Proc Natl Acad Sci USA* **103**, 18143–18147.
- 157 Ünsal-Kacmaz K, Chastain PD, Qu PP, Minoo P, Cordeiro-Stone M, Sancar A & Kaufmann WK (2007) The human Tim/Tipin complex coordinates an Intra-S checkpoint response to UV that slows replication fork displacement. *Mol Cell Biol* **27**, 3131–3142.
- 158 Gotter AL, Suppa C & Emanuel BS (2007) Mammalian TIMELESS and Tipin are evolutionarily conserved replication fork-associated factors. *J Mol Biol* **366**, 36–52.
- 159 Mao Y, Fu A, Leaderer D, Zheng T, Chen K & Zhu Y (2013) Potential cancer-related role of circadian gene TIMELESS suggested by expression profiling and in vitro analyses. *BMC Cancer* **13**, 498.
- 160 Yoshida K, Sato M, Hase T, Elshazley M, Yamashita R, Usami N, Taniguchi T, Yokoi K, Nakamura S, Kondo M, Girard L, Minna JD & Hasegawa Y (2013) TIMELESS is over expressed in lung cancer and its expression correlates with poor patient survival. *Cancer Sci* **104**, 171–177.
- 161 Bianco JN, Bergoglio V, Lin YL, Pillaire MJ, Schmitz AL, Gilhodes J, Lusque A, Mazières J, Lacroix-Triki M, Roumeliotis TI *et al.* (2019) Overexpression of Claspin and Timeless protects cancer cells from replication stress in a checkpoint-independent manner. *Nat Commun* **10**, 910.
- 162 Rageul J, Park JJ, Zeng PP, Lee EA, Yang J, Hwang S, Lo N, Weinheimer AS, Schärer OD, Yeo JE *et al.* (2020) SDE2 integrates into the TIMELESS-TIPIN complex to protect stalled replication forks. *Nat Commun* **11**, 5495.
- 163 Pan A, Schernhammer ES, Sun Q & Hu FB (2011) Rotating night shift work and risk of type 2 diabetes: two prospective cohort studies in women. *PLoS Medicine* **8**, e1001141.
- 164 Kinouchi K & Sassone-Corsi P (2020) Metabolic rivalry: circadian homeostasis and tumorigenesis. *Nat Rev Cancer* **20**, 645–661.
- 165 Engelen E, Janssens RC, Yagita K, Smits VA, van der Horst GTJ & Tamanini F (2013) Mammalian TIMELESS is involved in period determination and DNA damage-dependent phase advancing of the circadian clock. *PLoS One* **8**, e56623.
- 166 Pagueiro AM, Liu Q, Baker CL, Dunlap JC & Loros JJ (2006) The *Neurospora* checkpoint kinase 2: a regulatory link between the circadian and cell cycles. *Science* **313**, 644–649.
- 167 Oklejewicz M, Destici E, Tamanini F, Hut RA, Janssens R & van der Horst GT (2008) Phase resetting of the mammalian circadian clock by DNA damage. *Curr Biol* **18**, 286–291.
- 168 Papp SJ, Huber AL, Jordan SD, Kriebs A, Nguyen M, Moresco JJ, Yates JR & Lamia KA (2015) DNA damage shifts circadian clock time via Hausp-dependent Cry1 stabilization. *eLife* **4**, e04883.
- 169 Matsuo T, Yamaguchi S, Mitsui S, Emi A, Shimoda F & Okamura H (2003) Control mechanism of the circadian clock for the timing of cell division in vivo. *Science* **302**, 255–259.
- 170 Yang Q, Pando BF, Dong G, Golden SS & van Oudenaarden A (2010) Circadian gating of the cell cycle revealed in single cyanobacteria cells. *Science* **327**, 1522–1526.
- 171 Geyfman M, Kumar V, Liu Q, Ruiz R, Gordon W, Espitia F, Cam E, Millar SE, Smyth P, Ihler A *et al.* (2012) Brain and muscle Arnt-like protein-1 (BMAL1) controls circadian cell proliferation and susceptibility to UVB-induced DNA damage in the epidermis. *Proc Natl Acad Sci USA* **109**, 11758–11763.
- 172 Bouchard-Cannon P, Mendoza-Viveros L, Yuen A, Kærn M & Cheng HY (2013) The circadian molecular clock regulates adult hippocampal neurogenesis by controlling the timing of cell-cycle entry and exit. *Cell Rep* **5**, 961–973.
- 173 Karpowicz P, Zhang Y, Hogenesch JB, Emery P & Perrimon N (2013) The circadian clock gates the intestinal stem cell regenerative state. *Cell Rep* **3**, 996–1004.
- 174 Hong CI, Zámbořszky J, Baek M, Labiscsak L, Ju K, Lee H, Larrondo LF, Gotik A, Chong HS, Belden WJ *et al.* (2014) Circadian rhythms synchronize mitosis in *Neurospora crassa*. *Proc Natl Acad Sci USA* **111**, 1397–1402.
- 175 Miyagishima SY, Fuji T, Sumiya N, Hirooka S, Nakano A, Kabeya Y & Nakamura M (2014) Translation-independent circadian control of the cell cycle in a unicellular photosynthetic eukaryote. *Nat Commun* **5**, 3807.
- 176 Matsu-ura T, Dovzhenok A, Aihara E, Rood J, Le H, Ren Y, Rosselot AE, Zhang T, Lee C, Obrietan K *et al.* (2016) Intercellular coupling of the cell cycle and circadian clock in adult stem cell culture. *Mol Cell* **64**, 900–912.
- 177 Fung-Uceda J, Lee K, Seo PJ, Polen S, De Veylder L & Mas P (2018) The circadian clock sets the time of DNA replication licensing to regulate growth in *Arabidopsis*. *Dev Cell* **45**, 101–113.e4.
- 178 Liao Y & Rust MJ (2021) The circadian clock ensures successful DNA replication in cyanobacteria. *Proc Natl Acad Sci USA* **118**, e2022516118.
- 179 Gamsby JJ, Loros JJ & Dunlap JC (2009) A phylogenetically conserved DNA damage response resets the circadian clock. *J Biol Rhythms* **24**, 193–202.
- 180 Liu X, Dang Y, Matsu-Ura T, He Y, He Q, Hong CI & Liu Y (2017) DNA replication is required for

- circadian clock function by regulating rhythmic nucleosome composition. *Mol Cell* **67**, 203–213.e4.
- 181 Dakup P & Gaddameedhi S (2017) Impact of the circadian clock on UV-induced DNA damage response and photocarcinogenesis. *Photochem Photobiol* **93**, 296–303.
- 182 Shostak A (2017) Circadian clock, cell division, and cancer: from molecules to organism. *Int J Mol Sci* **18**, 873.
- 183 Gaucher J, Montellier E & Sassone-Corsi P (2018) Molecular cogs: interplay between circadian clock and cell cycle. *Trends Cell Biol* **28**, 368–379.
- 184 Ruby CL, Major RJ & Hinrichsen RD (2021) Regulation of tissue regeneration by the circadian clock. *Eur J Neurosci* **53**, 3576–3597.
- 185 Lubov JE, Cvammen W & Kemp MG (2021) The impact of the circadian clock on skin physiology and cancer development. *Int J Mol Sci* **22**, 6112.
- 186 Sancar A & Van Gelder RN (2021) Clocks, cancer, and chronochemotherapy. *Science* **371**. <https://doi.org/10.1126/science.abb0738>
- 187 Lam VL, Li YH, Liu X, Murphy KA, Diehl JS, Kwok RS & Chiu JC (2018) CK1 $\alpha$  collaborates with DOUBLETIME to regulate PERIOD function in the *Drosophila* circadian clock. *J Neurosci* **38**, 10631–10643.
- 188 Ousley A, Zafarullah K, Chen Y, Emerson M, Hickman L & Sehgal A (1998) Conserved regions of the *timeless (tim)* clock gene in *Drosophila* analyzed through phylogenetic and functional studies. *Genetics* **148**, 815–825.
- 189 Blackford AN & Jackson SP (2017) ATM, ATR, and DNA-PK, the trinity at the heart of the DNA damage response. *Mol Cell* **66**, 801–817.