



## Correction to: Patients with B-cell lymphoma receiving anti-CD20 monoclonal antibody-containing chemotherapies and seroreactive patterns in response to COVID-19 vaccination

Tohru Inaba<sup>1</sup> · Keita Okumura<sup>2</sup> · Chika Maekura<sup>3</sup> · Ayako Muramatsu<sup>3</sup> · Tsutomu Kobayashi<sup>3</sup> · Junya Kuroda<sup>3</sup> · Yoko Nukui<sup>1</sup>

Published online: 5 May 2022  
© Japanese Society of Hematology 2022

**Correction to: International Journal of Hematology**  
<https://doi.org/10.1007/s12185-022-03349-1>

In the original publication of the article, the table related to case study was missing. The case study table is given in this correction.

---

The original article can be found online at <https://doi.org/10.1007/s12185-022-03349-1>.

---

✉ Tohru Inaba  
[inaba178@koto.kpu-m.ac.jp](mailto:inaba178@koto.kpu-m.ac.jp)

- <sup>1</sup> Department of Infection Control and Laboratory Medicine, Kyoto Prefectural University of Medicine, Kawaramachi-Hirokoji, Kamigyo-ku, Kyoto 602-8566, Japan
- <sup>2</sup> Faculty of Clinical Laboratory, Kyoto Prefectural University of Medicine, Kawaramachi-Hirokoji, Kamigyo-ku, Kyoto 602-8566, Japan
- <sup>3</sup> Division of Hematology and Oncology, Kyoto Prefectural University of Medicine, Kawaramachi-Hirokoji, Kamigyo-ku, Kyoto 602-8566, Japan

Case	Disease	CD20 MoAb <sup>*1</sup>	SARS-CoV2		Anti-SARS-CoV-2 Antibodies (AU/mL) <sup>*2</sup>					Serum IgG (mg/dL) <sup>*3</sup>	Time <sup>*4</sup>
			Infection	Vaccine	IgG-N	IgM-N	IgG-S1	IgM-S1	Nab		
Case 1	FL	37	+	-	n.t. <sup>*5</sup>	n.t.	n.t.	n.t.	0.68	563	+2w
					0.27	0.14	0.39	0.21	3.48	441	+4w
					0.19	0.33	0.37	0.23	1.44	183	+10w
Case 2	FL	1	+	-	9.46	0.82	0.47	0.48	2.37	685	+2w
					<b><u>235.22</u></b>	<b><u>29.82</u></b>	<b><u>16.79</u></b>	2.23	<b><u>21.21</u></b>	576	+4w
Case 3	FL	12	+	+	0.32	1.12	7.09	<b><u>594.44</u></b>	<b><u>140.16</u></b>	492	+2w
					0.32	0.95	8.01	<b><u>578.75</u></b>	<b><u>213.00</u></b>	461	+4w
Case 4	Healthy	0	-	+	0.53	0.16	9.78	0.63	5.88	1,290	+2w
					0.50	0.17	<b><u>2145.96</u></b>	<b><u>18.97</u></b>	<b><u>2836.82</u></b>	1,291	+4w

\*1: Number of anti-CD20 monoclonal antibody-containing chemotherapy sessions administered in each case

\*2: Bold numbers with underline mean positive antibody titers (> 10 AU/mL)

\*3: Reference interval of serum IgG is 861–1747 mg/dL in our hospital

\*4: In Cases 1–3, time from the onset of SARS-CoV2 infection. In Case 4, time from the initial vaccination

\*5: n.t.: not tested

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.