



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Available online at
ScienceDirect
 www.sciencedirect.com

Elsevier Masson France
EM|consulte
 www.em-consulte.com



Editorial

Lung transplantation in the COVID-19 Era: A multi-faceted challenge

Abbreviations

ARDS	acute respiratory distress syndrome
COVID-19	coronavirus disease 2019
ECMO	extra-corporeal membrane oxygenation
ICU	intensive care unit
IQR	interquartile range
LTx	lung transplantation
SARS-CoV-2	severe acute respiratory syndrome coronavirus 2
SOT	solid organ transplantation
WHO	World Health Organisation

As of 17 September 2021, COVID-19 had affected 226,844,344 individuals worldwide, of whom 4666,334 had died [1]. This massive pandemic has inflicted an unprecedented and onerous burden on hospitals everywhere. To protect the quality of care, preparations were made [2, 3], guidelines developed and implemented [4–6], and local strategies devised [7, 8].

Although physicians performing solid-organ transplantation (SOT) suspected that the COVID-19 pandemic would adversely affect both recipients and organ procurement [9–11], their fears fell far short of reality. French lung transplantation (LTx) units observed devastating effects [12] as soon as one month after the World Health Organisation classified COVID-19 as a pandemic [13].

Now, after more than 18 months fighting COVID-19, reports of direct consequences on SOT recipients and of collateral damage to transplant programs have been published. Among SOT patients, LTx recipients have been the most severely affected. In a somewhat ironic turn of events, LTx is now suggested to treat patients with severe SARS-CoV-2 pneumonia.

COVID-19 pneumonia in lung-transplant recipients

A British epidemiological study that used the analytics programme OpenSAFELY to assess the primary-care records of 17,278,392 adults found that having a transplanted organ was amongst the comorbid conditions associated with the greatest increase in COVID-19 severity, the adjusted hazard ratio for death being 3.53 (95%CI, 2.77–4.49) [14]. Among SOT recipients, those with transplanted lungs seem the most at risk. In a nationwide Spanish study of 778 SOT or haematopoietic-stem-cell recipients who contracted COVID-19 during the first 6 months of the pandemic, 54 patients had LTx, which was associated with a 46% mortality rate compared to 27% for the overall cohort (odds ratio, 2.5; 95%CI, 1.4–4.6) [15]. Another nationwide study, conducted in France, identified 35 LTx recipients with confirmed or strongly suspected COVID-19 during the first wave [16]. Mortality was lower, at 14.3%, and overweight was significantly associated with death in the multivariate analysis.

Collateral damage on lung transplantation (LTx) programmes

That the COVID-19 pandemic would have surreptitious effects was suspected very early on [12]. A study sought to determine whether the number of SOT procedures in France and the US was associated with the number of patients who had COVID-19 during the first few months of the pandemic [17]. The number of transplantations from deceased donors dropped massively, by 90.6% in France and 51.1% in the US, as the number of COVID-19 infections increased. The decrease occurred for all organs but was greatest for kidneys and, to a lesser extent, lungs.

Recent retrospective work has provided valuable insights into the consequences of the pandemic on SOT in 22 countries in Europe, the Americas, and Japan [18]. Data from national or regional organ-procurement and transplantation networks collected in 2019 and 2020 were compared. Raw data on numbers of transplantations of each organ were obtained and the number of patient life-years lost because of the drop in transplantation activity was estimated. The findings are of the utmost interest. First, the overall SOT decrease was 15.92%, with the drop being greatest for kidneys (19.14%) and smallest for hearts (5.44%). The 15.51% reduction in LTx masked substantial geographical differences, with a 66.67% increase in Slovenia (plus 6 LTx procedures in 2020 vs. 2019) and a 85.71% drop in Chile (minus 6 LTx procedures in 2020). In France, the drop was –31.27% (101 fewer LTx procedures in 2020). Second, an important original feature of the study is the estimation of the number of patient life-years lost due to the drop in SOT activity, based on a comparison of patients who received transplants to those who remained on the waiting list. For LTx candidates, the loss was 1799 years.

Data collected by the French biomedicine agency (Agence de la Biomédecine) indicate a decrease in access to organ transplants of candidates listed in 2020 compared to those listed during the two previous years. This decrease occurred for all organ types except hearts. LTx candidates saw their cumulative incidence of access to transplantation fall from 55 [interquartile range 52–59] in 2018 and 2019 to 45 [IQR 39–51] in 2020 ($p < 0.01$). This major drop was not associated with significant changes in early post-LTx outcomes or in the 3-month post-listing cumulative incidence of death or delisting. Nevertheless, whether 3 months is sufficient to assess the consequences of the downturn in LTx activity is debatable.

Interestingly, similar findings were obtained in the US [19], where listing of LTx candidates and LTx procedures dropped by 40% and 28%, respectively, compared to 28% and 13% for the heart. Moreover, whereas heart transplantation activity recovered rapidly after the first COVID-19 wave, the same was not true for LTx.

Several reasons may explain the 31.27% decrease in French LTx activity during the pandemic [18]. In France, during the first

Table 1

The ten criteria required to consider lung transplantation in patients with end-stage lung disease due to COVID-19, adapted from Cypel and Keshavjee [34].

Candidates
Age under 65 years
Single organ dysfunction
Allow sufficient time for lung recovery (4 to 6 weeks)
Radiological evidence of irreversible lung disease
Patient awake and able to discuss transplantation
Patient participating in physical rehabilitation
Absence of typical contra-indication to LTx
Absence of SARS-CoV2 infectivity
Centre
Centre experienced in performing high-risk LT
Access to a broad-donor pool and low waiting-list mortality

COVID-19 wave in early 2020, the LTx task force of the French pulmonology society (Société de Pneumologie de Langue Française) decided that LTx should be performed only in patients on the high-priority waiting list and in those deemed to require the procedure urgently [12]. This decision was based on the need to ensure optimal use of the few operating rooms that had not been converted into COVID-19 ICUs, concern about shortage of critical-care devices such as ventilators and extra-corporeal membrane oxygenation pumps, and the potential risks associated with exposure of LTx candidates or recipients to COVID-19 infection in crowded ICUs. However, these concerns were not the main reason for the drop in LTx procedures. Donations fell in France by 16% from 1 March to 31 July 2020 compared to the same periods in 2018 and 2019. The decrease was similar in the US [20] and even higher in the UK [21]. A key reason for the decrease in donations is that ICU beds were not available for donors (as for recipients) [22, 23]. Also, the very tight first lockdown in France may have reduced the number of donors who died from traumatic injuries, as reported in the US [20]. Finally, the discarding of organs from 14 potential donors (1% of potential donors) between 1 March and 31 July due to positive SARS-CoV2 tests made a minor contribution [22].

Efforts were made to maintain LTx activity despite the pandemic. The UK set up COVID-light hospitals, where only patients requiring elective care and free of the SARS-CoV-2 virus were treated. One of these centres continued to offer care to all LTx candidates but performed only 7 LTx procedures (58% of all LTx procedures in the UK) between 1 March and 31 May 2020, and LTx activity dropped by 77% compared to 2019 [21]. Clearly, the COVID-light approach is insufficient to tackle the LTx challenges raised by the pandemic. Alternative strategies must be devised and tested to protect LTx candidates and recipients as we prepare for further COVID-19 waves or COVID-19 endemicity.

Lung transplantation (LTx) for end-stage COVID-19 pneumonia

LTx has recently emerged as a therapeutic option for refractory acute respiratory distress syndrome caused by COVID-19 [24–32]. We identified 26 reported cases. The disease was extremely severe, with patients having no evidence of lung healing, and 24 requiring ECMO for a median duration of 39.5 days [IQR, 21.3–65.5]. The pre-LTx ECMO durations were shortest for the earliest patients (range, 10–19 days) [24, 25]. In the following reports, more time was given to the possibility of improvement (pre-LTx ECMO duration range, 17–180 days) [26–28, 30–32]. Surgery was challenging, with dissection difficulties in 7 patients [26, 27, 33] and high blood-product requirements. The outcomes were encouraging, with a median post-

operative follow-up of 90 days [IQR, 52–144] in 23 patients and only 3 patients dying [25, 26, 30].

The hazards of LTx in this particular setting are still debated [34–36]. The healing potential of the lung is not predictable. A comprehensive pre-transplant assessment, notably of the patient's neurocognitive and psychological profile, is difficult to achieve. These patients cannot give informed consent to LTx. We do not know whether an acceptable strategy to ensure the optimal use of scarce organs is to perform LTx in patients with immediately life-threatening disease, after a pared-down pre-LTx evaluation, but not in those whose condition is severe and chronic but stable. Ten criteria for selecting patients with COVID-19 for consideration as LTx candidates have been suggested (Table 1) [34]. New data may prompt a modification of these criteria within the next few months.

Finally, acute respiratory distress due to any cause [37] including COVID-19 [38] has been reported to induce lung fibrosis. Time will tell whether sequelae from COVID-19 will represent a significant indication for LTx.

Thus, the COVID-19 pandemic has had a global and severe adverse effect on LTx recipients, candidates, and procedures. In the field of transplantation, LTx appears particularly vulnerable. Although the overwhelming of the entire healthcare system was unpredictable, LTx physicians must now use their newly acquired experience to develop master plans for addressing healthcare system crises, including future COVID-19 waves. Understanding how different countries and healthcare systems respond to COVID-19-related challenges may improve pandemic preparedness, including the strategies devised to maintain transplantation activity in order to prevent the loss of patient life-years.

Declarations of Competing Interest

None.

Jonathan Messika, MD, PhD*
 APHP, Nord-Université de Paris, Hôpital Bichat, Pulmonology B and Lung
 Transplantation Department, F-75018, Paris, France
 Université de Paris, INSERM PHERE UMRS 1152, F-75018 Paris, France
 Paris Transplant Group, Paris, France

Antoine Roux, MD, PhD
 Paris Transplant Group, Paris, France
 Pulmonology, Adult Cystic Fibrosis Centre, and Lung Transplantation
 Department, Foch Hospital, Suresnes, France
 Versailles-Saint-Quentin-en-Yvelines University, Versailles, France

Gaëlle Dauriat, MD
 Jérôme Le Pavec, MD, PhD
 for the Groupe de Transplantation Pulmonaire de la Société de Pneumologie de Langue Française
 Paris-Saclay University, School of Medicine, Le Kremlin-Bicêtre, France
 INSERM UMR_S 999, Pulmonary Hypertension: Pathophysiology and Novel Therapies Unit, Marie Lannelongue Hospital, Paris Saint-Joseph Hospital Group, Le Plessis-Robinson, France
 Marie Lannelongue Hospital, Paris Saint-Joseph Hospital Group, Department of transplantation and thoracic and vascular surgery, Pulmonary Hypertension National Referral Centre, Le Plessis Robinson, France

*Corresponding author at: Service de Pneumologie B et Transplantation Pulmonaire, Hôpital Bichat Claude-Bernard, 46 rue Henri Huchard, 75018 Paris, France.
 E-mail address: jonathan.messika@aphp.fr (J. Messika).

Received 7 October 2021
 Accepted 8 November 2021
 Available online 17 December 2021

References

[1] World Health Organization. WHO Coronavirus (COVID-19) Dashboard [Internet]. [cited 2021 Sep 20]. Available from: <https://covid19.who.int/>.

[2] Celarier T, Lafaie L, Goethals L, Barth N, Gramont B, Ojardias E, et al. Covid-19: adapting the geriatric organisations to respond to the pandemic. *Respir Med Res* 2020;78:100774.

[3] Lim J, Broughan J, Crowley D, O’Kelly B, Fawsitt R, Burke MC, et al. COVID-19’s impact on primary care and related mitigation strategies: a scoping review. *Eur J Gen Pract* 2021;27:166–75.

[4] Curigliano G, Banerjee S, Cervantes A, Garassino MC, Garrido P, Girard N, Haanen J, Jordan K, Lordick F, Machiels JP, Michielin O, Peters S, Tabernero J, Douillard JY, Pentheroudakis G, Addeo A, Albiges L, Ascierto PA, Banerjee S, Barlesi F, Caldas C, Cardoso F, Cervantes A, Chaberny IF, Cherny NI, Choueiri TK, Chua MLK, Crisciello C, Curigliano G, de Azambuja E, et al. Managing cancer patients during the COVID-19 pandemic: an ESMO multidisciplinary expert consensus. *Ann Oncol* 2020;31:1320–35.

[5] Global Initiative for Chronic Obstructive Lung Disease. GOLD COVID-19 Guidance [Internet]. [cited 2021 Sep 20]. Available from: <https://goldcopd.org/gold-covid-19-guidance/>.

[6] Girard N, Grenillier L, Zalcmán G, Cadranet J, Moro-Sibilot D, Mazières J, Audigier-Valette C, Bennouna J, Besse B, Cortot A, Couraud S, Duruisseaux M, Giroux-Leprieur E, Toffart A-C, Westeel V, Wislez M. Proposals for managing patients with thoracic malignancies during COVID-19 pandemic. *Respirat Med Res* 2020;78:100769.

[7] Peiffer-Smadja N, Lucet J-C, Bendjeloul G, Bouadma L, Gerard S, Choquet C, Jacques S, Khalil A, Maisani P, Casalino E, Descamps D, Timsit J-F, Yazdanpanah Y, Lescure F-X. Challenges and issues about organizing a hospital to respond to the COVID-19 outbreak: experience from a French reference centre. *Clin Microbiol Infect* 2020;26:669–72.

[8] Assistance Publique–Hôpitaux de Paris’ response to the COVID-19 pandemic. *Lancet North Am Ed* 2020;395:1760–1.

[9] Holm AM, Mehra MR, Courtwright A, Teuteberg J, Sweet S, Potena L, Singer LG, Farrero M, Shullo MA, Benza R, Ensminger S, Aslam S. Ethical considerations regarding heart and lung transplantation and mechanical circulatory support during the COVID-19 pandemic: an ISHLT COVID-19 Task Force statement. *J Heart Lung Transplant* 2020;39:619–26.

[10] Ahn C, Amer H, Anglicheau D, Ascher NL, Baan CC, Battsetset G, Bat-Iredui B, Berney T, Betjes MGH, Bichu S, Birn H, Brennan D, Bromberg J, Caillard S, Cannon RM, Cantarovich M, Chan A, Chen ZS, Chapman JR, Cole EH, Cross N, Durand F, Egawa H, Emond JC, Farrero M, Friend PJ, Geissler EK, Ha J, Haberal MA, Henderson ML, et al. Global Transplantation COVID Report March 2020. *Transplantation* 2020;104:1974–83.

[11] Fishman JA, Grossi PA. Novel Coronavirus-19 (COVID-19) in the immunocompromised transplant recipient: #Flatteningthecurve. *Am J Transplantat* [Internet] 2020 [cited 2020 Apr 4]. Available from <http://doi.wiley.com/10.1111/ajt.15890>.

[12] Picard C, Le Pavec J, Tissot A. Groupe Transplantation Pulmonaire de la Société de Pneumologie de Langue Française SPLF. Impact of the Covid-19 pandemic and lung transplantation program in France. *Respir Med Res* 2020;78:100758.

[13] World Health Organization. WHO director-general’s opening remarks at the media briefing on COVID-19. WHO; 2020 11 March.

[14] Williamson EJ, Walker AJ, Bhaskaran K, Bacon S, Bates C, Morton CE, et al. Factors associated with COVID-19-related death using OpenSAFELY. *Nature* 2020;584:430–6.

[15] Coll E, Fernández-Ruiz M, Sánchez-Álvarez JE, Martínez-Fernández JR, Crespo M, Gayoso J, et al. COVID-19 in transplant recipients: the Spanish experience. *Am J Transplant* [Internet] 2020. [cited 2020 Nov 28]. Available from <https://onlinelibrary.wiley.com/doi/10.1111/ajt.16369>.

[16] Messika J, Eloy P, Roux A, Hirschi S, Nieves A, Le Pavec J, et al. French group of lung transplantation. COVID-19 in lung transplant recipients. *Transplantation* 2021;105:177–86.

[17] Loupy A, Aubert O, Reese PP, Bastien O, Bayer F, Jacquelinet C. Organ procurement and transplantation during the COVID-19 pandemic. *Lancet* 2020;395:e95–6.

[18] Aubert O, Yoo D, Zielinski D, Cozzi E, Cardillo M, Dürr M, Domínguez-Gil B, Coll E, Da Silva MI, Sallinen V, Lemström K, Midtvedt K, Ulloa C, Immer F, Weissenbacher A, Vallant N, Basic-Jukic N, Tanabe K, Papatheodoridis G, Menoudakou G, Torres M, Soratti C, Hansen Krogh D, Lefaucheur C, Ferreira G, Silva HT, Hartell D, Forsythe J, Mumford L, Reese PP, et al. COVID-19 pandemic and worldwide organ transplantation: a population-based study. *Lancet Public Health* 2021;6:e709–19.

[19] Hallett A, Motter JD, Frey A, Higgins RS, Bush EL, Snyder J, Garonzik-Wang JM, Segev DL, Massie AB. Trends in heart and lung transplantation in the United States across the COVID-19 pandemic. *Transplant Direct* 2021;7:e759.

[20] Ahmed O, Brockmeier D, Lee K, Chapman WC, Doyle MBM. Organ donation during the COVID-19 pandemic. *Am J Transplant*. 2020;20:3081–8.

[21] Hardman G, Sutcliffe R, Hogg R, Mumford L, Grocott L, Mead-Regan S-J, et al. Transplant Cardiothoracic Advisory Group Clinical Audit Group. The impact of the SARS-CoV-2 pandemic and COVID-19 on lung transplantation in the UK: lessons learned from the first wave. *Clin Transplant* 2021;35:e14210.

[22] Legeai C, Malaquin G, Lamotte C, Antoine C, Averland B, Jasseron C, Bayer F, Bastien O, Kerbaul F. Impact of coronavirus disease 2019 on organ donation and transplantation in France. *Transpl. Int.* 2021;34:204–6.

[23] Angelico R, Trapani S, Manzia TM, Lombardini L, Tisone G, Cardillo M. The COVID-19 outbreak in Italy: initial implications for organ transplantation programs. *Am J Transplant* 2020;20:1780–4.

[24] Han W, Zhu M, Chen J, Zhang J, Zhu S, Li T, Cai H, Fang Q, Wei G, Liang T. Lung transplantation for elderly patients with end-stage COVID-19 pneumonia. *Ann Surg* 2020;272:e33–4.

[25] Chen J-Y, Qiao K, Liu F, Wu B, Xu X, Jiao G-Q, et al. Lung transplantation as therapeutic option in acute respiratory distress syndrome for coronavirus disease 2019-related pulmonary fibrosis. *Chin Med J* 2020;133:1390–6.

[26] Bharat A, Machuca TN, Querrey M, Kurihara C, Garza-Castillon R, Kim S, Manerikar A, Pelaez A, Pipkin M, Shahmohammadi A, Rackauskas M, Kg SR, Balakrishnan KR, Jindal A, Schaheen L, Hashimi S, Buddhdev B, Arjuna A, Rosso L, Palleschi A, Lang C, Jaksch P, Budinger GRS, Nosotti M, Hoetzenecker K. Early outcomes after lung transplantation for severe COVID-19: a series of the first consecutive cases from four countries. *Lancet Respir Med* 2021.

[27] Glorion M, De Wolf J, Zuber B, Cassiano F, Preau S, Al Brun, et al. Lung transplantation for COVID-19-associated acute respiratory distress syndrome: the first French patient. *Respirat Med Res* 2021;80:100851.

[28] Yeung JC, Cypel M, Chaparro C, Keshavjee S. Lung transplantation for acute COVID-19: the Toronto lung transplant program experience. *CMAJ* 2021;193:E1494–7.

[29] Sajid F, Ahmed T, Baz MA, Anstead MI. Lung transplantation in a patient with COVID-19-associated acute respiratory failure. *Cureus* 2021;13:e17152.

[30] Lindstedt S, Grins E, Larsson H, Nilsson J, Akbarshahi H, Silva I, Hyllen S, Wagner D, Sjögren J, Hansson L, Ederoth P, Gustafsson R. Lung transplant after 6 months on ECMO support for SARS-CoV-2-induced ARDS complicated by severe antibody-mediated rejection. *BMJ Open Respir Res* 2021;8.

[31] Domjan M, Harlander M, Knafelj R, Ribarič SF, Globokar MD, Gorjup V, et al. Lung transplantation for end-stage respiratory failure after severe COVID-19: a Report of 2 Cases. *Transplant Proc* 2021.

[32] Hawkins RB, Mehaffey JH, Charles EJ, Mannem HC, Roeser M. Lung transplantation for severe post–coronavirus disease 2019 respiratory failure. *Transplantation* 2021;105:1381–7.

[33] Lang C, Jaksch P, Hoda MA, Lang G, Staudinger T, Tschernko E, Zapletal B, Geleff S, Prosch H, Gawish R, Knapp S, Robak O, Thalhammer F, Indra A, Koestenberg M, Strassl R, Klikovits T, Ali K, Fischer G, Klepetko W, Hoetzenecker K, Schellongowski P. Lung transplantation for COVID-19-associated acute respiratory distress syndrome in a PCR-positive patient. *Lancet Respir Med* 2020;8:1057–60.

[34] Cypel M, Keshavjee S. When to consider lung transplantation for COVID-19. *Lancet Respir Med* 2020;8:944–6.

[35] Lepper PM, Langer F, Wilkens H, Schäfers H-J, Bals R. Lung transplantation for COVID-19-associated ARDS. *Lancet Respir Med* 2021(21) S2213–260000278–2.

[36] Messika J, Schmidt M, Tran-Dinh A, Mordant P. Lung transplantation for COVID-19-associated ARDS. *Lancet Respiratory Med* 2021;9:e89.

[37] Burnham EL, Janssen WJ, Riches DWH, Moss M, Downey GP. The fibroproliferative response in acute respiratory distress syndrome: mechanisms and clinical significance. *Eur Respir J* 2014;43:276–85.

[38] Spagnolo P, Balestro E, Aliberti S, Cocconcelli E, Biondini D, Casa GD, Sverzellati N, Maher TM. Pulmonary fibrosis secondary to COVID-19: a call to arms? *Lancet Respir Med* 2020;8:750–2.