



# HHS Public Access

Author manuscript

*Fertil Steril.* Author manuscript; available in PMC 2023 May 01.

Published in final edited form as:

*Fertil Steril.* 2022 May ; 117(5): 1083–1093. doi:10.1016/j.fertnstert.2022.01.013.

## Route of myomectomy and fertility: a prospective cohort study

**Lauren A. Wise, Sc.D.,**

Department of Epidemiology, Boston University School of Public Health, Boston, MA

**Laine Thomas, Ph.D.,**

Department of Biostatistics, Duke University, Durham, NC

**Sophia Anderson, M.P.H.,**

Department of Biostatistics, Duke University, Durham, NC

**Donna D. Baird, Ph.D.,**

Epidemiology Branch, National Institute of Environmental Health Sciences, Durham, NC

**Raymond M. Anchan, M.D.,**

Department of Obstetrics, Gynecology and Reproductive Biology, Brigham and Women's Hospital, Boston, MA

**Kathryn L. Terry, Sc.D.,**

Department of Obstetrics, Gynecology and Reproductive Biology, Brigham and Women's Hospital, Boston, MA

**Erica E. Marsh, M.D., M.S.C.I.,**

Department of Obstetrics and Gynecology, University of Michigan Medical School, Ann Arbor, MI

**Ganesa Wegienka, Ph.D.,**

Department of Public Health Sciences, Henry Ford Health System, Detroit, MI

**Wanda Kay Nicholson, M.D., M.P.H.,**

Center for Women's Health Research, Department of Obstetrics and Gynecology, UNC School of Medicine, Chapel Hill, NC

**Kedra Wallace, Ph.D.,**

Department of Obstetrics and Gynecology, University of Mississippi Medical Center, Jackson, MS

**Robert Bigelow, Ph.D.,**

Duke Clinical Research Institute, Duke University, Durham, NC.

**James Spies, M.D.,**

Department of Radiology, MedStar Georgetown University Hospital in Washington, D.C.

**George L. Maxwell, M.D.,**

---

**Address correspondence to:** Dr. Lauren A. Wise, Department of Epidemiology, Boston University School of Public Health, 715 Albany Street, Boston, MA 02118, 617-358-3424, lwise@bu.edu.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Department of Obstetrics and Gynecology and the Women's Health Integrated Research Center,  
Inova Fairfax Hospital, Falls Church, VA

**Vanessa Jacoby, M.D.,**

School of Medicine, University of California San Francisco, San Francisco CA

**Evan R. Myers, M.D., M.P.H.,**

Department of Obstetrics and Gynecology, Duke University School of Medicine, Durham, NC

**Elizabeth A. Stewart, M.D.**

Department of Obstetrics and Gynecology, Mayo Clinic, Rochester, MN

## Abstract

**Objective:** To assess prospectively the association between myomectomy route and fertility.

**Design:** Prospective cohort study.

**Setting:** The Comparing Treatments Options for Uterine Fibroids (COMPARE-UF) Study is a multi-site national registry of eight clinic centers across the United States ([NCT02260752](https://clinicaltrials.gov/ct2/show/study/NCT02260752), [clinicaltrials.gov](https://clinicaltrials.gov)).

**Patients:** Reproductive-aged women undergoing surgery for symptomatic uterine fibroids.

**Interventions:** N/A.

**Main Outcome Measures:** We used life-table methods to estimate cumulative probabilities and 95% confidence intervals (CI) of pregnancy and live birth by myomectomy route during 12, 24, and 36 months of follow-up (2015–2019). We also conducted 12-month interval-based analyses that used logistic regression to estimate odds ratios (ORs) and 95% confidence intervals for associations of interest. In all analyses, we used propensity score weighting to adjust for differences across surgical routes.

**Results:** Among 1,095 women who underwent myomectomy (abdominal=388, hysteroscopic=273, and laparoscopic=434), 202 reported pregnancy and 91 reported live birth during 36 months of follow-up. There was little difference in the 12-month probability of pregnancy or live birth by route of myomectomy overall, or among women intending pregnancy. In interval-based analyses, adjusted ORs for pregnancy were 1.28 (95% CI: 0.76–2.14) for hysteroscopic myomectomy and 1.19 (95% CI: 0.76–1.85) for laparoscopic myomectomy compared with abdominal myomectomy. Among women intending pregnancy, adjusted ORs were 1.27 (95% CI: 0.72–2.23) for hysteroscopic myomectomy and 1.26 (95% CI: 0.77–2.04) for laparoscopic myomectomy compared with abdominal myomectomy. Associations were slightly stronger but less precise for live birth.

**Conclusions:** The probability of conception or live birth did not differ appreciably by myomectomy route among women followed for 36 months post-surgery.

## Keywords

Fibroids; myomectomy; fertility; pregnancy

## Introduction

Uterine fibroids (UF) are the leading indication for hysterectomy in the United States (U.S.) (1, 2) and account for more than \$2.2 billion annually in health care costs (3). While the lifetime cumulative incidence of clinical diagnosis is approximately 30%, data from standardized screening of women aged 35–49 years estimated a cumulative incidence of ultrasound-detectable UF by age 50 of >70% (4). Studies show a greater burden of UF among Black than White women, with disproportionately higher incidence, earlier ages at diagnosis and surgery, and more severe symptoms at the time of initial diagnosis (4–7). UF symptoms include heavy menstrual bleeding, anemia, pain, pelvic pressure, and genitourinary symptoms. The peak incidence rate for UF symptoms occurs during the reproductive years (7, 8), when many women are attempting pregnancy. Given that Black women tend to experience earlier onset of UF and more severe disease, the impact of UF on their fertility may be greater.

Depending on their location within the uterus, UF may be associated with impaired fertility (9–12). However, studies of UF and fertility are inconsistent and the association may be due in part to uncontrolled or residual confounding (e.g., by age), referral bias (13), or detection bias (7, 14). About 10% of pregnant women have detectable UF in the first trimester (15), and UF presence has been associated with spontaneous abortion in some (16, 17) but not all studies (18). Furthermore, treatments for UF may affect fertility via adverse sequelae. In the U.S., health insurance typically covers UF treatment but not infertility treatment; thus, UF treatment to improve fertility is likely overused. However, research investigating the extent to which fertility outcomes differ based on route of myomectomy is limited (19–22).

Myomectomy, the most common uterine-preserving procedure performed for UF in the U.S. (23–30), accounts for about 22% of all UF surgeries (30). Considering both inpatient and outpatient procedures in the U.S., the most common surgical route for myomectomy is abdominal (via laparotomy) (>75%), followed by laparoscopic (with or without robotic assistance) (~15%) and hysteroscopic (~10%) routes (20, 30–33). According to data from the American College of Surgeons National Surgical Quality Improvement Program database, the percentage of abdominal myomectomies increased by 11 percentage points from 2012 through 2016 (while laparoscopic myomectomy decreased), likely due to concerns about morcellation and cancer (32, 34). Compared with laparoscopic myomectomy, abdominal myomectomy has been associated with longer hospitalizations, higher readmission rates, and greater morbidity (31, 32).

Abdominal myomectomy tends to be more commonly recommended for patients who have larger uterine volume, multiple UF, and UF that cannot be removed easily by other means. In contrast, laparoscopic myomectomy tends to be recommended for women with smaller uterine volume and subserosal/intramural UF (31). Hysteroscopic myomectomy is recommended for patients with symptomatic submucous UF. A 2020 review of surgical treatment of UF for subfertility, including four randomized controlled trials with 442 participants, concluded that there was very low-quality evidence on the association between surgical approach for myomectomy and subsequent subfertility overall and by UF location (20). Observational studies have also been limited by small sample size, retrospective study

design (which may be prone to selection bias), inconsistency of results across studies, and examination of only one type of surgical route (11–13, 16, 19–21, 31, 35–45).

Prospective cohort studies that compare fertility success across surgical approaches for myomectomy can fill important gaps in the literature. In this report, we examine prospectively the association between route of myomectomy (abdominal, hysteroscopic, and laparoscopic) for UF and the probability of conception and live birth during 36 months of follow-up, censoring women with varying lengths of follow-up and adjusting for potential confounding variables. We hypothesize that surgical route of myomectomy would not be strongly associated with fertility outcomes after accounting for differences in patient and UF characteristics across treatment groups. Evidence-based research is critical to generate the information necessary for patients to choose the surgical route for myomectomy that meets their individual needs, goals, and preferences.

## Materials and Methods

The Comparing Treatments Options for Uterine Fibroids (COMPARE-UF) Study is a multi-site national registry of women who were scheduled for treatment for symptomatic fibroids at one of eight clinic centers across the U.S. ([NCT02260752](https://clinicaltrials.gov/ct2/show/study/NCT02260752), [clinicaltrials.gov](https://clinicaltrials.gov)): Mayo Clinic, INOVA Health System, Brigham and Women's, University of Mississippi Medical Center, University of California Fibroid Network, Henry Ford Health System, University of Michigan, and University of North Carolina during 2015–2019. Duke Clinical Research Institute (DCRI) served as the Research Data and Coordinating Center. The primary objective of the registry was to compare prospectively the effectiveness of different surgical and interventional treatment options (hysterectomy, myomectomy, uterine artery embolization) on patient-reported outcomes postoperatively and during 3 years of follow-up using validated general and disease-specific surveys of quality of life. Details on the study design, protocol, and rationale for COMPARE have been published elsewhere (46). The registry protocol was reviewed and approved by the Duke Institutional Review Board for the coordinating center and by the review boards at each of the clinical recruitment sites.

Trained site coordinators screened all women for eligibility. Eligible participants then provided informed consent and HIPPA authorization, and completed an online baseline questionnaire through a secure, password protected, web-based study portal (SignalPath, LLC, Durham, NC) or through paper or phone interview with clinical site coordinators. The baseline questionnaire elicited self-reported data on patient socio-demographics, medical history, fibroid history, prior fibroid procedures, current and prior fibroid therapies, reproductive history, measures of financial distress, and childbearing plans. Per protocol, the baseline questionnaire was completed within the 30-day window before the procedure.

Follow-up questionnaires were completed 12, 24, and 36 months after the procedure. Participants completed questionnaires through the web-based portal, at in-person visits, or via telephone interview with the Research Call Center at the DCRI. The DCRI sent reminders to participants to complete the questionnaires. If a participant was lost to follow-up, coordinators at both the DCRI, and the local recruitment sites attempted to contact the participant using medical records to ascertain any new contact information.

### Assessment of uterine characteristics and myomectomy

Myomectomy was performed according to professional standards and institutional protocols at each clinical site. The choice of myomectomy and surgical route was made independently of COMPARE-UF study protocols. The routes of myomectomy examined in this study included abdominal, hysteroscopic, and laparoscopic. Details about the surgery were obtained from medical records. All participant records, including pelvic imaging reports, were reviewed by a single centralized team of abstractors to ensure consistency across sites. UF details were collected from the participants' imaging reports, which included uterine dimensions and the dimensions of each UF.

### Assessment of fertility and pregnancy outcomes

On annual follow-up questionnaires from 12 months through 36 months post-procedure, women were asked: "In the past year, have you had any pregnancies?" Those who responded "yes" were then asked about the number of pregnancies and the outcome of each pregnancy (up to three pregnancies), with the following response options: "pregnant and not yet delivered," "delivered a single baby," "delivered twins," "delivered triplets," "miscarriage (also known as spontaneous abortion)," "elective or therapeutic abortion," "still birth," or "tubal or ectopic pregnancy." We did not ascertain whether pregnancies were achieved with the use of assisted reproductive technologies (ART).

### Assessment of covariates

We collected self-reported data on socio-demographics at baseline. These factors included age, self-identified race ("How would you best describe your race?" with response options: Black/African-American, Asian, American Indian/Alaska Native, Native Hawaiian/Pacific Islander, White, Other), ethnicity (Hispanic/Latina, Non-Hispanic Latina), reproductive history (gravidity, parity), contraceptive history, body mass index (BMI, kg/m<sup>2</sup>, calculated using self-reported height and weight), marital status, educational level and insurance source. Additional baseline covariate data included clinical factors, such as smoking status, co-morbid conditions (e.g., diabetes, hypertension), gynecologic conditions (sexually transmitted infections, abnormal cervical cytology, polycystic ovarian syndrome), mental health history, and history of prior medical and surgical therapies for UF. Uterine and UF characteristics at baseline, including UF size, number, and location, and uterine volume (cm<sup>3</sup>) were derived from the pretreatment imaging reports.

On the baseline and annual follow-up questionnaires through 36 months, participants were asked about their intentions for pregnancy, specifically whether they were "trying to get pregnant now." If not, women were asked "Are you planning to become pregnant in the future?" with response options: "Yes, likely within the next 2 years," "Would like to keep as an option," and "No." To ascertain infertility history, the baseline questionnaire included the question: "Did you ever try for more than one year to get pregnant?" All follow-up questionnaires included the question: "Have you been trying to get pregnant for a year or more?" At all time points (baseline and follow-up), participants completed the Patient Health Questionnaire-2, a two-item measure to screen for clinical depression (47, 48); the Menopause Rating Scale, a measure of climacteric symptoms (49, 50); the Uterine Fibroid Symptom (UFS)-quality of life (QOL), a disease-specific instrument that assesses symptom

severity and health-related quality of life in women with UF (51), and the visual analog scale (VAS), which is a validated, subjective measure for acute and chronic pain (0=“no pain” and 100=“worst pain”). The post-procedure survey, completed within 11–18 months after the procedure, collected information about the time to resumption of usual activities, interim hospitalizations, procedural complications, and incidental cancer diagnoses.

## Exclusions

The present analysis included all COMPARE-UF eligible patients enrolled from December 28, 2015 through December 17, 2019. We excluded participants who underwent a procedure other than myomectomy because other treatments may have been contraindicated for patients desiring future fertility, and participants who received myomectomy but had missing data on surgical route (Supplemental Figure 1). The final analytic sample for analysis was 1,095 participants: 388 who underwent abdominal myomectomy, 273 who underwent hysteroscopic myomectomy, and 434 who underwent laparoscopic myomectomy.

All sites recruited women in each of these groups with the site-specific myomectomy percentages ranging from 19.2% (University of Mississippi Medical Center) to 71.7% (University of California Fibroid Network).

## Statistical Analysis

We assessed prospectively the association between surgical route for myomectomy (abdominal, hysteroscopic, laparoscopic) and self-reported pregnancy and live birth in each 12-month interval during 36 months of follow-up (2015–2019). We compared socio-demographic and clinical factors between the myomectomy surgical route groups. Our primary events of interest were time from index procedure to occurrence of 1) first pregnancy and 2) first live birth during the follow-up period. Time was measured as 12, 24, or 36 months corresponding to first, second, or third annual follow-up, respectively. Because of the observational design of COMPARE, we used inverse propensity weighting methods, specifically overlap weighting, to adjust for confounding (52, 53). The propensity score (probability of surgical route for myomectomy) was estimated using a multinomial regression model with myomectomy route as the dependent variable and each participant characteristic considered to be potential confounders as independent variables. Potential confounders for inclusion in the propensity model were identified *a priori* based on a review of the literature, clinical experience, and the drawing of a causal diagram, and included age, race/ethnicity, number of prior UF procedures, BMI, history of PCOS, contraception (combined oral contraception, progestin-only oral contraception, patch, vaginal ring, implant, hormone-containing intrauterine device (IUD), progestin-only injectable), number of children (0, 1, 2), fertility intent (currently trying, not currently trying but intend to try within 2 years, not currently trying but intend to try in the future, not interested in future pregnancy), history of difficulty becoming pregnant, and uterine volume. In sensitivity analyses, we further adjusted for the largest UF volume and UF with submucous location. We did not consider “number of UF” in the propensity score due to incomplete data on this variable. Covariates with missing data at baseline were imputed using the fully conditional specification method in SAS PROC MI (54), using all patient-reported variables available in the COMPARE-UF database (46). Given the low percentage of missing data for any



given covariate (<5%), we used a single imputation dataset in this analysis, consistent with previous publications from this registry and prior sensitivity analyses that showed no difference using multiple imputation.

First, we used life-table methods with propensity score weighting to estimate the probabilities of pregnancy and live birth and 95% confidence intervals (CIs) in each time interval (0–12 months, 0–24 months, or 0–36 months), after accounting for censoring. Women were censored at the first occurrence of any of the following events: report of natural or surgical menopause, loss to follow-up, or end of follow-up (36 months). We then conducted a 12-month interval-based analysis that updated pregnancy intent annually and used logistic regression to estimate odds ratios (ORs) and 95% confidence intervals for the association between myomectomy routes and annual outcomes of conception and live birth. The unit of analysis was study time (person-years). The analysis pooled results across all intervals. In all analyses, we included a fixed effect for treatment and used propensity score weighting to create comparable groups by procedure. In addition, we repeated analyses in subgroups restricted by pregnancy intent: actively trying to conceive or intending to conceive within the next 2 years. Potential correlation between patients from the same clinical center was handled by fitting a robust empirical variance estimator, with clustering by clinical center.

We performed sensitivity analyses that excluded women with hysteroscopic myomectomy as a comparison group, owing to the large differences in patient and UF characteristics between these women and all other participants. This involved re-running the propensity score weighting to balance the UF characteristics across the abdominal and laparoscopic myomectomy groups, the life-table analyses, and logistic regression models for associations with pregnancy and livebirth. A subsequent sensitivity analysis was conducted to account for additional UF characteristics: maximum UF volume and submucous location. These variables were not included in the primary propensity model because their method of collection was not standardized across clinical sites and they were thought to be captured less accurately than uterine volume. All analyses were conducted using SAS (Cary, North Carolina).

## Results

### Baseline socio-demographic and clinical characteristics

There were no appreciable differences in the percentages lost to follow-up by myomectomy group (data not shown). Participants who underwent abdominal myomectomy tended to be younger, nulliparous, have larger uterine volume at surgery, larger maximum UF volume, and were more likely to identify as Black or African American than women who underwent other routes of myomectomy (Table 1). They were also more likely to be currently trying (29.1%) or planning to try to conceive within the next two years (32.3%), relative to the other routes of myomectomy. Hysteroscopic myomectomy patients were substantially more likely than the other two myomectomy groups to have 2 or more prior UF procedures. There was little difference in history of infertility across the three groups. Likewise, the groups were not notably different with respect to UFS-QOL, EQ, or VAS scores. Comparability

between the treatment groups was achieved at baseline after propensity weighting (data not shown).

### **Probabilities of pregnancy and live birth, overall and by myomectomy route**

Among 1,095 women who underwent myomectomy, 202 reported pregnancy and 91 reported live birth during follow-up; some of these women were still pregnant at the end of follow-up. There was no appreciable difference in the probability of pregnancy or live birth by route of myomectomy overall, among women intending pregnancy within 2 years, or among women actively trying to conceive (Table 2).

Among women who had a myomectomy, the strongest predictors of reported conception were age and pregnancy intent at baseline (data not shown). Among women who reported currently trying to conceive at baseline, the cumulative probabilities of pregnancy within the first year of follow-up for women aged 30, 31–39, 40–44, and 45 years were: 0.37 (95% CI: 0.24–0.47), 0.29 (95% CI: 0.23–0.35), 0.16 (95% CI: 0.09–0.22), and 0.14 (95% CI: 0.03–0.23). After three years, those respective cumulative probabilities of pregnancy increased to: 0.67 (95% CI: 0.48–0.79), 0.57 (95% CI: 0.47–0.65), 0.34 (95% CI: 0.21–0.45), and 0.30 (95% CI: 0.08–0.47). This statistical model had a Harrell's C-index of 0.80 (55), indicating very good prediction. Other variables in this model that did not appreciably improve prediction included: myomectomy route, use of contraception at baseline, number of prior UF procedures, parity, and infertility history.

### **Odds ratios of pregnancy and live birth, by myomectomy route**

Overall, adjusted ORs for pregnancy were 1.28 (95% CI: 0.76–2.14) for hysteroscopic myomectomy and 1.19 (95% CI: 0.76–1.85) for laparoscopic myomectomy compared with abdominal myomectomy (Table 3). There was evidence of substantial confounding by patient and UF characteristics, as evidenced by the attenuation of unadjusted ORs following adjustment for confounding using propensity weighting.

Among women intending pregnancy, ORs were 1.27 (95% CI: 0.72–2.23) for hysteroscopic myomectomy and 1.26 (95% CI: 0.77–2.04) for laparoscopic myomectomy compared with abdominal myomectomy. Associations were similar for live birth.

When we repeated analyses after excluding women with hysteroscopic myomectomy and derived new propensity weights to account for patient and UF differences across laparoscopic versus abdominal myomectomy, results were similar when comparing these two surgical approaches (Table 4). Additional adjustment for maximum UF size and submucous UF also yielded consistent findings (Table 4).

## **Discussion**

In this prospective analysis of COMPARE-UF participants undergoing myomectomy for symptomatic UF, there was little association between surgical route for myomectomy and the probability of conception or livebirth during a 36-month follow-up period, after adjusting for patient demographics, reproductive history, and uterine volume. Among myomectomy patients, the strongest predictors of pregnancy success were age and pregnancy intent



at baseline. Among women who reported currently trying to conceive at baseline, the cumulative probabilities of pregnancy during three years of follow-up, after accounting for varying lengths of follow-up using life-table methods, were as high as 67% among women aged 30 or younger and 30% among women aged 45 years. The model with age and pregnancy intent showed excellent predictive probability for pregnancy (80%); after accounting for age and pregnancy intent, myomectomy route was not an important predictor of pregnancy. These results contribute to the sparse literature on the influence of surgical route of myomectomy and fertility outcomes.

Large differences in pre-treatment patient characteristics were observed across the different routes of myomectomy. These differences are not surprising given that procedures like abdominal myomectomy are typically recommended for women with larger uterine volumes, and larger and more numerous UF. Although we successfully adjusted for many of the observed differences using propensity weighting, this approach includes assumptions that may not fully capture the severity of UF characteristics among women who underwent abdominal myomectomy (e.g., setting mean uterine volume to 300 cm<sup>3</sup> for all subtypes of myomectomy, even though the mean volume for all women with abdominal myomectomy was ~900 cm<sup>3</sup>). To increase the generalizability of our findings, we repeated our analyses after excluding women with hysteroscopic myomectomy, for whom UF disease severity would be lower relative to women undergoing abdominal or laparoscopic myomectomy. The analyses restricted to abdominal and laparoscopic myomectomy focused on treatments with better covariate overlap. Such a comparison would better emulate the real-life situation where a given patient might be eligible for abdominal or laparoscopic myomectomy, but not hysteroscopic myomectomy. Again, these results showed little evidence for a difference in pregnancy or live birth comparing abdominal and laparoscopic myomectomy surgical routes. Thus, our results indicate that the choice of abdominal vs. laparoscopic myomectomy for women with UF that cannot be appropriately treated via the hysteroscopic route can be based on other considerations besides future fertility.

Although myomectomy is the treatment of choice for women desiring to preserve their fertility(24–29) and is frequently performed among women with unexplained infertility or recurrent spontaneous abortion, its effect on subsequent fertility is unclear (11–13, 16, 20, 21, 35–43). One study reported that the surgical route of myomectomy has little effect on fertility outcomes (44), but other studies have not directly compared surgical routes of myomectomy (19, 31, 45). One difficulty in comparing outcomes in nonrandomized participants is that important factors associated with pregnancy outcomes, particularly age and UF characteristics, may differ between groups prior to treatment (56, 57). For example, a recent study reported that women with greater than 6 UF removed were less likely to conceive; however, it was unclear whether this was related to the severity of UF itself or to the surgical procedure (19). Moreover, non-patient factors, including the skill of the surgeon and the availability and/or use of ART following UF treatment may have influenced study outcomes as well.

Strengths of the analysis include the prospective design, 3-year follow-up period, relatively large sample size, the availability of pre-treatment medical and operative notes, detailed covariate information, and the application of validated propensity weighting methods to

account for differences in pre-treatment characteristics (e.g., uterine anatomy—overall uterine size; number, size, and location of individual UF). There are limited comparative data on fertility outcomes among women undergoing myomectomy, and thus observational studies that are prospective in design can make an important contribution to the literature.

Limitations of the study include the restriction of analyses to women undergoing myomectomy only and potential unmeasured differences in the distribution of uterine anatomy characteristics across myomectomy procedures, which could have introduced residual confounding by indication. However, sensitivity analyses that included additional UF characteristics in the propensity score (e.g., location and size of largest UF) had little impact on the results. To the extent that confounding was not properly accounted for, we might expect lower fertility success among women undergoing abdominal myomectomy relative to the other types of myomectomy because women offered abdominal myomectomy tend to have more severe disease (e.g., larger and more numerous UF; submucous UF which could be more strongly associated with inhibition of implantation) (7, 8). Many of the demographic characteristics that are more common among women with severe UF (e.g., later reproductive age, African ancestry) are also risk factors for adverse reproductive outcomes such as infertility and spontaneous abortion (58, 59), and could confound the potential association between myomectomy route and these outcomes (7). This, in turn, limits our ability to compare fertility across different treatments. As mentioned above, propensity weighting may have made the results less generalizable to women with more severe UF who undergo abdominal myomectomy. Whether it is even appropriate to compare abdominal with laparoscopic and hysteroscopic myomectomy is debatable given a single patient may never be offered all three of these options. However, the extent to which differences in pre-operative uterine anatomy or other UF characteristics alone, independent of route of procedure, would have had a direct effect on fertility outcomes is unclear. Lack of data on specific types of reproductive failures such as fertilization, implantation, or post-implantation losses precluded the examination of potential mechanisms. We did not have data on whether women used fertility treatments to conceive or whether they conceived spontaneously, and differences in these factors may have obscured differences in fertility success among the surgical routes for myomectomy (60–64).

The COMPARE-UF data were collected from a convenience sample of patients undergoing UF procedures at 10 clinical sites across the U.S.; thus, the prevalence of myomectomy subtypes in this population is not representative of the general population. The primary eligibility criterion for inclusion in the COMPARE-UF registry was the presence of symptomatic UF, including subfertility as a syndrome. The proportion of women undergoing hysteroscopic resection reflects the distribution of women with UF suitable for hysteroscopic resection among our study population, the majority of whom were not actively trying to get pregnant. We also note that live birth rates were partly limited by varying lengths of follow-up. If patients were advised to wait 4–6 months post-procedure before attempting to conceive and had average fecundability, the first births would not take place until after 12 months of follow-up.

Another important limitation is that we relied on clinical imaging and operative reports at participating clinical sites to characterize the location of the UF being removed. Though

reports were abstracted using a standard form that included data on FIGO stage, fewer than 2% of COMPARE-UF reports used the FIGO classification. The general categorization of UF into submucous, intramural, and subserosal has been in practice for several decades, and there is some clinical and some basic science evidence to indicate that submucous UF are more likely to contribute to infertility given their ability to cause uterine cavity distortion (11). There is also evidence that removal of submucous UF increases subsequent pregnancy rates (11). However, controversy remains about the role of intramural UF in the pathogenesis of infertility (65). In a recently-published debate (65), experts cited several mechanisms by which intramural UF could influence fertility, including impaired endometrial and myometrial blood supply, reduced endometrial receptivity, greater myometrial contractility, thickening of the UF capsule, and hormonal and genetic alterations, all of which favored removal of intramural UF to improve fertility. Other experts argued against removal of intramural UF to improve fertility, citing concerns about surgical complications and challenges in the interpretation of published studies due to methodologic issues such as confounding, biologic heterogeneity (e.g., driver mutations; FIGO type 3 vs. 4), and selection bias related to differential referral patterns and insurance coverage for UF care (65–68). Conversely, there is general agreement that subserosal UF have limited, if any, impact on fertility although data are also limited, particularly for larger UF (11). Finally, comparing fertility in women with intramural UF surrounded by myometrium (FIGO type 4) with those that contact the endometrium (FIGO type 3) is a novel area of investigation (65, 66), but was beyond the scope of this report.

Results from the present study indicate that there is little difference in the probability of pregnancy or live birth during 36 months of follow-up according to surgical route of myomectomy, particularly when comparing abdominal vs. laparoscopic routes, after accounting for pre-treatment differences in patient characteristics. Additional follow-up may be needed to determine if the similarity in fertility outcomes across myomectomy groups persists over time. If confirmed, our results provide little reason for change in how current myomectomy route is chosen by patients in consultation with their providers in regard to a patient's desire for future fertility.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments:

This study was supported by grant number P50HS023418 from the Agency for Healthcare Research and Quality (AHRQ) with funding provided by the Patient-Centered Outcomes Research Institute (PCORI) under MOU number 2013-001. The content of this manuscript is solely the responsibility of the authors and readers should not interpret any statement in this product as an official position or the views of AHRQ, the U.S. Department of Health and Human Services, or PCORI.

## References

1. Wilcox LS, Koonin LM, Pokras R, Strauss LT, Xia Z, Peterson HB. Hysterectomy in the United States, 1988–1990. *Obstet Gynecol* 1994;83:549–55. [PubMed: 8134065]
2. Farquhar CM, Steiner CA. Hysterectomy rates in the United States 1990–1997. *Obstet Gynecol* 2002;99(2):229–34. [PubMed: 11814502]

3. Flynn M, Jamison M, Datta S, Myers E. Health care resource use for uterine fibroid tumors in the United States. *Am J Obstet Gynecol* 2006;195(4):955–64. [PubMed: 16723104]
4. Baird DD, Dunson DB, Hill MC, Cousins D, Schectman JM. High cumulative incidence of uterine leiomyoma in black and white women: ultrasound evidence. *Am J Obstet Gynecol* 2003;188(1):100–7. [PubMed: 12548202]
5. Marshall LM, Spiegelman D, Barbieri RL, Goldman MB, Manson JE, Colditz GA, et al. Variation in the incidence of uterine leiomyoma among premenopausal women by age and race. *Obstet Gynecol* 1997;90(6):967–73. [PubMed: 9397113]
6. Kjerulff KH, Langenberg P, Seidman JD, Stolley PD, Guzinski GM. Uterine leiomyomas: racial differences in severity, symptoms, and age at diagnosis. *J Reprod Med* 1996;41(7):483–90. [PubMed: 8829060]
7. Wise LA, Laughlin-Tommaso SK. Epidemiology of Uterine Fibroids: From Menarche to Menopause. *Clin Obstet Gynecol* 2016;59(1):2–24. [PubMed: 26744813]
8. Laughlin SK, Schroeder JC, Baird DD. New directions in the epidemiology of uterine fibroids. *Semin Reprod Med* 2010;28(3):204–17. [PubMed: 20414843]
9. Whynott RM, Vaught KCC, Segars JH. The Effect of Uterine Fibroids on Infertility: A Systematic Review. *Semin Reprod Med* 2017;35(6):523–32. [PubMed: 29100240]
10. Metwally M, Farquhar CM, Li TC. Is another meta-analysis on the effects of intramural fibroids on reproductive outcomes needed? *Reprod Biomed Online* 2011;23(1):2–14. [PubMed: 21549642]
11. Pritts EA, Parker WH, Olive DL. Fibroids and infertility: an updated systematic review of the evidence. *Fertil Steril* 2009;91(4):1215–23. [PubMed: 18339376]
12. Sunkara SK, Khairy M, El-Toukhy T, Khalaf Y, Coomarasamy A. The effect of intramural fibroids without uterine cavity involvement on the outcome of IVF treatment: a systematic review and meta-analysis. *Hum Reprod* 2010;25(2):418–29. [PubMed: 19910322]
13. Johnson G, MacLehose RF, Baird DD, Laughlin-Tommaso SK, Hartmann KE. Uterine leiomyomata and fecundability in the Right from the Start study. *Hum Reprod* 2012;27(10):2991–7. [PubMed: 22811308]
14. Schwartz SM, Marshall LM, Baird DD. Epidemiologic contributions to understanding the etiology of uterine leiomyomata. *Environ Health Perspect* 2000;108(Suppl 5):821–7. [PubMed: 11035989]
15. Laughlin SK, Baird DD, Savitz DA, Herring AH, Hartmann KE. Prevalence of Uterine Leiomyomas in the First Trimester of Pregnancy: An Ultrasound-Screening Study. *Obstetrics & Gynecology* 2009;113(3):630–5. [PubMed: 19300327]
16. Klatsky PC, Tran ND, Caughey AB, Fujimoto VY. Fibroids and reproductive outcomes: a systematic literature review from conception to delivery. *Am J Obstet Gynecol* 2008;198(4):357–66. [PubMed: 18395031]
17. Lai J, Caughey AB, Qidwai GI, Jacoby AF. Neonatal outcomes in women with sonographically identified uterine leiomyomata. *J Matern Fetal Neonatal Med* 2012;25(6):710–3. [PubMed: 22409539]
18. Hartmann KE, Velez Edwards DR, Savitz DA, Jonsson-Funk ML, Wu P, Sundermann AC, et al. Prospective Cohort Study of Uterine Fibroids and Miscarriage Risk. *Am J Epidemiol* 2017;186(10):1140–8. [PubMed: 28591761]
19. Shue S, Radeva M, Falcone T. Comparison of Long-Term Fertility Outcomes after Myomectomy: Relationship with Number of Myomas Removed. *J Minim Invasive Gynecol* 2018;25(6):1002–8. [PubMed: 29371170]
20. Metwally M, Raybould G, Cheong YC, Horne AW. Surgical treatment of fibroids for subfertility. *Cochrane Database Syst Rev* 2020;1:CD003857. [PubMed: 31995657]
21. Khaw SC, Anderson RA, Lui MW. Systematic review of pregnancy outcomes after fertility-preserving treatment of uterine fibroids. *Reprod Biomed Online* 2020;40(3):429–44. [PubMed: 32081542]
22. Dubuisson JB, Chapron C, Fauconnier A, Babaki-Fard K. Laparoscopic myomectomy fertility results. *Ann N Y Acad Sci* 2001;943:269–75. [PubMed: 11594546]
23. Management of Symptomatic Uterine Leiomyomas: ACOG Practice Bulletin, Number 228. *Obstet Gynecol* 2021;137(6):e100–e15. [PubMed: 34011888]

24. Malone LJ. Myomectomy: recurrence after removal of solitary and multiple myomas. *Obstet Gynecol* 1969;34(2):200–3. [PubMed: 5816311]
25. Wechter ME, Stewart EA, Myers ER, Kho RM, Wu JM. Leiomyoma-related hospitalization and surgery: prevalence and predicted growth based on population trends. *Am J Obstet Gynecol* 2011;205(5):492 e1–5. [PubMed: 22035951]
26. Jin C, Hu Y, Chen XC, Zheng FY, Lin F, Zhou K, et al. Laparoscopic versus open myomectomy--a meta-analysis of randomized controlled trials. *Eur J Obstet Gynecol Reprod Biol* 2009;145(1):14–21. [PubMed: 19398260]
27. Kotani Y, Tobiume T, Fujishima R, Shigeta M, Takaya H, Nakai H, et al. Recurrence of uterine myoma after myomectomy: Open myomectomy versus laparoscopic myomectomy. *J Obstet Gynaecol Res* 2018;44(2):298–302. [PubMed: 29227004]
28. Stewart EA, Faur AV, Wise LA, Reilly RJ, Harlow BL. Predictors of subsequent surgery for uterine leiomyomata after abdominal myomectomy. *Obstet Gynecol* 2002;99(3):426–32. [PubMed: 11864669]
29. Reed SD, Newton KM, Thompson LB, McCrummen BA, Warolin AK. The incidence of repeat uterine surgery following myomectomy. *J Womens Health (Larchmt)* 2006;15(9):1046–52. [PubMed: 17125423]
30. Barrett ML, Weiss AJ, Stocks C, Steiner CA, Myers ER. Procedures to Treat Benign Uterine Fibroids in Hospital Inpatient and Hospital-Based Ambulatory Surgery Settings, 2013. HCUP Statistical Brief #200. January 2016 Agency for Healthcare Research and Quality, Rockville, MD, 2016.
31. Hartmann KE, Fennesbeck C, Surawicz T, Krishnaswami S, Andrews JC, Wilson JE, et al. Management of Uterine Fibroids. Comparative Effectiveness Review No. 195. (Prepared by the Vanderbilt Evidence-based Practice Center under Contract No. 290–2015-00003-I.) AHRQ Publication No. 17(18)-EHC028-EF Rockville, MD: Agency for Healthcare Research and Quality, 2017.
32. Stentz NC, Cooney LG, Sammel M, Shah DK. Changes in Myomectomy Practice After the U.S. Food and Drug Administration Safety Communication on Power Morcellation. *Obstet Gynecol* 2017;129(6):1007–13. [PubMed: 28486366]
33. Barrett ML, Weiss AJ, Stocks C, Steiner CA, Myers ER. Procedures to Treat Benign Uterine Fibroids in Hospital Inpatient and Hospital-Based Ambulatory Surgery Settings, 2013: Statistical Brief #200. Healthcare Cost and Utilization Project (HCUP) Statistical Briefs Rockville (MD) 2006.
34. Matsushita T, Sekizawa A, Jacobs LK. Racial Disparities in Response to a US Food and Drug Administration Safety Communication Regarding the Use of Power Morcellation for the Treatment of Uterine Leiomyoma. *J Minim Invasive Gynecol* 2020;27(1):178–85 e1. [PubMed: 30936031]
35. Practice Committee of the American Society for Reproductive Medicine. Electronic address Aao, Practice Committee of the American Society for Reproductive M. Removal of myomas in asymptomatic patients to improve fertility and/or reduce miscarriage rate: a guideline. *Fertil Steril* 2017;108(3):416–25. [PubMed: 28865538]
36. Brady PC, Stanic AK, Styer AK. Uterine fibroids and subfertility: an update on the role of myomectomy. *Curr Opin Obstet Gynecol* 2013;25(3):255–9. [PubMed: 23562956]
37. Lebovitz O, Orvieto R, James KE, Styer AK, Brown DN. Predictors of reproductive outcomes following myomectomy for intramural fibroids. *Reprod Biomed Online* 2019;39(3):484–91. [PubMed: 31378690]
38. Gavai M, Berkes E, Lazar L, Fekete T, Takacs ZF, Urbancsek J, et al. Factors affecting reproductive outcome following abdominal myomectomy. *J Assist Reprod Genet* 2007;24(11):525–31. [PubMed: 18008158]
39. Kelly BA, Bright P, Mackenzie IZ. Does the surgical approach used for myomectomy influence the morbidity in subsequent pregnancy? *J Obstet Gynaecol* 2008;28(1):77–81. [PubMed: 18259905]
40. Pinto Pabon I, Magret JP, Unzurrunzaga EA, Garcia IM, Catalan IB, Cano Vieco ML. Pregnancy after uterine fibroid embolization: follow-up of 100 patients embolized using tris-acryl gelatin microspheres. *Fertil Steril* 2008;90(6):2356–60. [PubMed: 18339388]



41. Sinclair D, Gaither K, Mason TC. Fertility outcomes following myomectomy in an urban hospital setting. *J Natl Med Assoc* 2005;97(10):1346–8. [PubMed: 16353656]
42. Surrey ES, Minjarez DA, Stevens JM, Schoolcraft WB. Effect of myomectomy on the outcome of assisted reproductive technologies. *Fertil Steril* 2005;83(5):1473–9. [PubMed: 15866587]
43. Vimercati A, Scioscia M, Lorusso F, Laera AF, Lamanna G, Coluccia A, et al. Do uterine fibroids affect IVF outcomes? *Reprod Biomed Online* 2007;15(6):686–91. [PubMed: 18062866]
44. Metwally M, Cheong YC, Horne AW. Surgical treatment of fibroids for subfertility. *Cochrane Database Syst Rev* 2012;11:CD003857. [PubMed: 23152222]
45. Stewart EA, Laughlin-Tommaso SK, Catherino WH, Lalitkumar S, Gupta D, Vollenhoven B. Uterine fibroids. *Nat Rev Dis Primers* 2016;2:16043. [PubMed: 27335259]
46. Stewart EA, Lytle BL, Thomas L, Wegienka GR, Jacoby V, Diamond MP, et al. The Comparing Options for Management: PATient-centered REsults for Uterine Fibroids (COMPARE-UF) registry: rationale and design. *Am J Obstet Gynecol* 2018;219(1):95 e1–e10. [PubMed: 29750955]
47. Spitzer RL, Kroenke K, Williams JB. Validation and utility of a self-report version of PRIME-MD: the PHQ primary care study. *Primary Care Evaluation of Mental Disorders. Patient Health Questionnaire. JAMA* 1999;282(18):1737–44. [PubMed: 10568646]
48. Kroenke K Patients presenting with somatic complaints: epidemiology, psychiatric comorbidity and management. *Int J Methods Psychiatr Res* 2003;12(1):34–43. [PubMed: 12830308]
49. Heinemann K, Ruebig A, Potthoff P, Schneider HP, Strelow F, Heinemann LA, et al. The Menopause Rating Scale (MRS) scale: a methodological review. *Health Qual Life Outcomes* 2004;2:45. [PubMed: 15345062]
50. Zollner YF, Acquadro C, Schaefer M. Literature review of instruments to assess health-related quality of life during and after menopause. *Qual Life Res* 2005;14(2):309–27. [PubMed: 15892422]
51. Spies JB, Coyne K, Guaou G, Boyle D, Skyrnarz-Murphy K, Gonzalves SM. The UFSQ-OL, a new disease-specific symptom and health-related quality of life questionnaire for leiomyomata. *Obstet Gynecol* 2002;99(2):290–300. [PubMed: 11814511]
52. Li F, Morgan KL, Zaslavsky AM. Balancing covariates via propensity score weighting. *Journal of the American Statistical Association* 2018;113(521):390–400.
53. Li F, Thomas LE, Li F. Addressing Extreme Propensity Scores via the Overlap Weights. *Am J Epidemiol* 2019;188(1):250–7. [PubMed: 30189042]
54. Austin PC, White IR, Lee DS, van Buuren S. Missing Data in Clinical Research: A Tutorial on Multiple Imputation. *Can J Cardiol* 2020.
55. Harrell FE Jr., Califf RM, Pryor DB, Lee KL, Rosati RA. Evaluating the yield of medical tests. *JAMA* 1982;247(18):2543–6. [PubMed: 7069920]
56. American College of Obstetricians Gynecologists' Committee on Practice Bulletins-Gynecology. Management of Symptomatic Uterine Leiomyomas: ACOG Practice Bulletin, Number 228. *Obstet Gynecol* 2021;137(6):e100–e15. [PubMed: 34011888]
57. Burke CT, Funaki BS, Ray CE Jr., Kinney TB, Kostelic JK, Loesberg A, et al. ACR Appropriateness Criteria(R) on treatment of uterine leiomyomas. *J Am Coll Radiol* 2011;8(4):228–34. [PubMed: 21458760]
58. Berkowitz GS, Blackmore-Prince C, Lapinski RH, Savitz DA. Risk factors for preterm birth subtypes. *Epidemiology* 1998;9(3):279–85. [PubMed: 9583419]
59. Goldenberg RL, Culhane JF, Iams JD, Romero R. Epidemiology and causes of preterm birth. *Lancet* 2008;371(9606):75–84. [PubMed: 18177778]
60. Reddy UM, Wapner RJ, Rebar RW, Tasca RJ. Infertility, assisted reproductive technology, and adverse pregnancy outcomes: executive summary of a National Institute of Child Health and Human Development workshop. *Obstet Gynecol* 2007;109(4):967–77. [PubMed: 17400861]
61. Luke B, Brown MB. Elevated risks of pregnancy complications and adverse outcomes with increasing maternal age. *Hum Reprod* 2007;22(5):1264–72. [PubMed: 17289684]
62. Basso O, Baird DD. Infertility and preterm delivery, birthweight, and Caesarean section: a study within the Danish National Birth Cohort. *Hum Reprod* 2003;18(11):2478–84. [PubMed: 14585905]



63. Wise LA, Mikkelsen EM, Sorensen HT, Rothman KJ, Hahn KA, Riis AH, et al. Prospective study of time to pregnancy and adverse birth outcomes. *Fertil Steril* 2015;103(4):1065–73. [PubMed: 25707334]
64. Messerlian C, Maclagan L, Basso O. Infertility and the risk of adverse pregnancy outcomes: a systematic review and meta-analysis. *Hum Reprod* 2013;28(1):125–37. [PubMed: 23042798]
65. Dolmans MM, Isaacson K, Zhang W, Gordts S, Munro MG, Stewart EA, et al. Intramural myomas more than 3–4 centimeters should be surgically removed before in vitro fertilization. *Fertil Steril* 2021;116(4):945–58. [PubMed: 34579828]
66. Munro MG. Uterine polyps, adenomyosis, leiomyomas, and endometrial receptivity. *Fertil Steril* 2019;111(4):629–40. [PubMed: 30929720]
67. Tanos V, Balami S, Lingwood L. Junctional zone endometrium alterations in gynecological and obstetrical disorders and impact on diagnosis, prognosis and treatment. *Curr Opin Obstet Gynecol* 2019;31(6):418–27. [PubMed: 31573995]
68. Tanos V, Lingwood L, Balami S. The importance of the junctional zone of the endometrium in human reproduction. *Hum Fertil (Camb)* 2020:1–9.

**Table 1:**

Baseline characteristics of COMPARE-UF participants by surgical route of myomectomy

Characteristic	Myomectomy route			Total
	Abdominal	Laparoscopic	Hysteroscopic	
Number of women	388	434	273	1,095
Age (years), mean (SD)	36.8 (5.7)	37.3 (5.9)	40.9 (7.2)	38.0 (6.4)
30	49 (12.6%)	53 (12.2%)	25 (9.2%)	127 (11.6%)
31–39	218 (56.2%)	227 (52.3%)	88 (32.2%)	533 (48.7%)
40–44	85 (21.9%)	103 (23.7%)	73 (26.7%)	261 (23.8%)
45	36 (9.3%)	51 (11.8%)	87 (31.9%)	174 (15.9%)
Race				
Black or African American	198 (51.0%)	154 (35.8%)	104 (38.2%)	456 (41.8%)
White	118 (30.4%)	188 (43.7%)	127 (46.7%)	433 (39.7%)
Other	72 (18.6%)	88 (20.5%)	41 (15.1%)	201 (18.4%)
Hispanic or Latina	27 (7.0%)	21 (5.0%)	29 (10.9%)	77 (7.2%)
Body mass index (kg/m <sup>2</sup> ), mean (SD)	29.0 (7.1)	27.7 (7.2)	30.0 (8.9)	28.7 (7.7)
History of polycystic ovary syndrome (PCOS)	23 (6.1%)	26 (6.2%)	14 (5.1%)	63 (5.9%)
Contraception to prevent pregnancy				
Combined oral contraception, patch, or ring	38 (9.8%)	24 (5.5%)	23 (8.4%)	85 (7.8%)
Progestin-only implant	18 (4.6%)	8 (1.8%)	8 (2.9%)	34 (3.1%)
Progestin-only oral contraception	23 (5.9%)	11 (2.5%)	6 (2.2%)	40 (3.7%)
Hormone-containing intrauterine device	19 (4.9%)	7 (1.6%)	6 (2.2%)	32 (2.9%)
Progestin-only injectable	20 (5.2%)	8 (1.8%)	9 (3.3%)	37 (3.4%)
Fertility planning status				
Currently trying	113 (29.1%)	124 (28.6%)	52 (19.0%)	289 (26.4%)
Not currently trying, but within 2 years	125 (32.2%)	116 (26.7%)	41 (15.0%)	282 (25.8%)
Not currently trying, but keeping option open for future	109 (28.1%)	104 (24.0%)	46 (16.8%)	259 (23.7%)
Not currently trying, not interested in future pregnancy	41 (10.6%)	87 (20.0%)	133 (48.7%)	261 (23.8%)
Parity (number of births)				
0	315 (81.2%)	335 (77.2%)	143 (52.4%)	793 (72.4%)
1	46 (11.9%)	62 (14.3%)	43 (15.8%)	151 (13.8%)
2	27 (7.0%)	37 (8.5%)	87 (31.9%)	151 (13.8%)
History of difficulty conceiving	101 (27.1%)	117 (28.0%)	68 (25.2%)	286 (27.0%)
<b>Fibroid characteristics</b>				
Number of prior fibroid procedures				
0	317 (81.7%)	371 (85.5%)	215 (78.8%)	903 (82.5%)
1	66 (17.0%)	52 (12.0%)	49 (18.0%)	167 (15.3%)
2	5 (1.3%)	11 (2.5%)	9 (43.3%)	25 (2.3%)
Uterine volume (cm <sup>3</sup> ), mean (SD)	912.8 (737)	486.4 (390)	265.6 (275)	580.5 (585)
Maximum fibroid volume (cm <sup>3</sup> ), mean (SD)	483.5 (743.4)	265.3 (313.6)	72.9 (425.9)	295.2 (552.2)
Any submucous fibroid				
Yes	105 (27.1%)	96 (22.1%)	172 (63.0%)	373 (34.1%)

Characteristic	Myomectomy route			Total
	Abdominal	Laparoscopic	Hysteroscopic	
No/missing <sup>a</sup>	283 (72.9%)	338 (77.9%)	101 (37.0%)	722 (65.9%)
UFS-QOL, mean (SD)				
Concern	47.1 (32.2)	52.6 (33.9)	37.2 (28.4)	46.8 (32.6)
Activities	52.7 (28.8)	56.3 (29.2)	50.0 (29.1)	53.5 (29.1)
Energy/mood	50.3 (27.7)	52.8 (28.3)	49.2 (27.9)	51.0 (28.0)
Control	48.7 (27.0)	51.7 (27.6)	48.8 (26.8)	49.9 (27.2)
Self-conscious	39.8 (31.6)	50.2 (31.6)	52.7 (32.7)	47.3 (32.3)
Sexual function	54.3 (34.8)	55.9 (33.7)	49.8 (35.7)	53.8 (34.7)
Total summary of 6 subscale scores above	49.5 (25.8)	53.4 (25.9)	47.3 (26.0)	50.6 (26.0)
Symptom severity	51.6 (25.6)	48.6 (23.9)	54.4 (24.6)	51.1 (24.8)
EQ-5-Dimension Scale (% without problems)				
Mobility	314 (81.6%)	378 (87.7%)	228 (83.8%)	920 (84.6%)
Self-care	365 (95.1%)	416 (96.7%)	255 (93.4%)	1036 (95.3%)
Usual activities	250 (65.1%)	287 (66.9%)	192 (70.3%)	729 (67.1%)
Pain/discomfort	80 (20.8%)	108 (25.1%)	101 (37.0%)	289 (26.5%)
Anxiety/depression	156 (40.7%)	163 (38.0%)	112 (41.2%)	431 (39.8%)
Visual Analogue Pain Scale (0–100) (VAS), mean (SD)	72.5 (19.9)	74.7 (16.7)	73.3 (18.2)	73.6 (18.3)

Abbreviations: UFS-QOL=Uterine Fibroid Symptom and Quality of Life subscale. EQ=EuroQOL scale. SD=standard deviation.

<sup>a</sup>Unable to distinguish “no” from “missing” due to lack of standardization and inconsistent reporting of imaging data across clinical sites.

**Table 2:**

Cumulative probability of pregnancy and live birth during follow-up, by myomectomy route<sup>a</sup>

Myomectomy Route	Subgroup	Pregnancies/Total women (%)	Probability of pregnancy (95% CI) by follow-up time		
			12 months	24 months	36 months
Abdominal	All women	69/388 (17.8%)	0.13 (0.08–0.14)	0.20 (0.16–0.25)	0.24 (0.19–0.30)
	Trying to conceive or intending to conceive within the next 2 years <sup>b</sup>	64/238 (26.9%)	0.25 (0.17–0.34)	0.37 (0.27–0.48)	0.45 (0.33–0.59)
	Trying to conceive <sup>b</sup>	36/113 (31.9%)	0.28 (0.18–0.42)	0.41 (0.28–0.57)	0.47 (0.30–0.66)
Hysteroscopic	All women	37/273 (13.6%)	0.16 (0.11–0.22)	0.24 (0.17–0.32)	0.33 (0.23–0.45)
	Trying to conceive or intending to conceive within the next 2 years <sup>b</sup>	34/94 (36.2%)	0.26 (0.17–0.38)	0.41 (0.29–0.55)	0.56 (0.40–0.74)
	Trying to conceive <sup>b</sup>	22/53 (41.5%)	0.36 (0.24–0.53)	0.48 (0.33–0.66)	0.63 (0.42–0.83)
Laparoscopic	All women	96/434 (22.1%)	0.16 (0.13–0.19)	0.24 (0.20–0.29)	0.27 (0.23–0.34)
	Trying to conceive or intending to conceive within the next 2 years <sup>b</sup>	88/241 (36.5%)	0.28 (0.21–0.35)	0.40 (0.32–0.49)	0.50 (0.38–0.62)
	Trying to conceive <sup>b</sup>	60/125 (48.0%)	0.40 (0.30–0.52)	0.54 (0.42–0.67)	0.67 (0.50–0.83)
Myomectomy Route	Subgroup	Live births/Total women (%)	Probability of live birth (95% CI) by follow-up time		
			12 months	24 months	36 months
Abdominal	All women	28/388 (7.2%)	0.01 (0.00–0.05)	0.10 (0.06–0.17)	0.10 (0.06–0.17)
	Trying to conceive or intending to conceive within the next 2 years <sup>b</sup>	26/238 (10.9%)	0.01 (0.00–0.07)	0.20 (0.12–0.32)	0.20 (0.12–0.32)
	Trying to conceive <sup>b</sup>	16/113 (14.2%)	0.02 (0.00–0.12)	0.25 (0.14–0.43)	0.25 (0.14–0.43)
Hysteroscopic	All women	19/273 (7.0%)	0.04 (0.02–0.08)	0.13 (0.08–0.21)	0.19 (0.12–0.30)
	Trying to conceive or intending to conceive within the next 2 years <sup>b</sup>	17/94 (18.1%)	0.05 (0.02–0.14)	0.21 (0.12–0.35)	0.31 (0.18–0.50)
	Trying to conceive <sup>b</sup>	10/53 (18.9%)	0.06 (0.02–0.19)	0.27 (0.14–0.48)	0.30 (0.16–0.52)
Laparoscopic	All women	44/434 (10.1%)	0.02 (0.01–0.05)	0.12 (0.08–0.17)	0.14 (0.10–0.21)
	Trying to conceive or intending to conceive within the next 2 years <sup>b</sup>	39/241 (16.2%)	0.03 (0.01–0.08)	0.20 (0.13–0.29)	0.25 (0.16–0.37)
	Trying to conceive <sup>b</sup>	22/125 (17.6%)	0.03 (0.01–0.10)	0.26 (0.15–0.41)	N/A (no data)

Abbreviations: CI=confidence interval.

<sup>a</sup>Cumulative probability accounts for censoring using life-table methods and adjusts for confounding using propensity score weights.

<sup>b</sup>Based on self-report at baseline only.

**Table 3:**

Myomectomy route in relation to pregnancy and live birth, interval-based analysis

Myomectomy Route	All women Number of events/ Total (%)	Unadjusted		Adjusted <sup>a</sup>		Adjusted and restricted to women with pregnancy intent <sup>a,b</sup>	
		OR	95% CI	OR	95% CI	OR	95% CI
<i>Pregnancy</i>							
Abdominal	69/388 (17.8%)	1.00	reference	1.00	reference	1.00	reference
Hysteroscopic	37/273 (13.6%)	0.81	0.52, 1.24	1.28	0.76, 2.14	1.27	0.72, 2.23
Laparoscopic	96/434 (22.1%)	1.38	0.98, 1.94	1.19	0.76, 1.85	1.26	0.77, 2.04
<i>Live birth</i>							
Abdominal	28/388 (7.2%)	1.00	reference	1.00	reference	1.00	reference
Hysteroscopic	19/273 (7.0%)	1.03	0.57, 1.86	1.66	0.80, 3.43	1.71	0.81, 3.65
Laparoscopic	44/434 (10.1%)	1.47	0.90, 2.39	1.38	0.72, 2.62	1.49	0.77, 2.89

Abbreviations: OR=odds ratio, CI=confidence interval.

<sup>a</sup>Applies propensity score weights to account for differences in demographics, symptoms, and fibroid characteristics.<sup>b</sup>Restricted to women with fertility intent (actively trying to conceive or intending to conceive within the next two years) at the beginning of the 12-month interval (time-varying covariate).

**Table 4:**

Myomectomy route (laparoscopic vs. abdominal) in relation to pregnancy and live birth during follow-up, interval-based analysis

Myomectomy Route	Unadjusted		Adjusted <sup>a</sup>				Adjusted and restricted to women with pregnancy intent <sup>a,b</sup>			
	OR	95% CI	OR	95% CI	OR <sup>c</sup>	95% CI <sup>c</sup>	OR	95% CI	OR <sup>c</sup>	95% CI <sup>c</sup>
<i>Pregnancy</i>										
Abdominal	1.00	Reference	1.00	Reference	1.00	Reference	1.00	Reference	1.00	Reference
Laparoscopic	1.38	0.98, 1.94	1.15	0.81, 1.66	1.14	0.79, 1.64	1.25	0.85, 1.84	1.25	0.84, 1.84
<i>Live birth</i>										
Abdominal	1.00	Reference	1.00	Reference	1.00	Reference	1.00	Reference	1.00	Reference
Laparoscopic	1.47	0.90, 2.39	1.32	0.79, 2.19	1.24	0.75, 2.06	1.42	0.84, 2.39	1.35	0.80, 2.25

Abbreviations: OR=odds ratio, CI=confidence interval.

<sup>a</sup>Applies propensity score weights to account for differences in demographics, symptoms, and fibroid characteristics.

<sup>b</sup>Restricted to women with fertility intent (actively trying to conceive or intending to conceive within the next two years) at the beginning of the 12-month interval (time-varying covariate).

<sup>c</sup>Propensity score weights augmented by maximum fibroid volume and the presence of at least one submucous fibroid.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript