

# Utilization of cervical cancer screening service and associated factors among women of reproductive age group in Ilu Abba Bor zone, southwestern Ethiopia, 2019: Cross-sectional study

SAGE Open Medicine  
Volume 10: 1–8  
© The Author(s) 2022  
Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/20503121221094904  
journals.sagepub.com/home/smo



Dejene Edosa Dirirsa<sup>1</sup> , Eyosiyas Yeshialem Asefa<sup>2</sup>  
and Mukemil Awol Salo<sup>1</sup> 

## Abstract

**Introduction:** Cervical cancer is one of the severest risks to women's life. It is the main reason for more than half million of morbidity and 266,000 deaths worldwide annually.

**Objective:** The aim of this study was to assess the level of cervical cancer screening service utilization and associated factors among women of the reproductive age group coming for health care service to public health facilities of Ilu Abba Bor zone, southwestern Ethiopia, 2019.

**Methods:** An institutional-based cross-sectional study design was employed from 1 May to 30 May 2019. Participants were selected by systematic random sampling technique. Data were collected using a pretested questionnaire and entered into Epi data 3.1 and exported to SPSS version 23 for analysis. Multivariable logistic regression was used to observe the factors associated with cervical cancer screening utilization service. An adjusted odds ratio with a 95% confidence interval was used. Variables with a p-value of <0.05 were considered to be statistically significant.

**Results:** From a total of 259 women involved in the study, about 19 (7.3%) of the study participants had been screened for cervical cancer at least once in the past 3 years during the study period. Educational status: primary (grade 1–8) (adjusted odds ratio = 3; 95% confidence interval = (1.15–7.91)), secondary (grade 9–12) (adjusted odds ratio = 4.21; 95% confidence interval = (1.04–7.46)), and tertiary (grade 12+) (adjusted odds ratio = 6.01; 95% confidence interval = (3.01–11.35)), nulliparous (adjusted odds ratio = 0.256; 95% confidence interval = (0.105–0.642)), and looking for information about cervical cancer (adjusted odds ratio = 3.4; 95% confidence interval = (1.8–6.2)) were significant factors associated with utilization of cervical cancer screening service.

**Conclusion:** The study revealed that there was low cervical cancer screening service utilization in the study area. Educational status, parity, and looking for cervical cancer information were independently associated with cervical cancer screening utilization service. There is need to improve women's education, dissemination of information about importance of cervical cancer screening, and focus on nulliparous women to have good practice of cervical cancer screening.

## Keywords

Screening utilization, cervical cancer, Ethiopia

Date received: 5 August 2021; accepted: 29 March 2022

## Introduction

Cancer is a disease which is characterized by uncontrolled cell proliferation in the body. Cervical cancer refers to cancer that begins in the cervix. Seventy percent of cervical cancer cases worldwide are caused by two types of human papillomaviruses (HPV): HPV-16 and HPV-18.<sup>1</sup> Cervical cancer is

<sup>1</sup>Department of Midwifery, College of Health Sciences, Salale University, Fiche, Ethiopia

<sup>2</sup>Department of Midwifery, College of Health Science and Medicine, Debre Berhan University, Debre Berhan, Ethiopia

### Corresponding author:

Dejene Edosa Dirirsa, Department of Midwifery and Medicine, College of Health Science, Salale University, P.O.Box: 245, Fiche, Ethiopia.  
Email: dejenedosa43@gmail.com



one of the most serious threats to women's health, accounting for more than half a million cases of morbidity and 266,000 deaths globally each year. Most of the problems (85%) occur in low-income countries around the world.<sup>2</sup>

Cervical cancer screening has been recognized to be a very effective prevention strategy for cervical cancer. The recommendations of the United States Preventive Services Task Force (USPSTF), American Cancer Society (ACS), and World Health Organization (WHO) point out that cervical cancer screening once in a lifetime can significantly reduce the risk of death from the disease and the incidence of advanced cervical cancer.<sup>3</sup>

Despite government support for health policy, women in underdeveloped nations have a lower rate of cervical cancer screening than those in developed countries: for example, Kampala, Uganda 7%,<sup>4</sup> Bhutan 6%,<sup>5</sup> and Nigeria 8.3%,<sup>6</sup> respectively. Surprisingly, only 1% of Ethiopian women receive cervical cancer screening, with 90% of women receiving no screening at all.<sup>7</sup>

More than 80% of cervical cancer identified with advanced stage in low-income countries, resulting in low survival rates after surgery or radiotherapy. WHO investigated and reported a cervical cancer screening technique called visual inspection with acetic acid (VIA) in order to help Ethiopia and other resource-poor countries.<sup>8</sup> In low-income countries, using VIA for simultaneous diagnosis and treatment in a single visit is an effective prevention technique that can reduce complications and mortality.<sup>9</sup>

The WHO indicated that the number of health institutions in Ethiopia that provide cervical cancer screening are very low, and it is expected to be used by women greater than 30 years old and women at risk to acquire cervical cancer, such as women with multiple sexual partners, smoking, begin sexual intercourse earlier, have HIV, and other sexually transmitted infections (STIs).<sup>10</sup>

According to the Black Lion Specialized Hospital's Oncology Center, Ethiopia, the majority of cancer cases are diagnosed at a late stage. Treatment options for this condition are limited at this time. This is mainly due to a lack of screening service, early detection and treatment services, as well as inadequate diagnostic facilities and poorly structured referral system.<sup>11</sup>

Study conducted in Dire Dawa Ethiopia revealed that age > 40 years, attending private health institutions, occupation, visiting gynecology ward, having good knowledge about cervical cancer screening, counseling from health professionals, and using family planning methods were determinant factors of CC screening uptake.<sup>12</sup> Another study conducted in Arba Minch Ethiopia showed that only 9.6% women have had cervical cancer screening within the past 3 years. There was association between CC screening awareness and utilization of CC screening service.<sup>13</sup>

According to studies conducted in Bahir Dar, Saint Paul's Teaching Hospital, and Bishoftu, the rural residence, low

monthly income, history of STIs, frequent health facility visits, providers' recommendation, and poor knowledge about CC screening were determinants of cervical cancer screening service utilization.<sup>14-17</sup> The study conducted in Northern and southwestern Ethiopia also identified that level of education, lack of awareness about CC screening, occupation, having children, and visiting health facilities were the determinant factors associated with the uptaking of cervical cancer screening.<sup>18,19</sup>

Regular and early screening and treatment as part of targeted interventions is mandatory to reduce the incidence and mortality associated with the cervical cancer. But there are very limited studies conducted in the country, specifically in the study area to assess the utilization of cervical cancer screening service. The aim of this study was to assess the level of cervical cancer screening service utilization and associated factors among women of the reproductive age group coming for health care service to public health facilities of Ilu Abba Bor zone, southwestern Ethiopia, 2019.

## Methods

### Study setting and period

The study was implemented in Ilu Abba Bor zone, Oromia regional state from 1 May to 30 May 2019. There are 12 districts in the zone and based on the 2007 census conducted by the Central Statistical Agency of Ethiopia, the zone has a total population of 1,271,609 from which 44,124 (3.47%) were females of the reproductive age group. The capital town of the zone is Mettu, located in the southwestern of Ethiopia at 600 km from Addis Ababa. In the zone, there were 20 health centers and 2 public hospitals giving the health services. The study was done at 12 health centers and 1 hospital (Mettu Karl referral hospital). Mettu Karl referral hospital was the only health facility providing cervical cancer screening service using VIA for women of the reproductive age group who were referred from other health institutions and at high risk to develop cervical cancer.

### Study design

An institutional-based cross-sectional study design was implemented.

### Source Population

All women of the reproductive age group who visited public health facilities of Ilu Abba Bor zone for health care services were the source population.

### Study population

All sampled women of the reproductive age group who visited the selected public health facilities of Ilu Abba Bor zone

for health care services during the data collection period were the study population.

### **Inclusion criteria**

The women of reproductive age who visited selected public health facilities for health care services such as antenatal care (ANC), family planning, gynecological care, and adult outpatient departments for medical care during the data collection period were included.

### **Exclusion criteria**

Those women who were critically ill and unable to respond for the required information were excluded from the study.

### **Sample size determination**

Single population proportion formula was used to estimate the sample size required for this study with the following assumptions:  $P=19.8\%$ , taken from the study conducted in Mekelle<sup>8</sup> (which revealed the prevalence of cervical cancer screening utilization service was  $19.8\%$ ), at the marginal error of  $5\%$  with a  $95\%$  confidence level. Accordingly, the sample size was 244 and by considering  $10\%$  of calculated sample size for non-respondents, the final sample size was 268.

### **Sampling technique and procedures**

Twelve health centers and one public Hospital, namely Hurrumu, Yayo, Gore, Mettu, Nopa, Becho, Darimu, Uka, Bure, Sibbo, Burusa and Alge health centers, and Metu Karl Referral Hospital, were selected randomly by using the lottery method. The final sample size was proportionally allocated for each facility based on 3-month average client flow. Finally, we selected the participants using a systematic random sampling technique after calculating K-interval for each facility until we got the proportionally allocated sample size.

### **Study variables**

*Outcome variable.* Utilization of cervical cancer screening service measured by asking the respondent's action toward screening for a premalignant cervical lesion in the past 3 years. Those who ever screened at least once within the past 3 years were regarded as utilized cervical cancer screening service and those who had never screened in the past 3 years were regarded as having not utilized cervical cancer screening service.<sup>2</sup>

*Independent variables.* Age, marital status, religion, ethnicity, participant's educational status, husband's educational status, monthly family income, parity, participant's occupation, source of information, and knowledge about cervical cancer.

### **Data collection tools and procedures**

The data were collected by four female Midwives under the Supervision of investigators using a pretested structured questionnaire. The questionnaire was written in English first, translated into Afaan Oromoo and then back to English to ensure consistency. The Afaan Oromoo version of the questionnaire was pretested on  $5\%$  of women seeking health care at unselected facilities. The data collectors were given 2 days of training on the data collection technique, as well as the study's aim and objectives. The exit interview took place in a separate area set aside for this purpose near each health facility's waiting room.

### **Operational definitions**

*Cervical cancer screening service utilization.* Women who are eligible for cervical cancer screening and have ever been screened for cervical cancer during the last 3 years.

*Cervical cancer screening.* Application of acetic acid to detect cervical tissue abnormalities in an asymptomatic population.

*Good knowledge about cervical cancer screening.* The women who scored greater than the mean of knowledge measuring questions (9 questions).

*Poor knowledge about cervical cancer screening.* The women who scored less than the mean of knowledge measuring questions (9 questions).

### **Statistical analysis**

The Epi data 3.1 and SPSS version 23 were used for data analysis. Multivariable logistic regression was used to assess the relative effect of independent variables on the dependent variable to control the possible confounders and to select significant predictors of utilization of cervical cancer screening included in the final model. Finally, the adjusted odds ratio (AOR) with its  $95\%$  confidence interval (CI) and a  $p$ -value of  $0.05$  was used to identify significant associated factors of cervical cancer screening service utilization. The findings were presented using text and tables.

### **Ethical approval**

Ethical approval was obtained from Mettu University Research and Ethical Committee with the approval number of MU/67/566/19. The permission letter was provided from Ilu Abba Bor zonal Health Office, and other concerned bodies in the study area. During data collection, each respondent was informed about the purpose, scope, and expected outcome of the research. Informed written consent was obtained from each subject and legally authorized representatives of the subjects who were minors before proceeding data collection.

**Table 1.** Socio-demographic characteristics of the study participants in public health facilities of Ilu Abba Bor zone, Southwest Ethiopia, 2019.

Variables		Count	N %	Variables		Count	N %
Age in years	15–24	99	38.2	Husband's occupation	Employee (Governmental Organization/Non-governmental Organization)	18	6.9
	25–34	125	48.3		Merchant	84	32.4
	35–44	35	13.5		Student	66	25.5
Marital status	Married	157	60.6	Farmer	28	10.8	
	Single	54	20.8	Daily worker	47	18.1	
	Divorced	28	10.8	Others <sup>b</sup>	16	6.2	
	Widowed	16	6.2	Husband's educational status	No education	34	13.1
	Separated	4	1.5		Informal education	66	25.5
Religion	Orthodox	58	22.4		Primary (grade 1–8)	77	29.7
	Protestant	82	31.7	Secondary (grade 9–12)	54	20.8	
	Muslim	88	34.0	Tertiary (grade 12+)	28	10.8	
	Catholic	26	10.0	Monthly income	<\$72	93	39.2
Others <sup>a</sup>	5	1.9	\$72–\$143		90	38.0	
Ethnicity	Oromo	124	47.9		\$144–\$215	31	13.1
	Amhara	49	18.9		>\$215	23	9.7
	Guragie	40	15.4	Parity	Nulliparous	54	20.8
	Tigrie	28	10.8		1–2	108	41.7
	Others <sup>a</sup>	18	6.9	>2	97	37.5	
Participant's Educational status	No education	49	18.9	Participant's occupation	Employee (GO/NGO)	42	16.2
	Informal education	51	19.7		Merchant	86	33.2
	Primary (grade 1–8)	105	40.5		Student	57	22.0
	Secondary (grade 9–12)	33	12.7		Farmer	38	14.7
	Tertiary (12+)	21	8.1		Daily worker	24	9.3
				Others <sup>b</sup>	12	4.6	

<sup>a</sup>Wolayita, Sidama, and Afar.

<sup>b</sup>House wife, private, shopkeeper.

## Results

### *Socio-demographic characteristics of the study participants in public health facilities of Ilu Abba Bor zone, southwest Ethiopia, 2019*

Out of the total 268 participants included in this study, 259 completed the interview making a response rate of 96.6%. The mean age of the study subjects was found to be 27.34 years with a standard deviation,  $SD \pm 6.17$  and ranged from 17 to 40 years. Most of the respondents (60.6%,  $n = 157$ ) were currently married. The highest grade completed by the respondents was primary education (1–8) (40.5%,  $n = 105$ ) and informal education (19.7%,  $n = 51$ ) (see Table 1).

### *Health service-related characteristics among women of reproductive age group in public health facilities of Ilu Abba Bor zone, 2019*

The result of this study indicated that more than half (54.1%,  $n = 140$ ) of the respondents never received information from any health care providers about cervical cancer screening. The private health facility takes the major (65.6%,  $n = 78$ ) part to be the source of information. Almost all (92.7%,  $n = 240$ ) of the study participants have been looking for the

information about cervical cancer and nearly half (49.6%,  $n = 119$ ) of them searched for the information from health care providers (see Table 2).

### *Level of knowledge about cervical cancer screening services among women of reproductive age group in public health facilities of Ilu Abba Bor zone, 2019*

About 124 (47.9%) of the respondents (95% CI, 4.21–5.37) had poor knowledge, meaning they scored less than the mean of knowledge measuring scores (4.32;  $SD = 1.75$ ) and the minimum and maximum knowledge scores were 0 and 9, respectively. Nearly three-quarters of respondents (74.5%,  $n = 193$ ) were unaware of the causes of cervical cancer. The HPV was the most (41.3%,  $n = 107$ ) often stated cause of cervical cancer. One hundred nine of study participants (42.1%,  $n = 109$ ) said that early cervical cancer screening is useful in preventing the disease.

The majority of the study participants (53.7%,  $n = 139$ ) stated that beginning of sexual relations before age of 17 increased the risk of having cervical cancer for women. Only 13.5% of study participants mentioned that cervical cancer screening is important for women of childbearing age (see Table 3).

**Table 2.** Health service–related characteristics among women of reproductive age group in public health facilities of Ilu Abba Bor zone, 2019.

S. no	Variables		Count	N %
1	Have you ever received information from any health care providers about cervical cancer screening?	Yes	119	45.9
		No	140	54.1
2	If yes for question 1, from where did you receive?	Government health facility	38	31.9
		Private health facility	78	65.6
		Drug vendors	3	2.5
3	Have you been looking for the information about cervical cancer?	Yes	240	92.7
		No	19	7.3
4	If yes for question 3, from where have you searched for the information?	1. TV	53	22.1
		2. Radio	41	17.1
		3. Health care providers	119	49.6
		4. Internet	27	11.2

**Table 3.** Level of knowledge about cervical cancer screening among women of reproductive age group in public health facilities of Ilu Abba Bor zone, 2019.

Knowledge measurement tools		Frequency	N %
Do you know the causes of cervical cancer?	Yes	66	25.5
	No	193	74.5
What are the causes of cervical cancer if you answered yes to the above question?	HIV	72	27.8
	HPV	107	41.3
	Tuberculosis	58	22.4
	Do not know	22	8.5
What are the benefits of cervical cancer screening in its early stages?	To prevent cancer	109	42.1
	To control cancer	138	53.3
	Other <sup>a</sup>	12	4.6
Is it true that starting sexual relations before the age of 17 increases the risk of cervical cancer?	Yes	139	53.7
	No	120	46.3
Which category of women should be checked for cervical cancer, in your opinion?	Women with gynecological problems	100	38.6
	Pregnant women only	114	44.1
	All women of childbearing age	35	13.5
	Sexually active women	4	1.5
	Do not know	6	2.3
Is unprotected sexual activity related with an increased risk of cervical cancer?	Yes	91	35.1
	No	168	64.9
Do you believe that women who acquire cervical cancer symptoms should be treated?	Yes	125	48.3
	No	134	51.7
Do you believe cervical cancer can be prevented?	Yes	152	58.7
	No	107	41.3
	Do not know	69	45.4
Do you believe cervical cancer can be cured if it is diagnosed earlier?	Yes	150	57.9
	No	109	42.1

HPV: human papilloma virus.

<sup>a</sup>To identify cervical cancer.

### *Utilization of cervical cancer screening among reproductive age women coming to public health facilities of Ilu Abba Bor zone, 2019*

According to this study, only 7.3% (n=19, 95% CI (4.2–10.4)) of cervical cancer screening services were used in the previous 3 years. The main reasons for not being screened for cervical cancer were a lack of availability of the service (54.1%; n=129) and cost issues (24.3%; n=58). Other

reasons given were not knowing where to acquire the service (14.4%; n=34), fear of discrimination (3.6%; n=9), and others (see Table 4).

### *Factors associated with utilization of cervical cancer screening service*

Women's educational status, primary (grade 1–8), secondary (grade 9–12), and tertiary (grade 12+) level of education

**Table 4.** Utilization of cervical cancer screening among reproductive age women coming to public health facilities of Ilu Abba Bor zone, 2019.

S. no	Variables		Frequency	Percentage
1	Have you ever been screened for the cervical cancer in the last 3 years	Yes	19	7.3
		No	240	92.7
2	If not screened for the cervical cancer in the last 3 years, what was the reasons for not being screened?	Service is not available nearby	130	54.2
		Financial problem	58	24.2
		Unaware of where to get the service	34	14.4
		Fear of discrimination	9	3.6
		Other <sup>a</sup>	9	3.6

<sup>a</sup>Fear of test result.

**Table 5.** Factors associated with utilization of cervical cancer screening among the respondents in public health facilities, Ilu Abba Bor zone, southwest Ethiopia, 2019.

Variables	Categories	Count	%	COR (95% CI)	AOR (95% CI)	p value
Education	Cannot read and write	49	18.9	–	–	–
	Can read and write	51	19.7	0.9 (0.4–2)	0.51 (0.21–1.24)	0.136
	Primary (Grade 1–8)	105	40.5	7.2 (3–11.57)	3 (1.15–7.91)*	0.030*
	Secondary (9–12)	33	12.8	2.8 (1.11–2.73)	4.2 (1.04–7.46)*	0.001*
	Tertiary (12+)	21	8.1	1.76 (1.05–3.6)	6 (3.01–11.35)*	0.002*
Occupation of husband	Student	66	25.5	–	–	–
	Merchant	84	32.4	0.61 (0.32–1.17)	0.45 (0.19–1.1)	0.074
	Employed	18	6.9	2.45 (0.78–7.64)	1.8 (0.48–6.62)	0.394
	Farmer	28	10.8	0.52 (0.21–1.3)	0.63 (0.19–2.12)	0.451
	Daily worker	47	18.1	0.98 (0.47–2.1)	0.7 (0.26–1.98)	0.515
	Others	16	6.2	1.57 (0.51–4.81)	0.56 (0.13–2.4)	0.433
Education of husband	Cannot read and write	34	13.1	–	–	–
	Can read and write	66	25.5	1.72 (0.75–3.96)	2.25 (0.86–5.92)	0.099
	Primary (Grade 1–8)	77	29.7	1.69 (0.68–4.2)	2.28 (0.75–6.98)	0.148
	Secondary (Grade 9–12)	54	20.8	0.87 (0.37–2.07)	2.19 (0.72–6.66)	0.167
	Tertiary (Grade 12+)	28	10.8	0.82 (0.3–2.27)	1.76 (0.48–6.48)	0.395
Monthly income of family	<\$72	93	39.2	–	–	–
	\$72–\$143	90	38.0	0.58 (0.32–1.04)	0.6 (0.3–1.2)	0.147
	\$144–\$215	31	13.1	1.25 (0.54–2.87)	1.4 (0.54–3.4)	0.522
	>\$215	23	9.7	0.51 (0.2–1.29)	0.5 (0.18–1.57)	0.251
Parity of the respondent	>2	54	20.8	–	–	–
	1–2	108	41.7	1.51 (0.78–2.91)	0.8 (0.34–1.91)	0.615
	Nulliparous	97	37.5	0.61 (0.31–1.2)	0.256 (0.11–0.64)*	0.004*
Intention to be screened	No	139	53.7	–	–	–
	Yes	120	46.3	0.7 (0.44–1.16)	1.2 (0.62–2.32)	0.582
Looking for information	No	119	45.9	–	–	–
	Yes	140	54.1	2.6 (1.57–4.3)	3.4 (1.83–6.2)*	<0.001*

COR: crude odds ratio; AOR: adjusted odds ratio; CI: confidence interval; –: Reference.

\*Significant variables.

were 3 times (AOR=3; 95% CI (1.15–7.91)), 4.2 times (AOR=4.21; 95% CI (1.04–7.46)), and 6 times (AOR=6.01; 95% CI (3.01–11.35)) more likely to utilize cervical cancer screening service than women who cannot read and write, respectively. In addition, respondents who were nulliparous were 74.4% (AOR=0.26; 95% CI (0.105–0.642)) less likely to utilize cervical cancer screening service than women of parity more than two. The odds of utilization of cervical cancer screening was about 3.4 times more among women who

were looking for cervical cancer information than their counterparts (AOR=3.4; 95% CI (1.8–6.2)) (see Table 5).

## Discussion

The aim of this study was to assess the level of cervical cancer screening service utilization and associated factors among women of the reproductive age group coming for health care service to public health facilities of Ilu Abba Bor

zone, southwestern Ethiopia, 2019. Therefore, the study found that only 7.3% of women had received cervical cancer screening service at least once in the past 3 years. Factors such as education level, parity, and finding information about cervical cancer screening are significantly related to the use of cervical cancer screening.

The finding of cervical cancer screening service use (7.3%) in the current study was in line with the studies carried out in Debre Merkos (5.4%),<sup>3</sup> Arba Minchi (5.9%),<sup>10</sup> Bishoftu (5.8%),<sup>16</sup> and Ghana (7.8%)<sup>20</sup> of participants who had been screened for cervical cancer, respectively. Correspondingly, studies conducted in Kampala, Uganda (7%)<sup>4</sup> and Nigeria (8.3%)<sup>6</sup> support the result of this study.

This finding was found to be lower than results from studies carried out in Hawassa (40.1%),<sup>1</sup> Wolayita (18.6%),<sup>9</sup> Mekelle (19.8%),<sup>8</sup> Kenya (17.5%),<sup>21</sup> and Tanzania, Dare Salam (21%)<sup>22</sup> of participants being screened, respectively.

Compared with participants who cannot read or write, primary education (grades 1–8), secondary education (grades 9–12), and higher education (grades 12+) are 3 times, 4.2 times, and 6 times more likely to use cervical cancer screening services, respectively. Similar findings were observed in studies conducted by Hawassa,<sup>1</sup> Gondar,<sup>18</sup> Wolaita,<sup>9</sup> and Mekelle<sup>8</sup> where the level of education can predict the use of cervical cancer screening services.

Compared with respondents who had given birth twice, those who had never given birth were 74.4% less likely to be screened for cervical cancer. This may be because women who have not given birth before have less access to information, counseling, and cervical cancer screening services. This is consistent with a study conducted at Arba Minchi, which showed that multi-parities are closely related to the use of cervical cancer screening services.<sup>10</sup> Comparable studies conducted in Tanzania indicate that women who have given birth are more likely to use cervical cancer screening services than women who have never given birth.<sup>22</sup>

According to the result of this study, those who were looking for information about cervical cancer were about 3.4 times more likely to use cervical screening than those who were looking for information about cervical cancer. This finding was consistent with the result of a study done in Dare Salam, which indicated that the national health system factors influencing the early uptake of cervical cancer screening services include poor flow of information from national to lower level.<sup>23</sup>

### *Limitation of the study*

Since the study is facility-based study, it may not represent for all women not visited health institution during the study period. Therefore, we recommend future studies at the community level in the study area. Further research is needed to conduct qualitative research to find unknown reason why not being screened for cervical cancer.

## **Conclusion**

This study has shown that there was low cervical cancer screening service utilization in the study area. In addition, educational status, parity, and looking for cervical cancer information were independently associated with cervical cancer screening utilization. Therefore, stakeholders should improve women's education, dissemination of information about importance of cervical cancer screening, and focus on nulliparous women to have good practice of cervical cancer screening.

## **Acknowledgements**

We would like to thank Mettu University and colleagues for their valuable support throughout the research process. Our acknowledgment also extends to zonal health department and respective health facility officials for permitting us to do so as well as involved data collectors and supervisors for their commitment during the data collection process.

## **Author contributions**

D.E.D. and E.Y.A. originated the idea and participated in the proposal development, questionnaire development, data collection, and analysis. D.E.D. and M.A.S. participated in manuscript writing. All authors read the final version of the manuscript and approved this version of the manuscript to be considered for publication.

## **Availability of data and materials**

Data used to support the findings of this study are available from the corresponding author upon request.

## **Declaration of conflicting interests**

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## **Ethical approval**

Prior to data collection, ethical clearance was obtained from Research and Ethical committee of College of Health and Medical Sciences of Mettu University with the approval number MU/67/566/19. Written permission letter was produced from Ilu Abba Bor zonal Health Office and other concerned bodies in the study area. During data collection, each respondent was informed about the purpose, scope, and expected outcome of the research.

## **Funding**

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: The whole required (material and humanitarian) cost for this research work was covered by Mettu University.

## **Informed consent**

Informed written consent was obtained from each subject and legally authorized representatives of the subjects who were minors before proceeding data collection.

## ORCID iDs

Dejene Edosa Dirirsa  <https://orcid.org/0000-0003-2266-5006>

Mukemil Awol Salo  <https://orcid.org/0000-0001-6923-6650>

## Supplemental material

Supplemental material for this article is available online.

## References

- Assefa AA, Astawesegn FH and Eshetu B. Cervical cancer screening service utilization and associated factors among HIV positive women attending adult ART clinic in public health facilities, Hawassa town, Ethiopia: a cross-sectional study. *BMC Health Serv Res* 2019; 19: 847.
- Kasim J, Kalu A, Kamara B, et al. Cervical cancer screening service utilization and associated factors among women in the Shabadino District, Southern Ethiopia. *J Cancer Epidemiol* 2020; 2020: 6398394.
- Aynalem BY, Anteneh KT and Enyew MM. Utilization of cervical cancer screening and associated factors among women in Debremarkos town, Amhara region, Northwest Ethiopia: community based cross-sectional study. *PLoS ONE* 2020; 15(4): e0231307.
- Twinomujuni C, Nuwaha F and Babirye JN. Understanding the low level of cervical cancer screening in Masaka Uganda using the ASE model: a community-based survey. *PLoS ONE* 2015; 10(6): e0128498.
- Dhendup T and Tshering P. Cervical cancer knowledge and screening behaviors among female university graduates of year 2012 attending national graduate orientation program, Bhutan. *BMC Womens Health* 2014; 14: 44.
- Abiodun OA, Olu-Abiodun OO, Sotunsa JO, et al. Impact of health education intervention on knowledge and perception of cervical cancer and cervical screening uptake among adult women in rural communities in Nigeria. *BMC Public Health* 2014; 14: 814.
- Gakidou E, Nordhagen S and Obermeyer Z. Coverage of cervical cancer screening in 57 countries: Low average levels and large inequalities. *PLoS Med* 2008; 5: e132. doi: 10.1371/journal.pmed.0050132.
- Bayu H, Berhe Y, Mulat A, et al. Cervical cancer screening service uptake and associated factors among age eligible women in Mekelle Zone, Northern Ethiopia, 2015: a community based study using health belief model. *PLoS ONE* 2016; 11(3): e0149908.
- Bekele T, Arba A, Getahun D, et al. Cervical cancer screening service utilization and associated factors among women in reproductive age group attending public health facilities of Sodo Town, Wolaita Sodo. *EC Gynaecol* 2021; 10(2): 32–42.
- Gebru Z, Gerbaba M and Dirar A. Utilization of cervical carcinoma screening service and associated factors among currently married women in Arba Minch Town, Southern Ethiopia. *J Womens Health Care* 2016; 5: 297.
- Ayenew A, Zewdu B and Nigusie AA. Uptake of cervical cancer screening service and associated factors among age-eligible women in Ethiopia: systematic review and meta-analysis. *Infect Agents Cancer* 2020; 15: 67.
- Belay Y, Dheresa M, Sema A, et al. Cervical cancer screening utilization and associated factors among women aged 30 to 49 years in Dire Dawa, Eastern Ethiopia. *Cancer Control* 2020; 27(1): 1073274820958701.
- Seyoum T, Yesuf A, Kejela G, et al. Utilization of cervical cancer screening and associated factors among female health workers in governmental health institution of Arba Minch Town and Zuria District, Gamo Gofa Zone, Arba Minch, Ethiopia, 2016. *Arch Cancer Res* 2017; 5(4): 165.
- Muluneh BA, Atnafu DD and Wassie B. Predictors of cervical cancer screening service utilization among commercial sex workers in Northwest Ethiopia: a case-control study. *BMC Womens Health* 2019; 19: 162.
- Woldetsadik AB, Amhare AF, Bitew ST, et al. Socio-demographic characteristics and associated factors influencing cervical cancer screening among women attending in St. Paul's Teaching and Referral Hospital, Ethiopia. *BMC Womens Health* 2020; 20: 70.
- Ebo GG, Tilahun T and Heyi WD. Cervical cancer screening practice and associated factors in Bishoftu Town, Eastern Ethiopia. *Gynecol Obstet* 2020; 10: 538.
- Abate SM. Trends of cervical cancer in Ethiopia. *Cervical Cancer* 2015; 1: 103.
- Nega AD, Woldetsadik MA and Gelagay AA. Low uptake of cervical cancer screening among HIV positive women in Gondar University Referral Hospital, Northwest Ethiopia: cross-sectional study design. *BMC Womens Health* 2018; 18: 87.
- Chali K, Oljira D, Sileshi T, et al. Knowledge on cervical cancer, attitude toward its screening, and associated factors among reproductive age women in Metu Town, Ilu Aba Bor, South West Ethiopia, 2018: community-based cross-sectional study. *Cancer Rep* 2021; 4(5): e1382.
- Prempheh EAK. Women's awareness and factors affecting utilization of cervical cancer screening services in the Ejisu-Juaben municipality of Ghana. *J Cancer Tumor Int* 2018; 8(1): 1–11.
- Morema EN, Atieli HE, Onyango RO, et al. Determinants of cervical screening services uptake among 18–49 year old women seeking services at the Jaramogi Oginga Odinga Teaching and Referral Hospital, Kisumu, Kenya. *BMC Health Serv Res* 2014; 14: 335.
- Kileo NM, Michael D, Neke N, et al. Utilization of cervical cancer screening services and its associated factors among primary school teachers in Ilala Municipality, Dar es Salaam, Tanzania. *BMC Health Serv Res* 2015; 15: 552.
- Mugassa AM and Frumence G. Factors influencing the uptake of cervical cancer screening services in Tanzania: a health system perspective from national and district levels. *Nurs Open* 2020; 7(1): 345–354.