



# HHS Public Access

Author manuscript

*J Am Chem Soc.* Author manuscript; available in PMC 2022 September 01.

Published in final edited form as:

*J Am Chem Soc.* 2021 September 01; 143(34): 13759–13768. doi:10.1021/jacs.1c05769.

## Generation of Axially Chiral Fluoroallenes through a Copper-Catalyzed Enantioselective $\beta$ -Fluoride Elimination

**Thomas J. O'Connor,**

Department of Chemistry, University of California, Berkeley, California 94720, United States

**Binh Khanh Mai,**

Department of Chemistry, University of Pittsburgh, Pittsburgh, Pennsylvania 15260, United States

**Jordan Nafie,**

BioTools, Inc., Jupiter, Florida 33458, United States

**Peng Liu,**

Department of Chemistry, University of Pittsburgh, Pittsburgh, Pennsylvania 15260, United States

**F. Dean Toste**

Department of Chemistry, University of California, Berkeley, California 94720, United States

### Abstract

Herein we report the copper-catalyzed silylation of propargylic difluorides to generate axially chiral, tetrasubstituted monofluoroallenes in both good yields (27 examples >80%) and enantioselectivities (82–98% ee). Compared to previously reported synthetic routes to axially chiral allenes (ACAs) from prochiral substrates, a mechanistically distinct reaction has been developed: the enantiodiscrimination between enantiotopic fluorides to set an axial stereocenter. DFT calculations and vibrational circular dichroism (VCD) suggest that  $\beta$ -fluoride elimination from an alkenyl copper intermediate likely proceeds through a *syn*- $\beta$ -fluoride elimination pathway rather than an *anti*-elimination pathway. The effects of the C1-symmetric Josiphos-derived ligand on reactivity and enantioselectivity were investigated. Not only does this report showcase that alkenyl copper species (like their alkyl counterparts) can undergo  $\beta$ -fluoride elimination, but this elimination can be achieved in an enantioselective fashion.

### Graphical Abstract

---


**Corresponding Author Peng Liu** – Department of Chemistry, University of Pittsburgh, Pittsburgh, Pennsylvania 15260, United States; pengliu@pitt.edu; **F. Dean Toste** – Department of Chemistry, University of California, Berkeley, California 94720, United States; fdtoste@berkeley.edu.

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacs.1c05769>.

Synthesis procedures, characterization data for all new compounds, additional optimization data, computational details, and Cartesian coordinates of all computed structures (PDF)

The authors declare no competing financial interest.



## INTRODUCTION

The pursuit of synthetic methods to access structurally diverse allenes stems from their applications in both medicinal and materials chemistry as well as their ability to serve as a reactive functional group for further synthetic manipulations.<sup>1,2</sup> These cumulenes can exhibit axial chirality, and most of the naturally occurring allenic compounds that have been isolated are nonracemic.<sup>1a-c</sup> Although enantiopure bromoallenes have been discovered in nature (Scheme 1a) and axially chiral bromo-, chloro-, and iodoallenes have been synthesized, the corresponding axially chiral fluoroallenes are largely unknown.<sup>3,4</sup> Specifically, to the best of our knowledge, there is a single report where an enantioenriched tetrasubstituted, axially chiral monofluoroallene has been prepared, albeit in modest enantioselectivity.<sup>4c</sup> Due to a lack of general synthetic routes toward this chiral, fluorinated motif, their potential applications have remained unexplored.

An emerging route toward the catalytic synthesis of ACAs has been from prochiral substrates.<sup>5</sup> This strategy overcomes the requirement for a stoichiometric amount of a chiral auxiliary and/or enantioenriched substrates. Prochiral substrates that have been transformed into ACAs include propargylic electrophiles, 1,3-enynes, terminal alkynes, 1,3-dienes, racemic allenes, and vinyl triflates.<sup>6</sup> In the case of vinyl triflates, it was demonstrated that  $\beta$ -hydride elimination occurred in an enantioselective fashion, revealing a new mechanistic route for the synthesis of ACAs.<sup>6f</sup> Of the classes of ACAs (1,3-di-, tri-, and tetrasubstituted), tetrasubstituted ACAs remain difficult to synthesize in high enantiopurity.<sup>7</sup>

Moreover, the incorporation of a functional group directly attached to ACAs that permits further transformations has gained popularity as evidenced by recent reports of boryl and silyl substituted ACAs.<sup>8,9</sup> Access to tetrasubstituted, boryl, or silyl monofluoro ACAs would permit an array of further transformations that could generate quaternary, fluorine-containing stereocenters by way of axial-to-point chirality transfer.<sup>3a,10</sup>


The development of defluorination methods of (poly)-fluorinated compounds has emerged as a complementary strategy to access complex, fluorine-functionalized motifs that have typically been accessed from nonfluorinated substrates.<sup>11</sup> Of the metals that catalyze such

defluorination reactions, copper has been shown to be exceptionally competent, as evidenced by the numerous reports of catalytic hydrodefluorinations, defluoroborylations, and defluorosilylations of fluoroarenes and fluoroolefins.<sup>12</sup> An emerging trend in this field is the enantioselective defluorination of allylic CF<sub>3</sub> or CF<sub>2</sub>R groups, forging stereocenters adjacent to mono- and difluoroolefins (Scheme 1b).<sup>13</sup> Recently, a rare example of enantioselective defluorination via oxidative addition to a *gem*-difluoride has been reported as a method to generate products with a fluorine-containing stereogenic unit.<sup>14</sup> We envisioned that a mechanistically different approach to the desymmetrization of difluoromethylene groups, proceeding through an enantioselective  $\beta$ -fluoride elimination reaction, might be employed in the enantioselective synthesis of monofluoro ACAs (Scheme 1c).

To this end, we hypothesized that reaction of propargylic difluorides with a suitable chiral copper nucleophile would form an alkenyl copper species that could undergo an enantioselective  $\beta$ -fluoride elimination to generate tetrasubstituted monofluoro ACAs (Scheme 1c). Potential obstacles toward achieving this transformation included controlling both the regioselectivity<sup>15</sup> and enantioselectivity of the process, avoiding undesired reactivity of the alkenyl copper intermediate,<sup>16</sup> and preventing further silylation of the product.<sup>17</sup> Herein we demonstrate that  $\beta$ -fluoride elimination from an alkenyl copper species is possible and the discrimination of enantiotopic fluorides is a viable elementary process to achieve asymmetric synthesis of ACAs. DFT studies predict that this elimination proceeds through a *syn*-elimination pathway, which is in contrast to some studies of alkyl copper species that undergo  $\beta$ -fluoride elimination.<sup>12b,g-i,13b,c,f,g</sup>

## RESULTS AND DISCUSSION

We began our investigation by determining if  $\beta$ -fluoride elimination was feasible from a vinyl copper intermediate. In the presence of a suitable base, the borylation of **1a** with B<sub>2</sub>pin<sub>2</sub> generated the desired boryl monofluoroallene in 46% NMR yield (eq 1); however, attempts at isolating this product were



(1)

unsuccessful.<sup>8b</sup> We hypothesized that the corresponding silyl monofluoroallene of **1a** would be isolable and gratifyingly discovered that the silylation of **1a** with PhMe<sub>2</sub>SiBpin (**2**) led to **1b**. After a brief optimization of conditions, over 40 chiral ligands were examined for this transformation. Of the chiral ligands employed, only four gave the desired allene in greater than 20% ee.

Fortunately (*R,S*)-Josiphos afforded **1b** in a modest yield (71%) and promising enantioselectivity (20% ee). A series of Josiphos ligands were generated to evaluate their steric and electronic effects (see Supporting Information Tables S1 and S2) on the

reaction. Josiphos ligands containing aryl groups on both phosphorus atoms demonstrated good reactivity and enantioselectivity, specifically when the alkyldiarylphosphorus moiety possessed bulky 3,5-substituted arenes (Table 1, compare entries 2 and 3; Supporting Information Tables S2 and S3). However, once the 3,5-substituents became sterically too demanding, for example, with the TTB derivative, the reactivity dropped significantly (entry 5). Moreover, it appeared that steric rather than electronic factors played a decisive role in determining the enantioselectivity of the transformation (compare entries 3 and 8).

Further optimization led to simplified reaction conditions (Table 2). By lowering the temperature and changing the solvent, the enantiomeric excess of **1b** improved to 90%, but the overall conversion of **1a** dropped (entry 1). Increasing the amount of phenoxide base resulted in decoordination of the ligand, determined by <sup>31</sup>P NMR spectroscopy, and erosion of the ee of **1b**.<sup>18</sup> By use of an insoluble source of fluoride (CsF), an increase in the chemical yield of **1b** was achieved (83%, entry 2). Possible roles of CsF could be either trapping FBpin<sup>19</sup> and/or releasing CsOAr from ArOBpin.<sup>20</sup> Fortunately, the transformation proceeded with CsF as the sole base, removing the chance of phosphine decoordination due to excess phenoxide.<sup>21</sup> The reaction proceeded well in nonpolar, low coordinating solvents (entries 4–7), and the removal of MeCN led to an increase in catalytic activity (entries 8, 9). Switching to CuOTf·<sup>1</sup>/<sub>2</sub>C<sub>6</sub>H<sub>6</sub> as the copper source afforded the desired allene in almost quantitative yield (entry 10).

With the optimal conditions established, an alternative synthetic route employing readily available material to prepare propargylic difluorides was developed (Scheme 2). The decarboxylative bromination of difluorocarboxylic acids and the copper-catalyzed Sonogashira cross-coupling of terminal alkynes with difluorobenzyl bromides afforded difluoroalkynes, which were subjected to defluorosilylation under the optimized conditions. A range of functional groups were tolerated in the copper catalyzed transformation, affording the desired allenes **1–24b** in high yields (83–98%) and in good enantioselectivities (82–98%) after isolation (Table 3). Notably, alkynes (**19b**), alkenes (**20b**), enynes (**24b**), aldehydes (**8b**), ketones (**9b**), propargylic acetates (**23b**) as well as alkyl and aryl halides (**3b** and **6b**) were tolerated. Coordinating heterocycles (**14–16b**, **18b**, and **25b**), amides (**7b**, **11–13b**), and nitriles (**5b** and **10b**) also did not hamper catalysis. Although changing the electronics of the aryl ring slightly decreased the enantiomeric excess of the reaction (**1–6b**), increasing the steric bulk of the aryl group was well tolerated (**25b**). Although many functional groups were tolerated, the reaction proved more sensitive to alterations of the substituents directly attached to the allene, which could affect the barrier of silylation of the alkyne (**26–27b**) or impact the C—F bond strength (**28–29b**).

To explore the scalability of this method, **1b** was synthesized on a 6 mmol scale without a significant loss in enantioselectivity or chemical yield. It was discovered that other silylboranes could be utilized for this transformation when (*R,S*)-3,5-Trip-Josiphos was employed as the ligand (**1b-BnMe<sub>2</sub>Si**, **1b-CyMe<sub>2</sub>Si**, and **1b-Et<sub>3</sub>Si**). These allenyl silanes were synthesized on gram scales with comparable yields and enantioselectivities to **1b-PhMe<sub>2</sub>Si**.

Determination of the absolute configuration of allene **10b** was achieved using vibrational circular dichroism (VCD).<sup>22</sup> After a careful conformational search (see Supporting Information for details) at four levels of theory, B3LYP/6-31G(d), B3PW91/6-31G(d), B3LYP/cc-pVTZ, and B3PW91/cc-pVTZ,<sup>23</sup> the resulting conformers were Boltzmann averaged and plotted with a line width of 5 cm<sup>-1</sup> to produce the final theoretical spectra. The IR and VCD spectra were then frequency scaled<sup>23</sup> for comparison to the experimental data. Calculations at all four levels of theory matched well, proving the absolute configuration of **10b** to be *S*. Of the four methods employed, the best agreement with experimental data was from the B3PW91/cc-pVTZ level. The comparison of experimental and theoretical spectra was quantified<sup>24</sup> using BioTools (Jupiter, FL) CompareVOA software, with high neighborhood similarity for IR (90.4) and VCD (69.6), ESI (enantiomeric similarity index) for VCD (62.9), and a confidence level of 99%. Of particular note was the asymmetric allene C—C—C stretch observed at 1933 cm<sup>-1</sup>, which was one of many closely correlated bands between experiment and theory. In addition to allene **10b**, the absolute configuration of **1b-Et<sub>3</sub>Si**, **12b**, **17b**, and **25b** were also determined to be *S* by VCD analysis (see Supporting Information).

On the basis of previous reports regarding CuF<sup>12c,e,16a,21,25</sup> and copper silyl species,<sup>15b,f,17d,26</sup> we propose the following mechanism for the copper-catalyzed reaction (Figure 1). First, a complex between Josiphos and [CuOTf] undergoes salt metathesis with CsF to generate JosiphosCuF (**Cu1**).<sup>16a,21,25d</sup>  $\sigma$ -Bond metathesis with PhMe<sub>2</sub>SiBpin (**2**) generates JosiphosCuSiMe<sub>2</sub>Ph (**Cu2**) and releases FBpin.<sup>12c-e</sup> Subsequent coordination and silylation of the triple bond generates an alkenyl Cu species (**Cu3**).<sup>15a,e,27</sup> A  $\beta$ -fluoride elimination<sup>12g</sup> regenerates **Cu1**, which is trapped by FBpin,<sup>19,28</sup> **2**<sup>15c</sup> or decomposes the formed allenylsilane (**c**). As FBpin is more Lewis acidic than B<sub>2</sub>pin<sub>2</sub>,<sup>28</sup> the same is likely true with **2**. By using a judicious amount of CsF and a nonpolar solvent, we propose that the precipitation of Cs[F<sub>2</sub>Bpin]<sup>19,20</sup> drives this reaction forward. <sup>1</sup>H, <sup>19</sup>F, and <sup>11</sup>B NMR studies have identified **Cu1**, **Cu2**, LCuF<sub>2</sub>Bpin, as well as LCuOH and confirmed the generation of **Cu2** from both **Cu1** and LCuF<sub>2</sub>Bpin (see Supporting Information section 7). **Cu2** was also shown to react with alkyne **1a**, generating both allene **1b** and FBpin. It appears that LCuF<sub>2</sub>Bpin acts as a reservoir of Cu<sup>I</sup>F, and under catalytic conditions, a monomeric or dimeric CuF was not observed. Over the course of the reaction only **Cu2**, LCuF<sub>2</sub>Bpin, and LCuOH were observed, which converged to **Cu2** after the alkyne has been consumed (see Supporting Information section 7). Although the exact structure of **Cu1** is unknown, the speciation of **Cu1** appears to be both solvent and temperature dependent, in which the former has been observed for other Josiphos copper halide complexes (see Supporting Information section 7).<sup>29</sup> On the basis of our experiments, we propose that the silylation of the alkyne and the  $\beta$ -fluoride elimination reactions are the rate- and selectivity-determining steps, respectively. Using (*R,S*)-3,5-TES-JosiphosCuF<sub>2</sub>Bpin as a catalyst, the desired allene **1b** was obtained in a similar yield and enantiomeric excess, demonstrating its catalytic competence (see Supporting Information section 7).

Density functional theory (DFT) calculations were performed to investigate the reaction mechanism and origin of enantioselectivity of this Cu-catalyzed asymmetric silylation of propargylic difluorides. The DFT calculations were performed at the M06/

SDD(Cu,Fe,Cs)-6-311+G(d,p)/SMD-(toluene)//B3LYP-D3(zero)/SDD(Cu,Fe,Cs)-6-31G(d) level of theory using difluoroalkyne **27a** and PhMe<sub>2</sub>SiBpin (**2**) as model substrates. The (*R,S*)-3,5-TMS-Josiphos ligand was used in the DFT calculations for simplicity because the use of this ligand in the ligand screening provided only slightly lower ee than using (*R,S*)-3,5-*TES*-Josiphos (Table 1, entries 6 and 7). On the basis of the proposed catalytic cycle, the computed reaction energy profile is shown in Figure 2. The association of FBpin to monomeric LCuF (**26**) to form a heterodimer (**29**) is exergonic by 4.0 kcal/mol, suggesting that the more stable complex **29** can be an off-cycle reservoir of Cu<sup>I</sup>F. Although the dimerization of LCuF is exergonic by 7.3 kcal/mol, its formation is expected to be less favorable than forming **29** due to the low concentrations of LCuF under catalytic conditions (see Figure S1 for detailed discussions about the equilibrium of **26**, **29**, and the dimer of LCuF).

The  $\sigma$ -bond metathesis between monomeric LCuF (**26**) and PhMe<sub>2</sub>SiBpin<sup>30</sup> (**2**) takes place via a four-membered cyclic transition state (**TS-1**) to form silyl copper intermediate **30** and FBpin. This step requires a low activation barrier of 6.7 kcal/mol with respect to **26** and is exergonic by 24.7 kcal/mol. Migratory insertion of alkyne **27a** into the silyl copper (**TS-2**) gives alkenyl copper species **31**. This migratory insertion is highly regioselective for the formation of Cu—C bond at the alkyne terminus adjacent to the difluoromethylene. The transition state leading to the other regioisomer, *TS-2'*, is 10.9 kcal/mol higher in energy than **TS-2**. The high level of regioselectivity is due to steric repulsions between the silyl and the more hindered alkyne terminus (C1) in **TS-2'** as well as inductive effects of the difluoro substituents that stabilize the building of negative charge at C1 in **TS-2**. From **31**, both *syn*<sup>12g-i</sup> and *anti*<sup>13b</sup>  $\beta$ -fluoride elimination pathways were calculated. The *syn*-elimination of either of the two diastereotopic  $\beta$ -F in **31** (via **TS-3** and **TS-4**) involves a four-membered cyclic transition state, while the *anti*- $\beta$ -fluoride elimination is facilitated by CsF as a Lewis acid (via **TS-5** and **TS-6**). The FBPin-facilitated *anti*-elimination was also computed and is also less favorable than the *syn*-elimination (see Figure S3). The *syn*-elimination pathways require much lower barriers than the *anti*-elimination, which is in contrast to a computational study by Hoveyda and Torker that suggested the  $\beta$ -fluoride elimination from alkyl copper species favors the *anti*-pathway due to Lewis acid (i.e., Na<sup>+</sup>) coordination to the F<sup>-</sup> leaving group and the Bpin group on the substrate.<sup>13b</sup> In the present study, the lack of such chelating Lewis-acid coordination in the *anti*-pathway, the weaker Lewis acidity of CsF, and the strain release effect that alleviates steric repulsions between the SiMe<sub>2</sub>Ph group and the Cu in the *syn*-elimination transition state changed the reaction mechanism to favor the *syn*-elimination.<sup>31</sup>

Among the three key elementary steps in the catalytic cycle, the alkyne migratory insertion (**TS-2**) has the highest activation free energy ( $G^\ddagger = 19.9$  kcal/mol with respect to **30**). This finding is consistent with our experimental results that suggest this step being the rate-determining step (*vide supra*). The enantioselectivity-determining step is the *syn*- $\beta$ -fluoride elimination. **TS-3**, which leads to the (*S*)-enantiomer of the monofluoroallene product, is 1.9 kcal/mol more stable than **TS-4** that leads to the (*R*)-enantiomer. The predicted enantioselectivity is consistent with the absolute configuration of the product identified by the VCD analysis.

Next, we performed a detailed analysis to investigate the effects of the Josiphos ligand on the reactivity and enantioselectivity of the  $\beta$ -fluoride elimination. Because of the conformational flexibility of the (*R,S*)-3,5-TMS-Josiphos ligand,<sup>32</sup> a careful conformational search was performed for all intermediates and transition states in the catalytic cycle. These calculations revealed at least four different conformers of the 3,5-TMS-Josiphos-supported copper complexes. The two most stable and catalytically active ligand conformations **A** and **B** are shown in Table 4 (see Figures S4-S6 for all possible ligand conformations). Ligand conformation **A** involves a twist-boat-type six-membered ring and is more favorable in the copper fluoride (**26**), the  $\sigma$ -bond metathesis transition state (**TS-1**), and the silyl copper intermediate (**30**). In more sterically encumbered structures, including the migratory insertion transition state (**TS-2**), alkenyl copper (**31**), and the  $\beta$ -fluoride elimination transition states (**TS-3** and **TS-4**), ligand conformation **B** becomes more favorable. This ligand conformation involves a half-chair type six-membered ring, which points the Ar and Ph groups in quadrants I and II away from the Cu center. As such, the bulky SiMe<sub>2</sub>Ph group is placed between these unoccupied quadrants to minimize steric repulsions between the ligand and the SiMe<sub>2</sub>Ph group on the substrate.

Ligand conformation **B** not only stabilizes the silylalkenyl copper species but also plays a significant role in controlling the enantioselectivity of the  $\beta$ -fluoride elimination. The *P*-phenyl group in quadrant III and the *P*-3,5-TMS-phenyl group in quadrant IV point toward the Cu center and thus occupy these quadrants. The larger size of the 3,5-TMS-phenyl compared to phenyl indicates that the ligand–substrate repulsions in quadrant IV would be more pronounced than those in quadrant III. Indeed, quadrant diagrams of the  $\beta$ -fluoride elimination transition states (Figure 3) support this hypothesis. In the less favorable transition state **TS-4**, the phenyl group on the substrate is located in the more occupied quadrant IV, leading to steric repulsion with a TMS group on the ligand. By contrast, in the more favorable  $\beta$ -fluoride elimination transition state **TS-3**, the much smaller fluoro group is located in quadrant IV, and thus the ligand–substrate steric repulsions are diminished. Next, we performed energy decomposition analysis (EDA)<sup>33</sup> calculations to quantitatively analyze the ligand–substrate noncovalent interactions in **TS-3** and **TS-4** (see Supporting Information for computational details). The EDA calculations revealed that the dominant factor controlling the enantioselectivity is the Pauli repulsion (*i.e.*, steric repulsion) between the (*R,S*)-3,5-TMS-Josiphos ligand and the substrate. The Pauli repulsion energy ( $E_{\text{Pauli}}$ ) in **TS-4** is 1.7 kcal/mol higher than that in **TS-3** and thus destabilizes the former transition state.

Finally, we calculated the enantioselectivity-determining *syn*- $\beta$ -F elimination transition states of the reaction of the same substrate (**27a**) catalyzed by a SegPhos-supported Cu complex. The computed enantioselectivity is diminished ( $\Delta G^\ddagger = 0.2$  kcal/mol, Figure S7), indicating the C<sub>2</sub>-symmetric SegPhos ligand is not effective for asymmetric induction. This prediction is consistent with the low ee of 12% obtained experimentally at 65°C with 5 mol % CuCl, 40 mol % sodium phenoxide, and 6 mol % SegPhos ligand. Taken together, these ligand effect analyses revealed the unique roles of the conformationally flexible C<sub>1</sub>-symmetric Josiphos ligand, where it lowers the activation barrier for the rate-determining alkyne migratory insertion step and improves the enantioselectivity of the  $\beta$ -F elimination.

## CONCLUSION

The first copper-catalyzed, enantioselective  $\beta$ -fluoride elimination has been achieved. The resulting monofluoro ACAs represent the first examples of fluorine-containing, chiral tetrasubstituted allenes.<sup>34</sup> It is expected that such a motif will find value in pharmaceutical and agrochemical chemistry in addition to being a valuable building block for the generation of more elaborate fluorine-containing stereocenters. DFT calculations of the reaction mechanisms predicted that this elimination occurs in a *syn*-fashion, which is promoted by strain release of the *Z*- $\beta$ -silylalkenyl copper intermediate. The unique roles of the C1-symmetric Josiphos-derived ligand in promoting the reactivity and enantioselectivity were investigated. It is believed that lessons learned from this desymmetrization could be leveraged for the creation of other fluorine-containing stereocenters via defluorination pathways.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## ACKNOWLEDGMENTS

We gratefully acknowledge the National Institutes of Health (Grants R35 GM118190 and R35 GM128779) for financial support. T.J.O. thanks the NSF (Grant DGE 1752814) for a predoctoral fellowship. The College of Chemistry CheXray (NIH Shared Instrumentation Grant S10-RR027172) is acknowledged for X-ray crystallographic data, and we thank Dr. Hasan Celik and UC Berkeley's NMR facility in the College of Chemistry (CoC-NMR) for spectroscopic assistance. Instruments in the CoC-NMR are supported in part by Grant NIH S10OD024998. DFT calculations were carried out at the Center for Research Computing at the University of Pittsburgh, the Extreme Science and Engineering Discovery Environment (XSEDE), and the TACC Frontera Supercomputer.

## REFERENCES

- (1). (a)Modern Allene Chemistry; Krause N, Hashmi ASK, Eds.; Wiley-VCH: Weinheim, Germany, 2004.(b)Rivera-Fuentes P; Diederich F Allenes in molecular materials. *Angew. Chem., Int. Ed* 2012, 51, 2818–28.(c)Hoffmann-Roder A; Krause N Synthesis and properties of allenic natural products and pharmaceuticals. *Angew. Chem., Int. Ed* 2004, 43, 1196–216.(d)Cai F; Pu X; Qi X; Lynch V; Radha A; Ready JM Chiral allene-containing phosphines in asymmetric catalysis. *J. Am. Chem. Soc* 2011, 133, 18066–9. [PubMed: 21972824]
- (2). (a)Allen AD; Tidwell TT Ketenes and other cumulenes as reactive intermediates. *Chem. Rev* 2013, 113, 7287–342. [PubMed: 23767795] (b)Sutherland DR; Kinsman L; Angiolini SM; Rosair GM; Lee AL Gold(I)-Catalysed Hydroarylation of 1,3-Disubstituted Allenes with Efficient Axial-to-Point Chirality Transfer. *Chem. - Eur. J* 2018, 24, 7002–7009. [PubMed: 29542832] (c)Ma S Electrophilic addition and cyclization reactions of allenes. *Acc. Chem. Res* 2009, 42, 1679–88. [PubMed: 19603781] (d)Liu L; Ward RM; Schomaker JM Mechanistic Aspects and Synthetic Applications of Radical Additions to Allenes. *Chem. Rev* 2019, 119, 12422–12490. [PubMed: 31833759] (e)Krause N; Winter C Gold-catalyzed nucleophilic cyclization of functionalized allenes: a powerful access to carbo- and heterocycles. *Chem. Rev* 2011, 111, 1994–2009. [PubMed: 21314182] (f)Adams CS; Weatherly CD; Burke EG; Schomaker JM The conversion of allenes to strained three-membered heterocycles. *Chem. Soc. Rev* 2014, 43, 3136–63. [PubMed: 24647416] (g)Yu S; Ma S Allenes in catalytic asymmetric synthesis and natural product syntheses. *Angew. Chem., Int. Ed* 2012, 51, 3074–112.
- (3). (a)Yokobori U; Ohmiya H; Sawamura M Synthesis of Allenylsilanes through Copper-Catalyzed  $\gamma$ -Selective Coupling between  $\gamma$ -Silylated Propargylic Phosphates and Alkylboranes. *Organometallics* 2012, 31, 7909–7913.(b)Whitehead DC; Yousefi R; Jaganathan A; Borhan B An organocatalytic asymmetric chlorolactonization. *J. Am. Chem. Soc* 2010, 132, 3298–300.



[PubMed: 20170118] (c)Vaz B; Pereira R; Perez M; Alvarez R; de Lera AR Stereoselective Stille coupling of enantiopure haloallenes and alkenylstannanes for the synthesis of allenyl carotenoids. Experimental and computational studies. *J. Org. Chem* 2008, 73, 6534–41. [PubMed: 18681399] (d)Schade MA; Yamada S; Knochel P Synthesis of polyfunctional allenes by successive copper-mediated substitutions. *Chem. - Eur. J* 2011, 17, 4232–7. [PubMed: 21384450] (e)Ohno H; Ando K; Hamaguchi H; Takeoka Y; Tanaka T A highly cis-selective synthesis of 2-ethynylaziridines by intramolecular amination of chiral bromoallenes: improvement of stereoselectivity based on the computational investigation. *J. Am. Chem. Soc* 2002, 124, 15255–66. [PubMed: 12487601] (f)Marshall JA; Grant CM Formation of Transient Chiral Allenylindium Reagents from Enantioenriched Propargylic Mesylates through Oxidative Transmetalation. Applications to the Synthesis of Enantioenriched Homopropargylic Alcohols. *J. Org. Chem* 1999, 64, 696–697. [PubMed: 11674133] (g)Li H; Grassi D; Guenee L; Burgi T; Alexakis A Copper-catalyzed propargylic substitution of dichloro substrates: enantioselective synthesis of trisubstituted allenes and formation of propargylic quaternary stereogenic centers. *Chem. - Eur. J* 2014, 20, 16694–706. [PubMed: 25327947] (h)Kim MJ; Sohn TI; Kim D; Paton RS Concise substrate-controlled asymmetric total syntheses of dioxabicyclic marine natural products with 2,10-dioxabicyclo-[7.3.0]dodecene and 2,9-dioxabicyclo[6.3.0]undecene skeletons. *J. Am. Chem. Soc* 2012, 134, 20178–88. [PubMed: 23194584] (i)Evans PA; Murthy VS; Roseman JD; Rheingold AL Enantioselective Total Synthesis of the Nonisoprenoid Sesquiterpene (–)-Kumausallene. *Angew. Chem., Int. Ed* 1999, 38, 3175–3177. (j)Caporusso AM; Zampieri A; Aronica LA; Banti D Stereoselective synthesis of chiral 3-aryl-1-alkynes from bromoallenes and heterocuprates. *J. Org. Chem* 2006, 71, 1902–10. [PubMed: 16496974] (k)Caporusso AM; Filippi S; Barontini F; Salvadori P Coupling of chiral 1-bromo-1,2-dienes with zinc-based cuprates: a new procedure for the regio and stereoselective synthesis of functionalized acetylenic compounds. *Tetrahedron Lett.* 2000, 41, 1227–1230.

- (4). (a)Wang X; Wu Z; Wang J alpha-Fluoroallenoate Synthesis via N-Heterocyclic Carbene-Catalyzed Fluorination Reaction of Alkynals. *Org. Lett* 2016, 18, 576–9. [PubMed: 26771540] (b)Mae M; Hong JA; Xu B; Hammond GB Highly regioselective synthesis of gem-difluoroallenes through magnesium organocuprate SN2' substitution. *Org. Lett* 2006, 8, 479–82. [PubMed: 16435864] (c)Li T; Zhou C; Yan X; Wang J Solvent-Dependent Asymmetric Synthesis of Alkynyl and Monofluoroalkenyl Isoindolinones by CpRh(III) -Catalyzed C-H Activation. *Angew. Chem., Int. Ed* 2018, 57, 4048–4052. (d)Hammond GB Functionalized Fluorinated Allenes. In *Fluorine-Containing Synthons*; American Chemical Society, 2005; Vol. 911, pp 204–215. (e)Castelhana AL; Krantz A A novel route to allenyl fluorides. Synthesis of 4-amino-7-fluorohepta-5,6-dienoic acid, the first fluoroallenyl amino acid. *J. Am. Chem. Soc* 1987, 109, 3491–3493. (f)Lan Y; Hammond GB Functionalization of monofluoroallene and the synthesis of aryl-substituted conjugated fluorodienes. *Org. Lett* 2002, 4, 2437–9. [PubMed: 12098266] (g)Pacheco MC; Purser S; Gouverneur V The chemistry of propargylic and allylic fluorides. *Chem. Rev* 2008, 108, 1943–81. [PubMed: 18543877] (h)Zapata AJ; Gu Y; Hammond GB The first alpha-fluoroallenylphosphonate, the synthesis of conjugated fluoroenynes, and the stereoselective synthesis of vinylfluorophosphonates using a new multifunctional fluorine-containing building block. *J. Org. Chem* 2000, 65, 227–34. [PubMed: 10813920]
- (5). (a)Neff RK; Frantz DE Recent Advances in the Catalytic Syntheses of Allenes: A Critical Assessment. *ACS Catal.* 2014, 4, 519–528. (b)Chu W-D; Zhang Y; Wang J Recent advances in catalytic asymmetric synthesis of allenes. *Catal. Sci. Technol* 2017, 7, 4570–4579. (c)Brummond K; DeForrest J Synthesizing Allenes Today (1982–2006). *Synthesis* 2007, 2007, 795–818. (d)Ogasawara M Catalytic enantioselective synthesis of axially chiral allenes. *Tetrahedron: Asymmetry* 2009, 20, 259–271.
- (6). (a)Wang Y; Zhang W; Ma S A room-temperature catalytic asymmetric synthesis of allenes with ECNU-Phos. *J. Am. Chem. Soc* 2013, 135, 11517–20. [PubMed: 23869750] (b)Trost BM; Zell D; Hohn C; Mata G; Maruniak A Enantio- and Diastereoselective Synthesis of Chiral Allenes by Palladium-Catalyzed Asymmetric [3+2] Cycloaddition Reactions. *Angew. Chem., Int. Ed* 2018, 57, 12916–12920. (c)Trost BM; Schultz JE; Chang T; Maduabum MR Chemo-, Regio-, Diastereo-, and Enantioselective Palladium Allylic Alkylation of 1,3-Dioxaboroles as Synthetic Equivalents of alpha-Hydroxyketones. *J. Am. Chem. Soc* 2019, 141, 9521–9526. [PubMed: 31180647] (d)Liu H; Leow D; Huang KW; Tan CH Enantioselective synthesis of chiral allenoates by guanidine-catalyzed isomerization of 3-alkynoates. *J. Am. Chem. Soc*

2009, 131, 7212–3. [PubMed: 19422238] (e)Jiang Y; Diagne AB; Thomson RJ; Schaus SE Enantioselective Synthesis of Allenes by Catalytic Traceless Petasis Reactions. *J. Am. Chem. Soc* 2017, 139, 1998–2005. [PubMed: 28121128] (f)Crouch IT; Neff RK; Frantz DE Pd-catalyzed asymmetric beta-hydride elimination en route to chiral allenenes. *J. Am. Chem. Soc* 2013, 135, 4970–3. [PubMed: 23488914] (g)Chu WD; Zhang L; Zhang Z; Zhou Q; Mo F; Zhang Y; Wang J Enantioselective Synthesis of Trisubstituted Allenes via Cu(I)-Catalyzed Coupling of Diazoalkanes with Terminal Alkynes. *J. Am. Chem. Soc* 2016, 138, 14558–14561. [PubMed: 27788320] (h)Bayeh-Romero L; Buchwald SL Copper Hydride Catalyzed Enantioselective Synthesis of Axially Chiral 1,3-Disubstituted Allenes. *J. Am. Chem. Soc* 2019, 141, 13788–13794. [PubMed: 31423768] (i)Adamson NJ; Jeddi H; Malcolmson SJ Preparation of Chiral Allenes through Pd-Catalyzed Intermolecular Hydroamination of Conjugated Enynes: Enantioselective Synthesis Enabled by Catalyst Design. *J. Am. Chem. Soc* 2019, 141, 8574–8583. [PubMed: 31070902]

- (7). (a)Wu S; Huang X; Wu W; Li P; Fu C; Ma S A C-H bond activation-based catalytic approach to tetrasubstituted chiral allenenes. *Nat. Commun* 2015, 6, 7946. [PubMed: 26246391] (b)Scheipers I; Muck-Lichtenfeld C; Studer A Palladium-Catalyzed Decarboxylative gamma-Arylation for the Synthesis of Tetrasubstituted Chiral Allenes. *Angew. Chem., Int. Ed* 2019, 58, 6545–6548. (c)Qian D; Wu L; Lin Z; Sun J Organocatalytic synthesis of chiral tetrasubstituted allenenes from racemic propargylic alcohols. *Nat. Commun* 2017, 8, 567. [PubMed: 28924216] (d)Hayashi T; Tokunaga N; Inoue K Rhodium-catalyzed asymmetric 1,6-addition of aryltitanates to enynones giving axially chiral allenenes. *Org. Lett* 2004, 6, 305–7. [PubMed: 14723554] (e)Hashimoto T; Sakata K; Tamakuni F; Dutton MJ; Maruoka K Phase-transfer-catalysed asymmetric synthesis of tetrasubstituted allenenes. *Nat. Chem* 2013, 5, 240–4. [PubMed: 23422567] (f)Liao Y; Yin X; Wang X; Yu W; Fang D; Hu L; Wang M; Liao J Enantioselective Synthesis of Multisubstituted Allenes by Cooperative Cu/Pd-Catalyzed 1,4-Arylboration of 1,3-Enynes. *Angew. Chem., Int. Ed* 2020, 59, 1176–1180. (g)Armstrong RJ; Nandakumar M; Dias RMP; Noble A; Myers EL; Aggarwal VK Enantiodivergent Synthesis of Allenes by Point-to-Axial Chirality Transfer. *Angew. Chem., Int. Ed* 2018, 57, 8203–8208.
- (8). (a)Sang HL; Yu S; Ge S Copper-catalyzed asymmetric hydroboration of 1,3-enynes with pinacolborane to access chiral allenylboronates. *Org. Chem. Front* 2018, 5, 1284–1287. (b)Huang Y; Del Pozo J; Torker S; Hoveyda AH Enantioselective Synthesis of Trisubstituted Allenyl-B(pin) Compounds by Phosphine-Cu-Catalyzed 1,3-Enyne Hydroboration. Insights Regarding Stereochemical Integrity of Cu-Allenyl Intermediates. *J. Am. Chem. Soc* 2018, 140, 2643–2655. [PubMed: 29417810] (c)Gao DW; Xiao Y; Liu M; Liu Z; Karunananda MK; Chen JS; Engle KM Catalytic, Enantioselective Synthesis of Allenyl Boronates. *ACS Catal.* 2018, 8, 3650–3654. [PubMed: 30740261]
- (9). (a)Wang M; Liu ZL; Zhang X; Tian PP; Xu YH; Loh TP Synthesis of Highly Substituted Racemic and Enantioenriched Allenylsilanes via Copper-Catalyzed Hydrosilylation of (Z)-2-Alken-4-ynoates with Silylboronate. *J. Am. Chem. Soc* 2015, 137, 14830–3. [PubMed: 26560851] (b)Nishimura T; Makino H; Nagaosa M; Hayashi T Rhodium-catalyzed enantioselective 1,6-addition of arylboronic acids to enynamides: asymmetric synthesis of axially chiral allenylsilanes. *J. Am. Chem. Soc* 2010, 132, 12865–7. [PubMed: 20806919] (c)Liu ZL; Yang C; Xue QY; Zhao M; Shan CC; Xu YH; Loh TP Copper-Catalyzed Asymmetric Silylation of Propargyl Dichlorides: Access to Enantioenriched Functionalized Allenylsilanes. *Angew. Chem., Int. Ed* 2019, 58, 16538–16542.
- (10). (a)Danheiser RL; Carini DJ; Fink DM; Basak A Scope and stereochemical course of the (trimethylsilyl)cyclopentene annulation. *Tetrahedron* 1983, 39, 935–947. (b)Brawn RA; Panek JS Preparation and use of enantioenriched allenylsilanes for the stereoselective synthesis of homopropargylic ethers. *Org. Lett* 2007, 9, 2689–92. [PubMed: 17559219] (c)Marshall JA; Maxson K Stereoselective synthesis of stereotriad subunits of polyketides through additions of nonracemic allenylsilanes to (R)- and (S)-2-methyl-3-oxygenated propanals. *J. Org. Chem* 2000, 65, 630–3. [PubMed: 10813988] (d)Brawn RA; Panek JS Synthesis of enantioenriched homopropargylic sulfonamides by a three component reaction of aldehydes, sulfonamides, and chiral allenylsilanes. *Org. Lett* 2009, 11, 4362–5. [PubMed: 19775184] (e)Danheiser RL; Kwasigroch CA; Tsai YM Application of allenylsilanes in [3+2] annulation approaches to oxygen and nitrogen heterocycles. *J. Am. Chem. Soc* 1985, 107, 7233–7235. (f)Masse CE; Panek JS Diastereoselective Reactions of Chiral Allyl and Allenyl Silanes with Activated


- C:X .pi.-Bonds. Chem. Rev 1995, 95, 1293–1316.(g)Fleming I; Terrett NK Stereospecific syntheses and reactions of allyl- and allenyl-silanes. J. Organomet. Chem 1984, 264, 99–118. (h)Danheiser RL; Carini DJ (Trimethylsilyl)allenes as propargylic anion equivalents: synthesis of homopropargylic alcohols and ethers. J. Org. Chem 1980, 45, 3925–3927.(i)Brawn RA; Panek JS Stereoselective C-glycosidations with achiral and enantioenriched allenylsilanes. Org. Lett 2010, 12, 4624–7. [PubMed: 20839812] (j)Danheiser RL; Carini DJ; Basak A (Trimethylsilyl)-cyclopentene annulation: a regiocontrolled approach to the synthesis of five-membered rings. J. Am. Chem. Soc 1981, 103, 1604–1606.(k)Spiteri C; Moses JE Copper-catalyzed azide-alkyne cyclo-addition: regioselective synthesis of 1,4,5-trisubstituted 1,2,3-triazoles. Angew. Chem., Int. Ed 2010, 49, 31–3.(l)Cai B; Evans RW; Wu J; Panek JS Total Synthesis of Nuclear Factor of Activated T-Cells-68 (NFAT-68): Sequential Use of Chiral Allenylsilane and Titanium Alkoxide-Mediated Reductive Coupling Bond Construction. Org. Lett 2016, 18, 4304–7. [PubMed: 27513364] (m)Danheiser RL; Carini DJ; Kwasigroch CA Scope and stereochemical course of the addition of (trimethylsilyl)-allenes to ketones and aldehydes. A regiocontrolled synthesis of homopropargylic alcohols. J. Org. Chem 1986, 51, 3870–3878.(n)Weinreb SM Thermal and Lewis Acid Catalyzed Intramolecular Ene Reactions of Allenylsilanes. Synthesis 1998, 1998, 509–521.(o)Ohmiya H; Ito H; Sawamura M General and functional group-tolerable approach to allenylsilanes by rhodium-catalyzed coupling between propargylic carbonates and a silylboronate. Org. Lett 2009, 11, 5618–20. [PubMed: 19908884] (p)Curtis-Long MJ; Aye Y Vinyl-, propargyl-, and allenylsilicon reagents in asymmetric synthesis: a relatively untapped resource of environmentally benign reagents. Chem. - Eur. J 2009, 15, 5402–16. [PubMed: 19421982]
- (11). (a)Wang H; Jui NT Catalytic Defluoroalkylation of Trifluoromethylaromatics with Unactivated Alkenes. J. Am. Chem. Soc 2018, 140, 163–166. [PubMed: 29256607] (b)Vogt DB; Seath CP; Wang H; Jui NT Selective C-F Functionalization of Unactivated Trifluoromethylarenes. J. Am. Chem. Soc 2019, 141, 13203–13211. [PubMed: 31369264] (c)Thornbury RT; Toste FD Palladium-Catalyzed Defluorinative Coupling of 1-Aryl-2,2-Difluoroalkenes and Boronic Acids: Stereoselective Synthesis of Monofluorostilbenes. Angew. Chem., Int. Ed 2016, 55, 11629–11632.
- (12). (a)Hu J; Zhao Y; Shi Z Highly tunable multi-borylation of gem-difluoroalkenes via copper catalysis. Nature Catal. 2018, 1, 860–869.(b)Zhang J; Dai W; Liu Q; Cao S Cu-Catalyzed Stereoselective Borylation of gem-Difluoroalkenes with B<sub>2</sub>pin<sub>2</sub>. Org. Lett 2017, 19, 3283–3286. [PubMed: 28581762] (c)Tan D-H; Lin E; Ji W-W; Zeng Y-F; Fan W-X; Li Q; Gao H; Wang H Copper-Catalyzed Stereoselective Defluorinative Borylation and Silylation of gem-Difluoroalkenes. Adv. Synth. Catal 2018, 360, 1032–1037.(d)Sakaguchi H; Uetake Y; Ohashi M; Niwa T; Ogoshi S; Hosoya T Copper-Catalyzed Regioselective Monodefluoroborylation of Polyfluoroalkenes en Route to Diverse Fluoroalkenes. J. Am. Chem. Soc 2017, 139, 12855–12862. [PubMed: 28849929] (e)Sakaguchi H; Ohashi M; Ogoshi S Fluorinated Vinylsilanes from the Copper-Catalyzed Defluorosilylation of Fluoroalkene Feedstocks. Angew. Chem., Int. Ed 2018, 57, 328–332.(f)Niwa T; Ochiai H; Hosoya T Copper-Catalyzed ipso-Borylation of Fluoroarenes. ACS Catal. 2017, 7, 4535–4541.(g)Hu J; Han X; Yuan Y; Shi Z Stereoselective Synthesis of Z Fluoroalkenes through Copper-Catalyzed Hydrodefluorination of gem-Difluoroalkenes with Water. Angew. Chem., Int. Ed 2017, 56, 13342–13346.(h)Andrella NO; Xu N; Gabidullin BM; Ehm C; Baker RT Selective Copper Complex-Catalyzed Hydrodefluorination of Fluoroalkenes and Allyl Fluorides: A Tale of Two Mechanisms. J. Am. Chem. Soc 2019, 141, 11506–11521. [PubMed: 31305996] (i)Kojima R; Kubota K; Ito H Stereodivergent hydrodefluorination of gem-difluoroalkenes: selective synthesis of (Z)- and (E)-monofluoroalkenes. Chem. Commun 2017, 53, 10688–10691.
- (13). (a)Wang M; Pu X; Zhao Y; Wang P; Li Z; Zhu C; Shi Z Enantioselective Copper-Catalyzed Defluoroalkylation Using Arylboronate-Activated Alkyl Grignard Reagents. J. Am. Chem. Soc 2018, 140, 9061–9065. [PubMed: 29989802] (b)Paioti PHS; Del Pozo J; Mikus MS; Lee J; Koh MJ; Romiti F; Torker S; Hoveyda AH Catalytic Enantioselective Boryl and Silyl Substitution with Trifluoromethyl Alkenes: Scope, Utility, and Mechanistic Nuances of Cu-F beta-Elimination. J. Am. Chem. Soc 2019, 141, 19917–19934. [PubMed: 31809041] (c)Kojima R; Akiyama S; Ito H A Copper(I)-Catalyzed Enantioselective gamma-Boryl Substitution of Trifluoromethyl-Substituted Alkenes: Synthesis of Enantioenriched gamma,gamma-gem-Difluoroallylboronates. Angew. Chem., Int. Ed 2018, 57, 7196–7199.(d)Jang YJ; Rose D;

Mirabi B; Lautens M Rhodium-Catalyzed Enantioselective Defluorinative alpha-Arylation of Secondary Amides. *Angew. Chem., Int. Ed* 2018, 57, 16147–16151.(e)Huang Y; Hayashi T Rhodium-Catalyzed Asymmetric Arylation/Defluorination of 1-(Trifluoromethyl)alkenes Forming Enantioenriched 1,1-Difluoroalkenes. *J. Am. Chem. Soc* 2016, 138, 12340–3. [PubMed: 27627581] (f)Gao P; Yuan C; Zhao Y; Shi Z Copper-Catalyzed Asymmetric Defluoroborylation of 1-(Trifluoromethyl)Alkenes. *Chem.* 2018, 4, 2201–2211.(g)Akiyama S; Kubota K; Mikus MS; Paioti PHS; Romiti F; Liu Q; Zhou Y; Hoveyda AH; Ito H Catalytic Enantioselective Synthesis of Allylic Boronates Bearing a Trisubstituted Alkenyl Fluoride and Related Derivatives. *Angew. Chem., Int. Ed* 2019, 58, 11998–12003.


- (14). (a)Butcher TW; Yang JL; Amberg WM; Watkins NB; Wilkinson ND; Hartwig JF Desymmetrization of difluoromethylene groups by C-F bond activation. *Nature* 2020, 583, 548–553. [PubMed: 32480398]
- (15). (a)Garcia-Rubia A; Romero-Revilla JA; Mauleon P; Gomez Arrayas R; Carretero JC Cu-catalyzed silylation of alkynes: a traceless 2-pyridylsulfonyl controller allows access to either regioisomer on demand. *J. Am. Chem. Soc* 2015, 137, 6857–65. [PubMed: 25955333] (b)Hazra CK; Fopp C; Oestreich M Copper(I)-catalyzed regioselective addition of nucleophilic silicon across terminal and internal carbon-carbon triple bonds. *Chem. - Asian J* 2014, 9, 3005–10. [PubMed: 25065952] (c)Zhao M; Shan CC; Wang ZL; Yang C; Fu Y; Xu YH Ligand-Dependent-Controlled Copper-Catalyzed Regio- and Stereoselective Silaboration of Alkynes. *Org. Lett* 2019, 21, 6016–6020. [PubMed: 31318220] (d)Meng F; Jang H; Hoveyda AH Exceptionally E- and beta-selective NHC-Cu-catalyzed proto-silyl additions to terminal alkynes and site- and enantioselective proto-boryl additions to the resulting vinylsilanes: synthesis of enantiomerically enriched vicinal and geminal borosilanes. *Chem. - Eur. J* 2013, 19, 3204–14. [PubMed: 23325733] (e)Fujihara T; Tani Y; Semba K; Terao J; Tsuji Y Copper-catalyzed silacarboxylation of internal alkynes by employing carbon dioxide and silylboranes. *Angew. Chem., Int. Ed* 2012, 51, 11487–90.(f)Wang P; Yeo XL; Loh TP Copper-catalyzed highly regioselective silylcupration of terminal alkynes to form alpha-vinylsilanes. *J. Am. Chem. Soc* 2011, 133, 1254–6. [PubMed: 21208005]
- (16). (a)Uehling MR; Suess AM; Lalic G Copper-catalyzed hydroalkylation of terminal alkynes. *J. Am. Chem. Soc* 2015, 137, 1424–7. [PubMed: 25621888] (b)Uehling MR; Rucker RP; Lalic G Catalytic anti-Markovnikov hydrobromination of alkynes. *J. Am. Chem. Soc* 2014, 136, 8799–803. [PubMed: 24896663] (c)Shi SL; Buchwald SL Copper-catalysed selective hydroamination reactions of alkynes. *Nat. Chem* 2015, 7, 38–44. [PubMed: 25515888] (d)Mailig M; Hazra A; Armstrong MK; Lalic G Catalytic Anti-Markovnikov Hydroallylation of Terminal and Functionalized Internal Alkynes: Synthesis of Skipped Dienes and Trisubstituted Alkenes. *J. Am. Chem. Soc* 2017, 139, 6969–6977. [PubMed: 28449580] (e)Chemla F; Ferreira F The Chemistry of Organocopper Compounds; Rappoport Z, Marek I, Eds.; Patai Series; Wiley-Chichester, 2009; 527 pp, DOI: 10.1002/9780470682531.pat0444.(f)Whittaker AM; Lalic G Monophasic catalytic system for the selective semireduction of alkynes. *Org. Lett* 2013, 15, 1112–5. [PubMed: 23427913] (g)Barbero A; Pulido FJ Allylsilanes and vinylsilanes from silylcupration of carbon-carbon multiple bonds: scope and synthetic applications. *Acc. Chem. Res* 2004, 37, 817–25. [PubMed: 15491128]
- (17). (a)Yoshida H; Hayashi Y; Ito Y; Takaki K Inverse regioselectivity in the silylstannylation of alkynes and allenes: copper-catalyzed three-component coupling with a silylborane and a tin alkoxide. *Chem. Commun* 2015, 51, 9440–2.(b)Xu YH; Wu LH; Wang J; Loh TP Synthesis of multi-substituted vinylsilanes via copper(I)-catalyzed hydrosilylation reactions of allenes and propiolate derivatives with silylboronates. *Chem. Commun* 2014, 50, 7195–7. (c)Tani Y; Yamaguchi T; Fujihara T; Terao J; Tsuji Y Copper-catalyzed Silylative Allylation of Ketones and Aldehydes Employing Allenes and Silylboranes. *Chem. Lett* 2015, 44, 271–273.(d)Tani Y; Fujihara T; Terao J; Tsuji Y Copper-catalyzed regiodivergent silacarboxylation of allenes with carbon dioxide and a silylborane. *J. Am. Chem. Soc* 2014, 136, 17706–9. [PubMed: 25469703] (e)He ZT; Tang XQ; Xie LB; Cheng M; Tian P; Lin GQ Efficient Access to Bicyclo[4.3.0]nonanes: Copper-Catalyzed Asymmetric Silylative Cyclization of Cyclohexadienone-Tethered Allenes. *Angew. Chem., Int. Ed* 2015, 54, 14815–8.(f)Fujihara T; Sawada A; Yamaguchi T; Tani Y; Terao J; Tsuji Y Boraformylation and Silaformylation of Allenes. *Angew. Chem., Int. Ed* 2017, 56, 1539–1543.(g)Chang XH; Liu ZL; Luo YC; Yang C; Liu XW; Da BC; Li JJ; Ahmad T; Loh TP; Xu YH Copper-catalyzed silylation reactions of

- propargyl epoxides: easy access to 2,3-allenols and stereodefined alkenes. *Chem. Commun* 2017, 53, 9344–9347.
- (18). Lee J; Radomkit S; Torker S; Del Pozo J; Hoveyda AH Mechanism-based enhancement of scope and enantioselectivity for reactions involving a copper-substituted stereogenic carbon centre. *Nat. Chem* 2018, 10, 99–108. [PubMed: 29256506]
- (19). Tian YM; Guo XN; Kuntze-Fechner MW; Krummenacher I; Braunschweig H; Radius U; Steffen A; Marder TB Selective Photocatalytic C-F Borylation of Polyfluoroarenes by Rh/Ni Dual Catalysis Providing Valuable Fluorinated Arylboronate Esters. *J. Am. Chem. Soc* 2018, 140, 17612–17623. [PubMed: 30474979]
- (20). Somerville RJ; Hale LVA; Gomez-Bengoia E; Bures J; Martin R Intermediacy of Ni-Ni Species in sp(2) C-O Bond Cleavage of Aryl Esters: Relevance in Catalytic C-Si Bond Formation. *J. Am. Chem. Soc* 2018, 140, 8771–8780. [PubMed: 29909614]
- (21). Dang H; Mailig M; Lalic G Mild copper-catalyzed fluorination of alkyl triflates with potassium fluoride. *Angew. Chem., Int. Ed* 2014, 53, 6473–6.
- (22). (a)He Y; Wang B; Dukor RK; Nafie LA Determination of absolute configuration of chiral molecules using vibrational optical activity: a review. *Appl. Spectrosc* 2011, 65, 699–723. [PubMed: 21740631] (b)Merten C; Golub TP; Kreienborg NM Absolute Configurations of Synthetic Molecular Scaffolds from Vibrational CD Spectroscopy. *J. Org. Chem* 2019, 84, 8797–8814. [PubMed: 31046276] (c)Polavarapu PL; Santoro E Vibrational optical activity for structural characterization of natural products. *Nat. Prod. Rep* 2020, 37, 1661–1699. [PubMed: 32608433] (d)Chamberlain BT; Vincent M; Nafie J; Muller P; Greka A; Wagner FF Multigram Preparation of BRD4780 Enantiomers and Assignment of Absolute Stereochemistry. *J. Org. Chem* 2021, 86, 4281–4289. [PubMed: 33618515]
- (23). Devlin FJ; Stephens PJ; Cheeseman JR; Frisch MJ Ab Initio Prediction of Vibrational Absorption and Circular Dichroism Spectra of Chiral Natural Products Using Density Functional Theory: Camphor and Fenchone. *J. Phys. Chem. A* 1997, 101, 6322–6333.
- (24). (a)Polavarapu PL; Covington CL Comparison of experimental and calculated chiroptical spectra for chiral molecular structure determination. *Chirality* 2014, 26, 539–52. [PubMed: 24644231] (b)Debie E; De Gussem E; Dukor RK; Herrebout W; Nafie LA; Bultinck P A confidence level algorithm for the determination of absolute configuration using vibrational circular dichroism or Raman optical activity. *ChemPhysChem* 2011, 12, 1542–1549. [PubMed: 21542094]
- (25). (a)Cirriez V; Rasson C; Hermant T; Petriguet J; Diaz Alvarez J; Robeyns K; Riant O Copper-catalyzed addition of nucleophilic silicon to aldehydes. *Angew. Chem., Int. Ed* 2013, 52, 1785–8. (b)Gibbons SK; Hughes RP; Glueck DS; Royappa AT; Rheingold AL; Arthur RB; Nicholas AD; Patterson HH Synthesis, Structure, and Luminescence of Copper(I) Halide Complexes of Chiral Bis(phosphines). *Inorg. Chem* 2017, 56, 12809–12820. [PubMed: 29064687] (c)Gurung SK; Thapa S; Kafle A; Dickie DA; Giri R Copper-catalyzed Suzuki-Miyaura coupling of arylboronate esters: transmetalation with (PN)CuF and identification of intermediates. *Org. Lett* 2014, 16, 1264–7. [PubMed: 24499358] (d)Lee M; Nguyen M; Brandt C; Kaminsky W; Lalic G Catalytic Hydroalkylation of Allenes. *Angew. Chem., Int. Ed* 2017, 56, 15703–15707. (e)Mita T; Chen J; Sugawara M; Sato Y One-Pot Synthesis of  $\alpha$ -Amino Acids from CO<sub>2</sub> Using a Bimetal Reagent with Si–B Bond. *Org. Lett* 2012, 14, 6202–6205. [PubMed: 23205912] (f)Shibasaki M; Kanai M Copper(I) Fluoride and Copper(II) Fluoride. In *Encyclopedia of Reagents for Organic Synthesis*; John Wiley & Sons, Ltd.: Hoboken, NJ, 2007; DOI: 10.1002/047084289X.rm00744. (g)Suess AM; Uehling MR; Kaminsky W; Lalic G Mechanism of Copper-Catalyzed Hydroalkylation of Alkynes: An Unexpected Role of Dinuclear Copper Complexes. *J. Am. Chem. Soc* 2015, 137, 7747–53. [PubMed: 26042355] (h)Wyss CM; Tate BK; Bacsa J; Wieliczko M; Sadighi JP Dinuclear  $\mu$ -fluoro cations of copper, silver and gold. *Polyhedron* 2014, 84, 87–95.
- (26). (a)Meng F-F; Xie J-H; Xu Y-H; Loh T-P Catalytically Asymmetric Synthesis of 1,3-Bis(silyl)propenes via Copper-Catalyzed Double Proto-Silylations of Polar Enynes. *ACS Catal.* 2018, 8, 5306–5312. (b)Plotzitzka J; Kleeberg C [(NHC)Cu(I)-ER3] Complexes (ER3 = SiMe<sub>2</sub>Ph, SiPh<sub>3</sub>, SnMe<sub>3</sub>): From Linear, Mononuclear Complexes to Polynuclear Complexes with Ultrashort Cu(I)–Cu(I) Distances. *Inorg. Chem* 2016, 55, 4813–23. [PubMed: 27145039] (c)Plotzitzka J; Kleeberg C [(18-C-6)K][(N identical with C)Cu(I)-SiMe<sub>2</sub>Ph], a Potassium Silylcyanocuprate as a Catalyst Model for Silylation Reactions with Silylboranes: Syntheses, Structures, and Catalytic Properties. *Inorg. Chem* 2017, 56, 6671–6680. [PubMed: 28509537]

- (27). (a) Ohmiya H; Yokobori U; Makida Y; Sawamura M General approach to allenes through copper-catalyzed gamma-selective and stereospecific coupling between propargylic phosphates and alkylboranes. *Org. Lett* 2011, 13, 6312–5. [PubMed: 22054013] (b) Zhong C; Sasaki Y; Ito H; Sawamura M The synthesis of allenes by Cu(I)-catalyzed regio- and stereoselective reduction of propargylic carbonates with hydrosilanes. *Chem. Commun* 2009, 5850–2.
- (28). (a) Kuehn L; Stang M; Wurtemberger-Pietsch S; Friedrich A; Schneider H; Radius U; Marder TB FBpin and its adducts and their role in catalytic borylations. *Faraday Discuss.* 2019, 220, 350–363. [PubMed: 31513207] (b) Pietsch S; Neeve EC; Apperley DC; Bertermann R; Mo F; Qiu D; Cheung MS; Dang L; Wang J; Radius U; Lin Z; Kleeberg C; Marder TB Synthesis, Structure, and Reactivity of Anionic sp(2)-sp(3) Diboron Compounds: Readily Accessible Boryl Nucleophiles. *Chem. - Eur. J* 2015, 21, 7082–98. [PubMed: 25877472]
- (29). Harutyunyan SR; Lopez F; Browne WR; Correa A; Pena D; Badorrey R; Meetsma A; Minnaard AJ; Feringa BL On the mechanism of the copper-catalyzed enantioselective 1,4-addition of Grignard reagents to alpha,beta-unsaturated carbonyl compounds. *J. Am. Chem. Soc* 2006, 128, 9103–18. [PubMed: 16834384]
- (30). The formations of several possible off-cycle adducts of CsF with LCuF and PhMe2SiBpin were calculated (see Figure S1). The DFT calculations suggest that although these adducts are slightly more stable than the monomeric species, the dissociation of these adducts are facile. For simplicity, PhMe2SiBpin and FBpin were used as the reference in the computed reaction energy profile.
- (31). A Lewis base-promoted antielimination pathway, proposed in ref 13b, is unlikely in the present system because a Lewis base will not be able to coordinate to the highly sterically encumbered Cu in 31.
- (32). (a) Kobayashi K; Yamamoto Y; Miyaura N Pd/Josiphos-Catalyzed Enantioselective  $\alpha$ -Arylation of Silyl Ketene Acetals and Mechanistic Studies on Transmetalation and Enantioselection. *Organometallics* 2011, 30, 6323–6327. (b) Li C; Liu RY; Jesikiewicz LT; Yang Y; Liu P; Buchwald SL CuH-Catalyzed Enantioselective Ketone Allylation with 1,3-Dienes: Scope, Mechanism, and Applications. *J. Am. Chem. Soc* 2019, 141, 5062–5070. [PubMed: 30817137]
- (33). (a) Lu G; Liu RY; Yang Y; Fang C; Lambrecht DS; Buchwald SL; Liu P Ligand-Substrate Dispersion Facilitates the Copper-Catalyzed Hydroamination of Unactivated Olefins. *J. Am. Chem. Soc* 2017, 139, 16548–16555. [PubMed: 29064694] (b) Kang T; Erbay TG; Xu KL; Gallego GM; Burtea A; Nair SK; Patman RL; Zhou R; Sutton SC; McAlpine IJ; Liu P; Engle KM Multifaceted Substrate-Ligand Interactions Promote the Copper-Catalyzed Hydroboration of Benzylidenecyclobutanes and Related Compounds. *ACS Catal.* 2020, 10, 13075–13083. [PubMed: 33791144] (c) Xi Y; Su B; Qi X; Pedram S; Liu P; Hartwig JF Application of Trimethylgermyl-Substituted Bisphosphine Ligands with Enhanced Dispersion Interactions to Copper-Catalyzed Hydroboration of Disubstituted Alkenes. *J. Am. Chem. Soc* 2020, 142, 18213–18222. [PubMed: 32962336]
- (34). (a) Bizet V; Cahard D Asymmetric Fluorination Methods: Application in the Stereoselective Synthesis of Fluorinated Drugs. In *Stereoselective Synthesis of Drugs and Natural Products*; John Wiley & Sons, Inc., 2013. DOI: 10.1002/9781118596784.ssd044. (b) Wang J; Sánchez-Roselló M; Aceña JL; del Pozo C; Sorochinsky AE; Fustero S; Soloshonok VA; Liu H Fluorine in pharmaceutical industry: fluorine-containing drugs introduced to the market in the last decade (2001–2011). *Chem. Rev* 2014, 114, 2432–2506. [PubMed: 24299176] (c) Gillis EP; Eastman KJ; Hill MD; Donnelly DJ; Meanwell NA Applications of fluorine in medicinal chemistry. *J. Med. Chem* 2015, 58, 8315–8359. [PubMed: 26200936] (d) Zhu Y; Han J; Wang J; Shibata N; Sodeoka M; Soloshonok VA; Coelho JAS; Toste FD Modern Approaches for Asymmetric Construction of Carbon-Fluorine Quaternary Stereogenic Centers: Synthetic Challenges and Pharmaceutical Needs. *Chem. Rev* 2018, 118, 3887–3964. [PubMed: 29608052]




**Figure 1.**  
Proposed catalytic cycle.






**Figure 2.** Computed reaction energy profiles of the Cu-catalyzed silylation and asymmetric  $\beta$ -fluoride elimination. Gibbs free energies and enthalpies (in kcal/mol) are with respect to the monomeric copper fluoride **26**.







**Figure 3.** Origin of enantioselectivity in *syn*- $\beta$ -fluoride elimination. Most hydrogen atoms are omitted for clarity. Gibbs free energies and enthalpies are with respect to **31**.  $E_{\text{Pauli}}$  is the Pauli repulsion energy between the substrate and the Josiphos ligand from EDA calculations.

## (a) Naturally Occurring bromoallenes


(b) Literature Precedented  $\beta$ -Fluoride Elimination(c) This Work: Enantioselective  $\beta$ -Fluoride Elimination**Scheme 1.**

Allenyl Halide Natural Products and Synthesis of Monofluoro ACAs *via* Enantioselective  $\beta$ -Fluoride Elimination


**(a) Decarboxylative Bromination****(b) Copper-Catalyzed Sonogashira Cross-Coupling**

**Scheme 2.**  
Improved Synthesis of Propargylic Difluorides

Table 1.

Structural Effect of Josiphos on Transformation<sup>a</sup>

entry	R; R'	yield (SM %) <sup>b</sup>	ee (%) <sup>c</sup>
1	Me; H	59% (23)	45
2	Me; OMe	31% (14)	44
3	<sup>t</sup> Bu; OMe	41% (<1)	70
4	Mes; H	58% (38)	75
5	TTB; H	3% (79)	nd
6	TMS; H	73% (<1)	75
7	TES; H	66% (<1)	83
8	CF <sub>3</sub> ; H	43% (<2)	71
9	<sup>t</sup> Pr <sup>(F7)</sup> ; H	44% (43)	86




<sup>a</sup>Standard conditions: **1a** (0.10 mmol, 1.0 equiv), **2** (0.135 mmol, 1.35 equiv), CuCl (5 mol %), L (6 mol %), THF (1.0 mL), 65 °C, 24 h.

<sup>b</sup>Yield was determined by <sup>19</sup>F NMR of crude reaction, using PhF as an internal standard.

<sup>c</sup>Determined by HPLC with a chiral stationary phase.

Table 2.

Reaction Optimization<sup>a</sup>


entry	Cu	base (mol %)	solvent	yield (SM %) <sup>b</sup>	ee (%) <sup>c</sup>
1	Cu(MeCN) <sub>4</sub> BF <sub>4</sub>	A (20)	PhMe	50% (49)	90
2	"	A (20), CsF (100)	"	83% (13)	90
3	"	CsF (150)	"	75% (11)	91
4	Cu(MeCN) <sub>4</sub> BF <sub>4</sub>	"	THF	39% (48)	74
5	"	"	dioxane	65% (20)	91
6	"	"	cyclohexane	50% (31)	87
7	"	"	MTBE	75% (10)	92
8	" <sup>d</sup>	"	"	74% (<1)	91
9	Cu(MeCN) <sub>4</sub> OTf <sup>d</sup>	"	"	82% (4)	90
10	CuOTf·1/2C <sub>6</sub> H <sub>6</sub>	CsF (160)	"	98% (-)	90


<sup>a</sup>Standard conditions: **1a** (0.10 mmol, 1.0 equiv), **2** (0.135 mmol, 1.35 equiv), [Cu] (5 mol %), L (6 mol %), solvent (1.5 mL), 32 °C, 24 h. A = NaO(2-OMeC<sub>6</sub>H<sub>4</sub>).

<sup>b</sup>Yield was determined by <sup>19</sup>F NMR of crude reaction, using PhF as an internal standard.

<sup>c</sup>Determined by HPLC with a chiral stationary phase.

<sup>d</sup>MeCN removed before reaction.

Table 3.

Scope of Asymmetric  $\beta$ -Fluoride Elimination<sup>a</sup>

<sup>a</sup> Standard conditions: **1-25a** (0.20 mmol, 1.0 equiv), **2** (0.27 mmol, 1.35 equiv), CuOTf•1/2C<sub>6</sub>H<sub>6</sub> (6 mol %), L (7 mol %), CsF (1.6–1.8 equiv), 9:1 PhMe:MTBE or MTBE (3.0 mL), 35–45 °C, 24 h. Reported yields are of isolated allene. Enantiomeric excess was determined by HPLC with a chiral stationary phase.

Author Manuscript

Author Manuscript

Author Manuscript


Author Manuscript

<sup>b</sup> **1a** (6.0 mmol), CsF(25%)-CaF<sub>2</sub> (1.6 equiv), 30 h.

<sup>c</sup> **1a** (2.5–3.0 mmol), R<sub>3</sub>SiBpin (1.35–1.45 equiv), Cu (8–9 mol %), 3,5-TripJosphos (9–10 mol %), CsF(25%)-CaF<sub>2</sub> (1.8–2.5 equiv), MTBE, 28–45 °C, 30–48 h.

<sup>d</sup> NaO(2-OMeC<sub>6</sub>H<sub>4</sub>) (30 mol %), CsF (1.0 equiv), PhMe, 27 °C.

Table 4.

Relative Free Energies of Conformers of (*R,S*)-3,5-TMS-Josiphos-Supported Copper Complexes and Transition States<sup>a</sup>


ligand conformation	L <sub>2</sub> CuF (26)	$\sigma$ -bond metathesis (TS-1)	silyl copper (30)	migratory insertion (TS-2)	alkenyl copper (31)	<i>syn</i> - $\beta$ -fluoride elimination	
						TS-3	TS-4
A	<b>0.0</b>	<b>6.7</b>	<b>-24.7</b>	2.9	-42.8	-26.7	-28.4
B	4.8	10.8	-23.5	<b>-4.8</b>	<b>-47.1</b>	<b>-32.5</b>	<b>-30.6</b>

<sup>a</sup> All Gibbs free energies are in kcal/mol with respect to the monomeric copper fluoride **26**. Bold numbers indicate the favorable ligand conformation.