

Correction: *Heart failure medication dosage and survival in women and men seen at outpatient clinics*

Bots SH, Onland-Moret NC, Tulevski II, *et al.* Heart failure medication dosage and survival in women and men seen at outpatient clinics. *Heart* 2021;107:1748–55.

This article has been corrected since it was first published to correct . The rows were misaligned in the ‘Ejection fraction’ section; please see below for the corrected portion of the table.

	Women	Men
Ejection fraction (n, %)		
<40%	162 (28.9)	291 (47.3)
40%–49%	81 (15.3)	123 (20.0)
>50%	272 (48.5)	155 (25.2)
Not recorded	41 (7.3)	46 (7.5)



OPEN ACCESS

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

Heart 2022;108:e4. doi:10.1136/heartjnl-2021-319229corr1

