


ORIGINAL ARTICLE

Phosphorylation of Yun is required for stem cell proliferation and tumorigenesis

Xuejing Ren¹ | Hang Zhao¹ | Lin Shi¹ | Zhengran Li¹ | Ruiyan Kong¹ |
Rui Ma² | Lemei Jia³ | Shan Lu³ | Jian-Hua Wang³ | Meng-qiu Dong³ |
Yingchun Wang^{4,5} | Zhouhua Li¹ 

¹College of Life Sciences, Capital Normal University, Beijing, China

²Department of Neurology, Capital Medical University, Beijing, China

³National Institute of Biological Sciences, Beijing, China

⁴State Key Laboratory of Plant Genomics, Institute of Genetics and Developmental Biology, The Innovative Academy of Seed Design, Chinese Academy of Sciences, Beijing, China

⁵College of Advanced Agricultural Sciences, University of Chinese Academy of Sciences, Beijing, China

Correspondence

Zhouhua Li, College of Life Sciences, Capital Normal University, Beijing 100048, China.
Email: zhli@cnu.edu.cn

Funding information

Beijing Municipal Commission of Education, Grant/Award Number: KZ201910028040; National Natural Science Foundation of China, Grant/Award Numbers: 31972893, 31471384, 92054109

Abstract

Stem cells maintain adult tissue homeostasis under physiological conditions. Uncontrolled stem cell proliferation will lead to tumorigenesis. How stem cell proliferation is precisely controlled is still not fully understood. Phosphorylation of Yun is essential for ISC proliferation. Yun is essential for the proliferation of normal and transformed intestinal stem cells. Our mass spectrometry and biochemical data suggest that Yun can be phosphorylated at multiple residues in vivo. Interestingly, we show that the phosphorylation among these residues is likely interdependent. Furthermore, phosphorylation of each residue in Yun is important for its function in ISC proliferation regulation. Thus, our study unveils the important role of post-translational modification of Yun in stem cell proliferation.

1 | INTRODUCTION

Tissue homeostasis is maintained by residential stem cells, which proliferate and produce differentiated progeny to replenish lost cells. Thus, the proliferation (self-renewal) and differentiation of adult stem cells must be tightly balanced under physiological conditions. Disruption of this balance will result in excessive stem cell or precocious stem cell differentiation and finally stem cell depletion, eventually leading to various diseases, such as cancer and precocious aging.^{1–3} In particular, it has been proposed that various tumours possess cancer stem cells (CSCs), which are the driving force of tumour development and progression, dormancy, and recurrence.^{4–6} Therefore, illustrating

the underlying mechanisms of stem cell proliferation control is critical for understanding homeostasis regulation and for the development of potential therapeutics to treat human diseases including cancer.

The adult *Drosophila* intestine is an excellent model to study the regulation of stem cell proliferation/differentiation and tumorigenesis, which shares marked similarities in terms of development, cellular makeup, and genetic control with its mammalian counterparts.^{7–11} *Drosophila* intestinal stem cells (ISCs) are distributed along the basement membrane of the adult intestinal epithelium^{12,13} and divided asymmetrically to produce differentiating enteroblasts (EBs) or EE progenitors (EEPs). One of the Notch ligands, Delta, is specifically expressed in ISCs, while the Notch receptor is expressed in ISCs, EBs,

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *Cell Proliferation* published by John Wiley & Sons Ltd.

and EEPs (termed progenitors collectively). Notch signalling activation in EBs promotes their differentiation into absorptive enterocytes (ECs).^{12,14-17} Recent studies show that EE cells may not be generated

from EBs, but directly from ISCs or EEPs, which divide once to produce two EEs.¹⁸⁻²⁰ The proliferation and differentiation of ISCs are regulated by multiple signalling pathways such as the Notch, Wntless




FIGURE 1 Legend on next page.

(Wg), JAK/STAT, EGFR, Hippo, Insulin, Hedgehog, and BMP signalling to maintain tissue homeostasis under physiological and stressed conditions (see reviews²¹⁻²⁴ and references therein).

In our recent study, we identified a novel intrinsic factor Yun (Luck in Chinese) which sustains the proliferation of normal and transformed ISCs.²⁵ Yun is a novel protein without any known domains or motifs. During the course of our study, another group showed that it is implicated in cell proliferation in larval brain and spermatogenesis with unknown mechanisms and named it as *diamond (dind)*.²⁶ Yun is required for the proliferation of ISCs under physiological conditions and tissue regeneration under stress conditions.²⁵ The identification of Yun as an essential stem cell proliferation regulator has important applications, which can be used as a target to treat intestinal malignancies. However, it remains unexplored whether its activity in ISC proliferation is regulated by any post-translational modifications (PTMs).

In this study, we investigate whether PTMs exist in Yun and whether these PTMs are required for Yun's function in ISC proliferation. We find that Yun is phosphorylated *in vivo*. Several phosphorylation residues are identified by mass spectrometry. Site-directed mutagenesis analyses indicate that the phosphorylation of these residues may be interdependent and is important for the proper function of Yun in ISC proliferation regulation.

2 | RESULTS

2.1 | Yun is a positive regulator of ISC proliferation

To identify regulators involved in ISC maintenance and proliferation/differentiation, we carried out a genome-wide RNAi screen in *Drosophila* adult posterior midgut using an *esgGal4, UAS-GFP, tubGal80^{ts}* (*esg^{ts}*) driver, which is expressed in the progenitors (ISCs, EBs, and EEPs).^{25,27,28} From the screen, we identified many known and novel ISC regulators, including Yun (encoded by *CG7705*).²⁵ Yun encodes a novel protein without any known domains or motifs. We examined its expression pattern by immunostaining using a Yun-specific antibody and found that in adult intestines, Yun was mainly expressed in progenitors and EEs (Figure S1). Compared to the control, knocking down *yun* by expressing an effective shRNA RNAi line (*yun^{shRNA}*) showed a


significant reduction in the number of *esg⁺* cells (Figures 1A-C and S2). Consistently, the number of ISCs undergoing mitosis was significantly reduced in *esg^{ts} > yun^{shRNA}* intestines (Figures 1D-F). However, the observed ISC proliferation defects in the absence of Yun were not caused by cell death, indicating that Yun is required for the proliferation of normal ISCs under physiological conditions (Figure S3). We further examined whether Yun is required in ISCs for ISC proliferation control by depleting *yun* using an ISC-specific driver, *DI-Gal4*. Depletion of Yun specifically in ISCs also resulted in a significant reduction in the number of ISCs (Figures 1G-I). To further confirm the role of Yun played in ISC proliferation control, we generated *yun* null mutant ISC clones using MARCM clone technique.^{25,29} Compared with the control ISC clones which contained multiple cells, most *yun* null mutant ISC clones failed to proliferate and contained only 1 cell per clone, supporting the notion that Yun sustains ISC proliferation under normal conditions (Figures 1J-L). Conversely, ectopic expression of *yun* resulted in significant increase in progenitors (Figures 1M-O). Altogether, these data indicate that Yun is required for ISC proliferation under physiological conditions.

As Yun is also expressed in EE cells, we further examined what will happen in the absence of *yun* in EE cells. We depleted Yun specifically in EE cells using the EE-specific driver *prosGal4*. No obvious defects were observed when Yun is depleted in EEs, with no significant change of progenitors and EE cells, indicating that Yun in EE cells is dispensable for ISC proliferation (Figure S4).

2.2 | Yun is required for tumorigenesis

We then explored whether *yun* is also required for tumorigenesis. Ectopic expression of a constitutively active Raf (*Raf^{off}*, an activated component of the EGFR/MAPK signalling pathway) in progenitors causes rapid ISC proliferation and eventually leads to tumorigenesis (Figure 2A).^{30,31} We found that depletion of *yun* effectively suppressed the growth and proliferation of *Raf^{off}* tumours (Figures 2A-C), indicating that Yun is essential for tumorigenesis. Consistently, the number of mitotic cells observed in *esg^{ts} > Raf^{off}* intestines was dramatically suppressed by Yun depletion (Figures 2D-F). We further explored whether Yun specifically functions in ISCs to sustain the growth and

FIGURE 1 Yun is required for ISC proliferation. (A) *esg⁺* cells (green, by *esg > GFP*) in Control intestines at 29°C for 7 days (white arrowheads). (B) The number of *esg⁺* cells (green) is significantly decreased in *esg^{ts} > yun^{shRNA}* intestines at 29°C for 7 days compared to control (white arrowheads). (C) Quantification of the relative number of *esg⁺* cells in Control and *esg^{ts} > yun^{shRNA}* intestines. Mean ± SD is shown. *n* = 15. *****p* < 0.0001. (D) pH 3 staining (red) in Control intestines at 29°C for 7 days (white arrowheads). (E) pH 3 staining (red) in *esg^{ts} > yun^{shRNA}* intestines at 29°C for 7 days. (F) Quantification of the number of pH 3 cells per gut in Control and *esg^{ts} > yun^{shRNA}* intestines. Mean ± SD is shown. *n* = 20. *****p* < 0.0001. (G) ISCs (green, by *DI > GFP*) in Control intestines at 29°C for 7 days (white arrowheads). (H) Compared with Control, the number of ISCs (green) is significantly decreased in *DI^{ts} > yun^{shRNA}* intestines at 29°C for 7 days (white arrowheads). (I) Quantification of the relative number of ISCs in Control and *DI^{ts} > yun^{shRNA}* intestines. Mean ± SD is shown. *n* = 15. *****p* < 0.0001. (J) MARCM clones of FRT Control (green) at room temperature for 6 days after clone induction (ACI). (K) The clone size of *yun^{Δ9}* is significantly decreased (ACI 6 days). (L) Quantification of the clone size in Control and *yun^{Δ9}* mutant. Mean ± SD is shown. *n* = 15. *****p* < 0.0001. (M) *esg⁺* cells (green, by *esg > GFP*) in Control intestines at 29°C for 7 days (white arrowheads). Note that this panel is same as panel A for easy comparison. (N) The number of *esg⁺* cells (green) is significantly increased in *esg^{ts} > flag-yun* intestines at 29°C for 7 days compared to control (white arrowheads). (O) Quantification of the relative number of *esg⁺* cells in Control and *esg^{ts} > flag-yun* intestines. Mean ± SD is shown. *n* = 18. ***p* < 0.01. In all panels except graphs, GFP is in green and blue indicates DAPI staining of DNA. Scale bars: 20 μm



proliferation of *Raf^{gof}* tumour. Ectopic expression of *Raf^{gof}* in ISCs led to significant increase of *esg⁺* cells, forming tumours (Figure 2G). Consistently, depletion of *yun* completely suppressed the proliferation of *Raf^{gof}* tumours (Figures 2G–I). To further confirm the notion that Yun is

required for the proliferation of transformed ISCs derived from *Raf^{gof}* activation (or cancer stem cells, CSCs), we used MARCM technique to examine the roles of Yun in tumorigenesis. The size of *Raf^{gof}* MARCM clones is significantly increased compared to that of control, whereas




FIGURE 3 S450, S530, S532, S534, and S536 sites of endogenous Yun protein may be phosphorylated in vivo. (A–E) Five serine residues are likely to be phosphorylated identified by MS analysis in Yun: S450 (A), S530 (B), S532 (C), S534 (D), and S536 (E)




FIGURE 4 Yun is phosphorylated in vivo. (A) Detection of the phosphorylation status of Yun (pYun) using anti-phosphorylated serine/threonine (pS/T) and anti-phosphorylated serine (pS) antibodies by IP and western blot, respectively. NC: negative control. The 50 KDa band is likely the IgG heavy chain (IgG HC). (B) Detection of pYun with phos-tag and it can be dephosphorylated by λ -PP. (C) Quantification of the grayscale of pYun with or without λ -PP. Mean \pm SD is shown. $n = 3$. * $p < 0.05$. (D) Detection of pYun with phos-tag in flies expressing different Flag-tagged Yun forms by *tub^{ts}*. (E) Quantification of pYun/total Yun ratio of different Yun forms. Mean \pm SD is shown. $n = 3$. *** $p < 0.001$; **** $p < 0.0001$

the proliferation of *Raf^{gof}* MARCM clones was completely suppressed in the absence of *yun* (Figures 2K,L). Collectively, these data demonstrate that Yun is essential for tumorigenesis (and CSC proliferation).

2.3 | Yun is post-translationally modified by phosphorylation in vivo

To explore the mechanism of how Yun regulates ISC proliferation and tumorigenesis, we immunoprecipitated endogenous Yun proteins with a

3XFlag tag at its C-terminus under the control of its endogenous promoter.²⁵ Interestingly, double bands of endogenous Yun proteins which were close to each other could be detected in co-IP experiments, indicating that some kind of post-translational modification (PTM) may occur on endogenous Yun protein in vivo (Figure S5). We then performed IP-MS experiments to search for the possible PTM occurred on endogenous Yun protein. Detailed analysis of our mass spectrometry results showed that five serine residues of Yun (S450, S530, S532, S534, and S536) were likely to be phosphorylated (Figure 3). These data indicate that Yun is likely post-translationally modified by phosphorylation in vivo.




FIGURE 5 Legend on next page.

2.4 | Phosphorylation of different residues of Yun may be interdependent

To further confirm that Yun protein is phosphorylated *in vivo*, we pulled down Yun by IP experiments and detected it with phosphorylated serine and/or threonine (pS/T)- and phosphorylated serine (pS)-specific antibodies, respectively. The results showed that Yun could be detected by both antibodies, supporting the notion that Yun is phosphorylated at serine residue(s) *in vivo* (Figure 4A). Consistently, phosphorylation of Yun could also be detected when analysed with phos-tag (Figure 4B). Next, we treated the immunoprecipitated Yun proteins with λ -phosphatase (λ -PP), which could de-phosphorylate the phosphorylated proteins. The results showed that treatment with λ -PP could effectively diminish phosphorylation of Yun proteins (Figure 4B,C). Together, these data show that Yun protein is post-translationally modified by phosphorylation *in vivo*.

We further examined the contribution of phosphorylation status of Yun by these candidate phosphorylation residues and whether the phosphorylation status of individual residue can affect phosphorylation of the other residues. We performed site-directed mutagenesis of these residues, changing each of them to alanine to make phosphorylation dead mutants, and generated transgenic flies carrying these mutants individually (Figure 4D). We then examined the effects of these single sites on the phosphorylation status of Yun using phos-tag. We found that mutation of each single serine residue significantly affected, but did not completely abolish the phosphorylation of Yun protein, supporting the notion that Yun protein is phosphorylated at multiple serine residues (Figure 4D). Of note, mutation of each single serine residue affected the phosphorylation status of Yun at different extents, implying that phosphorylation of individual residue may affect the phosphorylation status of the other residue(s), thus the phosphorylation of these residues may be interdependent (Figure 4D,E).

2.5 | Phosphorylation of Yun is critical for ISC proliferation

Protein phosphorylation plays an important roles in regulating protein activity and function.^{32,33} To collectively address the role of phosphorylation of these serine residues in ISC proliferation regulation, we constructed two transgenes: one expressed a phosphorylation dead form of all five serine residues (*yun[5SA]*), while the other expressed a

phosphorylation mimic form of all five serine residues (*yun[5SD]*), and examined their function. Compared with control MARCM ISC clones, the ISC MARCM clones of *yun* null mutant failed to proliferate (Figure 5A,B). Ectopic expression of wildtype *yun* fully restored the proliferation defects observed in *yun* null mutant (Figures 5A–C and 5F). Whereas ectopic expression of phosphorylation dead *yun(5SA)* failed to restore the proliferation defects observed in *yun* null mutant, suggesting that the phosphorylation of these five serine residues collectively is essential for Yun function in ISC proliferation control (Figure 5D,F). Interestingly, ectopic expression of phosphorylation mimic *yun(5SD)* only partially restored the proliferation defects observed in *yun* null mutant, suggesting that the proper phosphorylation status of each single residue in Yun is also important for its function in ISC proliferation control (Figure 5E,F). Taken together, these data suggest that proper phosphorylation status of these five serine residues collectively is critical for Yun to regulate ISC proliferation.

We then examined the role of phosphorylation of individual serine residue in ISC proliferation regulation *in vivo* by performing individual rescue experiments using mutated Yun carrying either phosphorylation dead form (SA) or phosphorylation mimic form (SD) of a single residue, respectively. Consistently, ectopic expression of wildtype *yun* could restore the proliferation defects observed in *yun* null mutant (Figures 5G–I and 5T). Interestingly, ectopic expression of phosphorylation dead *yun(S450A)* failed to restore the proliferation defects observed in *yun* null mutant, suggesting that the phosphorylation of this single serine residue is critical for Yun's function in ISC proliferation control (Figure 5J and T). Whereas ectopic expression of phosphorylation mimic *yun(S450D)* significantly restored the proliferation defects observed in *yun* null mutant (Figure 5K and T). These data show that the phosphorylation of the serine 450 residue is important for Yun to regulate ISC proliferation. Similar results were obtained when either *yun(S530A)* or *yun(S530D)* was ectopically expressed, indicating that the phosphorylation of serine 530 residue is also important for Yun's function in ISC proliferation control (Figure 5L, M, and T). Furthermore, ectopic expression of phosphorylation dead *yun(S532A)*, *yun(S534A)*, and *yun(S536A)* failed to restore the proliferation defects observed in *yun* null mutant, while ectopic expression of phosphorylation mimic *yun(S532D)*, *yun(S534D)*, and *yun(S536D)* did not statistically restore the proliferation defects observed in *yun* null mutant, although ISC clones containing 2–3 cells could be occasionally observed, indicating that that the phosphorylation of serine 532, 534, and 536 residues is also necessary for Yun's function in

FIGURE 5 The phosphorylation of the five serine sites in Yun is collectively required for ISC proliferation. (A) MARCM clones of FRT control (green, ACI 6 days), DI and Pros are in red. (B) The clone size of *yun*^{Δ9} is significantly decreased (green, ACI 6 days). (C) Overexpressing Yun can rescue the clone size of *yun*^{Δ9} (green, ACI 6 days). (D) Expression of *yun(5SA)* mutant cannot rescue the proliferation defects observed in *yun*^{Δ9} (green, ACI 6d). (E) Expression of *yun(5SD)* mutant partially rescues the proliferation defects observed in *yun*^{Δ9} (green, ACI 6 days). (F) Quantification of the clone size of indicated genotypes. Mean ± SD is shown. $n = 20$. ^{ns} $p > 0.05$; * $p < 0.05$. (G) MARCM clones of FRT control (red, ACI 6 days). (H) The clone size of *yun*^{Δ9} is significantly decreased (red, ACI 6 days). (I) Overexpressing *yun* can rescue the clone size of *yun*^{Δ9} (red, ACI 6 days). (J, L, N, P, and R) Expression of different single phospho-dead point mutants of *yun(1SA)* cannot rescue the proliferation defects observed in *yun*^{Δ9} (red, ACI 6 days). (K, M, O, Q, and S) Expression of different single phospho-mimic point mutants of *yun(1SD)* partially rescue the proliferation defects observed in *yun*^{Δ9} (red, ACI 6 days). (T) Quantification of the clone size of indicated genotypes. Mean ± SD is shown. $n \geq 20$. ^{ns} $p > 0.05$; **** $p < 0.0001$. In all panels except graphs, GFP is in green and blue indicates DAPI staining of DNA. Scale bars: 20 μ m

ISC proliferation control (Figures 5N–T). Collectively, these data suggest that phosphorylation of all the five serine residues identified is important for Yun to regulate ISC proliferation, but the role(s) played by individual serine residue may be different and phosphorylation statuses among these residues are likely interdependent.

3 | DISCUSSION

The proliferation and differentiation of adult stem cells must be tightly balanced in order to maintain tissue homeostasis and prevent tumorigenesis. However, how stem cell proliferation is properly controlled, in particular how post-translational modifications of intrinsic regulators affect their functions in stem cell proliferation control, remains not fully understood. Here, we find that novel intrinsic Yun is phosphorylated at multiple serine residues *in vivo* and that the phosphorylation of these serine residues in Yun protein is critical for its function in ISC proliferation control. Interestingly, the phosphorylation of these serine residues may be interdependent and phosphorylation of Yun on the function of Yun in sustaining ISC proliferation seems to be complicated and the role(s) played by individual serine residue may be different.

Protein phosphorylation is one of the most widely observed PTMs, which plays important roles in regulating protein abundance, subcellular localization, function, and activity.^{32,33} Functional analysis of phosphorylated proteins is particularly important for deciphering the causes of diseases like cancer and neurodegenerative diseases, as well as for the development of therapeutic applications.^{34,35} Protein phosphorylation and de-phosphorylation reactions catalysed by protein kinases and protein phosphatases (PPases) are dynamic which change over time and space. The complex network of protein phosphorylation/de-phosphorylation provides rapid regulation of the phosphorylated status and supports many fundamental biological functions of cells. Thus identification of the protein kinases and possible PPases of a phosphorylated protein will be critical for unveiling the regulatory network of this protein and elucidating its molecular mechanism. Interestingly, two casein kinases (CKI α and CKII β) were found to associate with Yun in our IP-MS results, and their association was further confirmed by co-IP experiments, implying that they may be the kinases responsible for Yun phosphorylation (Figure S5). Consistent with this possibility, ectopic expression of either CKI α or CKII β could not restore the proliferation defects observed in the absence of Yun (Figure S6). CKI and CKII are members of Ser/Thr kinases; they interact with many other proteins, such as β -catenin, other kinases like PIP4K and S6KII, the RNA-binding protein Orb, circadian proteins (Tim (Timeless) and Per (Period)), and are involved in multiple signalling pathways, including Wnt, Hh, and Hpo.^{36–41} The two casein kinases are essential for ISC proliferation. Depletion of them completely blocks ISC proliferation. Importantly, ectopic expression of both wildtype *yun* and phosphorylation mimic *yun(5SD)* could not restore the proliferation defects in the absence of either casein kinase, suggesting that multiple downstream substrates are required for ISC proliferation control and expression of

single substrate is not sufficient to restore ISC proliferation blockage upon their depletion (Figure S5).

Our identification of Yun as a phosphorylated protein provides an interesting cutting point to elucidate the underlying mechanism of how Yun regulates ISC proliferation and tumorigenesis. It will be interesting to systematically investigate how these serine residues of Yun are differentially phosphorylated in ISCs, whether any interdependence(s) of phosphorylation status exists among these residues, and how the phosphorylation of these residues is coordinated to regulate the function of Yun in ISC proliferation control using site-specific phospho-Yun antibodies and corresponding knock-in flies carrying single and/or combined phospho-dead and phospho-mimic *yun* mutant in future work.

4 | EXPERIMENTAL PROCEDURES

4.1 | Fly lines and husbandry

Flies were maintained on standard media at 25°C. Two–three–days-old flies were selected and transferred to 29°C, unless otherwise specified. Flies were transferred to new vials with fresh food every day and dissected at specific time points as indicated. In all experiments, only female posterior midguts were analysed. Information about alleles and transgenes used in this study can be found either in FlyBase or as noted: *esgGal4*, *UAS-GFP*, *tubGal80^{ts}* (*esg^{ts}*, gift from N. Perrimon), *tubGal80^{ts}, tubGal4*, *UAS-Ra^{gof}* (BL2033), *yun^{shRNA}*, *yun^{Δ9}*, *UAS-flag-yun*, and *hsFlp*, *ActGal4*, *UAS-GFP*, *FRT82B-tubGal80* (for MARCM clonal analysis), *hsFlp*, *ActGal4*, *UAS-RFP*, *FRT82B-tubGal80* (for MARCM clonal analysis),²⁵ *DIGal4*, *tubGal80^{ts}* (*Dl^{ts}*, gifts from S. Hou and R. Xi), *esgGal4*, *Gal80^{ts}*, *UAS-CC3Ai*,⁴² *prosGal4*, *UAS-GFP*, *tubGal80^{ts}* (*pros^{ts}*, gift from Xiaohang. Yang), *Cklα^{v20}* (THU5314 and HMS02276), *FRT19A-Cklα^{8B12}* (BL63802), *CklIIβ^{v20}* (THU0596, BL34939, and HMS00058), *FRT19A-Ck2β^A* (BL67713), *UAS-Cklα-HA* (BL55067 and BL55068), and *UAS-CklIIβ* (F001372).

4.2 | RNAi knockdown and overexpression experiments

Crosses (unless stated otherwise) were maintained at 18°C to bypass potential requirements during early developmental stages. Two–three–days-old progeny with the desired genotypes were collected after eclosion and maintained at 29°C to inactivate Gal80^{ts} before dissection and immunostaining. The flies were transferred to new vials with fresh food every day.

4.3 | Constructs and transgenic flies

pUAST-flag-yun (CG7705) was constructed by cloning *yun* ORF (from cDNA clone LP22035, BDGP) into *pUAST-nflag*. *attB-pUAST-CklIIβ-myc* was constructed by cloning the ORF of *UAS-CklIIβ* (F001372) into the EcoRI and XbaI site of *attB-pUAST-myc*. All point mutation

constructs were made by Hieff Mut™ Multi Site-Directed Mutagenesis Kit (YEASEN, China). *attB-pUAST-yun(5SA:S450/530/532/534/536A)-flag-GFP* and *attB-pUAST-yun(5SD:S450/530/532/534/536D)-flag-GFP* vectors were constructed by cloning the mutated *yun(5SA)* and *yun(5SD)* into the EcoRI site of *attB-pUAST-flag-GFP* respectively. *attB-pUAST-flag-yun(1SA:S450A, S530A, S532A, S534A, or S536A, respectively)* and *attB-pUAST-flag-yun(1SD:S450D, S530D, S532D, S534D, or S536D, respectively)* vectors were constructed by cloning the mutated *yun(1SA)* and *yun(1SD)* into the KpnI site of *attB-pUAST-nflag*. Transgenic flies were obtained by standard P-element-mediated germline transformation carrying attP site at 36B, 75B, or 86F.

4.4 | MARCM clone analysis

The clonal analyses were achieved using the MARCM system.²⁹ The ISC clones were induced by heat shocking Two–three–days-old adult flies at 37°C for 60 min. The flies were maintained at a 25°C incubator and transferred to new vials with fresh food every day. The sizes of the marked clones were assayed at 6 days after clone induction (6D ACI, at least 10 midguts for each genotype were assayed).

4.5 | Immunostainings and fluorescence microscopy

For standard immunostainings, intestines were dissected in 1 × PBS (10 mM NaH₂PO₄/Na₂HPO₄, 175 mM NaCl, pH 7.4), and fixed in 4% paraformaldehyde for 25 min at room temperature. Samples were washed with 1 × PBT (0.1% Triton X-100 in 1 × PBS) and blocked in 3% BSA in 1 × PBT for 45 min. Primary antibodies were added to the samples and incubated at 4°C overnight. The following primary antibodies were used: mouse mAb anti-DI (C594.9B, 1:50, developed by S. Artavanis-Tsakonas, DSHB), mouse mAb anti-Prospero (MR1A, 1:100, developed by Chris Doe, DSHB), rabbit anti-pH 3 (1:2000, Millipore), rabbit anti-active caspase 3 (1:1000, Cell Signalling), and rabbit anti-Yun (1:1000).²⁵ Primary antibodies were detected by fluorescent-conjugated secondary antibodies from Jackson ImmunoResearch Laboratories. Secondary antibodies were incubated for 2 h at room temperature. DAPI (Sigma-Aldrich; 0.1 µg/ml) was added after secondary antibody staining. The samples were mounted in mounting medium (70% glycerol containing 2.5% DABCO). All images were captured using a Zeiss inverted confocal microscope and were processed in Adobe Photoshop and Illustrator.

4.6 | Co-immunoprecipitation and western blotting

Fly tissues were lysed in RIPA buffer [50 mM Tris.HCl, pH 8.0, 150 mM NaCl, 5 mM EDTA, pH 8.0, 0.5% Triton X-100, 0.5% NP-40, 0.5% sodium deoxycholate, and complete protease inhibitor cocktail tablets (Roche)] on ice for 30 min. After centrifugation, lysates were then diluted tenfold with RIPA buffer and subjected to immunoprecipitation using anti-FLAG M2 affinity gel (A2220; Sigma-Aldrich, USA).

The immunocomplexes were collected by centrifugation and washed with 1 ml of RIPA buffer three times. For western blotting, immunoprecipitated proteins were separated in SDS-PAGE and then blotted onto PVDF membranes. The membranes were stained with primary antibody overnight at 4°C. Followed by washing, PVDF membranes were incubated with secondary antibodies conjugated with HRP, then the membranes were scanned using Luminescent Image Analyser (GE, Sweden). Rabbit anti-Yun (1:1000),²⁵ mouse anti-Flag (1:1000, Sigma-Aldrich, USA), Rabbit anti-pS (1:1000, Abcam), Rabbit anti-pS/T (1:1000, Abcam), and mouse monoclonal anti-βTubulin (1:1000, 3G6, Abbkine, USA) antibodies were used. Phos-tag (Wako, Japan) was added into SDS-PAGE gel to exhibit phosphorylated Yun. λ-phosphatase (NEB) was used to dephosphorylate pYun.

4.7 | MS sample preparation

Proteins were precipitated with 25% trichloroacetic acid (TCA) for at least 30 min on ice. The protein pellets were washed twice with 500 µl ice-cold acetone, air dried, and then resuspended in 8 M urea, 100 mM Tris, and pH 8.5. After reduction (5 mM TCEP, RT, 20 min) and alkylation (10 mM iodoacetamide, RT, 15 min in the dark), the samples were diluted to 2 M urea with 100 mM Tris, pH 8.5 and digested with trypsin at 1/50 (w/w) enzyme/substrate ratio at 37°C for 16–18 h. The digestion was then stopped by addition of formic acid to 5% (final concentration).

4.8 | LC-MS/MS analysis

All the samples were analysed using an EASY-nLC 1000 system (Thermo Fisher Scientific) interfaced with a Q-Exactive mass spectrometer (Thermo Fisher Scientific). Peptides were loaded on a trap column (75 µm ID, 4 cm long, packed with ODS-AQ 12 nm-10 µm beads) and separated on an analytical column (75 µm ID, 12 cm long, packed with Luna C18 1.9 µm 100 Å resin) with a 60 min linear gradient at a flow rate of 200 µl/min as follows: 0–5% B in 2 min, 5–30% B in 43 min, 30–80% B in 5 min, and 80% B for 10 min (A = 0.1% FA, B = 100% ACN, 0.1% FA). Spectra were acquired in data-dependent mode: the top ten most intense precursor ions from each full scan (resolution 70,000) were isolated for HCD MS2 (resolution 17,500; NCE 27) with a dynamic exclusion time of 30 s. The AGC targets for the MS1 and MS2 scans were 3e6 and 1e5, respectively, and the maximum injection times for MS1 and MS2 were both 60 ms. Precursors with 1+, more than 7+ or unassigned charge states were excluded.

4.9 | Database search

The MS data were searched against a Uniprot *Drosophila melanogaster* protein database (database ID number of UP000000803) using ProLuCID with the following parameters: precursor mass tolerance, 3 Da; fragment mass tolerance 20 ppm; peptide length, minimum

6 amino acids and maximum 100 amino acids; enzyme, Trypsin, with up to three missed cleavage sites.⁴³ The results were filtered by DTASelect requiring FDR <1% at the peptide level and spectra count ≥ 2.⁴⁴ The proteins identified from the negative control and Flag-Yun IP were contrasted by Contrast.⁴⁴

4.10 | Data analysis

The number of intestines scored is indicated in the text. To determine the relative number of *esg*⁺ cells, confocal images of 40× lens/1.0 zoom from a defined posterior midgut region of different genotypes indicated were acquired. The number of *esg*⁺ cells from each confocal image was determined using Image-Pro Plus software, manually selecting the “filter” depending on the respective cell size to filter out background signals (referred to as the relative number of *esg*⁺ cells). The clone sizes were scored manually under Zeiss Imager Z2/LSM780 microscope for indicated genotypes. At least 10 different guts were analysed for each set. Statistical analysis was done using the Student's *t*-test. PEMS 3.1 software was used for SEM analyses and Sigma plot and GraphPad prism software for graph generation. The graphs were further modified using Adobe Photoshop and Illustrator. ⁿ*p* > 0.05; **p* < 0.05; ***p* < 0.01; *** *p* < 0.001; and **** *p* < 0.0001.

ACKNOWLEDGEMENTS

We are grateful to Yu Cai, Steven Hou, Rongwen Xi, Norbert Perrimon, Xiaohang Yang, DSHB, Bloomington *Drosophila* Stock Center, VDRC, NIG-FLY stock center, and TRIP for reagents and stocks, and DGRC for cDNA clones. This work is supported by grants from Beijing Municipal Commission of Education (No. KZ201910028040 to Zhouhua Li) and the National Natural Science Foundation of China (Nos. 92054109, 31972893, and 31471384 to Zhouhua Li).

CONFLICT OF INTEREST

The authors declare no conflicts of interest.

AUTHOR CONTRIBUTIONS

Conceptualization, Zhouhua Li and Hang Zhao; investigation, Xuejing Ren, Hang Zhao, Lin Shi, Zhengran Li, Ruiyan Kong, Rui Ma, Lemei Jia, Shan Lu, Jian-Hua Wang, Meng-qiu Dong, Yingchun Wang, Zhouhua Li; formal analysis, Xuejing Ren, Zhouhua Li, Hang Zhao; methodology, Rui Ma; validation, Xuejing Ren, Hang Zhao, and Zhouhua Li; writing—original draft preparation, Xuejing Ren, Hang Zhao, and Zhouhua Li; writing—review and editing, Hang Zhao and Zhouhua Li; supervision, Zhouhua Li; project administration, Zhouhua Li; funding acquisition, Zhouhua Li. All authors have read and agreed to the published version of the manuscript.

DATA AVAILABILITY STATEMENT

The data that supports the findings of this study are available in the supplementary material of this article.

ORCID

Zhouhua Li  <https://orcid.org/0000-0002-7660-8579>

REFERENCES

- Radtke F, Clevers H. Self-renewal and cancer of the gut: two sides of a coin. *Science*. 2005;307:1904-1909.
- Lin H. Cell biology of stem cells: an enigma of asymmetry and self-renewal. *J Cell Biol*. 2008;180:257-260.
- Morrison SJ, Spradling AC. Stem cells and niches: mechanisms that promote stem cell maintenance throughout life. *Cell*. 2008;132:598-611.
- Chaffer CL, Weinberg RA. A perspective on cancer cell metastasis. *Science*. 2011;331:1559-1564.
- Kleffel S, Schatton T. Tumor dormancy and cancer stem cells: two sides of the same coin? *Adv Exp Med Biol*. 2013;734:145-179.
- Kreso A, Dick JE. Evolution of the cancer stem cell model. *Cell Stem Cell*. 2014;14:275-291.
- Casali A, Batlle E. Intestinal stem cells in mammals and *drosophila*. *Cell Stem Cell*. 2009;4:124-127.
- Edgar BA. Intestinal stem cells: no longer immortal but ever so clever. *EMBO J*. 2012;31:2441-2443.
- Gervais L, Bardin AJ. Tissue homeostasis and aging: new insight from the fly intestine. *Curr Opin Cell Biol*. 2017;48:97-105.
- Jasper H. Intestinal stem cell aging: origins and interventions. *Annu Rev Physiol*. 2020;82:203-226.
- Stainier DYR. No organ left behind: tales of gut development and evolution. *Science*. 2005;307:1902-1904.
- Micchelli CA, Perrimon N. Evidence that stem cells reside in the adult *drosophila* midgut epithelium. *Nature*. 2006;439:475-479.
- Ohlstein B, Spradling A. The adult *drosophila* posterior midgut is maintained by pluripotent stem cells. *Nature*. 2006;439:470-474.
- Beebe K, Lee WC, Michelli CA. JAK/STAT signaling coordinates stem cell proliferation and multilineage differentiation in the *drosophila* intestinal stem cell lineage. *Dev Biol*. 2010;338:28-37.
- Ohlstein B, Spradling A. Multipotent *drosophila* intestinal stem cells specify daughter cell fates by differential notch signaling. *Science*. 2007;315:988-992.
- Perdigoto CN, Schweisguth F, Bardin AJ. Distinct levels of notch activity for commitment and terminal differentiation of stem cells in the adult fly intestine. *Development*. 2011;138:4585-4595.
- Yeung T, Chia L, Kosinski C, Kuo C. Regulation of self-renewal and differentiation by the intestinal stem cell niche. *Cell Mol Life Sci*. 2011;68:2513-2523.
- Biteau B, Jasper H. Slit/Robo signaling regulates cell fate decisions in the intestinal stem cell lineage of *drosophila*. *Cell Rep*. 2014;7:1867-1875.
- Chen J, Xu N, Wang C, et al. Transient Scute activation via a self-stimulatory loop directs enteroendocrine cell pair specification from self-renewing intestinal stem cells. *Nat Cell Biol*. 2018;20:152-161.
- Zeng X, Han L, Singh SR, et al. Genome-wide RNAi screen identifies networks involved in intestinal stem cell regulation in *drosophila*. *Cell Rep*. 2015;10:1226-1238.
- Colombani J, Andersen DS. The *drosophila* gut: a gatekeeper and coordinator of organism fitness and physiology. *WIREs Dev Biol*. 2020;9:e378.
- Guo Z, Lucchetta E, Rafel N, Ohlstein B. Maintenance of the adult *drosophila* intestine: all roads lead to homeostasis. *Curr Opin Genet Dev*. 2016;40:81-86.
- Jiang H, Tian A, Jiang J. Intestinal stem cell response to injury: lessons from *drosophila*. *Cell Mol Life Sci*. 2016;73:3337-3349.
- Joly A, Rousset R. Tissue adaptation to environmental cues by symmetric and asymmetric division modes of intestinal stem cells. *Int J Mol Sci*. 2020;21:6362.
- Zhao H, Shi L, Li Z, et al. The Yun/Prohibitin complex regulates adult *drosophila* intestinal stem cell proliferation through the transcription factor E2F1. *Proc Natl Acad Sci U S A*. 2022;119:e2111711119.
- Graziadio L, Palumbo V, Cipressa F, et al. Phenotypic characterization of diamond (*dind*), a *drosophila* gene required for multiple aspects of cell division. *Chromosoma*. 2018;127:489-504.

27. Liu F, Zhao H, Kong R, Shi L, Li Z, Ma R, Zhou H, Li Z. Derlin-1 and TER94/VCP/p97 are required for intestinal homeostasis. *J Genet Genomics*. 2022;49(3):195-207.
28. Ma H, Zhao H, Liu F, et al. Heparan sulfate negatively regulates intestinal stem cell proliferation in *drosophila* adult midgut. *Biol Open*. 2019;8: 047126.
29. Lee T, Luo L. Mosaic analysis with a repressible cell marker (MARCM) for *drosophila* neural development. *Trends Neurosci*. 2001;24:251-254.
30. Ma M, Zhao H, Binari R, Perrimon N, Li Z. Wildtype adult stem cells, unlike tumor cells, are resistant to cellular damages in *drosophila*. *Dev Biol*. 2016;411:207-216.
31. Zhao H, Shi L, Kong R, et al. Autophagy induction in tumor surrounding cells promotes tumor growth in adult *drosophila* intestines. *Dev Biol*. 2021;476:294-307.
32. Liu M, Su Y, Peng J, Zhu AJ. Protein modifications in hedgehog signaling: cross talk and feedback regulation confer divergent hedgehog signaling activity. *Bioessays*. 2021b;43:5.
33. Seok SH. Structural insights into protein regulation by phosphorylation and substrate recognition of protein kinases/phosphatases. *Life*. 2021;11:957.
34. Brognard J, Hunter T. Protein kinase signaling networks in cancer. *Curr Opin Genet Dev*. 2011;21:4-11.
35. Gross S, Rahal R, Stransky N, Lengauer C, Hoeflich KP. Targeting cancer with kinase inhibitors. *J Clin Invest*. 2015;125:1780-1789.
36. Akten B, Tangredi MM, Jauch E, et al. Ribosomal s6 kinase cooperates with casein kinase 2 to modulate the *drosophila* circadian molecular oscillator. *J Neurosci*. 2009;29:466-475.
37. Chen Y, Sasai N, Ma G, et al. Sonic hedgehog dependent phosphorylation by CK1 α and GRK2 is required for ciliary accumulation and activation of smoothened. *PLoS Biol*. 2011;9:14.
38. Jia H, Liu Y, Xia R, et al. Casein kinase 2 promotes hedgehog signaling by regulating both smoothened and Cubitus interruptus. *J Biol Chem*. 2010;285:37218-37226.
39. Szabó A, Papin C, Zorn D, et al. The CK2 kinase stabilizes CLOCK and represses its activity in the *drosophila* circadian oscillator. *PLoS Biol*. 2013;11:27.
40. Tsai CR, Galko MJ. Casein kinase 1 α decreases β -catenin levels at adherens junctions to facilitate wound closure in *drosophila* larvae. *Development*. 2019;146:175133.
41. Yanagawa S, Matsuda Y, Lee JS, et al. Casein kinase I phosphorylates the armadillo protein and induces its degradation in *drosophila*. *EMBO J*. 2002;21:1733-1742.
42. Schott S, Ambrosini A, Barbaste A, et al. A fluorescent toolkit for spatiotemporal tracking of apoptotic cells in living *drosophila* tissues. *Development*. 2017;144:3840-3846.
43. Xu T, Park SK, Venable JD, et al. ProLuCID: an improved SEQUEST-like algorithm with enhanced sensitivity and specificity. *J Proteomics*. 2015;129:16-24.
44. Tabb DL, McDonald WH, Yates JR. DTASelect and contrast: tools for assembling and comparing protein identifications from shotgun proteomics. *J Proteome Res*. 2002;1:21-26.

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: Ren X, Zhao H, Shi L, et al. Phosphorylation of Yun is required for stem cell proliferation and tumorigenesis. *Cell Prolif*. 2022;55(5):e13230. doi:[10.1111/cpr.13230](https://doi.org/10.1111/cpr.13230)