

Carceral Amplification of COVID-19

Impacts for Community, Corrections Officer, and Incarcerated Population Risks

Eric T. Lofgren,^a Kristian Lum,^b Aaron Horowitz,^c Brooke Mabubuonwu,^c Kellen Meyers,^d and Nina H. Fefferman^{e,f}

Abstract: COVID-19 is challenging many societal institutions, including our criminal justice systems. Some have proposed or enacted (e.g., the State of New Jersey) reductions in the jail and/or prison populations. We present a mathematical model to explore the epidemiologic impact of such interventions in jails and contrast them with the consequences of maintaining unaltered practices. We consider infection risk and likely in-custody deaths, and estimate how within-jail dynamics lead to spill-over risks, not only affecting incarcerated people but increasing exposure, infection, and death rates for both corrections officers and the broader community beyond the justice system. We show that, given a typical jail-community dynamic, operating in a business-as-usual way results in substantial, rapid, and ongoing loss of life. Our results are consistent with the hypothesis that large-scale reductions in arrest and speeding of releases are likely to save the lives of incarcerated people, jail staff, and the wider community.

Keywords: COVID-19; SARS-CoV-2; Carceral amplification; Jails
(*Epidemiology* 2022;33: 480–492)

Submitted February 10, 2021; accepted February 28, 2022
From the ^aPaul G. Allen School for Global Animal Health, Washington State University, Pullman, WA; ^bDepartment of Computer and Information Science, University of Pennsylvania, Philadelphia, PA; ^cACLU Analytics, American Civil Liberties Union, New York, NY; ^dDepartment of Mathematics, Tusculum University, Greeneville, TN; ^eDepartment of Mathematics, University of Tennessee, Knoxville, TN; and ^fDepartment of Ecology and Evolutionary Biology, University of Tennessee, Knoxville, TN. E.T.L. was supported by the CDC Cooperative Agreement RFA-CK-17-001-Modeling Infectious Diseases in Healthcare Program (MInD-Healthcare). A.H. and B.M. are employed as researchers at an advocacy organization, in collaboration with coauthors who are unpaid members of that same advocacy organization (E.T.L., K.L., K.M., N.F.F.). That organization had no role in determining how the study was conducted, or the results were obtained. Further, E.T.L. and N.F.F. are currently acting as pro bono expert witnesses for legal cases based on the results of this work.

All data used were gathered from publicly available, published sources, as referenced in the text and tables. Model source code is freely available on GitHub.

SDC Supplemental digital content is available through direct URL citations in the HTML and PDF versions of this article (www.epidem.com).

This manuscript was previously posted to medRxiv: doi: <https://doi.org/10.1101/2020.04.08.20058842>

Correspondence: Eric T. Lofgren, WSU Paul G. Allen School for Global Health, Allen Center, PO Box 647090, 1155 College Ave, Pullman, WA 99164. E-mail: eric.lofgren@wsu.edu.

Copyright © 2022 The Author(s). Published by Wolters Kluwer Health, Inc. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

ISSN: 1044-3983/22/334-480

DOI: 10.1097/EDE.0000000000001476

As the coronavirus disease 2019 (COVID-19) pandemic sweeps the globe, one of the critical functions of epidemiology is to consider how society can transform current practice to increase the health and safety of the public. Given the widespread risk of infection and the high case fatality rates, especially in older or medically compromised populations, the most effective strategies to reduce the impact of the disease may require that we be willing to consider structural reforms to our institutions to promote an overall greater good. To these ends, we have already seen systemic shifts in institutional practices that would be unthinkable under normal conditions: shelter-in-place orders closing businesses and restricting freedom of individual movement,¹ school closures to limit transmission compromising the ongoing education of children,² domestic travel restrictions and international border closures,³ suspension of visa processing,⁴ etc. Another clearly important institution that affects a substantial portion of the public directly⁵ and an even greater portion indirectly^{6–9} is our criminal legal system. The currently unfolding public health crisis makes clear the urgent need for rigorous analyses of the impact of maintaining current practices within these institutions, including both the costs to incarcerated people and their families as well as the community at large. We therefore explore the epidemiologic costs associated with our current system's functions as a necessary part of the policy conversations to decide whether or not these practices should be maintained or altered in response to a growing global crisis, especially with recurrent waves of new variants and incomplete vaccination coverage of the US population.

Analyzing incarcerated populations poses a unique epidemiologic problem for several reasons. The population experiences high rates of movement and turnover.^{10,11} Incarcerated people are responsible for purchasing their own hygiene products with limited resources.^{12,13} It is difficult or impossible for incarcerated people to practice recommended precautions from the US Centers for Disease Control and Prevention (CDC) for limiting transmission for a variety of factors including overcrowding and insufficient access to personal protective equipment.^{14–20} The incarcerated population has a higher expected rate of existing health conditions than the community from which they come.^{21–23} Jails are dependent completely on a

workforce that moves in and out of the jail and the community, including vendors, lawyers, corrections officers, medical staff, etc. There is strong evidence that incarceration itself is associated with profound adverse effects on the health of incarcerated people.^{24–26} These descriptors make jails highly likely not only to place detained people at increased risk of infection and resulting severe outcomes but also to function as a driver for increased infectivity, adversely impacting attempts to contain and mitigate disease spread in the broader communities in which jails are located.

To study the dynamics of this system and provide quantitative metrics for risk to incarcerated populations and the populations with which incarcerated people necessarily interact, we construct and tailor an epidemiologic model of COVID-19 transmission, and then use that model to consider how some possible reforms to the system will alter these baseline risks. In doing so, we focus only on interventions that do not rely on the suspension of any individual rights guaranteed to individuals by either the United States or individual state constitutions, but instead rely on elements of the criminal justice system that are already at the discretion of law enforcement, departments of corrections, and the court system (i.e., reduction in arrest intake, increased rates of returning incarcerated people to their homes, and improvement of conditions within the jails—indeed, these actions have already been undertaken by isolated, individual jails). We parameterize our model with data available from the Allegheny county jail system and perform a broad sensitivity analysis and comparison of relevant metrics to demonstrate how such models may apply to jail systems throughout the United States.

METHODS

Transmission Model

We begin by tailoring a standard Susceptible-Exposed-Infectious-Recovered (SEIR) model to the specific dynamics of COVID-19. We first split our total population into four categories of risk: children under 18 (denoted with the subscript K), low-risk adults (denoted with the subscript L), high-risk adults (denoted with the subscript H), and elderly adults (denoted with the subscript E). We also designate a separate population category for jail staff, O (note: while O was the selected notation, it is meant to capture all staff working at the jail, not only the corrections officers). These populations are then assigned into disease-related health status compartments: susceptible (S), exposed (in which individuals are pre-symptomatic, but do already produce low levels of infection transmission to others, E), infected (in which individuals are both symptomatic themselves and fully infectious to others, I), medically treated (those infectious individuals whose disease severity and healthcare access results in removal from the population into a medical care facility that prevents any further transmission of infection back into the population, M), and removed (those who have either recovered from the

infection and are now immune or those who have died, R). We also allow for the possibility that an infected person with sufficient disease severity to warrant medical treatment is unable to obtain care, and designate rates associated with this case, as designated by the subscript U . For clarity of the results, we do not consider death from any non-COVID-19 cause; this is done to highlight the COVID-19-specific dynamics. Additionally, as a simplifying assumption due to their low rates of both infections and complications, we do not model hospitalizations or deaths in children. Similarly, once hospitalized, patients are assumed not to spread COVID-19 further, as additionally modeling the impact of healthcare-associated COVID-19 cases is well beyond the scope of this model. Last, we split our population into segments depending on the subsection of the community or jail system in which they are currently functioning: the community at large, C , the processing system for the jail, P , the court system T , and the jail system, J .

A schematic for the whole model can be seen in Figure 1, and the differential equations comprising the model are in eAppendix 1; <http://links.lww.com/EDE/B908>. Figure 2 shows separately the model for transitions between locations (left) and the model for transitions between disease states (right). In this figure, we give the model parameters used to describe the rate of transition between locations and states. The model was implemented in R 3.6.3 using the *deSolve* package, with the visualization of results primarily using *ggplot2*.²⁷ Statistical analysis of one parameter (see below) was performed using the *flexsurv* package. The code and data used in this analysis are available at <https://www.github.com/epimodels/COVID19-Jails>. As this study used only publicly available data and does not involve human subjects, Institutional Review Board approval was not required.

Population Movement Into, Within, and Out of the Jail System

Our model captures movement between the community, processing, jail, and court—though this is meant to encompass all court appointments. It assumes staff move only between the community and jail; they are not arrested in our model. We base the parameters of movement into, within, and out of the jail on Allegheny County, PA, where detailed data on the jail population and facilities are available, including an automatically updating dashboard giving statistics for the jail population (<https://perma.cc/93RG-4WZ8>). In Allegheny County, the population at large is approximately 1.2 million people. The size of the jail population hovers around 2,500. We use these population figures to initialize our model.

In our model, individuals in the community are arrested at a rate of approximately 100 people per day (<https://perma.cc/9DSP-9CTY>). Based on discussion with experts, arrest rates were calibrated such that approximately 40% of those arrested were at high risk for COVID-19, though the underlying conditions this represents may be very different from those prevalent in the community at large (e.g., high levels




FIGURE 1. Schematic for a mathematical model of COVID-19 in a linked urban community-jail system. The population is represented in one of five possible compartments: S, E, I, needing M, and R. In addition, the population is divided into five distinct subpopulations: children under 18 years of age, elderly adults over 65 years of age, low-risk adults between 18 and 65, high-risk adults between 18 and 65 and jail staff (assumed to be between 18 and 65 years of age). Arrested adults move between the community, processing, and the court system and jail, while jail staff move between the community and jail. Children are assumed not to be eligible for arrest. E indicates exposed; I, infected; M, needing medical care; R, recovered/removed; S, susceptible.

of immune suppression from drug use).²⁸ Arrested individuals are brought to processing. From processing, individuals can either be released back into the community (60%) or taken to jail (40%). This results in an in-flow to the jail of approximately 40 individuals per day, which is consistent with Allegheny County's reporting. While in jail, individuals transition back and forth between the jail and court appointments. The daily number of movements between jail and court appointments is described as "well over 100" on the jail's web site. We assume movement of approximately 150 people per day between the jail and court. For each court appointment, we assume individuals spend half a day on average at the court facility. Importantly, we also assume that there is mixing in the court facility between those who are there for processing after arrest and those that are present there for court appointments. From the jail, individuals are released back into the community at a rate that is consistent with the reported 62-day average length of stay. One limitation of our model is that we do not account for

postjail destinations that are not the community, i.e., we do not model people moving from jail to prison. According to Wagner and Sakala,²⁹ the yearly number of admissions to prison is about 600,000 while the yearly number of admissions to jail is around 10.6 million. So, assuming that all prison admissions first had one jail admissions, around 95% of all jail admissions do not go on to prison; they are released back into the community as in our model. Thus, we expect that the omission of prison from our model does not substantially impact the overall findings.

An online database of public employees salaries in Allegheny County shows a population of 384 people whose job title is corrections officer, whose job location is the jail, and who are listed as active. Although this is certainly an underestimate of the total number of the jail's staff, which includes other types of employees, we think this is a useful approximation to the total number of staff. We use this figure as the number of staff members moving between community and jail. Staff transition between community and jail at a rate




FIGURE 2. Schematic for a mathematical model of COVID-19 in a linked urban community-jail system with parameter annotations. Left: Depiction of model emphasizing movement between locations. Here “Court” stands for both processing and trials. Right: Depiction of the model emphasizing transition between disease states. The parameter family is shown for each type of transition. For example, although we show just one α between community and court, in our model, there are separate subscripted α parameters corresponding to different rates of arrest for different age groups. Similar logic applies to the other parameters. E indicates exposed; I, infected; M, needing medical care; R, recovered/removed; S, susceptible.

that assumes 8-hour shift lengths in the jail per day with the remaining 16 hours per day spent in the community.

Estimation Population Mixing and Contact Rates

To estimate the β parameters governing movement between disease states, we break the problem into two parts: estimating an unscaled matrix of transmission rates between age groups and calculating appropriate scaling factors based on contact rates between different age strata. Because each location (community, jail, processing, and trial) comprises different numbers of people, this matrix must be rescaled by the population size in each location so that the mixing parameters have comparable meanings. We further scale β according to the assumption that people in jail, processing, and trial mix at a rate that is approximately three, six, and six times that of people in the community, respectively. This is based on qualitative evidence of increased mixing rates in those locations. Finally, we calibrate c_0 , the common scale parameter across all locations, using data on reported deaths in Allegheny County, PA, through the month after the relaxation of shelter-in-place conditions. Below we go into detail for each of the steps of this calculation. We first estimate parameters that describe the relative rate of transmission in the community between each of the community categories: children (under the age of 18, who are unable to enter into the jail model), adults 18–64 years of age, and elderly adults over the age of 65. Though we split low-risk adults and high-risk adults into different compartments in our model, we assume low-risk and high-risk adults have identical contact patterns. We denote the rate of transmission to category q from category r in the

community by β_{qr}^C . We decompose β_{qr}^C into a scale parameter β^C and matrix of scale-free relative transmission rates, β_{qr}^* .

To estimate the relative transmission rates, β_{qr}^* , we follow the methodology outlined in for modeling the age-stratified spread of COVID-19 in the United States. We define $\beta_{q,r}^* = u_q E_{q,r}$, where u_q is the susceptibility of people in age compartment q , and $E_{q,r}$ describes the mean number of effective contacts between individuals in the q th and r th age compartments of the contact matrix. As in Miller et al.,³⁰ we use consensus estimates of age-stratified susceptibility from Davies et al.,³¹ and use the contact matrix for a moderately aged population (the United Kingdom) as a stand-in for contact rates in Allegheny County, PA. Our model has fewer age groups than reported in Davies et al.³¹; for simplicity, we define only three age categories: We thus define u_q as a population-weighted average of the values reported in Davies et al.³¹ For example, Davies et al.³¹ reports u_{0-9} and u_{10-19} , the average susceptibility of individuals in the 0–9 and 10–19 age groups, respectively. To create our u_K , we set $u_K = \frac{n_{0-9}u_{0-9} + n_{10-17}u_{10-19}}{n_{0-9}n_{10-17}}$,

where n_{0-9} is the number of people in Allegheny County aged 0–9 and n_{10-17} is the number of people in Allegheny County aged 10–17. We use the same method to aggregate the average susceptibility for the 18- to 65-year-old age bracket and the 65 and over age bracket. We obtained age-stratified population totals for Allegheny County from the 2018 American Community Survey, conducted by the US Census Bureau. We use the socialmixr package R package³² to obtain the values of $E_{q,r}$. The resulting values of β_{qr}^* is shown in Table 1.

For each of the noncommunity locations, we assume that mixing patterns are not category-dependent. That is, we assume that all incarcerated people mix equally, regardless of age. Mathematically, this amounts to the assumption that $\beta_{qr}^j = \beta^j \beta^*$, $\beta_{qr}^p = \beta^p \beta^*$, and $\beta_{qr}^t = \beta^t \beta^*$, where β^* is defined to be consistent with the β^* ’s in the community, and β^j , β^p , and β^t are scaling parameters. Because most of the people in jail, processing, and trial are adults, we take $\beta^* = \beta_{LL}^*$.

This leaves setting the scale parameters, β^C , β^j , β^p , and β^t . There are two considerations to account for here: differences in the size of the population in each of the four locations, and differences in contact rates within the four locations. We define $\beta^C = \frac{c_0}{n_c}$, $\beta^j = \frac{c_0 c_j}{n_j}$, $\beta^p = \frac{c_0 c_p}{n_p}$, and $\beta^t = \frac{c_0 c_t}{n_t}$, where $n_c = 1,200,000$, $n_j = 2,600$, $n_t = 150$, and $n_p = 100$ are the size of the population in the community, jail, trial, and processing, respectively. Scaling each of the β terms in the model by the population sizes amounts to the assumption that transmission is contact-based. This assumption leads to conservative estimates of the speed of the spread in the jail system relative to a fomite-based transmission model. The parameters c_j , c_p , and c_t are factors that denotes how many times more contacts per day a person in jail or processing, respectively, has than a person in the community. We set these values to be $c_j = 3$ and $c_p = c_t = 6$, corresponding to an assumption of three and six times more contact in jail and processing and trial, respectively, than take place in the community. These values were chosen to reflect the conditions for both health and crowding of populations within jail facilities,^{21–26} and the understood gathering and transportation protocols associated with intake and release processing and court appearances.

The sensitivity of the model to these and other parameter choices is explored in eTable 1; <http://links.lww.com/EDE/B908>. An analysis of the sensitivity of the model to perturbations of the parameter values allows us to determine how changes in parameters (due to variation over time, uncertainty in measurement, or otherwise) impacts the quantitative outcomes of the model.³³ It is possible to use such a sensitivity analysis to identify the more influential parameters.³⁴

To calibrate the model for our baseline scenario, we then find a c_0 such that approximately 80% of the population is ultimately infected by the time the spread dies out in our model.³⁵ We selected an 80% final infection rate for consistency with predictions of the spread of COVID-19 under the assumption of no mitigation measures in place from an

influential microsimulation model. To calibrate the model for our shelter-in-place scenario, we select a c_0 that matches the trajectory of COVID-19 in Allegheny County under shelter-in-place. Specifically, we calibrate our model to the time series of deaths in Allegheny County as reported in data released by the New York Times. Because very few infections or deaths were reported before the initial shelter-in-place order on 23 March 2020, we assume the beginning of the death time series all took place under shelter-in-place conditions. On 15 May, the county entered the “yellow phase” in which businesses began reopening. Using previously published estimates of the incubation time and time to death of COVID-19, we assume a 30-day period between initial infection and death,^{36–38} so we take the total number of deaths that occurred under shelter-in-place to be the total number of deaths as of 15 June 2020, 30 days after shelter-in-place conditions were relaxed. There were 174 deaths that occurred as of this date. We set the start-date of the epidemic to be 1 month before the first death, 20 February 2020. We then perform a grid search to find a shelter-in-place scalar for c_0 such that over the course of approximately 115 days between 20 February 20 and 15 June, approximately 174 deaths occurred in our simulation and the trajectories are similar. eFigure 1; <http://links.lww.com/EDE/B908> shows a comparison between the time series of COVID-19 deaths in Allegheny County and the trajectory produced by the c_0 scalar that corresponds to the best fit.

Estimation of Other Model Parameters

Parameters concerning the natural history of COVID-19, patient progression, etc., were primarily obtained from existing estimates in the modeling literature, where possible using estimates from as close to the modeled catchment area as possible (i.e., CHIME from UPenn Medicine, <https://penn-chime.phl.io/>). Citations for specific parameter values may be found in Table 1.

In one case, $\hat{\gamma}$ or the asymptomatic period⁻¹, the original source reported that their estimate was likely an underestimation due to censoring. However, given that the authors provided the data within their manuscript,³⁹ the data were re-estimated to account for censoring using a parametric survival model assuming an exponential distribution (the distribution typically implied by the uniform hazard of transitioning from one compartment to another within a compartmental model). The fit for this exponential model may be found in eFigure 2; <http://links.lww.com/EDE/B908>.

Modeled Scenarios and Interventions

We represented the effects of several policy interventions or failures as changes to various parameters in this model. We consider four categories of scenarios that could vary the rate of spread: in addition to modeling shelter-in-place (reduced mixing) conditions in the community, we modeled scenarios related to reductions in arrest rates, increases in release rates, and changes to within-jail conditions. These scenarios are detailed in Table 2. Most scenarios are additive; that is, all arrest reduction interventions assume a baseline scenario of

TABLE 1. Relative Transmission Rates of COVID-19 From an Individual per Row to an Individual per Column

	Child	Adult	Elderly
Child	2.72	1.60	0.93
Adult	1.92	6.22	4.39
Elderly	0.09	0.49	1.38

TABLE 2. Parameter Values, Meanings, and Sources for a Community-jail Transmission Model of COVID-19

Parameter	Value	Description
σ	0.50	Percent reduction in transmission during asymptomatic period (compared with symptomatic) ⁴⁰
γ^{-1}	5.1 days	Incubation period ³⁷
$\hat{\gamma}^{-1}$	6.7 days	Asymptomatic period ³⁹
δ^{-1}	10 days	Symptomatic period ⁴⁰
$\delta_{\text{discharge}}^{-1}$	9 days	Hospitalization length of stay (discharged alive) ^{41a}
$\delta_{\text{death}}^{-1}$	4.2 days	Hospitalization length of stay (discharge dead) ⁴¹
$\delta_{\text{deathU}}^{-1}$	4.2 days	Time to death for unhospitalized critical cases
ω_L	$5.9\text{days}^{-1} \times 0.0625$	Time from symptom onset to hospitalization \times probability of needing hospitalization (low risk) ^{41,42}
ω_H	$5.9\text{days}^{-1} \times 0.118$	Time from symptom onset to hospitalization \times probability of needing hospitalization (high risk) ^{41,42}
ν	95%	Hospitalized case survival rate (low risk) ⁴³
ν_H	66.6%	Hospitalized case survival rate (high risk) ⁴³
ν_U	0.0%	Case survival rate for untreated cases in need of hospitalization (low risk) ⁴³
ν_{UH}	0.0%	Case survival rate for untreated cases in need of hospitalization (high risk)
α_L	3.57e-06	Per capita hourly arrest rate (low risk). Equates to 60 arrests per day
α_E	7.35e-06	Per capita hourly arrest rate (elderly). Equates to 1 arrest per day
α_H	1.11e-03	Per capita hourly arrest rate (high risk). Equates to 40 arrests per day
ψ_C	$12\text{hours}^{-1} \times 0.60$	Processing time from arrest to returning to community \times probability of release after arrest
ψ_J	$12\text{hours}^{-1} \times 0.40$	Processing time from arrest to jail \times probability of jail after arrest
κ	2.60e-03	Per capita hourly probability of scheduled court appearance
τ	12 hours	Time from scheduled court appearance to return to jail
ρ^{-1}	62 days	Length of stay in jail ^b
μ_C^{-1}	8 hours	Shift length for jail staff
μ_J^{-1}	16 hours	Time spent in the community for jail staff
ζ	1.00	Probability an incarcerated person needing treatment will receive it
pop	1.22 million people	Population of jail catchment area

^a<https://penn-chime.phl.io/>.

^b<https://www.alleghenycountyanalytics.us/index.php/category/topics/crime-and-justice/>.

shelter-in-place in the community. The scenarios involving faster release of individuals in jail all assume both shelter-in-place in the community, and were each run under each of the “Arrest Reduction” scenarios to determine the cumulative effects of arrest reduction, increased release rates, and community shelter-in-place conditions. The mixing reduction scenario in the jail assumes shelter-in-place in the community as well as a 25% reduction in arrests (equivalent to the “Bail Eligible” Arrest Reduction scenario), as it is unlikely that jails will be able to effectively reduce contact rates without reducing their average daily population. Finally, in the “Reduced Detection” scenario, we assume shelter-in-place, but vary the likelihood that serious cases of COVID-19 within the incarcerated population are caught and treated in a timely manner.

Below, vulnerable populations are defined as individuals over the age of 65 or at increased risk of complications from COVID-19 due to other comorbidities. We estimate that 40% of the jail population is vulnerable by this definition, based on reported rates of comorbidities among incarcerated people.⁴⁴ We estimate that around 25% of those arrested are bail eligible, based on information from Allegheny county on the rate at which cash bail was used between February and June of 2019.⁴⁵

RESULTS

Unsurprisingly given the epidemiological dynamics of COVID-19, absent any intervention our models showed a substantial outbreak in the community, causing 1,051,238

infections (both symptomatic and asymptomatic) as well as requiring 71,735 hospitalizations and ultimately resulting in 11,203 fatalities over the 180 days of the simulation, with the peak of infections occurring 81 days after the first infective case appeared in the population. Among those incarcerated, the outbreak was considerably more severe, causing a cumulative 4,779 cases requiring 312 hospitalizations and 58 deaths among those incarcerated, the 2,500 person jail being 0.2% the size of the wider community (Figure 3). The peak of this within-jail epidemic was also considerably earlier, with the peak of the epidemic occurring 29 days after the first infective case appeared in the community.

Given the dominant approach to controlling COVID-19 in the community and the widespread calls to “flatten the curve,” for the remaining results, we assume the presence of a shelter-in-place order or similar social distancing intervention only in the community as the comparator scenario, represented as a reduction in the mixing frequency of all age groups in the community. In line with the experience of communities

undergoing such distancing interventions, this decrease in overall contacts results in a substantially delayed epidemic, with 39,965 infections in the community as well as far lower burdens in terms of both hospitalizations and fatalities. In contrast, the early dynamics of the COVID-19 outbreak within the incarcerated population are identical, while in the latter half of the simulation, the outbreak dynamics in the incarcerated population are markedly worse, resulting in 8,339 infections after 180 days and proportionately more hospitalizations and COVID-19–related fatalities (Table 3). Shelter-in-place orders had no discernible estimated impact on the health outcomes of the staff of the jail.

All of the four considered arrest deferral scenarios had substantial estimated impacts on the course of the epidemic in the incarcerated population, while also lessening the impact of the epidemic on the community and, to a lesser extent, the jail’s staff. Discontinuing the arrest of bail-eligible individuals, which corresponds to a $\approx 25\%$ reduction of admissions into the jail, resulted in a 24% reduction in infections in the




FIGURE 3. Epidemic curves from a simulated COVID-19 epidemic in an urban community (right) and the connected population of persons in a jail (left). The curves demonstrate the expected magnitude and timing of the outbreak in the different populations, broken into the different etiologically relevant categories (i.e. exposed, infected, and hospitalized).

TABLE 3. Scenarios and Parameter Adjustments for a Number of Policy-based Interventions to Curtail COVID-19 in Jail and the Community

Scenario Name	Parameters	Multiplier of Baseline	Scenario Description
Shelter in place Arrest reduction	β_C^*	0.4	Effective contact rate in the community is reduced by a factor of 1/2.5
Bail eligible	$\alpha_L, \alpha_H, \alpha_E$	0.75	Divert all bail-eligible arrests (estimated at 25% of all arrests)
Vulnerable only	α_H, α_E	0.10	Divert arrests of 90% of vulnerable populations ¹¹
Low level	$\alpha_L, \alpha_H, \alpha_E$	0.17	Divert all low-level arrests (estimated at 83% of arrests)
Arrest fewer people Faster release	$\alpha_L, \alpha_H, \alpha_E$	0.10	Divert 90% of current arrests
Increase release speed	ρ^{-1}	2	2× rate of release from jail
Vulnerable only In-jail scenarios	ρ_H^{-1}	2	2× rate of release for vulnerable only
Mixing reduction	$\beta_I, \beta_P, \beta_J$	0.4	Reduction of baseline contact rates in jails by the same factor as the community under shelter-in-place
Reduced detection	ζ	0.99, 0.95, 0.90	Reduction in infection detection and timely hospitalization in jails by $1 - \zeta$

incarcerated population, and a 20% reduction in infections within the community. Note that, as discussed above, this model seeks to examine the role of jails in a population otherwise doing relatively well in controlling their outbreak. As such, these numbers arise in a population with a sustained, 6-month, broad-based shelter-in-place order, and as such, the cases arising from the jail population as a percentage of total cases were expected to be higher than they would be in a population with less stringent controls.

Broader, more sweeping arrest deferral programs resulted in correspondingly larger impacts in both the incarcerated population and the community as a whole. The discontinuation of arresting individuals for low level offenses ($\approx 83.4\%$ reduction) and the blanket reduction of arrests by 90% resulted in a 77% and 80% reduction in infections within the incarcerated population (with correspondingly fewer hospitalizations and deaths), respectively. These strategies also resulted in the greatest decrease in infections among staff (11% and 14%) and in the community at large (62% and 66%). Finally, a strategy built on deferring the arrest of individuals at high risk of developing COVID-19-related complications by 90%, tailoring the intervention to groups of epidemiologic importance rather than the nature of their offense, resulted in a 30% decrease in infections within the incarcerated population and a 61% decrease in deaths among the same population (Table 4).

In comparison, a strategy deferring the same number of people with no regard to their underlying risk, had smaller estimated effects on the number of infections and deaths. Specifically, releasing the same number of individuals as in the above scenario without regard to their risk resulted in only a 19% decrease in infections within the incarcerated population

TABLE 4. Cumulative Simulated COVID-19 Infections, Hospitalizations, and Deaths in the Community, Among Incarcerated Individuals, and Among Jail Staff During a 6-month Period With and Without a Shelter-in-place Social Distancing Intervention

	Without Social Distancing	With Social Distancing
Community		
Infections	926,108	450,621
Hospitalizations	51,497	22,892
Deaths	12,133	4,546
Incarcerated		
Infections	4,949	7,421
Hospitalizations	264	362
Deaths	79	106
Staff		
Infections	370	370
Hospitalizations	21	21
Deaths	2	2

and a 18% reduction in deaths. The deferral strategy targeting individuals for high-risk outcomes caused 0.9% more infections in the community relative to the same proportionately large but broader strategy, and the decreased number of deaths among persons in jail was partially offset by this increase. The targeted strategy resulted in a combined number of COVID-19 fatalities in both the population of persons in jail and in the community of 293 compared with 372 fatalities under the broader strategy.

Pairing increased arrest deferral with a more rapid release of persons who were already incarcerated enhanced the impact of those interventions, reducing infections, hospitalizations,

and deaths overall (Figure 4). The rate of decrease was less dramatic in the community and staff populations, especially at lower levels of accelerated release schedules.

When accompanied by the deferred arrest of bail-eligible individuals to reduce the incarcerated population, supplementary measures to reduce transmission among incarcerated persons had a marked benefit in both reducing the amplitude of the epidemic curve in incarcerated people and jail staff, as well as shifting the overall community epidemic curve later (Figure 5). These interventions may be thought of as either measures to reduce mixing—such as allowing greater space between individuals in common areas or the staggering of the use of shared facilities—or the provision of supplies such as soap and hand sanitizer that reduce the level of viral contamination of patient’s hands, physical surfaces, etc. Compared with the baseline mixing rate among persons in jail, a reduction to an equivalent level of mixing as the community while sheltering in place would reduce infections in this population by 37% as well as delay the peak of the epidemic by approximately 40 days.

An increase in the detection of severe COVID-19 cases among incarcerated persons from 95% to 100% (equivalent

to the same detection of the need for medical treatment available in the community) unsurprisingly increased the number of hospitalizations, as 5 of every 100 incarcerated persons needing hospitalization were no longer missed, either for lack of access to care, insufficient diagnostic capacity, or other reasons. Similarly, owing to the vast reduction in the case fatality rate (CFR) between hospitalized (5% for low risk and 33.3% for high risk) severe cases and unhospitalized severe cases (CFR = 100% for both groups), the number of deaths dropped by 91% when the detection of severe cases rose to the same level as the community. Between these scenarios, the number of infections rose slightly with better detection, increasing by 0.2% (Table 5). This is likely due to the slightly longer time an untreated severe case spends in the incarcerated population before they are removed due to death versus when a treated case is transferred for hospitalization. This effect will only be present if the level of viral shedding is constant (or increasing) over the course of a clinical infection. If instead the mechanism by which a severe COVID-19 patient dies is a cytokine storm or other process not involving the virus overwhelming the immune system, we would not expect this effect to be observed. However, even in the pessimistic case wherein viral




FIGURE 4. Cumulative infections, hospitalizations, and deaths in the community (first row), among persons in jail (second row) and among jail staff (third row) for several combinations of incarceration deferment and accelerated release.

Outcomes with Reducing Mixing

Bail Eligible Scenario with Mixing Reduction from Baseline to General Population Level Mixing




FIGURE 5. Epidemic curves for the community (top panel), persons in jail (middle panel), and jail staff (bottom panel) under a shelter in place order as well as the deferment of bail-eligible persons. The curves show the impact of increased reduction in mixing (e.g., from the ability to physically distance persons in jail while in common areas) from baseline (dark blue) to identical to the community’s shelter-in-place order (green).

shedding is constant throughout the clinical course of infection, the slight rise in infections is offset by the decrease in the number of COVID-19–related fatalities.

DISCUSSION

Using a computational simulation approach, our study estimated the impact of the COVID-19 epidemic among the incarcerated population, as well as among workers within the judicial system and the community within which these

facilities reside. Our models estimate that, in the absence of community mitigation such as strict social distancing, by only 30 days after the introduction of the first infection to the community, we can expect 2,628 infections among incarcerated people, resulting in two in-custody deaths. These results clearly follow from the features of the jail system themselves in challenging ways. While only 1% of the population entering into the jail system are elderly,⁴⁶ incarceration in jail is itself associated with degraded health of incarcerated people^{24–26}

TABLE 5. Cumulative Simulated COVID-19 Infections, Hospitalizations, and Deaths in the Community, Among Incarcerated Individuals, and Among Jail Staff During a 6-month Period Under Four Arrest Reduction Scenarios

	Community Social Distancing, No Arrest Reduction	Reduce Bail Eligible Arrests by 90%	Reduce Arrests in Groups Vulnerable to COVID-19 by 90%	Reduce Arrests for Low Level Offenses by 90%	Blanket 90% Arrest Reduction
Community					
Infections	450,621	433,262	429,257	362,055	349,306
Hospitalizations	22,892	21,604	21,320	16,855	16,079
Deaths	4,546	4,228	4,163	3,135	2,967
Incarcerated					
Infections	7,421	5,721	5,334	1,931	1,682
Hospitalizations	362	285	235	103	91
Deaths	106	84	45	31	27
Staff					
Infections	370	368	368	332	323
Hospitalizations	21	21	21	19	18
Deaths	2	2	2	2	2

leaving them more vulnerable to infection and severe outcomes from infection.⁴⁷ As individual robustness to disease decreases, the epidemiologic result is the increased vulnerability of the whole jail population.

Beyond the direct implications for the health of incarcerated people, jail populations have high rates of re-entry into the general community and they depend on people who regularly mix with the outside community. It is worth noting that, even as some court systems transitioned to videoconferencing and remote hearings, incarcerated populations do not have access to teleconferencing capabilities where they are housed, and must still be transported to, and staffed while using, communications facilities. Jail populations are largely composed of individuals who have not been convicted of a crime, and therefore will be released quickly back into the general community rather than to further incarceration within the carceral system. Jails with disease prevalence higher than the general populations they serve will therefore act as sources of infection, reseeding infection into communities that may be striving to contain or mitigate ongoing outbreaks, or even reintroducing infection into otherwise disease-free populations. Dynamics consistent with these predictions have already been observed^{48,49} but can and should be considered as an ongoing challenge. It is important to note that this would happen even if no one were released given the volume of people coming in and out of jails in staff and vendor roles, so should therefore not be construed as an implication that releases should be suspended or impeded. As COVID-19 continues to spread throughout the United States, tracking data within jails and the communities they serve will be critical in validating studies such as this one and in shaping best practices to limit jail-driven spread going forwards.

These estimated impacts are not, however, inevitable, and may be mitigated through a number of policy changes. Some obvious potential courses of action suggest themselves

immediately. New arrests mean that people of unknown disease status may be regularly brought into jails, increasing the likely severity of outbreaks both by the plausible continuous introduction of new sources of infection and by the maintenance of higher rates of contact among susceptible incarcerated people due to the density and structure of jail housing arrangements. If jurisdictions across the country reduce their intake by significant percentages, our models predict that we could meaningfully directly reduce the disease incidence in the incarcerated population (as seen in Table 4). Moreover, these same strategies also clearly produced a reduction in the source of risk to incarcerated people's families, jail staff, and the broader community (Table 4). These strategies could be enacted in a number of ways, such as (but not limited to) replacing misdemeanor arrests with citations, avoiding recommendations for jail time or prohibitive terms for bail conditions, or refusing to detain anyone for nonpayment of fines or fees during the course of the outbreak. Some of these reductions may be accomplished as a result of shelter-in-place orders themselves, though our model assumes no decrease in crime or arrests as a direct result of these policies.

Having considered these potential strategies for categorical reduction in intake into jails, we also considered the case in which the categorical consideration for reduction in intake stemmed instead from the health of the arrested person. Although this resulted in a smaller within-jail outbreak and reduced the resulting fatalities, we failed to achieve any noteworthy reduction in disease burden in the broader community by taking this action. It is therefore more effective to more aggressively reduce the intake rate across the entire population than to attempt to single out particular categories of individuals due to their likely susceptibility to severe morbidity or mortality from infection. The larger the reduction in overall intake, the greater the reduction in disease achieved for all populations (incarcerated people, the broader community, and

jail staff, in decreasing proportion of effect). These broader interventions are also likely to be relatively straightforward to implement administratively, without knowledge of an individual's underlying comorbidities, if any.

Our model's results predict that, in addition to reducing rates of intake into the jail system, increasing the rate of release from jails may help to mitigate the impacts of the epidemic. This increase however needed to be coupled with a decreased rate of intake rather than being enacted in isolation. Increasing release rates while maintaining the same rate of intake increased the predicted number of infections in the community and among jail staff. Rapid release in isolation does theoretically reduce the population size of the incarcerated population and the chance of transmission. However, the risk of transmission in scenarios with rapid releases but little change in jail intake remained high enough that, when coupled with rapid releases, incarcerated individuals are effectively returned to their community having been infected and with ample chance of transmitting to the community. In some ways, one can consider rapid release-only policies to be analogous to an accidental and well-intentioned form of patient dumping. This phenomenon may even still occur when expedited release is coupled with decreased rates of intake if the rate of release is insufficient (Figure 4). Again, our results clearly demonstrate that the greater the proportion of the incarcerated population we can include in such a policy, the more effective the intervention is at mitigating the outbreak.

Critically, the factors that cause these outbreak dynamics and drive the resulting efficacy of proposed interventions are features implicit in the nature of the jail system itself. The living conditions foster disease spread. Incarcerated people are shuttled back and forth to court or, where court proceedings are halted due to this pandemic, forced to remain in their cells or dorms. Incarcerated people occupy shared spaces in which physical distancing is impossible due to space, overcrowding, or the requirement of constant supervision. Incarcerated people are often not provided with the means to disinfect their surroundings or practice all of the hygiene guidelines suggested by the CDC. Building from what is known from COVID-19 transmission at this point in the epidemic, we believe that improved facility sanitation, access to free personal hygienic care, such as warm water, free soap, free hand sanitizer, and free cleaning products, increased time spent outside, increased physical/social distancing measures, increased access to free medical care, and improved nutrition are all factors that are likely to result in the improvement of individual health outcomes for people incarcerated within the jail system. Alterations to function and practice of the jail system that can correct for these challenges are unlikely to occur quickly enough or substantially enough to improve the epidemiologic risks for the incarcerated people within the jail system. As our results have shown, even when the within-jail transmission rates were improved by interventions such as reduction in intake from new arrests leading to a decrease in the size of the

incarcerated population, we could not effectively reduce the outbreak of infection in either the staff or incarcerated people down to the levels of the broader community.

As with all models, the conclusions of this study depend on an accurate representation of the flow of individuals between the jail system and the wider community, either due to arrests or due to their employment as jail staff, as well as the values of the parameters used to determine how swiftly this flow occurs. The inherent nature of emerging epidemics makes both of these things uncertain—the clinical and biological aspects of the pathogen might not be fully understood, and the data needed to parameterize these models are often sparse and incomplete. This problem is especially acute in models of this sort, which seek to present a “what-if” scenario to stave off a public health crisis, rather than analyze how that crisis unfolded after the fact. Nevertheless, while the exact projected magnitudes may be sensitive to these unknowns, in truth, the greatest utility of models such as these in determining best courses of action and likely magnitudes of the effects that can be gained from those actions, rather than exact predictions of precise numbers of individuals.⁵⁰ Due to the logical nature of the processes studied, so long as errors in the parameters used are consistent across scenarios, they will not impact the understanding that results from our projections about which courses of action achieve the best outcomes, even if those errors would alter our understanding of the precise amount of effect achieved by each intervention.

The focus of this model is on the early, emerging stages of COVID-19, and on attempting to prevent the large-scale outbreaks and super-spreading events seen in jails, prisons, and detention centers. Importantly, it is also focused on a period before the widespread availability of effective vaccines, which are currently and indisputably the most effective means of mitigating the pandemic. While the results of this model may be thought of as lessons learned and suggestions for an inevitable future pandemic, the combination of more transmissible variants of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and the relatively modest uptake of vaccines both in developing countries and in many geographic areas and subpopulations within the United States (including among corrections officers), many of the interventions here remain critical as we continue to experience the familiar “wave” patterns that characterize the COVID-19 pandemic.

TABLE 6. Infections, Hospitalizations, and Deaths Among Incarcerated Individuals Depending on the Probability that a Severe Case Within the Simulated Jail Is Detected and Appropriately Treated

Probability a Severe Case Is Detected and Treated	Infections	Hospitalizations	Deaths
0.95	7,361	316	383
0.99	7,409	353	165
0.00	7,421	362	106

REFERENCES

- Mervosh S, Lu D, Swales V. See Which States and Cities Have Told Residents to Stay at Home. *New York Times*. April 2020. Available at: <https://www.nytimes.com/interactive/2020/us/coronavirus-stay-at-home-order.html>.
- Bayham J, Fenichel EP. Impact of school closures for COVID-19 on the US health-care workforce and net mortality: a modelling study. *Lancet Public Health*. 2020;5:e271–e278.
- Parmet WE, Sinha MS. COVID-19—the law and limits of quarantine. *New Engl J Med*. 2020;382:e28.
- Bernstein H, González J, Gonzalez D, Jagannath J. *Immigrant-Serving Organizations Perspectives on the COVID-19 Crisis*. Urban Institute; 2020.
- Sharp R. *The Incarceration Nation: Interpreting the United States Imprisonment Rate*. The University of Akron; 2018.
- Sykes BL, Pettit B. Measuring the exposure of parents and children to incarceration. In: Mark Eddy J, Poehlmann-Tynan J, eds. *Handbook on Children with Incarcerated Parents*. Springer; 2019:11–23.
- Weidner RR, Schultz J. Examining the relationship between US incarceration rates and population health at the county level. *SSM-Popul Health*. 2019;9:100466.
- Kelley S. Mass incarceration. *Hum Ecol*. 2019;47:15.
- Enns PK, Yi Y, Comfort M, et al. What percentage of Americans have ever had a family member incarcerated?: evidence from the family history of incarceration survey (FamHIS). *Socius*. 2019;5:2378023119829332.
- Lester HD, Miller MJ. Discrete event simulation of jail operations in pursuit of organizational culture change. In: *Computer Security*. Springer; 2019:307–322.
- Minton TD. Jail Inmates at Midyear 2010 - Statistical Tables. Bureau of Justice Statistics; 2011.
- Raher S. The Company Store: A Deeper Look at Prison Commissaries. 2018. Available at: <https://www.prisonpolicy.org/reports/commissary.html>. Accessed 6 April 2020.
- Bick JA. Infection control in jails and prisons. *Clin Infect Dis*. 2007;45:1047–1055.
- Hoge CW, Reichler MR, Dominguez EA, et al. An epidemic of pneumococcal disease in an overcrowded, inadequately ventilated jail. *N Engl J Med*. 1994;331:643–648.
- Rubin R. The challenge of preventing COVID-19 spread in correctional facilities. *JAMA*. 2020;323:1760–1761.
- Reinhart E, Chen D. Incarceration and its disseminations: COVID-19 Pandemic Lessons From Chicago's Cook County Jail: Study examines how arrest and pre-trial detention practices may be contributing to the spread of COVID-19. *Health Aff*. 2020;39:1412–1418.
- Akiyama MJ, Spaulding AC, Rich JD. Flattening the curve for incarcerated populations—COVID-19 in jails and prisons. *New Engl J Med*. 2020;382:2075–2077.
- Hawks L, Woolhandler S, McCormick D. COVID-19 in Prisons and Jails in the United States. *JAMA Intern Med*. 2020;180:1041–1042.
- Thigpen M, Beauclair T, Hutchinson V, Persons V, Paul C. *Jail Design Guide*. National Institute of Corrections; 2011.
- Centers for Disease Control and Prevention. Interim guidance on management of coronavirus disease 2019 (COVID-19) in correctional and detention facilities. Centers for Disease Control and Prevention, 2020.
- Nowotny KM. Health care needs and service use among male prison inmates in the United States: a multi-level behavioral model of prison health service utilization. *Health Justice*. 2017;5:9.
- Mignon S. Health issues of incarcerated women in the United States. *Cien Saude Colet*. 2016;21:2051–2060.
- Barnert ES, Perry R, Morris RE. Juvenile incarceration and health. *Acad Pediatr*. 2016;16:99–109.
- McClelland DC, Alexander C, Marks E. The need for power, stress, immune function, and illness among male prisoners. *J Abnorm Psychol*. 1982;91:61–70.
- Jacobs ET, Mullany CJ. Vitamin D deficiency and inadequacy in a correctional population. *Nutrition*. 2015;31:659–663.
- Kouyoumdjian FG, Andreev EM, Borschmann R, Kinner SA, McConnon A. Do people who experience incarceration age more quickly? Exploratory analyses using retrospective cohort data on mortality from Ontario, Canada. *PLoS One*. 2017;12:e0175837.
- Soetaert K, Petzoldt T, Setzer RW. Solving differential equations in R: package deSolve. *J Stat Softw*. 2010;33:1–25.
- Nnorom-Dike O, Ekwebelem O, Ofielu E, Attah M, Ekwe D. Long term immunologic consequences of illicit drug abuse. *J Alcohol Drug Abuse Subst Depend*. 2020;12. Available at: <https://www.heraldopenaccess.us/openaccess/long-term-immunologic-consequences-of-illicit-drug-abuse>.
- Wagner P, Sakala L. Mass Incarceration: The Whole Pie. Prison Policy Initiative. 2014;12.
- Miller IF, Becker AD, Grenfell BT, Metcalf CJE. Disease and healthcare burden of COVID-19 in the United States. *Nat Med*. 2020;26:1212–1217.
- Davies NG, Klepac P, Liu Y, et al. Age-dependent effects in the transmission and control of COVID-19 epidemics. *MedRxiv*. Preprint posted online March 27, 2020. doi: 10.1101/2020.03.24.20043018.
- Funk S. socialmixr: Social Mixing Matrices for Infectious Disease Modelling. R package version 0.1.8.2020. Available at: <https://CRAN.R-project.org/package=socialmixr>.
- Wu J, Dhingra R, Gambhir M, Remais JV. Sensitivity analysis of infectious disease models: methods, advances and their application. *J R Soc Interface*. 2013;10:20121018.
- Lurette A, Touzeau S, Lamboni M, Monod H. Sensitivity analysis to identify key parameters influencing Salmonella infection dynamics in a pig batch. *J Theor Biol*. 2009;258:43–52.
- Ferguson N, Laydon D, Nedjati Gilani G, et al. Report 9: Impact of non-pharmaceutical interventions (NPIs) to reduce COVID19 mortality and healthcare demand. Imperial College; 2020.
- Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*. 2020;395:1054–1062.
- Lauer SA, Grantz KH, Bi Q, et al. The incubation period of Coronavirus Disease 2019 (COVID-19) from publicly reported confirmed cases: estimation and application. *Ann Int Med*. 2020;172:577–582.
- Johndrow J, Ball P, Gargiulo M, Lum K. Estimating the Number of SARS-CoV-2 Infections and the Impact of Mitigation Policies in the United States. *Harvard Data Science Review*. 2020;11. Available at: <https://hdr.mitpress.mit.edu/pub/9421kmzi>.
- Hu Z, Song C, Xu C, et al. Clinical characteristics of 24 asymptomatic infections with COVID-19 screened among close contacts in Nanjing, China. *Sci China Life Sci*. 2020;63:706–711.
- Beckett SJ, Dominguez-Mirazo M, Lee S, Andris C, Weitz JS. Spread of COVID-19 through Georgia, USA. Near-term projections and impacts of social distancing via a metapopulation model. 2020. Available at: https://github.com/WeitzGroup/MAGEmodel_covid19_GA/blob/master/Report/GA_COVID19_assessment_21Apr2020.pdf. Accessed 6 April 2020.
- Verity R, Okell LC, Dorigatti I, et al. Estimates of the severity of COVID-19 disease. *MedRxiv*. Preprint posted online March 13, 2020. doi: 10.1101/2020.03.09.20033357
- Tindale L, Coombe M, Stockdale JE, et al. Transmission interval estimates suggest pre-symptomatic spread of COVID-19. *MedRxiv*. Preprint posted online March 06, 2020. doi: 10.7554/eLife.57149
- COVID C. Severe Outcomes Among Patients with Coronavirus Disease 2019 (COVID-19)—United States, February 12–March 16, 2020. *CDC*. 2020.
- Maruschak LM, Berzofsky M, Unangst J. Medical problems of state and federal prisoners and jail inmates, 2011–12. US Department of Justice, Office of Justice Programs, Bureau of Justice Statistics; 2015.
- Pennsylvania ACLU. Punishing Poverty: Cash Bail In Allegheny County. 2019. Available at: https://aclupa.org/sites/default/files/field_documents/allegheny_county_report_final.pdf. Accessed 6 April 2020.
- Allegheny County Department of Human Services. Allegheny County Jail Population Management: Interactive Dashboards. 2020. Available at: <https://www.alleghenycountyanalytics.us/index.php/2019/11/04/allegheny-county-jail-population-management-dashboards-2/>. Accessed 6 April 2020.
- Binswanger IA, Blatchford PJ, Forsyth SJ, Stern MF, Kinner SA. Epidemiology of infectious disease-related death after release from prison, Washington State, United States, and Queensland, Australia: a cohort study. *Public Health Rep*. 2016;131:574–582.
- Williams T, Seline L, Griesbach R. Coronavirus cases rise sharply in prisons even as they plateau nationwide. *New York Times*. June 2020;18.
- Ransom J. Virus Raged at City Jails, leaving 1,259 guards infected and 6 dead. *New York Times*. May 2020;20.
- Lofgren ET, Halloran ME, Rivers CM, et al. Opinion: mathematical models: a key tool for outbreak response. *Proc Natl Acad Sci USA*. 2014;111:18095–18096.