

ERRATA

Combined Microautoradiography–16S rRNA Probe Technique for Determination of Radioisotope Uptake by Specific Microbial Cell Types In Situ

CLEBER C. OUVENEY AND JED A. FUHRMAN

Department of Biological Sciences, University of Southern California, Los Angeles, California 90089-0371

Volume 65, no. 4, p. 1746–1752, 1999. Page 1748, legend to Fig. 1: lines 4 and 5 should read as follows: “The arrow in panel E indicates a cluster of silver grains with no corresponding cell in either the DAPI-stained (D) or the probe (F) field. Such clusters of silver grains with no cells represented less than 2% of the ³H-labeled cells and were thought to be sites where cells were lost during the procedure. Bar = 5 μm.”

Microbial Desulfurization of a Crude Oil Middle-Distillate Fraction: Analysis of the Extent of Sulfur Removal and the Effect of Removal on Remaining Sulfur

M. J. GROSSMAN, M. K. LEE, R. C. PRINCE, K. K. GARRETT,
G. N. GEORGE, AND I. J. PICKERING

*Exxon Research and Engineering Co., Annandale, New Jersey 08801, and Stanford
Synchrotron Radiation Laboratory, Stanford, California 94309*

Volume 65, no. 1, p. 181–188, 1999. Page 183: Table 1 should read as shown below.

TABLE 1. Sulfur removal from OB oil by ECRD-1

Culture	% Sulfur (± SD)		% Sulfur removed
	Sterile control	Biodesulfurized culture	
Undiluted OB oil	2.17 (0.01)	1.99 (0.0002)	8.1
OB oil diluted 10-fold in decane (20 ppm sulfur)	2.05 (0.005)	1.44 (0.004)	29.8
OB oil diluted 20-fold in decane (10 ppm sulfur)	2.05 (0.005)	1.45 (0.008)	29.3