



Published in final edited form as:

*Biochem Biophys Res Commun.* 2022 June 30; 611: 146–150. doi:10.1016/j.bbrc.2022.04.069.

## Inactivation of axon guidance molecule netrin-1 in human colorectal cancer by an epigenetic mechanism

Hironao Nakayama<sup>a,\*†</sup>, Hidetaka Ohnuki<sup>b,\*</sup>, Masako Nakahara<sup>a</sup>, Hisayo Nishida-Fukuda<sup>c</sup>, Tomohisa Sakaue<sup>d,e</sup>, Shinji Fukuda<sup>d,f</sup>, Shigeki Higashiyama<sup>d,g</sup>, Yuki Doi<sup>h</sup>, Masahiro Mitsuyoshi<sup>h,i</sup>, Takashi Okimoto<sup>h,j</sup>, Giovanna Tosato<sup>b</sup>, Chiaki Kusumoto<sup>a,k,†</sup>

<sup>a</sup>Department of Medical Science and Technology, Hiroshima International University, Higashi-hiroshima, Hiroshima 739-2695, Japan

<sup>b</sup>Laboratory of Cellular Oncology, Center for Cancer Research, National Cancer Institute, National Institutes of Health, Bethesda, MD 20892, USA

<sup>c</sup>Department of Genome Editing, Institute of Biomedical Science, Kansai Medical University, 2-5-1 Shin-machi, Hirakata, Osaka, 5731010, Japan

<sup>d</sup>Division of Cell Growth and Tumor Regulation, Proteo-Science Center, Ehime University, Toon, Shitsukawa, Ehime 791-0295, Japan

<sup>e</sup>Department of Cardiovascular and Thoracic Surgery, Ehime University Graduate School of Medicine, Toon, Shitsukawa, Ehime 791-0295, Japan

<sup>f</sup>Department of Biochemistry, School of Dentistry, Aichi Gakuin University, Nagoya, Aichi 464-8650, Japan

<sup>g</sup>Department of Molecular and Cellular Biology, Osaka International Cancer Institute, Chuo-ku, Osaka 541-8567, Japan

<sup>h</sup>Department of Surgery, Nippon Kokan Fukuyama Hospital, Fukuyama, Hiroshima 721-0927, Japan

<sup>i</sup>Department of Surgery I, School of Medicine, University of Occupational and Environmental Health, Kita-kyushu, Fukuoka 807-0804, Japan

<sup>j</sup>Department of Surgery, Kyushu Rosai Hospital, Kita-kyushu, Fukuoka 800-0296, Japan

<sup>k</sup>Department of Gastroenterology, Nippon Kokan Fukuyama Hospital, Fukuyama, Hiroshima 721-0927, Japan

<sup>†</sup>**Corresponding authors:** To whom correspondence should be addressed: Hironao Nakayama, Department of Medical Science and Technology, Hiroshima International University, Higashi-hiroshima, Hiroshima 739-2695, Japan, hironao@hirokoku-u.ac.jp; Tel.: +81 (823) 70-4640 Or Chiaki Kusumoto, Department of Medical Science and Technology, Hiroshima International University, Higashi-hiroshima, Hiroshima 739-2695, Japan, kusumoto@hirokoku-u.ac.jp; Tel.: +81 (823) 70-4594.

\*These authors contributed equally to this work.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conflict of Interest

The authors declare that there are no conflicts of interest.

## Abstract

Netrin-1, the protein product of the *NTN1* gene, is an axon guidance molecule implicated in regulation of cell survival and tumorigenesis. Expression of the netrin-1 receptors deleted in colorectal cancer (*DCC*) and uncoordinated 5 homolog (*UNC5H*) is frequently silenced in colorectal cancer (CRC) by either loss of heterozygosity or epigenetic mechanisms. However, netrin-1 expression and regulation in CRC are mostly unknown. Here, we report that *NTN1* expression is significantly reduced in most CRC tissues compared to the adjacent normal intestinal mucosa, and that *NTN1* DNA methylation is significantly higher in CRCs (24.6%) than in the adjacent normal intestinal mucosa (4.0%). In 6 CRC cell lines, *NTN1* expression is low. Treatment with 5-Aza-2'-deoxycytidine increased expression of *NTN1* in CRC cell lines, indicating that DNA methylation represses *NTN1* transcription in CRCs. *NTN1* DNA hypermethylation was significantly associated with advanced CRC disease. Median netrin-1 serum levels were significantly decreased in CRC patients (330.1 pg/mL) compared with normal individuals (438.6 pg/mL). Our results suggest that netrin-1 is a candidate biomarker for CRC.

## Keywords

netrin-1; axon guidance; colorectal cancer; DNA methylation; VEGF-A

## 1. Introduction

Colorectal cancer (CRC) is the second most deadly and third most commonly diagnosed cancer in the world, according to GLOBOCAN 2020 [1]. Currently, early diagnosis of CRC is achieved by screening colonoscopy, which is extremely effective, but invasive. It would be useful to detect CRC with a non-invasive procedure. Like other epithelial cancers, CRC is believed to arise through the successive accumulation of genetic and epigenetic alterations in oncogenes, tumor suppressor genes, and DNA repair pathway genes [2,3]. DNA hypermethylation of promoter-associated CpG islands is a frequent and early mechanism of gene inactivation in CRC and other cancers, but no DNA methylation marker is a clinically applicable biomarker of CRC [4,5].

Netrins are laminin-like proteins, first identified as axonal guidance molecules in *Caenorhabditis elegans* [6]. The netrin family consists of four secreted molecules, netrin-1, netrin-3, netrin-4, and netrin-5, and two membrane-anchored members, netrin-G1 and netrin-G2. Netrin-1, secreted from floor plate during brain development, establishes a gradient that attracts to the midline the commissural axons that express netrin receptors [6,7]. Several receptors mediate the function of netrin family proteins, including uncoordinated-5-Homologs (*UNC5H1-4/A-D*), deleted in colorectal cancer (*DCC*), the *DCC* orthologue neogenin (*NEO1*), and Down syndrome cell adhesion molecule (*DSCAM*). These receptors are “dependence receptors” as they promote cell proliferation, migration, and survival when activated by the ligand, but trigger apoptosis when the ligand is limiting [8]. In the gut, netrin-1 is predominantly expressed at the bottom of the intestinal crypt, whereas *DCC* is expressed all along the villus. Within the crypt, netrin-1 binding to *DCC* induces crypt cell survival. At the villus tips, where netrin-1 levels are low, *DCC* promotes apoptotic cell death [3,9]. *DCC* was identified as a candidate tumor suppressor in CRC

with a deletion in chromosome 18q21 [10,11], and loss of netrin-1 receptor expression was associated with a poor prognosis in patients with CRC [3,9]. Also, expression of *UNC5H* genes is silenced in CRC, either by loss of heterozygosity or epigenetic mechanisms [12].

Netrin-1 mRNA and protein levels are increased in metastatic breast cancer, non-small cell lung cancer, and pancreatic adenocarcinoma, where netrin-1 functions as an oncogene and may be associated with loss of netrin-1 receptors [13–16]. Recently, we have shown that netrin-1 promotes invasiveness and angiogenesis in glioblastoma and medulloblastoma by activating the cysteine protease cathepsin B [17–19]. However, the function of netrin-1 in CRC is not clear. In this report, we demonstrate that netrin-1 mRNA and protein levels are significantly reduced in most CRC tissues compared to the adjacent normal intestinal mucosa, and that *NTN1* methylation is significantly greater in CRC tissues compared to the adjacent normal mucosa. The demethylating drug 5-Aza-2'-deoxycytidine restores *NTN1* expression in CRC cell lines, suggesting that *NTN1* expression is silenced by DNA methylation in CRC. We also find that serum levels of netrin-1 are significantly decreased in patients with CRC compared with normal individuals and are predictors of disease severity. Together, our results suggest that netrin-1 reduction is a candidate biomarker of CRC.

## 2. Materials and Methods

### 2.1. CRC samples

Tissue samples of CRC and matched normal intestinal mucosa were collected during a colonoscopy from 25 patients at Nippon Kokan Fukuyama Hospital (Fukuyama, Japan), prior to initiation of cancer therapy, and were stored at  $-80^{\circ}\text{C}$  until analyzed. Peripheral blood samples, collected from the CRC patients in endotoxin-free silicone-coated tubes (Sekisui Medical Co., Tokyo, Japan) a few days prior to surgery, were allowed to clot at room temperature for 30 min before centrifugation (3,000 rpm, 10 min); serum was removed and stored at  $-80^{\circ}\text{C}$  until analyzed. To establish a normal reference range, serum netrin-1 concentrations were measured in 30 healthy volunteer blood donors. This study was approved by the Institutional Review Board of Hiroshima International University (approved IRB protocol number: 18–015) and Nippon Kokan Fukuyama Hospital (approved IRB protocol number: 2018–01).

### 2.2. NTN1 expression analysis with TCGA datasets

Paired RNA-seq datasets of rectum adenocarcinoma (TCGA-READ)/colon adenocarcinoma (TCGA-COAD) and adjacent normal tissue were downloaded from The Cancer Genome Atlas (TCGA) using the data transfer software, GDC-client (version 1.5.0). Paired samples of solid tumor and normal tissue from 9 patients of TCGA-READ and 46 patients of TCGA-COAD are listed in Supplementary Table 4. Datasets were unzipped using the Linux command “*gzip -cd*” on a high-performance computer, NIH Biowulf. *NTN1* expression levels (Fragments Per Kilobase pair per Million reads, FPKM) were extracted from each dataset using the Linux command “*grep -w NTN1*”. Extracted data files were combined by the Linux command “*cat*”. The results were visualized using Microsoft EXCEL.

### 2.3. DNA methylation analyses with TCGA datasets

Datasets of DNA methylation arrays (Illumina Human Methylation array 450k or 27k) from paired rectum/colon solid tumor and paired normal tissue were downloaded from TCGA-READ and TCGA-COAD using the GDC-client through NIH Biowulf. Individual datasets are listed in Supplementary Table 5. Datasets were unzipped using “*gzip -cd*”, and DNA methylation status in a CpG island near the promoter of *NTN1* (probe cg19564877) was extracted using the Linux command “*grep -w cg19564877*”. The extracted data files were combined using the Linux command “*cat*”. The results were visualized using Microsoft EXCEL.

For correlation analysis between *NTN1* DNA methylation and TNM stage (Fig. 2D), datasets of DNA methylation arrays from normal tissues of 51 patients and tumors of 617 patients were downloaded from TCGA-READ and TCGA-COAD (see Supplementary Table 6). Extracted data files from each patient were combined as a CSV file on NIH Biowulf. The combined CSV file was opened by Microsoft EXCEL, and data were summarized based on TNM stage.

### 2.4. Correlation analysis of DNA methylation and mRNA expression of NTN1

Datasets of RNA-seq and DNA methylation arrays from the same patients were downloaded from TCGA-READ and TCGA-COAD using the GDC-client. The used datasets are listed in Supplementary Table 7. Datasets were unzipped using the Linux command “*gzip -cd*”. *NTN1* gene expression and the “sample\_submitter\_id” were extracted from RNA-seq datasets as described in the previous method section. DNA methylation levels and the “sample\_submitter\_id” were extracted from datasets of DNA methylation arrays described in the previous method section. Extracted data files of *NTN1* gene expression and DNA methylation of *NTN1* CpG island proximal to the promoter were combined into a CSV file using the Linux command “*cat*”. A table is generated, which has 3 columns; 1st column is “sample\_submitter\_id”, 2nd column is *NTN1* FPKM values, and 3rd column is DNA methylation level of *NTN1* CpG island proximal to the promoter. The data were visualized as a dot plot shown in Fig. 2C.

### 2.5. Bisulfite modification and sequence of NTN1 promoter region

Bisulfite modification of genomic DNA from cultured cells was performed by EpiTect Bisulfite Kits (Qiagen, Chatsworth, CA). We first identified tentative CpG sites in the *NTN1* promoter region (chr17: 9,020,851–9,021,600). There were 68 CpG sites within the promoter region (nucleotides –1523 to –774) analyzed in this study. The nucleotide number +1 is defined as the start codon of *NTN1*. Bisulfite-modified DNA was amplified using the primer set listed in Supplementary Table 1. Each of the PCR products was subsequently cloned using the pT7Blue T-vector (Sigma-Aldrich). To quantify the percentage of methylated CpG sites, 8 clones were sequenced and identified the bisulfite converted DNA.

### 2.6. Statistical analysis

GraphPad Prism 6 statistical software (GraphPad Software; San Diego, CA) was used for statistical analyses. The results are represented as medians and interquartile ranges were used to summarize the tumor patient data, and controls were compared using the

nonparametric Mann-Whitney *U* test. Diagnostic accuracy was assessed with receiver operating characteristic (ROC) curve analysis, and the Youden Index was used to identify cut-off values.  $P < 0.05$  was considered statistically significant.

### 3. Results and Discussion

We first evaluated netrin-1 expression in CRC. Levels of *NTN1* mRNA were derived from TCGA data sets of paired samples of CRC tissues and normal adjacent intestinal mucosa ( $n = 55$ ). *NTN1* is expressed at significantly ( $P = 3.67 \times 10^{-13}$ ) lower levels in CRC tissue samples compared to the adjacent normal intestinal mucosa (Fig. 1A and B). We confirmed this observation in additional 25 paired samples of CRC and normal adjacent intestinal mucosa ( $P < 0.05$ , Fig. 1C), by using qRT-PCR (Supplementary Table 1 and 2).

Using these 25 paired samples of CRC and adjacent normal intestinal mucosa, we examined expression of netrin-1 receptors. Netrin-1 has multiple receptors [6]. By qRT-PCR, netrin-1 receptors *DCC*, *NEO1*, *UNC5B*, *UNC5C* and *UNC5D* were expressed in the normal mucosa, whereas *DSCAM* and *UNC5A* were not expressed. *NEO1* and *UNC5B* were expressed at the highest levels in the normal mucosa. Expression of all 5 receptors was significantly decreased in CRCs compared with the normal mucosa (Fig. 1C). Angiogenesis contributes to tumor progression and metastasis. Vascular endothelial growth factor (VEGF) is a potent angiogenic factor. Circulating VEGF-A levels are abnormally increased in patients with CRC and anti-VEGF therapy is effective at reducing tumor growth in patients with CRC [20–22]. Unlike netrin-1 and its receptors, expression of *VEGFA* was elevated in CRC samples compared to the normal adjacent intestinal mucosa ( $P < 0.05$ , Fig. 1D), consistent with previous reports [23,24]. Western blot analysis in 3 randomly selected CRC samples showed that protein levels of netrin-1 and *UNC5C* are reduced in CRCs compared with the corresponding normal colonic mucosa (Fig. 1E). Together, these results show that netrin-1 mRNA and protein levels are reduced in CRC tissues compared to the normal adjacent mucosa.

Methylation-induced gene silencing frequently occurs in cancer [12,25,26]. We examined the status of *NTN1* methylation in CRC samples and paired samples from the normal adjacent intestinal mucosa ( $n=92$ ) using available TCGA datasets derived from the Human Methylation 450K or 27K Array (Illumina). This comparison, focused on the level of DNA methylation within the CpG islands at the 5' untranslated region of *NTN1* (position –692 to +2,061 relative to the transcription start site), revealed that the region was hypermethylated in tumor samples compared with the paired normal intestinal samples (mean  $\beta$  values: 24.7% in tumors and 4.01% in normal tissue,  $P = 7.98 \times 10^{-13}$ , Fig. 2A and B). The mean percentage of CpG methylation of *NTN1* was inversely correlated with *NTN1* expression level (Pearson's  $r = -0.30$ ), suggesting that DNA methylation represses *NTN1* transcription in CRCs (Fig. 2C). In addition, hypermethylation of *NTN1* was frequently observed in patients with TNM stages II-IV (Fig. 2D), indicating that DNA hypermethylation of *NTN1* was significantly associated with advanced CRC.

By qRT-PCR, we examined *NTN1* expression in normal colonic epithelial cells (ECs) and 6 CRC cell lines. Consistent with the result of CRC patient samples (Fig. 1), all CRC cell lines

showed significantly ( $P < 0.01$ ) reduced *NTN1* levels compared to normal colon ECs (Fig. 3A). Since methylation-induced gene silencing is frequently observed in human cancers, we examined the DNA methylation status in normal colon ECs and HT29 cells. By bisulfite sequencing, the *NTN1* promoter region (nucleotides -1523 to -1250) was hypermethylated in 64.1% (15 of 23) CpG sites, compared to 5.4% (1 of 23) CpG sites in normal colon ECs (Fig. 3B and Supplementary Fig. 1). We cultured CRC cell lines and normal colon ECs with decitabine (5-Aza-2'-deoxycytidine, DAC). Treatment with DAC resulted in a significant increase of *NTN1* in 5 CRC cell lines. However, DAC had a minimal effect on *NTN1* expression in normal colonic ECs and in SW480 cells (Fig. 3C). Western blot analysis showed that DAC treatment increased netrin-1 protein levels in the HT29 and HCT116 cell lines (Fig. 3D).

Netrin-1 receptors were also measured (Supplementary Fig. 2A). Five netrin-1 receptors were detected in CRC cell lines, whereas *UNC5D* and *DSCAM* were not detected. *DCC* and *UNC5A-C* were decreased in CRC cell lines compared to the normal colonic ECs (Supplementary Fig. 2A). Previous studies reported that netrin-1 receptors *DCC* and *UNC5A-C* are down-regulated in multiple cancers, including CRC, and suggested that loss of netrin-1 receptors is caused by loss of heterozygosity or epigenetic processes [12,27]. Consistent with these previous results, DAC treatment restored mRNA expression in *DCC* and *UNC5A-C* receptors in the CRC cell lines (Supplementary Fig. 2B). Altogether, these results suggest that DNA hypermethylation regulates the transcriptional repression of netrin-1 ligand and receptors in human CRC.

At present, it is not clear whether netrin-1 expression plays a role in colorectal cancer development [28,29]. Several lines of evidence supported a crucial role of netrin-1 during colorectal tumorigenesis by regulating apoptosis [28]. Overexpression in the mouse gut of netrin-1, under the control of the gut-specific liver fatty acid-binding protein gene (*Fabpl*) promoter, inhibited intestinal epithelial cell death and promoted formation of hyperplastic and neoplastic lesions [28]. Additionally, inhibition of netrin-1-mediated apoptosis was associated with tumor progression in mice carrying the adenomatous polyposis coli (*APC*) mutation [28]. Inflammatory bowel disease in a mouse model was associated with increased netrin-1 expression in the colonic epithelial cells via the NF- $\kappa$ B pathway and with colorectal cancer development [29]. These apparently inconsistent results might be attributable to forced expression of netrin-1 in the mouse model, and on netrin-1 affecting various cell types, including endothelial cells and immune cells, all contributing to tumor development [6,7]. Further work is needed to dissect the mechanisms underlying the pro-tumorigenic effects of ectopic netrin-1 expression mice. Another potential explanation for these discrepant results is that reduction of netrin-1 expression may inhibit the survival of highly proliferative cells at the base of the intestinal crypt because netrin-1/DCC interactions are essential for tight regulation of epithelial cells proliferation and differentiation in the intestine. Thus, we can speculate that the loss of the proximal-to-distal netrin-1 gradient in a netrin-1-deprived environment impairs intestinal cellular homeostasis, eventually initiating malignant transformation.

Recent studies show that serum netrin-1 levels may be clinically useful as a biomarker for human cancers [30]. We measured serum levels of netrin-1 by ELISA in patients with CRC

and healthy blood donors. The baseline demographic, clinical, and CRC characteristics of the participants are presented in Supplementary Table 2. Accuracy of the netrin-1 ELISA was determined by spiking and recovery testing (Supplementary Table 3). The median serum netrin-1 concentration was 438.6 pg/mL (IQR, 332.1–666.1) for control and 330.1 pg/mL (IQR, 181.9–472.3) for patients with CRC ( $P = 0.0028$ ), indicating that serum netrin-1 levels are significantly decreased in CRCs (Fig. 4A). A receiver operating characteristic (ROC) curve analysis indicated that serum netrin-1 provided moderate diagnostic accuracy in differentiating tumor from control groups [area under the curve (AUC): 0.710,  $P = 0.0065$ ]. The Youden Index revealed that the optimal cut-off value is  $< 293.8$  pg/mL (Fig. 4B). We also measured serum VEGF-A levels which served as an experimental positive control [24]. Serum VEGF-A levels were elevated in patients with CRC (median 271.5 pg/mL, IQR, 161.5–474.3) compared to controls (median 178.1 pg/mL, IQR, 97.3–347.4,  $P = 0.010$ ) (Fig. 4C).

In summary, we show here that CRC patients samples exhibit reduced expression of *NTN1* attributable to DNA methylation. Our data further indicate that detection of *NTN1* DNA hypermethylation and serum levels of netrin-1 may provide biomarkers for predicting CRC progression. Further studies are needed to define the function of netrin-1 inactivation on tumor cell proliferation, invasiveness, and angiogenesis in CRC. However, these data provide a basis for investigating netrin-1 as a potential new target for treatment of CRC.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

Research reported in this publication was supported in part by Japan Society for the Promotion of Science (JSPS) KAKENHI Grant Numbers 18K08984 and 21K09192 (H.N.), and the Chugoku Occupational Health Association (C.K.). H.O. and G.T. are supported by the Intramural Program of the Center for Cancer Research, National Cancer Institute, National Institutes of Health. This study was supported by Ehime University Proteo-Science Center (PROS) and Advanced Research Support Center. This work used the computational resources of the NIH High-Performance Computing (HPC) Biowulf cluster (<http://hpc.nih.gov>).

## Abbreviations:

<b>DCC</b>	deleted in colorectal cancer
<b>UNC5H</b>	uncoordinated 5 homolog
<b>CRC</b>	colorectal cancer
<b>DSCAM</b>	down syndrome cell adhesion molecule
<b>VEGF</b>	vascular endothelial growth factor
<b>EC</b>	epithelial cell
<b>DAC</b>	5-Aza-2'-deoxycytidine
<b>APC</b>	adenomatous polyposis coli

## References


- [1]. Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, Bray F, Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries, *CA Cancer J Clin* 71 (2021) 209–249. 10.3322/caac.21660. [PubMed: 33538338]
- [2]. Hanahan D, Weinberg RA, The hallmarks of cancer, *Cell* 100 (2000) 57–70. 10.1016/s0092-8674(00)81683-9. [PubMed: 10647931]
- [3]. Mehlen P, Fearon ER, Role of the dependence receptor DCC in colorectal cancer pathogenesis, *J Clin Oncol* 22 (2004) 3420–3428. 10.1200/jco.2004.02.019. [PubMed: 15310786]
- [4]. Draht MXG, Goudkade D, Koch A, Grabsch HI, Weijnenberg MP, van Engeland M, Melotte V, Smits KM, Prognostic DNA methylation markers for sporadic colorectal cancer: a systematic review, *Clin Epigenetics* 10 (2018) 35. 10.1186/s13148-018-0461-8. [PubMed: 29564023]
- [5]. Yi JM, Dhir M, Van Neste L, Downing SR, Jeschke J, Glöckner SC, de Freitas Calmon M, Hooker CM, Funes JM, Boshoff C, Smits KM, van Engeland M, Weijnenberg MP, Iacobuzio-Donahue CA, Herman JG, Schuebel KE, Baylin SB, Ahuja N, Genomic and epigenomic integration identifies a prognostic signature in colon cancer, *Clin Cancer Res* 17 (2011) 1535–1545. 10.1158/1078-0432.Ccr-10-2509. [PubMed: 21278247]
- [6]. Lai Wing Sun K, Correia JP, Kennedy TE, Netrins: versatile extracellular cues with diverse functions, *Development* 138 (2011) 2153–2169. 10.1242/dev.044529. [PubMed: 21558366]
- [7]. Mehlen P, Delloye-Bourgeois C, Chédotal A, Novel roles for Slits and netrins: axon guidance cues as anticancer targets?, *Nat Rev Cancer* 11 (2011) 188–197. 10.1038/nrc3005. [PubMed: 21326323]
- [8]. Llambi F, Causeret F, Bloch-Gallego E, Mehlen P, Netrin-1 acts as a survival factor via its receptors UNC5H and DCC, *Embo j* 20 (2001) 2715–2722. 10.1093/emboj/20.11.2715. [PubMed: 11387206]
- [9]. Mehlen P, Llambi F, Role of netrin-1 and netrin-1 dependence receptors in colorectal cancers, *Br J Cancer* 93 (2005) 1–6. 10.1038/sj.bjc.6602656. [PubMed: 15956977]
- [10]. Fearon ER, Cho KR, Nigro JM, Kern SE, Simons JW, Ruppert JM, Hamilton SR, Preisinger AC, Thomas G, Kinzler KW, et al. , Identification of a chromosome 18q gene that is altered in colorectal cancers, *Science* 247 (1990) 49–56. [PubMed: 2294591]
- [11]. Vogelstein B, Fearon ER, Kern SE, Hamilton SR, Preisinger AC, Nakamura Y, White R, Allelotype of colorectal carcinomas, *Science* 244 (1989) 207–211. [PubMed: 2565047]
- [12]. Shin SK, Nagasaka T, Jung BH, Matsubara N, Kim WH, Carethers JM, Boland CR, Goel A, Epigenetic and genetic alterations in Netrin-1 receptors UNC5C and DCC in human colon cancer, *Gastroenterology* 133 (2007) 1849–1857. 10.1053/j.gastro.2007.08.074. [PubMed: 18054557]
- [13]. Dumartin L, Quemener C, Laklai H, Herbert J, Bicknell R, Bousquet C, Pyronnet S, Castronovo V, Schilling MK, Bikfalvi A, Hagedorn M, Netrin-1 mediates early events in pancreatic adenocarcinoma progression, acting on tumor and endothelial cells, *Gastroenterology* 138 (2010) 1595–1606, 1606 e1591–1598. 10.1053/j.gastro.2009.12.061. [PubMed: 20080097]
- [14]. Fitamant J, Guenebeaud C, Coissieux MM, Guix C, Treilleux I, Scoazec JY, Bachelot T, Bernet A, Mehlen P, Netrin-1 expression confers a selective advantage for tumor cell survival in metastatic breast cancer, *Proceedings of the National Academy of Sciences of the United States of America* 105 (2008) 4850–4855. 10.1073/pnas.0709810105. [PubMed: 18353983]
- [15]. Kaufmann S, Kuphal S, Schubert T, Bosserhoff AK, Functional implication of Netrin expression in malignant melanoma, *Cellular oncology : the official journal of the International Society for Cellular Oncology* 31 (2009) 415–422. 10.3233/clo-2009-0491. [PubMed: 19940358]
- [16]. Rodrigues S, De Wever O, Bruyneel E, Rooney RJ, Gespach C, Opposing roles of netrin-1 and the dependence receptor DCC in cancer cell invasion, tumor growth and metastasis, *Oncogene* 26 (2007) 5615–5625. 10.1038/sj.onc.1210347. [PubMed: 17334389]
- [17]. Akino T, Han X, Nakayama H, McNeish B, Zurakowski D, Mammoto A, Klagsbrun M, Smith E, Netrin-1 promotes medulloblastoma cell invasiveness and angiogenesis, and demonstrates elevated expression in tumor tissue and urine of patients with pediatric medulloblastoma, *Cancer research* 74 (2014) 3716–3726. 10.1158/0008-5472.can-13-3116. [PubMed: 24812271]



- [18]. Nakayama H, Kusumoto C, Nakahara M, Fujiwara A, Higashiyama S, Semaphorin 3F and Netrin-1: The Novel Function as a Regulator of Tumor Microenvironment, *Front Physiol* 9 (2018) 1662. 10.3389/fphys.2018.01662. [PubMed: 30532711]
- [19]. Shimizu A, Nakayama H, Wang P, Konig C, Akino T, Sandlund J, Coma S, Italiano JE Jr., Mammoto A, Bielenberg DR, Klagsbrun M, Netrin-1 Promotes Glioblastoma Cell Invasiveness and Angiogenesis by Multiple Pathways Including Activation of RhoA, Cathepsin B, and cAMP-response Element-binding Protein, *The Journal of biological chemistry* 288 (2013) 2210–2222. 10.1074/jbc.M112.397398. [PubMed: 23195957]
- [20]. George ML, Tutton MG, Janssen F, Arnaout A, Abulafi AM, Eccles SA, Swift RI, VEGF-A, VEGF-C, and VEGF-D in colorectal cancer progression, *Neoplasia* 3 (2001) 420–427. 10.1038/sj.neo.7900186. [PubMed: 11687953]
- [21]. Hanrahan V, Currie MJ, Gunningham SP, Morrin HR, Scott PA, Robinson BA, Fox SB, The angiogenic switch for vascular endothelial growth factor (VEGF)-A, VEGF-B, VEGF-C, and VEGF-D in the adenoma-carcinoma sequence during colorectal cancer progression, *J Pathol* 200 (2003) 183–194. 10.1002/path.1339. [PubMed: 12754739]
- [22]. McIntyre A, Harris AL, Metabolic and hypoxic adaptation to anti-angiogenic therapy: a target for induced essentiality, *EMBO Mol Med* 7 (2015) 368–379. 10.15252/emmm.201404271. [PubMed: 25700172]
- [23]. Werther K, Christensen IJ, Brünner N, Nielsen HJ, Soluble vascular endothelial growth factor levels in patients with primary colorectal carcinoma. The Danish RANX05 Colorectal Cancer Study Group, *Eur J Surg Oncol* 26 (2000) 657–662. 10.1053/ejso.2000.0977. [PubMed: 11078612]
- [24]. Werther K, Christensen IJ, Nielsen HJ, Prognostic impact of matched preoperative plasma and serum VEGF in patients with primary colorectal carcinoma, *Br J Cancer* 86 (2002) 417–423. 10.1038/sj.bjc.6600075. [PubMed: 11875709]
- [25]. Grandin M, Mathot P, Devailly G, Bidet Y, Ghantous A, Favrot C, Gibert B, Gadot N, Puisieux I, Herceg Z, Delcros JG, Bernet A, Mehlen P, Dante R, Inhibition of DNA methylation promotes breast tumor sensitivity to netrin-1 interference, *EMBO Mol Med* 8 (2016) 863–877. 10.15252/emmm.201505945. [PubMed: 27378792]
- [26]. Li VS, Yuen ST, Chan TL, Yan HH, Law WL, Yeung BH, Chan AS, Tsui WY, So S, Chen X, Leung SY, Frequent inactivation of axon guidance molecule RGMA in human colon cancer through genetic and epigenetic mechanisms, *Gastroenterology* 137 (2009) 176–187. 10.1053/j.gastro.2009.03.005. [PubMed: 19303019]
- [27]. Thiebault K, Mazelin L, Pays L, Llambi F, Joly MO, Scoazec JY, Saurin JC, Romeo G, Mehlen P, The netrin-1 receptors UNC5H are putative tumor suppressors controlling cell death commitment, *Proceedings of the National Academy of Sciences of the United States of America* 100 (2003) 4173–4178. 10.1073/pnas.0738063100. [PubMed: 12655055]
- [28]. Mazelin L, Bernet A, Bonod-Bidaud C, Pays L, Arnaud S, Gespach C, Bredesen DE, Scoazec JY, Mehlen P, Netrin-1 controls colorectal tumorigenesis by regulating apoptosis, *Nature* 431 (2004) 80–84. 10.1038/nature02788. [PubMed: 15343335]
- [29]. Paradisi A, Maisse C, Coissieux MM, Gadot N, Lépinasse F, Delloye-Bourgeois C, Delcros JG, Svrcek M, Neufert C, Fléjou JF, Scoazec JY, Mehlen P, Netrin-1 up-regulation in inflammatory bowel diseases is required for colorectal cancer progression, *Proceedings of the National Academy of Sciences of the United States of America* 106 (2009) 17146–17151. 10.1073/pnas.0901767106. [PubMed: 19721007]
- [30]. Ramesh G, Berg A, Jayakumar C, Plasma netrin-1 is a diagnostic biomarker of human cancers, *Biomarkers* 16 (2011) 172–180. 10.3109/1354750x.2010.541564. [PubMed: 21303223]


**Highlights:**

- Netrin-1 mRNA and protein levels are down-regulated in most CRCs.
- Netrin-1 DNA methylation is higher in CRCs than in corresponding normal mucosa.
- Hypermethylation of netrin-1 is associated with CRC patients with advanced TNM stage.
- Serum levels of netrin-1 are significantly decreased in patients with CRC.




**Figure 1.**

*NTN1* expression is reduced in CRC compared to normal colon. A,B, *NTN1* expression in paired samples of CRC and normal colonic tissues by RNA-seq from TCGA database (n = 55). Results for individual CRC and paired normal colon tissue samples are displayed as dots at the same position in the x axis (A). Mean mRNA levels in CRC and paired normal colonic epithelium tissues are displayed as box-and-whisker plots (B). The horizontal line indicated the median (B). C,D, Relative mRNA levels of *NTN1* and its receptors and *VEGFA* measured by qRT-PCR in paired samples of CRC and normal colon tissues (n = 25). mRNA levels were normalized to *GAPDH* mRNA. E, Netrin-1 and UNC5C protein levels in three paired samples of CRC and normal colon tissues (normal: N, tumor: T) were analyzed by Western blot analysis.




**Figure 2.**

*NTN1* DNA is frequently hypermethylated in CRCs. A,B, DNA methylation levels in *NTN1* CpG island (chr17: 8,865,316–8,866,185) located proximal to the *NTN1* promoter were analyzed from the Human Methylation 450K Array (HM450, Illumina) in TCGA datasets of paired samples of CRC and normal colon tissues (n = 96). Results from individual pairs are shown as dots (A) and cumulative group results are shown as box-and-whisker plots (B). C, Correlation between *NTN1* gene expression and DNA methylation in colorectal cancer (n=95) and adjacent normal colon tissues (n=39). D, Correlation between *NTN1* DNA methylation and TNM cancer stage. Datasets were from TCGA database (see Supplementary Tables 5-7). Data represent the mean  $\pm$  SD [normal: T1 (n=1), T2 (n=10), T3 (n=69), T4 (n=7); tumor: T1 (n=20), T2 (n=108), T3 (n=430), T4 (n=72)], \*  $P < 0.05$ .



**Figure 3.**

Effects of DNA hypermethylation on *NTN1* expression in CRC cell lines. A, *NTN1* mRNA levels were measured in normal colon EC and CRC cell lines. mRNA levels were normalized to *GAPDH* mRNA. \*  $P < 0.001$ , \*\*  $P < 0.01$ . B, Bisulfite sequencing of normal EC and HT29 cells. Empty circles indicate unmethylated CpG sites (#1–23), filled circles represent methylated CpG sites (#1–23). C, Cells incubated with (w) or without (wo) DAC (10  $\mu$ M) for 3 days were analyzed by qRT-PCR (fold change, w / wo DAC). *NTN1* mRNA levels were normalized to *GAPDH* mRNA. D, CRC cell lines (HT29 and HCT116) were treated with DAC (10  $\mu$ M) for 3 days and cell lysates were analyzed by Western blotting. Normal EC and HCT116 cells after CRISPR/Cas9-mediated netrin-1 knockout are used as controls for Western blotting.



**Figure 4.** Serum netrin-1 levels are reduced in patients with CRCs compared to blood donors. Serum netrin-1 (A) and VEGF-A (C) levels were quantified by ELISA in patients with CRCs (n=31) and normal individuals (n=30). B, Specificity of and sensitivity of netrin-1 serum levels at predicting CRC. ROC curve analysis for determining the predictive value of serum netrin-1 concentration for CRC diagnosis.