

ORIGINAL ARTICLE

The efficacy and safety of adjunctive perampanel for the treatment of refractory focal-onset seizures in patients with epilepsy: A meta-analysis

Yiming Li¹  | Ya Zeng²  | Jie Mu³  | Dong Zhou³ 

¹West China Clinical Medical School, West China Hospital, Sichuan University, Chengdu, China

²Department of Pharmacy, West China Hospital, Sichuan University, Chengdu, China

³Department of Neurology, West China Hospital, Sichuan University, Chengdu, China

Correspondence

Jie Mu, Department of Neurology, West China Hospital, Sichuan University, Chengdu, China.

Emails: mujie2010@foxmail.com

Abstract

Objective: The last decade has seen an increase in the use of anti-seizure medications (ASMs); however, the burden of treating drug-resistant epilepsy has not fallen. We performed this meta-analysis to evaluate the optimal dose of Perampanel (PER) as a new adjunctive treatment for drug-resistant seizures.

Methods: We searched for studies published from inception to February 1, 2021 from PubMed, Central Register of Controlled Trials (CENTRAL), and ScienceDirect. Research characteristics, patients' characteristics, and treatment regimen, concomitant ASMs, clinical outcomes were extracted. The practical outcome included a reduction in seizures frequency $\geq 50\%$, $\geq 75\%$, and $\geq 100\%$ from baseline convulsive seizure frequency, and the safety outcome included the proportion of drug withdrawal and adverse reactions. Odds ratios (OR) for 95% confidence intervals (95% CI) were estimated by the inverse variance method.

Results: Four trials which enrolled 2187 participants (1569 in the PER group and 618 in the placebo group) were included. Results showed that 8 or 12 mg per day had the best effect on all three outcomes, with no significant difference between 8 and 12 mg per day ($\geq 50\%$ reduction, 35.5% vs 36.1%, $P = .84$; $\geq 75\%$ reduction, 17.8% vs 19.1%, $P = .64$; seizure-free, 3.5% vs 3.7%, $P = .85$). In addition, 12-mg PER compared to 8 mg had a higher proportion of trial withdrawal (8.7% vs 17.0%; $P < .00001$) and treatment-emergent adverse event (TEAE) resulting in dose reduction/discontinuation (18.5% vs 32.0%; $P < .00001$). The adverse events (AEs) significantly associated with adjunctive PER were dizziness, somnolence, fatigue, and irritability.

Significance: Adjunctive treatment of PER was associated with a more significant reduction in the frequency of seizures in patients with refractory epilepsy than placebo, but with a higher frequency of AEs. PER at a daily dose of 8 mg appears to have the best ratio between efficacy and tolerance in most study participants.

KEYWORDS

efficacy, perampanel, refractory focal-onset seizures, safety

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2021 The Authors. *Epilepsia Open* published by Wiley Periodicals LLC on behalf of International League Against Epilepsy

1 | INTRODUCTION

It is estimated that the incidence of epilepsy is about 80/100 000, and the prevalence is 5/1000, with about 70 million people suffering from epilepsy globally.^{1,2} Treatment of epilepsy is symptom based. Although many patients with epilepsy can control their seizures, there are still more than one-third of the cases of seizures that cannot be controlled.³ Uncontrolled epileptic seizures are likely to be a risk factor of impaired quality of life, disability, and premature death with severe physical and psychological dysfunction.⁴ However, with the increasing appearance and use of anti-seizure medications (ASMs) over the past few years, the burden of treating drug-resistant epilepsy has not been reduced. New and effective treatment options are still needed.⁵ The safety and effectiveness of newer drugs also require periodic evaluation.

Perampanel (PER) is a selective, non-competitive, and orally active alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionic acid receptor antagonist. This receptor plays a vital role in mediating rapid excitatory synaptic transmission, generation, and transmission of epileptic activity.⁶ In 2014, the European Union and the United States approved of PER (for patients 18 years of age or older), and to date, more than 40 countries around the world have approved of this drug.^{7,8} All anti-seizure medications (ASMs) were associated with the risk of treatment-emergent adverse events (TEAEs); Therefore, minimizing TEAEs is an important consideration when using ASMs. This meta-analysis assesses the efficacy and safety of adjunctive PER for the treatment of refractory focal-onset seizures in patients with epilepsy.

2 | METHODS

2.1 | Search strategy

We (Yiming Li and Ya Zeng) independently searched PubMed, ScienceDirect, and Central Register of Controlled Trials (CENTRAL) from their earliest dates up to February 1, 2021. The final search string was “perampanel” [Mesh] or “3-(2-cyanophenyl)-5-(2-pyridyl)-1-phenyl-1,2-dihydropyridin-2-one” AND “refractory partial-onset seizures” [Mesh]AND randomized controlled trial [ptyp]. No other filters were used. Studies were selected with the following entry criteria: Phase III, randomized, double-blind, placebo-controlled, parallel-group design with 6-week observational baseline and 19-week double-blind treatment phase (6-week titration period and 13-week maintenance period). Participants who meet the following criteria are included: age > 12, diagnosed with focal-onset seizures (partial-onset seizures) at least two ASMs failures in the

Key points

- Perampanel (PER) as a new adjunctive treatment for drug-resistant seizures is efficacy.
- Adjunctive treatment of PER was associated with a higher frequency of adverse effects.
- The optimal dose of Perampanel is 8 mg per day as a new adjunctive treatment for drug-resistant seizures between efficacy and tolerance.

previous 2 years, at least five focal seizures in the 6-week baseline phase and was taking stable doses of 1-4 approved concomitant ASMs. Exclusion criteria were as follows: not English language studies, RCT articles using the same experimental data, articles without recording 50% reduction in the frequency of seizures, and articles using indeterminate drug doses. No PROSPERO registration number for the moment. The specific screening process is shown in [Figure 1](#).

2.2 | Outcome measures

The primary outcome was the responder rate, defined as the percentage of patients who experienced a $\geq 50\%$ reduction in seizure frequency in the maintenance period relative to baseline in the European Union. Responder rate is a single point in the entirety of response and not a continuous variable. Thus, the 50% responder rate provides less information from all possible responses; it may be less sensitive. Increasing the gauge of responder rate to 75% showed more significant improvements compared with placebo. So, we defined a $\geq 75\%$ reduction in seizure frequency and seizure free as the secondary outcome.

Safety outcomes were as follows: 1. the proportions of patients who drop out of the treatment for any reason; 2. all TEAEs related to PER (ie, dizziness, somnolence, headache, fatigue, upper respiratory tract infection, nasopharyngitis, gait disturbance, or irritability).

2.3 | Data and assessment of the risk of bias

To ensure the consistency of the data collection of each study, we conform to the conditions of the study of the following information into a structured Excel data table: research characteristics (such as sample size, titration time, and maintenance time), the patient's characteristics (such as age, gender, etiology, duration of disease,




FIGURE 1 Flow diagram of study selection process

and complications), and treatment regimen, concomitant ASMs, and clinical outcomes (evaluation forms from the Cochrane Manual were used to assess the quality of selection, performance, detection, attrition, and reporting biases for each qualifying study [2011]).⁹

2.4 | Statistical analysis

To develop the meta-analysis, we used Cochrane Collaboration's Review Manager software (RevMan 5.4) to derive pooled effect estimates, such as odd ratios (OR) and associated 95% confidence intervals (CI). Statistical significance was assessed at a nominal α level of .05. The heterogeneity index I^2 determined the choice of fixed-effect model or random-effect model.

3 | RESULTS

3.1 | Study characteristics

Two hundred and twenty-four records were initially identified by searching the database and trial register. Four randomized controlled trials (RCTs) were retrieved for detailed evaluation.¹⁰⁻¹³ The essential characteristics of the studies are provided in [Table 1](#).

3.2 | Risk of bias assessment

Four studies were multicenter, randomized, double-blinded, placebo-controlled, parallel-group trials, which resulted in a low risk of bias ([Table 2](#)).

TABLE 1 Basic characteristics of the included studies

Article	Inclusion criteria	Race	Female gender, n (%)	Mean age (years)	Intervention group				
					Placebo	PER 2 mg	PER 4 mg	PER 8 mg	PER 12 mg
Jacqueline A. French 2013 ¹²	Diagnosis of simple or complex focal seizures permitted only one inducer ASM and must have been on a stable dose of any concomitant benzodiazepines	White (322) Asian (42) Others (22)	65 (47.8)/64 (49.6)/71 (58.7)	34.4 (13.6)/36.7 (14.4)/35.5 (14.1)	n = 136	n = 129	n = 129	n = 129	n = 121
Jacqueline A. French 2012 ¹⁴	Diagnosed with focal-onset seizures, with stable doses of 2-3 approved ASMs	White (334) Asian (4) Others (50)	67 (55.4)/68 (51.1)/65 (48.5)	35.6 (14.7)/35.8 (14.2)/36.7 (14.6)	n = 121	n = 133	n = 133	n = 133	n = 134
T. Nishida 2017 ¹³	/	/	90 (48.6)/95 (52.8)/84 (48.8)/92 (54.4)	33.4 (12.6)/33.8 (13.6)/33.6 (12.2)/34.6 (12.8)	n = 185	n = 180	n = 172	n = 169	n = 169
GL Krauss 2012 ¹⁵	Diagnosed with simple or complex focal-onset seizures, with stable doses of 2-3 approved ASMs	White (459) Asian (244) Others (3)	89 (50.9)/94 (54.0)/84 (48.0)/93 (51.7)	34.5 (13.2)/33.1 (13.2)/33.6 (14.1)/32.3 (12.3)	n = 175	n = 174	n = 174	n = 175	n = 180

3.3 | Efficacy

We compared four dosage types of PER (2, 4, 8, or 12 mg) with placebo; pooled data from the four RCTs showed that PER 4, 8, 12-mg groups all had a superior response compared with the placebo group, and the PER doses of 8 and 12 mg appeared to be more effective than 4-mg dose of PER (8 mg: $\geq 50\%$ reduction, 25.6% vs 35.5%, $P = .002$; $\geq 75\%$ reduction, 12.4% vs 16.8%, $P = .07$; seizure-free, 3.5% vs 3.6%, $P = .99$; 12 mg: $\geq 50\%$ reduction, 25.6% vs 36.0%, $P = .002$; $\geq 75\%$ reduction, 12.4% vs 19.1%, $P = .01$; seizure-free, 3.5% vs 3.7%, $P = .86$). We compared the PER doses of 8 mg with 12 mg in three efficacy outcomes ($\geq 50\%$ reduction, 35.5% vs 36.1%, $P = .84$; $\geq 75\%$ reduction, 17.8% vs 19.1%, $P = .64$; seizure-free, 3.5% vs 3.7%, $P = .85$), the data showed that there is no significant difference between the doses of 8 and 12 mg. The details of 50% reduction in seizure can be found in Figure 2. To sum up, the efficacy of different doses of PER is in the following order: 8 = 12 mg > 4 mg, the minimum effective dose of PER may be 4 mg/d as no statistically significant difference in the 2 mg/d dose compared with placebo.

3.4 | Treatment withdrawal and adverse events

In all trials, 152 (9.52%) and 27 (4.3%) patients withdrew from the study due to drug-related TEAEs in the PER-added and placebo groups, respectively (OR 2.50, 95% CI 1.64-3.82; $I^2 = 0\%$; $P < .0001$). The incidences of patients' withdrawal from the study due to drug-related TEAEs were higher in PER supplementation with 2, 8, and 12 mg compared with the placebo group (2 mg, 6.7% vs 3.8%, $P < .02$; 8 mg, 8.4% vs 4.4%, $P < .004$; 12 mg, 17.0% vs 4.6%, $P < .00001$). PER supplementation with 4 mg showed no significant difference (3.7% vs 3.6%, $P < .92$) (Table 3).

In addition, 12-mg PER compared to 8 mg had a higher proportion of trial withdrawal (8.7% vs 17.0%; $P < .00001$) and TEAEs resulting in dose reduction/discontinuation (18.5% vs 32.0%; $P < .00001$) without a significant increase in efficacy. To sum up, the safety of different doses of PER is in the following order: 4 > 8 > 12 mg. Two milligrams was not included in this comparison because 2 mg had no efficacy; thus, we considered the safety of 2 mg to be meaningless.

The incidence of treatment-related TEAEs was higher in the PER-added group (59.6% vs 37.9%; $P < .00001$) (Table 3), which was also related to dosage. TEAEs were higher in PER supplementation with 4, 8, and 12 mg compared with the placebo group (4 mg, 45.4% vs 30.7%, $P = .0001$; 8 mg, 61.2% vs 34.8%, $P < .00001$; 12 mg, 75.2%

TABLE 2 Analysis of efficacy and the association between perampanel (PER) and placebo

Outcome or subgroup	Number of studies	Participants	I ² , %	Odds ratio (95% CI)	P
1.1 50% reduction in the seizure frequency	4	2187	37%	1.96 [1.56, 2.45]	<.00001
1.1.1 PER 2 mg/d vs placebo	1	364	/	1.18 [0.70, 2.00]	.53
1.1.2 PER 4 mg/d vs placebo	2	751	0	1.45 [1.02, 2.08]	.04
1.1.3 PER 8 mg/d vs placebo	4	1222	0	2.12 [1.63, 2.75]	<.00001
1.1.4 PER 12 mg/d vs placebo	3	866	41	2.53 [1.87, 3.44]	<.00001
1.2 75% reduction in the seizure frequency	4	2178	0%	2.74 [1.93, 3.89]	<.00001
1.2.1 PER 2 mg/d vs placebo	1	265	/	1.94 [0.87, 4.34]	.1
1.2.2 PER 4 mg/d vs placebo	2	709	0%	1.73 [1.04, 2.88]	<i>P</i> = .03
1.2.3 PER 8 mg/d vs placebo	4	1224	0%	3.01 [2.04, 4.43]	<i>P</i> < .00001
1.2.4 PER 12 mg/d vs placebo	3	868	0%	3.29 [2.10, 5.15]	<i>P</i> < .00001
1.3 Seizure freedom during the treatment	4	2178	0%	3.24 [1.42, 7.83]	.005
1.3.1 PER 2 mg/d vs placebo	1	265	/	1.55 [0.26, 9.39]	<i>P</i> = .63
1.2.2 PER 4 mg/d vs placebo	2	709	0%	3.20 [1.02, 10.00]	<i>P</i> = .05
1.2.3 PER 8 mg/d vs placebo	4	1224	0%	3.51 [1.45, 8.51]	<i>P</i> = .005
1.2.4 PER 12 mg/d vs placebo	3	868	0%	3.88 [1.35, 11.14]	<i>P</i> = .01

vs 37.0%, *P* = .0002). There was no significant difference in PER supplementation with 2 mg (37.2% vs 31.9%, *P* < .29) (Table 3).

There was no statistical difference in severe TEAEs (5.1% vs 5.1%; *P* < .88). The most common TEAE was dizziness and there is a significant difference between the PER-added group and the placebo group (29.1% vs 8.1%; *P* < .00001). In addition, the incidence of major TEAEs in the PER-added group and the placebo group was as follows: somnolence (16.0% vs 7.2%; *P* < .001), headache (8.6% vs 10.0%; *P* = .94), fatigue (7.2% vs 4.4%; *P* = .03), upper respiratory tract infection (5.0% vs 3.6%; *P* = .28), nasopharyngitis (8.4% vs 8.0%; *P* = .86), gait disturbance (3.8% vs 2.2%; 0.06), irritability (7.1% vs 2.3%; 0.003), rash (2.4% vs 1.1%; *P* = .31), nausea (3.3% vs 2.8%; 0.72), and falls (11.2% vs 6.6%; *P* = .12) (Table 4).

4 | DISCUSSION

Epilepsy remains uncontrolled in one-third of patients despite appropriate medical therapy. PER has been marketed as third-generation ASMs and adjunctive treatments for focal-onset seizures in China in 2019. Guideline of American Academy of Neurology and the American Epilepsy Society 2018 recommended PER for treatment-resistant adult focal epilepsy (level A).¹⁴ Oral PER has superficial pharmacological characteristics: it can be rapidly absorbed from the gastrointestinal tract; Steady-state plasma concentrations can be reached within 14 days of oral administration, with a terminal half-life of

approximately 70-120 hours.¹⁵ PER does not affect plasma concentrations of ASM taken simultaneously.¹⁶

Our meta-analysis results indicate that adjunctive treatment of PER at a daily dose of 4, 8, or 12 mg significantly reduced the number of seizures in patients with refractory focal seizures that were rarely discontinued due to unacceptable TEAEs. The minimum effective dosage of PER maybe 4 mg/d, because there is no statistically significant difference in the 2 mg/d dose compared with placebo. However, since 2 mg was included in only one experiment, its effectiveness needs to be further verified. The PER doses of 8 mg and 12 mg are more effective than 4 mg, and the PER doses of 8 and 12 mg had no significant difference in efficacy. In addition, a small number of patients experienced seizures, which may be a dose-dependent phenomenon; although the number was small, this trend was statistically significant.

Perampanel increased the incidence of TEAEs compared with placebo, which was higher in PER supplementation with 8 and 12 mg. The vast majority of TEAEs were mild/moderate at standard doses. Dizziness is the most common AE and may have a dose-response; however, only a small number of those patients (8.3%) discontinued treatment due to this symptom, and no safety issues have been identified.¹⁰ In addition, PER may increase the incidence of somnolence, fatigue, and irritability.

Patients experienced more TEAEs and had a higher proportion of trial withdrawal after taking the 12-mg dose, and the efficacy is less significant than that of 8 mg, so PER 8 mg/d may be the perfect option. However,




FIGURE 2 Effect of perampanel on 50% reduction in refractory focal-onset seizure

there are still additional benefits for a significant number of patients to accept the 12 mg.¹⁷ The 12-mg dose may be an essential option to achieve the goal of a more significant reduction of seizures and free seizures in patients who can tolerate and do not achieve optimal response at an 8-mg dose.

In addition, although the maintenance period is twice if the titration period, the frequency of TEAEs during the maintenance period is lower than titration, indicating that they are transient, with no increase in the incidence of TEAE over time, and no potential tolerance.^{10,18} The low or nonexistent incidence of the first occurrence of

TABLE 3 Treatment dropout between perampanel (PER) and placebo

Outcome or subgroup	Studies	Events/participants (%)		I ² (%)	Odd ratio (95% CI)	P
		PER	Placebo			
Treatment dropout						
PER any dose	4	152/1569 (9.5%)	27/618 (4.4%)	0	2.50 [1.64, 3.82]	<.001
PER 2 mg/d	1	12/180 (6.7%)	7/185 (3.8%)	0	1.82 [0.70, 4.72]	.02
PER 4 mg/d	2	13/384 (3.7%)	13/361 (3.6%)	0	1.04 [0.47, 2.27]	.92
PER 8 mg/d	4	53/633 (8.4%)	27/617 (4.4%)	15	2.00 [1.24, 3.22]	.004
PER 12 mg/d	3	74/435 (17.0%)	20/433 (4.6%)	0	4.23 [2.53, 7.08]	<.001
Any TEAE leading to dose reduction/interruption						
PER any dose	4	286/1569 (18.2%)	24/618 (3.9%)			
PER 2 mg/d	1	3/180 (1.7%)	6/185 (3.2%)		0.51 [0.12, 2.05]	.34
PER 4 mg/d	2	32/348 (9.2%)	13/361 (3.6%)	0	2.13 [1.20, 3.79]	.01
PER 8 mg/d	4	112/606 (18.5%)	24/618 (3.9%)	0	5.41 [3.57, 8.21]	<.001
PER 12 mg/d	3	139/435 (32.0%)	18/433 (4.2%)	0	9.87 [6.24, 15.62]	<.001
Any TEAE						
PER any dose	4	1026/1569 (75.6%)	411/618 (66.5%)		0.98 [0.92, 1.05]	.62
PER 2 mg/d	1	111/180 (61.7%)	101/185 (54.6%)		1.13 [0.95, 1.35]	.17
PER 4 mg/d	2	232/348 (66.7%)	218/361 (60.4%)	27%	1.10 [0.99, 1.23]	.09
PER 8 mg/d	4	479/570 (84.0%)	411/682 (60.3%)	91%	1.25 [1.02, 1.55]	.03
PER 12 mg/d	3	383/435 (88.0%)	310/433 (71.6%)	61%	1.21 [1.09, 1.35]	.0004
Any treatment-related TEAE						
PER any dose	4	936/1569 (59.7%)	234/618 (37.9%)	62	2.72 [2.23, 3.31]	.00001
PER 2 mg/d	1	67/180 (37.2%)	59/185 (31.9%)		1.17 [0.88, 1.55]	.29
PER 4 mg/d	2	158/348 (45.4%)	111/361 (30.7%)	0	1.48 [1.22, 1.79]	.0001
PER 8 mg/d	3	292/477 (61.2%)	169/485 (34.8%)	0	1.70 [1.49, 1.95]	<.00001
PER 12 mg/d	2	236/314 (75.2%)	110/297 (37.0%)	80%	2.00 [1.40, 2.86]	.0002

Note: Risk ratios are from a fixed-effects model.

TABLE 4 TEAEs between perampanel (PER) and placebo

Outcome	Studies	Events/participants (%)		I ² (%)	Odd Ratio (95% CI)	P
		PER	Placebo			
All TEAEs	4	936/1569 (59.7%)	234/618 (37.9%)	62	2.72 [2.23, 3.31]	<.001
Dizziness	4	458/1569 (29.2%)	50/618 (8.1%)	80	4.83 [3.55, 6.58]	<.001
Somnolence	4	251/1569 (16.0%)	45/618 (7.3%)	64	2.45 [1.75, 3.41]	<.001
Headache	4	135/1569 (8.6%)	62/618 (10.0%)	0	1.01 [0.74, 1.39]	.94
Fatigue	3	86/1188 (7.2%)	22/497 (4.4%)	0	1.74 [1.07, 2.83]	.03
Upper respiratory tract infection	3	53/1052 (5.0%)	13/361 (3.6%)	0	1.41 [0.76, 2.62]	.28
Nasopharyngitis	2	89/1052 (8.5%)	29/361 (8.0%)	0	0.96 [0.62, 1.48]	.86
Gait disturbance	2	51/1319 (3.9%)	11/482 (2.3%)	0	1.88 [0.97, 3.66]	.06
Irritability	2	56/788 (7.1%)	7/297 (2.4%)	41	3.48 [1.54, 7.83]	.003
Rash	1	13/531 (2.4%)	2/176 (1.1%)		2.18 [0.49, 9.77]	.31
Nausea	1	18/531 (3.4%)	5/176 (2.8%)		1.20 [0.44, 3.28]	.72
Fall	1	30/267 (11.2%)	8/121 (6.6%)		1.79 [0.79, 4.02]	.16

these TEAEs after 6 months to 1 year of treatment is further evidence that long-term treatment with PER is safe and well tolerated.¹⁸

Three patients died in these four RCTs; one died of sudden cardiac death in the placebo group, one died of an unknown cause in the PER 8-mg group and one died of convulsion during baseline. Because of the minor frequency of events, it remains unclear whether drugs caused the deaths. In addition, three patients from the placebo group, one from the PER 2-mg group, two from the PER 8-mg group, and two from PER 12 mg patients appeared with suicidal tendencies. Again, we do not know for sure whether the suicidal tendencies were related to the effects of the drugs. The data showed no statistically significant difference in severe TEAEs between the PER and the placebo groups. Overall, the incidence of psychotic serious TEAEs was higher in the 12-mg group than in the other dose or placebo groups. Although there was no statistically significant difference in severe adverse reactions between the placebo and PER groups, the proportion of TEAEs leading to discontinuation and dose reduction/interruption was higher in the PER group than in the placebo group. Dose reduction rather than PER withdrawal was used in most cases to deal with TEAEs.

5 | CONCLUSIONS

Perampanel is an optional adjunctive method for refractory focal epilepsy. Adjunctive treatment of PER was associated with a more significant reduction in the frequency of seizures in patients with refractory epilepsy than placebo, but with a higher frequency of AEs. A daily dose of PER 8 mg is considered the best dosing option. To enhance patient tolerance, we suggest increasing and reducing the dose gradually when starting or discontinuing. More research will be forthcoming to explain further the true therapeutic potential and clinical significance of this latest ASM.

6 | STRENGTHS AND LIMITATIONS OF THIS STUDY

- This is a meta-analysis of the efficacy and safety of different doses of adjunctive PER in patients with focal-onset seizures.
- In this meta-analysis, efficacy and safety analyses are mainly based on daily doses of PER estimates of seizure response during the maintenance phase, the most accurate phase to represent steady-state drug levels in the entire treatment period.
- This meta-analysis included only four RCTs. The literature did not rate "refractory" (for example, patients with focal epilepsy who had previously been resistant to

more than three different drugs were significantly more resistant than patients who had been previously resistant to one drug).

- This meta-analysis inherited the general limitations of the four RCTs, such as the short duration of maintenance and the potential impact of concomitant drugs.
- This meta-analysis has no information about PER monotherapy's efficacy, tolerability, and safety PER during pregnancy and lactation.

CONFLICT OF INTEREST

Neither of the authors has any conflict of interest to disclose. We confirm that we have read the Journal's position on issues involved in ethical publication and affirm that this report is consistent with those guidelines.

AUTHOR CONTRIBUTIONS

Dong Zhou, Jie Mu, and Yiming Li proposed and designed the purpose; Yiming Li and Ya Zeng developed and revised the Searching strategy. All co-authors have contributed to the revision of the manuscript.

ORCID

Yiming Li  <https://orcid.org/0000-0002-2022-6811>

Ya Zeng  <https://orcid.org/0000-0001-8379-9003>

Jie Mu  <https://orcid.org/0000-0002-9773-3838>

Dong Zhou  <https://orcid.org/0000-0001-7101-4125>

REFERENCES

1. Hirtz D, Thurman DJ, Gwinn-Hardy K, Mohamed M, Chaudhuri AR, Zalutsky R. How common are the "common" neurologic disorders? *Neurology*. 2007;68(5):326–37.
2. Cagnetti C, Lattanzi S, Foschi N, Provinciali L, Silvestrini M. Seizure course during pregnancy in catamenial epilepsy. *Neurology*. 2014;83(4):339–44.
3. Chen Z, Brodie MJ, Liew D, Kwan P. Treatment outcomes in patients with newly diagnosed epilepsy treated with established and new antiepileptic drugs: a 30-year longitudinal cohort study. *JAMA Neurol*. 2018;75(3):279–86.
4. Lattanzi S, Zaccara G, Giovannelli F, Grillo E, Nardone R, Silvestrini M, et al. Antiepileptic monotherapy in newly diagnosed focal epilepsy. A Network meta-analysis. *Acta Neurol Scand*. 2019;139(1):33–41.
5. Laxer KD, Trinka E, Hirsch LJ, Cendes F, Langfitt J, Delanty N, et al. The consequences of refractory epilepsy and its treatment. *Epilepsy Behav*. 2014;37:59–70.
6. Hanada T. The discovery and development of perampanel for the treatment of epilepsy. *Expert Opin Drug Discov*. 2014;9(4):449–58.
7. FYCOMPA (Perampanel). US Prescribing Information. Eisai Inc.; 2014
8. Schmidt LA, Myers JL, McHugh JB. Napsin A is differentially expressed in sclerosing hemangiomas of the lung. *Arch Pathol Lab Med*. 2012;136(12):1580–4.
9. Wang QB, Chen YQ, Shen JJ, Zhang C, Song B, Zhu XJ, et al. Sixteen cases of pulmonary sclerosing haemangioma: CT

- findings are not definitive for preoperative diagnosis. *Clin Radiol.* 2011;66(8):708–14.
10. Pulido-Moran M, Moreno-Fernandez J, Ramirez-Tortosa C, Ramirez-Tortosa M. Curcumin and health. *Molecules.* 2016;21(3):264.
 11. Solheim TS, Laird BJA, Balstad TR, Stene GB, Bye A, Johns N, et al. A randomized phase II feasibility trial of a multimodal intervention for the management of cachexia in lung and pancreatic cancer. *J Cachexia Sarcopenia Muscle.* 2017;8(5):778–88.
 12. French JA, Krauss GL, Steinhoff BJ, Squillacote D, Yang H, Kumar D, et al. Evaluation of adjunctive perampanel in patients with refractory partial-onset seizures: results of randomized global phase III study 305. *Epilepsia.* 2013;54(1):117–25.
 13. Nishida T, Lee SK, Inoue Y, Saeki K, Ishikawa K, Kaneko S. Adjunctive perampanel in partial-onset seizures: Asia-Pacific, randomized phase III study. *Acta Neurol Scand.* 2018;137(4):392–9.
 14. Kanner AM, Ashman E, Gloss D, Harden C, Bourgeois B, Bautista JF, et al. Practice guideline update summary: efficacy and tolerability of the new antiepileptic drugs II: treatment-resistant epilepsy: report of the Guideline Development, Dissemination, and Implementation Subcommittee of the American Academy of Neurology and the American Epilepsy Society. *Neurology.* 2018;91(2):82–90.
 15. Diaz JA, Kim WH, Fernandez CP, Jeong J, Afrin F, Lillehoj HS, et al. Identification and expression analysis of duck interleukin-17D in *Riemerella anatipestifer* infection. *Dev Comp Immunol.* 2016;61:190–7.
 16. Yamamoto Y, Shiratani Y, Asai S, Usui N, Nishida T, Imai K, et al. Pharmacokinetics, tolerability, and clinical effectiveness of perampanel in Japanese patients with epilepsy. *Seizure.* 2020;83:181–6.
 17. French JA, Krauss GL, Biton V, Squillacote D, Yang H, Laurenza A, et al. Adjunctive perampanel for refractory partial-onset seizures: randomized phase III study 304. *Neurology.* 2012;79(6):589–96.
 18. Ko D, Yang H, Williams B, Xing D, Laurenza A. Perampanel in the treatment of partial seizures: time to onset and duration of most common adverse events from pooled Phase III and extension studies. *Epilepsy Behav.* 2015;48:45–52.

How to cite this article: Li Y, Zeng Y, Mu J, Zhou D. The efficacy and safety of adjunctive perampanel for the treatment of refractory focal-onset seizures in patients with epilepsy: A meta-analysis. *Epilepsia Open.* 2022;7:271–279. <https://doi.org/10.1002/epi4.12574>