


# Hsa\_circ\_0017639 regulates cisplatin resistance and tumor growth via acting as a miR-1296-5p molecular sponge and modulating sine oculis homeobox 1 expression in non-small cell lung cancer

Feiyun Chang, Jiali Li, Quan Sun, Shuqing Wei, and Yongming Song 

Department of Thoracic Surgery, Shanxi Province Cancer Hospital, Shanxi Hospital Affiliated to Cancer Hospital, Chinese Academy of Medical Sciences, Cancer Hospital Affiliated to Shanxi Medical University, Taiyuan, Shanxi, China

## ABSTRACT


Cisplatin (DDP)-induced chemoresistance is an important reason for the failure of non-small cell lung cancer (NSCLC) treatment. Circular RNAs (circRNAs) participate in the chemoresistance of diverse cancers. However, the function of hsa\_circ\_0017639 (circ\_0017639) in the DDP resistance of NSCLC is unclear. Forty-one NSCLC samples (21 DDP-resistant samples and 20 DDP-sensitive samples) were utilized in the research. The relative expression levels of some genes were determined by real-time quantitative polymerase chain reaction (RT-qPCR). 3-(4,5-Dimethylthiazol-2-yl)-2,5-Diphenyltetrazolium Bromide (MTT) assay for half-maximal inhibitory concentration (IC<sub>50</sub>) value of DDP and cell viability, colony formation and 5-ethynyl-2'-deoxyuridine (EDU) assays for cell proliferation, flow cytometry assay for cell apoptosis, transwell assay for cell invasion and wound-healing assay for cell migration were performed. The regulation mechanism of circ\_0017639 was demonstrated by a dual-luciferase reporter assay. We observed higher levels of circ\_0017639 in DDP-resistant NSCLC samples and cells. Functionally, circ\_0017639 silencing decreased tumor growth and elevated DDP sensitivity *in vivo* and induced apoptosis, repressed proliferation, invasion, and migration of DDP-resistant NSCLC cells *in vitro*. Mechanically, circ\_0017639 modulated sine oculis homeobox 1 (SIX1) expression via sponging microRNA (miR)-1296-5p. Also, miR-1296-5p inhibitor restored circ\_0017639 knockdown-mediated impacts on cell DDP resistance in DDP-resistant NSCLCs. Furthermore, SIX1 overexpression counteracted the inhibiting impact of miR-1296-5p upregulation on DDP resistance and malignant phenotypes of DDP-resistant NSCLC cells. In conclusion, circ\_0017639 conferred DDP resistance and promoted tumor growth via elevating SIX1 expression through sequestering miR-1296-5p in NSCLC, providing a new mechanism for understanding the chemoresistance and progression of NSCLC.

## ARTICLE HISTORY

Received 4 January 2022  
Revised 8 March 2022  
Accepted 10 March 2022

## KEYWORDS




DDP;  
IC<sub>50</sub>; NSCLC; circ\_0017639;  
miR-1296-5p; SIX1



## Introduction

Lung cancer causes a higher mortality rate than any other malignant tumor [1]. Despite the remarkable development of immunotherapy and molecular-targeted therapy, the 5-year overall survival rate of lung cancer patients is still very

low [2,3]. Non-small cell lung cancer (NSCLC) is the major subtype of lung cancer (approximately 85% of lung cancer cases) [4,5]. Cisplatin (DDP) is a first-line chemotherapeutic drug for advanced NSCLC. Normally, patients with NSCLC respond to the initial treatment of

**CONTACT** Yongming Song  [Songyongming0303@163.com](mailto:Songyongming0303@163.com)  Department of Thoracic Surgery, Shanxi Province Cancer Hospital, Shanxi Hospital Affiliated to Cancer Hospital, Chinese Academy of Medical Sciences, Cancer Hospital Affiliated to Shanxi Medical University, Taiyuan 030013, Shanxi, China  
 Supplemental data for this article can be accessed [here](#).

© 2022 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.  
This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

DDP, but resistance often develops, resulting in poor survival [6].

Recently, changes in circular RNAs (circRNAs) levels have been observed in DDP-resistant NSCLC cells [7]. CircRNAs are mainly produced through the process of back-splicing, in which the 3' and 5' ends of the precursor RNA are cleaved and joined by the splicing mechanism [8]. Moreover, circRNAs have the advantages of specific expression in different cell types, tissue types, and developmental stages [9]. Researchers uncovered that circRNAs participate in miscellaneous pathological processes, including tumorigenesis, tumor metastasis, and drug resistance [10]. For instance, circRNA circ-fibroblast growth factor receptor 1 accelerated anti-programmed death-1 resistance and tumor growth through upregulating C-X-C motif chemokine receptor 4 in NSCLC [11]. Another study exposed that decreased circRNA circ\_0001946 expression in NSCLC was related to DDP insensitivity [12]. By analyzing the GSE112214 microarray, we found that circRNA hsa\_circ\_0017639 (circ\_0017639) is a differentially expressed circRNA in NSCLC tissues. Li and colleagues revealed that circ\_0017639 plays an oncogenic function in gastric cancer [13]. At present, the action of circ\_0017639 in NSCLC is indistinct, so it aroused our interest.

Researchers have demonstrated that circRNAs exert a crucial regulatory action as microRNA (miR) decoys/sponges in tumors [14]. MiRs are important post-transcriptional regulators of gene expression, and they control many cellular processes and developmental in eukaryotic organisms [15]. MiR-1296-5p exerts a tumor-suppressing impact in osteosarcoma [16], breast cancer [17], and hepatocellular cancer [18]. However, the action of miR-1296-5p in NSCLC resistance to DDP has not been verified. Transcription factor sine oculis homeobox 1 (SIX1) is a member of the sine oculis (So/Six) homeobox protein family and a powerful regulator of organogenesis [19]. SIX1 is considered to be a cancer fetoprotein, and its inappropriate re-expression can lead to genome instability, malignant transformation, and metastasis [20]. Moreover, SIX1 participates in the chemoresistance of many cancers [21–23]. Upregulation of SIX1 can confer paclitaxel resistance in breast cancer [24], DDP resistance in

NSCLC [25], and 5-fluorouracil resistance in hepatocellular cancer [26]. However, the mechanism related to SIX1 dysregulation in DDP resistance of NSCLC remains unclear.

Thus, the research was to explore the function of circ\_0017639 in NSCLC resistance to DDP and its regulatory mechanism. Herein, circ\_0017639 was verified as an adverse circRNA in NSCLC. Mechanically, circ\_0017639 upregulation endowed DDP resistance in NSCLC by elevating the SIX1 expression via serving as a miR-1296-5p decoy.

## Materials and methods

### Biological specimens

Forty-one human NSCLC samples and matched adjacent normal samples were collected under the authorization of the Ethical Committee of Shanxi Province Cancer Hospital, Shanxi Hospital Affiliated to Cancer Hospital, Chinese Academy of Medical Sciences, Cancer Hospital Affiliated to Shanxi Medical University. According to the DDP treatment response, 41 NSCLC samples were divided into two parts: DDP-resistant NSCLC group (21 samples) and DDP-sensitive NSCLC group (20 samples). All registered NSCLC patients provided informed consent and underwent surgical resection at Shanxi Province Cancer Hospital, Shanxi Hospital Affiliated to Cancer Hospital, Chinese Academy of Medical Sciences, Cancer Hospital Affiliated to Shanxi Medical University.

### Cell culture

Human bronchial epithelial cells (HBE), HEK-293 T cells, and lung cancer cells (H460, H1650, A549, and H1299) (Procell, Wuhan, China) were maintained under the right conditions (5% carbon dioxide and 37°C). HEK-293 T and A549 cells were, respectively, cultured in DMEM (PM150210, Procell) and Ham's F-12 K (PM150910, Procell), whereas other cells were cultured in Roswell Park Memorial Institute-1640 Medium (PM150110, Procell). All media utilized for cell growth were supplemented with 10% Fetal Bovine Serum (FBS) (164,210–500, Procell) and 1% Penicillin/Streptomycin (PB180120, Procell).

### Establishment of DDP-resistant NSCLC cells

DDP-resistant NSCLC cells (A549/DDP and H1299/DDP) were established by a gradual increase in DDP concentration as described previously [27]. Briefly, A549 cells were initially exposed to DDP (0.5 µg/mL) DDP (Sigma, St Louis, MO, USA) for 3 days and then recovered within 3 days. After three cycles, the concentration of DDP was gradually increased to 1, 2, 4, 8 and 10 µg/mL to generate DDP-resistant NSCLC cells.

### Oligonucleotides and plasmids

Small interference (si) RNA against circ\_0017639 (si-circ\_0017639), miR-1296-5p inhibitor (anti-miR-1296-5p), miR-1296-5p mimic (miR-1296-5p), as well as the negative controls of oligonucleotide sequences mentioned above (si-NC, anti-miR-NC, and miR-NC) were synthesized by AoKe Biotech (Beijing, China). For pCD5-ciR-circ\_0017639 (circ\_0017639) and pcDNA-SIX1 (SIX1) plasmids, the cDNA sequences of circ\_0017639 and SIX1 were inserted into empty pCD5-ciR (vector) (Geneseeed, Guangzhou, China) and pcDNA (Thermo Fisher, Burlington, ON, Canada) vectors, respectively. Transfection of DDP-resistant NSCLC cells with specific oligonucleotides and/or plasmids was performed using Lipofectamine 3000 (Thermo Fisher) or Lipofectamine RNAiMax (Thermo Fisher).

### RNA isolation and RNase R digestion

Total RNA was isolated using an RNA Isolation Kit (Qiagen, Hilden, Germany) following the manufacturer's steps. RNA from the nucleus or cytoplasm was extracted using the Cytoplasmic & Nuclear RNA Purification Kit (Norgen Biotek, Thorold, Canada) based on the manufacturer's procedures. To validate the circular structure of circ\_0017639, total RNA from DDP-resistant NSCLC cells was incubated with diethylpyrocarbonate (DEPC)-treated water (Sigma) or RNase R (20 U/µL, Sigma) [28].

### Real-time quantitative polymerase chain reaction (RT-qPCR)

The synthesis of complementary DNA from total RNA was conducted with Prime Script™ RT reagent

kit (TaKaRa, Dalian, China) or miScript II RT Kit (Qiagen). Quantitative PCR was set up with the SYBR Green kit (Qiagen) in triplicate, and data were calculated using the  $2^{-\Delta\Delta C_t}$  method [29]. Primer sequences are exhibited in Table 1. U6 or  $\beta$ -Actin was used as an endogenous reference.

### 3-(4,5-Dimethylthiazol-2-yl)-2,5-Diphenyltetrazolium Bromide (MTT) assay

The half-maximal inhibitory concentration (IC<sub>50</sub>) value and viability of DDP-resistant NSCLC cells were evaluated using the MTT assay following the previous description [30]. In brief, cells with or without DDP treatment were incubated for 48 h, followed by co-incubating with the MTT solution (10 µL) (Roche, Basel, Switzerland) for 4 h. Following this, the purple crystals were dissolved, and the optical density was analyzed using an absorbance reader (BioTek, Winooski, VT, USA).

### Colony formation assay

A previous report was referenced to perform this experiment [31]. Approximately  $2 \times 10^2$  DDP-resistant NSCLC cells were seeded into 6-well plates. After culture for 12 days, the cloned cells were stained with 0.1% crystal violet (Beyotime, Shanghai, China) and then photographed and counted using a microscope (Olympus, Tokyo, Japan).

### 5-ethynyl-2'-deoxyuridine (EDU) assay

The Cell-Light™ EDU Apollo 567 In Vitro Imaging Kit (Thermo Fisher) was utilized to observe the EDU-positive cells following the manufacturer's instructions. Nuclei were labeled with blue fluorescence using 4,6-diamidino-2-phenylindole

**Table 1.** Primer sequences used for RT-qPCR.

Genes	Primer sequences (5'-3')
circ_0017639	Forward (F): 5'-CAGATTTGCGAAGCCATTTC-3' Reverse (R): 5'-TCCTCGAACCCAGTCAAGTCA-3'
SIX1	F: 5'-AGGTCAGCAACTGGTTAAGAACC-3' R: 5'-GAGGAGAGAGTTGGTCTGCTTG-3'
$\beta$ -Actin	F: 5'-CAGCCATGTACGTTGCTATCCA-3' R: 5'-TCACCGGAGTCCATCACGAT-3'
miR-1296-5p	F: 5'-CGTTAGGGCCCTGGCTCC-3' R: 5'-CAGTGCCTGTCGTGGAGT-3'
U6	F: 5'-CTCGCTTCGGCAGCAC-3' R: 5'-AACGCTTCACGAATTTGCGT-3'

(DAPI). Images were captured using a microscope (Olympus).

### **Flow cytometry assay**

DDP-resistant NSCLC cells were harvested and then trypsinized gently. Following this, the cells were re-suspended with  $1 \times$  binding buffer and then stained with Annexin V-fluorescein isothiocyanate (FITC) (5  $\mu$ L) and phosphatidylinositol (PI) (5  $\mu$ L) using the Annexin V-FITC/PI apoptosis detection kit (Becton Dickinson, Santa Cruz, California, USA). After incubation for 5 min on ice, the cells were analyzed using a FACS Verse flow cytometer (Becton Dickinson).

### **Transwell invasion assay**

Cell invasion was analyzed using transwell chambers with Matrigel (Costar, Cambridge, MA, USA) as previously depicted [31]. Approximately  $1 \times 10^5$  DDP-resistant NSCLC cells in 200  $\mu$ L serum-free medium were seeded into the upper compartment, and 600  $\mu$ L completed media containing 10% FBS (Procell) was added to the lower compartment. 24 h later, the invaded cells were stained with 0.1% crystal violet (Beyotime), followed by counting using a microscope (Olympus) in 5 random fields.

### **Wound-healing assay**

The wound-healing assay was utilized for migration analysis [32]. In short, about  $1 \times 10^3$  DDP-resistant NSCLC cells were seeded into 6-well plates. Wounds were scratched in the cell monolayer using a pipette tip when the cells reach 90–100% confluence. After rinsing with phosphate-buffered saline, the cells were incubated for 24 h under the right conditions. A microscope (Olympus) was utilized to capture the pictures at 0 and 24 h.

### **Dual-luciferase reporter assay**

The fragments of wild type (WT) circ\_0017639 and WT SIX1 3'UTR and their mutant (MUT) sequences were ligated into the pMIR-REPORT vector (Thermo Fisher). Luciferase assay was

performed by co-transfecting with the luciferase vector along with miR-1296-5p or miR-NC and pRL-TK plasmids (Thermo Fisher) into DDP-resistant NSCLC cells. Analysis of luciferase activity was done with the Pierce™ Renilla-Firefly Luciferase Dual Assay Kit (Thermo Fisher).

### **Western blotting**

Total protein was extracted using the RIPA buffer (Beyotime). Western blotting was done as described by Chen *et al.* [33] with specific primary antibodies against SIX1 (ab243247, Abcam, USA), matrix metalloproteinase 9 (MMP9) (ab58803, Abcam), and Cleaved-caspase-3 (Cleaved-caspase-3, Abcam), and  $\beta$ -Actin (ab8226, Abcam). Development of the blots was done with the SuperSignal West Pico PLUS Chemiluminescent Substrate (Thermo Fisher).

### **In vivo experiments**

The animal assay was manipulated under the authorization of the Animal Care Committee of Shanxi Province Cancer Hospital, Shanxi Hospital Affiliated to Cancer Hospital, Chinese Academy of Medical Sciences, Cancer Hospital Affiliated to Shanxi Medical University. For xenograft assay, 32 BALB/c nude mice (Vital River Laboratory, Beijing, China) were randomly divided into 4 groups (8 mice in each group) and then processed as follows: (1) mice were injected with sh-NC-transduced H1299/DDP cells ( $4 \times 10^6$ ) and administered with phosphate-buffered saline or DDP; (2) mice were injected with sh-circ\_0017639-transduced H1299/DDP cells ( $4 \times 10^6$ ) and administered with phosphate-buffered saline or DDP. For DDP treatment, mice were intraperitoneally administered with DDP (5 mg/kg) every 3 days from day 8. Tumor volume was measured every 3 days ( $\text{Volume} = (\text{length} \times \text{width}^2)/2$ ) from the beginning of DDP administration. Xenograft tumors were excised, weighed, and paraffin-embedded after injection for 23 days. Paraffin-embedded tissues were cut (5.0  $\mu$ m sections) and subjected to immunohistochemistry (IHC) to detect the level of SIX1 as described previously [34]. An antibody against SIX1 (PA5-51,654, 1:500, Thermo Fisher) was used for IHC analysis.

## Statistical analysis

At least three biological repeats were performed for each experiment. Data were analyzed using GraphPad Prism 8 software (GraphPad, La Jolla, CA, USA) and presented as mean  $\pm$  standard deviation. Correlation was evaluated using Pearson's correlation analysis. Significance was determined using the Student's *t*-test or analysis of variance. If the *P*-value was below 0.05, the results were considered statistically significant.

## Results

The current research aimed to investigate the action of circ\_0017639 in the resistance of NSCLC to DDP. Our findings demonstrated that circ\_0017639 promoted DDP resistance and tumor progression via sequestering miR-1296-5p and subsequent elevating SIX1 expression. This study provided new insights into DDP resistance and highlighted a potential target for improving DDP therapy in NSCLC.

### Identification of circ\_0017639 expression in DDP-resistant and DDP-sensitive NSCLC samples

To identify the potential circRNAs associated with the DDP resistance of NSCLC, we analyzed the GSE112214 microarray. Heatmap showed the 12 observably upregulated circRNAs in GSE112214 ( $P < 0.001$ ) (Figure 1(a)). The difference of has\_circRNA\_100542 (circ\_0017639) was more obvious than other circRNAs and it had been reported to play an oncogenic role in gastric cancer [13]. Therefore, we speculated that circ\_0017639 might be involved in the regulation of lung cancer progression. A diagram showed the level of circ\_0017639 in 3 NSCLC samples and 3 matching adjacent normal samples in the GSE112214 microarray (Figure 1(b)). We also observed that circ\_0017639 was overexpressed in 41 NSCLC samples (with respect to adjacent normal samples) and 4 NSCLC cell lines (with respect to HBE cells) (Figure 1(c-d)). Notably, circ\_0017639 expression was significantly elevated in 21 DDP-resistant samples compared to 20 DDP-sensitive samples (Figure 1(e)). We then constructed DDP-resistant NSCLC cell lines and

their IC<sub>50</sub> values were about 3 times greater than that of their parental cells (Figure 1(f-h)). There was an apparent elevation in circ\_0017639 expression in DDP-resistant cells respective to their parental cells (Figure 1(i)). As exhibited in Figure 1(j)-1 K, circ\_0017639 was resistant to RNase R treatment in contrast to linear scm like with four mbt domains 2 (SFMBT2). Subcellular fractionation assay showed that the cellular distribution of circ\_0017639 was mainly distributed in the cytoplasm of DDP-resistant NSCLC cells, suggesting that circ\_0017639 might function as a miRNA molecular sponge (Figure 1l-1 M). Overall, these results manifested that higher levels of circ\_0017639 might be related to the DPP resistance of NSCLC.


### Circ\_0017639 silencing reduced the resistance and malignancy of DDP-resistant NSCLC cells

To test the potential role of circ\_0017639 in the DDP resistance of NSCLC, we transfected si-circ\_0017639 into DDP-resistant NSCLC cells to silence circ\_0017639 expression. The results exhibited a lower level in circ\_0017639 expression after si-circ\_0017639 transfection than the control group (Figure 2(a)). Moreover, circ\_0017639 inhibition resulted in a marked decrease in the IC<sub>50</sub> value and viability of DDP-resistant cells (Figure 2(b-c)). We also observed that the number of colonies and EDU-positive cells was less in the circ\_0017639-inhibiting DDP-resistant cells (Figure 2(d,e)). Also, the apoptotic rate of circ\_0017639-inhibiting DDP-resistant NSCLC cells was elevated (Figure 2(f)). Furthermore, the invasion and migration abilities of DDP-resistant cells were repressed after circ\_0017639 knockdown (Figure 2(g-h)). In addition, circ\_0017639 inhibition decreased the MMP9 protein level, whereas elevated the Cleaved-caspase-3 protein level in DDP-resistant NSCLC cells (Figure 2(i-j)). Collectively, these results manifested that circ\_0017639 inhibition decreased the resistance and malignancy of DDP-resistant NSCLC cells.

### Circ\_0017639 acted as a miR-1296-5p sponge

It has been proved that circRNAs can exert functions via sponging miRs [35]. Bioinformatics






**Figure 2. Influence of circ\_0017639 inhibition on the sensitivity and malignancy of DDP-resistant NSCLC cells.** (a) RT-qPCR analysis of circ\_0017639 in DDP-resistant NSCLC cells transfected with si-circ\_0017639 or si-NC. (b and c) MTT assay assessed the IC<sub>50</sub> value and viability of circ\_0017639-inhibiting DDP-resistant NSCLC cells and control cells. (d-h) The colony formation, proliferation, apoptosis, invasion, and migration of circ\_0017639-inhibiting DDP-resistant NSCLC cells and control cells were analyzed by colony formation, EDU, flow cytometry, transwell invasion, and wound-healing assays. (i and j) Western blotting detected MMP9 and Cleaved-caspase-3 protein levels in DDP-resistant NSCLC cells transfected with si-circ\_0017639 or si-NC. \*\**P* < 0.01, \*\*\**P* < 0.001, and \*\*\*\**P* < 0.0001.

analysis (circinteractome) predicted that there were many miRNAs that might bind to circ\_0017639. Through literature review, seven miRNAs (miR-1208 [36], miR-1305 [37], miR-188-3p [38], miR-198-5p [39], miR-224-5p [40], miR-885-3p [41] and miR-1296-5p [42]) that are lowly expressed in lung cancer and inhibit the progression of lung cancer were selected for further analysis. Also, silenced circ\_0017639 expression markedly increased the expression levels of miR-1305, miR-224-5p, miR-885-3p and miR-1296-5p, and miR-1296-5p with the largest differential change was selected as the candidate miRNA (supplementary Fig. 1A). The binding sites between miR-1296-5p and circ\_0017639 are provided in Figure 3(a). We then performed

a dual-luciferase reporter assay to examine whether miR-1296-5p target circ\_0017639. The overexpression efficiency of miR-1296-5p mimic is displayed in Figure 3(b). Also, miR-1296-5p mimic repressed the luciferase activity of WT-circ\_0017639 in DDP-resistant NSCLC cells, but the luciferase activity of MUT-circ\_0017639 was not affected (Figure 3(c-d)). We also observed lower levels of miR-1296-5p in DDP-resistant samples, and its expression had a negative correlation with circ\_0017639 (Figure 3(e,f)). Similar results were noted in DDP-resistant cell lines (in contrast to their parental cells) and NSCLC cells (in contrast to HBE cells) (Figure 3(g)). The overexpression efficiency of circ\_0017639 was confirmed by RT-qPCR, as displayed in Figure 3(h). Moreover,



**Figure 3. Identification of circ\_0017639 as a miR-1296-5p sponge.** (a) Schematic illustrating the sequence alignment of circ\_0017639 with miR-1296-5p. (b) Analysis of miR-1296-5p in DDP-resistant NSCLC cells with miR-1296-5p mimic or miR-NC. (c and d) Dual-luciferase reporter assay analysis of the luciferase activities of WT-circ\_0017639 and MUT-circ\_0017639 plasmids in the presence of miR-1296-5p mimic or miR-NC in DDP-resistant NSCLC cells. (e) Expression of miR-1296-5p in 21 DDP-resistant NSCLC samples and 20 NSCLC DDP-sensitive NSCLC samples was analyzed by RT-qPCR. (f) Pearson's correlation analysis uncovered the correlation between miR-1296-5p and circ\_0017639 in DDP-resistant NSCLC samples. (g) RT-qPCR analysis of miR-1296-5p in DDP-resistant NSCLC cells (in contrast to their parent cells) and NSCLC cells (in contrast to HBE cells). (h) RT-qPCR analysis of circ\_0017639 in DDP-resistant NSCLC cells with circ\_0017639 or pCD5-ciR. (i) Effects of circ\_0017639 inhibition and overexpression on miR-1296-5p expression in DDP-resistant NSCLC cells. \*\* $P < 0.01$ , \*\*\* $P < 0.001$ , and \*\*\*\* $P < 0.0001$ .



circ\_0017639 inhibition elevated miR-1296-5p expression, but circ\_0017639 overexpression had the opposing effect (Figure 3(i)). Together, circ\_0017639 served as a miR-1296-5p sponge.

### ***Circ\_0017639 regulated the resistance and malignancy of DDP-resistant NSCLC cells through sponging miR-1296-5p***

Given that circ\_0017639 acted as a miR-1296-5p sponge, we further investigated whether circ\_0017639 regulated DDP-resistant NSCLC cell resistance through binding to miR-1296-5p. Transfection with anti-miR-1296-5p caused an observable decrease in miR-1296-5p expression in DDP-resistant cells (Figure 4(a)). Moreover, the elevation of miR-1296-5p in circ\_0017639-inhibiting DDP-resistant NSCLC cells was substantially restored after miR-1296-5p knockdown (Figure 4(b)). Silenced miR-1296-5p expression restored circ\_0017639 knockdown-mediated decrease of the IC<sub>50</sub> value and viability of DDP-resistant cells (Figure 4(c-d)). Furthermore, the inhibiting effect of circ\_0017639 knockdown on DDP-resistant cell colony formation and proliferation was offset after anti-miR-1296-5p introduction (Figure 4(e-f)). As expected, miR-1296-5p inhibition counteracted the accelerating impact of circ\_0017639 silencing on DDP-resistant cell apoptosis (Figure 4(g)). Additionally, decreased miR-1296-5p expression effectively abolished the inhibitory impact of circ\_0017639 knockdown on DDP-resistant cell invasion and migration (Figure 4(h-i)). In addition, protein levels of MMP9 and Cleaved-caspase-3 in DDP-resistant NSCLC cells mediated by circ\_0017639 inhibition were restored after miR-1296-5p inhibitor introduction (Figure 4(j-k)). In sum, circ\_0017639 sponged miR-1296-5p to regulate DDP-resistant NSCLC cell resistance.


### ***MiR-1296-5p directly targeted SIX1***

To seek the downstream targets of miR-1296-5p, an online software targetScan was utilized. Among the many predicted targets, seven genes (SOX4 [43], ZEB2 [44], AVL9 [45], PTPN9 [46], RAB22A [47], ITGB8 [48] and SIX1 [49]) that have been reported to play an oncogenic role in


lung cancer were further analyzed. In addition, miR-1296-5p overexpression significantly reduced the expression levels of ZEB2 and SIX1, especially SIX1 (Supplementary Fig. 1B). Figure 5(a) exhibits the potential binding between miR-1296-5p and SIX1. Moreover, miR-1296-5p mimic decreased the luciferase activity of the WT-SIX1 3'UTR plasmid, but there was no difference in the MUT-SIX1 3'UTR plasmid (Figure 5(b-c)). By consulting the gene expression profiling interactive analysis (GEPIA) database, we found that SIX1 was highly expressed in NSCLC patients (Figure 5(d)). Consistently, SIX1 was overexpressed in DDP-resistant samples (Figure 5(e)). Furthermore, the expression of SIX1 mRNA was negatively correlated with miR-1296-5p in DDP-resistant NSCLC samples (Figure 5(f)). We also observed a higher level of SIX1 protein in three DDP-resistant NSCLC samples than that in DDP-sensitive NSCLC samples (Figure 5(g)). In addition, higher levels of protein SIX1 were gained in NSCLC cells and DDP-resistant cells in contrast to their corresponding control cells (Figure 5(h)). As expected, miR-1296-5p overexpression decreased SIX1 protein levels in DDP-resistant cells, but miR-1296-5p inhibition had an opposing effect (Figure 5(i)). Overall, SIX1 acted as a miR-1296-5p target.

### ***MiR-1296-5p decreased DDP-resistant NSCLC cell resistance and malignancy via targeting SIX1***

We then further addressed whether miR-1296-5p performs its function by targeting SIX1. Protein levels of SIX1 were prominently increased in DDP-resistant NSCLC cells after SIX1 transfection (Figure 6(a)). The decreased protein level of SIX1 induced by miR-1296-5p mimic was reversed by forcing the expression of SIX1 (Figure 6(b)). Moreover, elevated SIX1 expression abolished the decreased IC<sub>50</sub> value and viability of DDP-resistant cells caused by miR-1296-5p overexpression (Figure 6(c-d)). Also, miR-1296-5p mimic repressed DDP-resistant cell colony formation and proliferation, and induced DDP-resistant cell apoptosis, but these trends induced by miR-1296-5p mimic were restored after SIX1 introduction (Figure 6(e-g)). In addition, miR-1296-5p elevation mediated repression on DDP-resistant cell



**Figure 4. Circ\_0017639 sponged miR-1296-5p to regulate DDP-resistant NSCLC cell sensitivity and malignancy.** (a) Analysis of miR-1296-5p in DDP-resistant NSCLC cells transfected with anti-miR-NC or anti-miR-1296-5p by RT-qPCR. (b) Influence of miR-1296-5p inhibitor on circ\_0017639-inhibiting DDP-resistant NSCLC cells. (c and d) Effects of miR-1296-5p silencing on the IC<sub>50</sub> value and viability of DDP-resistant NSCLC cells were assessed by MTT assay. (e-i) Impacts of miR-1296-5p knockdown on colony formation, proliferation, apoptosis, invasion, and migration of circ\_0017639-inhibiting DDP-resistant NSCLC cells were analyzed by colony formation, EDU, flow cytometry, transwell invasion, and wound-healing assays. (j and k) Influence of miR-1296-5p silencing on MMP9 and Cleaved-caspase-3 protein levels in circ\_0017639-inhibiting DDP-resistant NSCLC cells. \**P* < 0.05, \*\**P* < 0.01, \*\*\**P* < 0.001, \*\*\*\**P* < 0.0001.



**Figure 5. SIX1 as a miR-1296-5p target was verified.** (a) Schematic illustrating the sequence alignment of SIX1 3'UTR with miR-1296-5p. (b and c) Dual-luciferase reporter assay assessed the luciferase activity of MUT-SIX1 3'UTR and MUT-SIX1 3'UTR plasmids in DDP-resistant NSCLC cells transfected with miR-1296-5p mimic or miR-NC. (d) The level of SIX1 mRNA in NSCLC through consulting the GEPIA database. (e) Analysis of SIX1 mRNA in 21 DDP-resistant NSCLC samples and 20 DDP-sensitive NSCLC samples by RT-qPCR. (f) Correlation of SIX1 mRNA and miR-1296-5p in DDP-resistant NSCLC samples was analyzed by Pearson's correlation analysis. (g and h) Protein level of SIX1 in DDP-resistant NSCLC tissues, NSCLC cells, and DDP-resistant NSCLC cells was detected. (i) Impacts of miR-1296-5p mimic and inhibitor on SIX1 protein levels in DDP-resistant NSCLC cells. \*\*\* $p < 0.001$  and \*\*\*\* $p < 0.0001$ .

migration and invasion was counteracted after SIX1 overexpression (Figure 6(h-i)). As expected, forced SIX1 expression overturned miR-1296-5p overexpression-mediated effects on protein levels of MMP9 and Cleaved-caspase-3 (Figure 6(j-k)). Collectively, miR-1296-5p targeted SIX1 to decrease the resistance and malignancy of DDP-resistant NSCLC cells.


#### **Circ\_0017639 regulated SIX1 expression via adsorbing miR-1296-5p**

Next, we further surveyed whether circ\_0017639 regulates SIX1 expression via adsorbing miR-1296-5p. Circ\_0017639 silencing reduced SIX1 at


mRNA and protein levels, but this decrease was reversed after anti-miR-1296-5p introduction (Figure 7(a-b)). These data manifested that Circ\_0017639 regulated SIX1 expression via sponging miR-1296-5p.

#### **Circ\_0017639 inhibition decreased xenograft tumor growth and sensitized cells to DDP in vivo**

To further support the above findings, we intraperitoneally administered DDP or PBS into nude mice injected with H1299/DDP cells carrying sh-circ\_0017639 or sh-NC. Injection with H1299/DDP cells carrying sh-circ\_0017639 decreased



**Figure 6. MiR-1296-5p decreased DDP-resistant NSCLC cell resistance and malignancy via targeting SIX1.** (a) After SIX1 or pcDNA transfection, the SIX1 protein level was analyzed. (b) Influence of SIX1 introduction on the SIX1 protein level in miR-1296-5p-overexpressed DDP-resistant NSCLC cells was evaluated using western blotting. (c and d) Influence of SIX1 overexpression on the IC<sub>50</sub> value and viability of miR-1296-5p-elevated DDP-resistant NSCLC cells was evaluated by MTT assay. (e-i) Impacts of SIX1 upregulation on colony formation, proliferation, apoptosis, invasion, and migration of miR-1296-5p-overexpressed DDP-resistant NSCLC cells were analyzed by colony formation, EDU, flow cytometry, transwell invasion, and wound-healing assays. (j and k) Effects of SIX1 upregulation on protein levels of MMP9 and Cleaved-caspase-3 in miR-1296-5p-overexpressed DDP-resistant NSCLC cells were evaluated by western blotting. \*\**P* < 0.01, \*\*\**P* < 0.001, and \*\*\*\**P* < 0.0001.




**Figure 7. Circ\_0017639 regulated SIX1 expression via adsorbing miR-1296-5p.** (a and b) Analysis of SIX1 at mRNA and protein levels in DDP-resistant NSCLC cells transfected with si-NC, si-circ\_0017639, si-circ\_0017639+ anti-miR-NC, or si-circ\_0017639+ anti-miR-1296-5p by RT-qPCR or western blotting. \*\* $P < 0.01$ , \*\*\* $P < 0.001$ , and \*\*\*\* $P < 0.0001$ .

xenograft tumor growth and sensitized cells to DDP treatment (Figure 8(a-b)). Furthermore, the levels of circ\_0017639 and SIX1 protein were decreased in sh-circ\_0017639+ PBS and sh-circ\_0017639+ DDP groups compared to their matched control group, but miR-1296-5p expression had an opposite tendency (Figure 8(c-d)). Also, the number of SIX1-positive cells was less in sh-circ\_0017639+ PBS and sh-circ\_0017639+ DDP groups with respect to their matched control group (Figure 8(e)). Together, circ\_0017639 inhibition decreased xenograft tumor growth and sensitized cells to DDP *in vivo*.

## Discussion

Since DDP-induced resistance leads to treatment failure of NSCLC, it is essential to continue to explore the mechanism of DDP resistance in NSCLC. Our research described that circ\_0017639 conferred DDP resistance through modulating the miR-1296-5p/SIX1 pathway in NSCLC.

CircRNAs have been demonstrated to exert essential roles in the response of cancer cells to DDP treatment [50]. Herein, we searched for a differentially expressed circRNA circ\_0017639 in NSCLC samples from the GSE112214 dataset.



**Figure 8. Knockdown of circ\_0017639 decreased xenograft tumor growth and sensitized cells to DDP *in vivo*.** (a and b) The average tumor volume and weight of mice in sh-NC+PBS, sh-circ\_0017639+ PBS, sh-NC+DDP, sh-circ\_0017639+ DDP groups. (c and d) Analysis of circ\_0017639, miR-1296-5p, and SIX1 protein in xenograft tumors in the above groups was done. (e) Analysis of SIX1 in xenograft tumors in the above groups by IHC. \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ , and \*\*\*\* $P < 0.0001$ .

The report of Li *et al.* exposed that circ\_0017639 sponged miR-224-5p to elevate ubiquitin-specific protease 3 expression, resulting in elevating cell metastasis and proliferation in gastric cancer [13]. Nevertheless, circ\_0017639 exerted an inhibiting impact on glioma cell metastasis and proliferation, which might be related to the tissue-specific expression [51]. A recent report showed

that circ\_0017639 was highly expressed in NSCLC samples and cell lines, and circ\_0017639 overexpression promoted NSCLC cell proliferation, invasion, and migration via the PI3K/AKT signaling [52]. The difference between our study and the previous report [52] was to explore the function of circ\_0017639 in DDP resistance in NSCLC. Our data showed higher levels of circ\_0017639 in

DDP-resistant samples and cell lines. Loss-of-function assays indicated that circ\_0017639 silencing elevated DDP sensitivity of DDP-resistant NSCLC cells in xenograft models and facilitated DDP-resistant NSCLC cell apoptosis and decreased DDP-resistant NSCLC cell proliferation, invasion, and migration *in vitro*. These results suggested the contribution of circ\_0017639 in NSCLC resistance to DDP.

Considering that circRNAs can function as miR sponges [35]. We observed that circ\_0017639 was mainly distributed in the cytoplasm of DDP-resistant NSCLC cells through subcellular fractionation assay, implying that circ\_0017639 might function as a miRNA molecular sponge. Furthermore, circ\_0017639 as a miR-1296-5p sponge was identified through bioinformatics analysis and dual-luciferase reporter assay. Researchers had proved that miR-1296-5p could function as an inhibitor in gastric cancer by targeting cyclin-dependent kinase 6 and epidermal growth factor receptor [53], ERBB2-positive breast cancer by downregulating ERBB2 [17], in osteosarcoma through repressing notch receptor 2 expression [16]. A recent study described that miR-1296 repressed NSCLC cell invasion by blocking the Wnt signaling [54]. Our data exhibited that miR-1296-5p inhibitor could restore circ\_0017639 silencing-mediated impact on DDP-resistant NSCLC cell resistance. Thus, we concluded that circ\_0017639 regulated DDP resistance and tumor growth through sponging and repressing miR-1296-5p in NSCLC.

Subsequently, we discovered SIX1 as a miR-1296-5p target through bioinformatics analysis and dual-luciferase reporter assay. SIX1 could induce vascular endothelial growth factor-C expression, leading to cell migration in breast cancer [55]. It has been reported that cyclin A1 and cyclin D1 were transcriptional targets of SIX1 and that SIX1 promoted cell proliferation in breast cancer by increasing cyclin A1 expression [56] and in pancreatic cancer via increasing cyclin D1 expression [57]. Xia *et al.* revealed that the upregulation of miR-204 constrained NSCLC cell invasion and proliferation through targeting SIX1 [58]. Moreover, miR-186-5p decreased DDP resistance via downregulating

SIX1 in NSCLC [49]. Also, FOXD2 adjacent opposite strand RNA 1 increased SIX1 expression by adsorbing miR-185-5p, thereby conferring DDP resistance of NSCLC [25]. Our data showed the upregulation of SIX1 in DDP-resistant NSCLC samples and cell lines, and SIX1 overexpression lessened the repressive effect of miR-1296-5p overexpression on DDP resistance in DDP-resistant NSCLC cells, manifesting that miR-1296-5p could regulate DDP resistance via targeting SIX1 in DDP-resistant NSCLC cells. Notably, circ\_0017639 could regulate SIX1 expression via serving as a decoy of miR-1296-5p. Thus, we inferred that circ\_0017639 regulated DDP resistance by regulating the miR-1296-5p/SIX1 pathway in NSCLC.

## Conclusion

Our findings highlighted the promoting function of circ\_0017639 in DDP resistance of NSCLC. Moreover, circ\_0017639 upregulation elevated SIX1 expression through adsorbing miR-1296-5p via acting as a miR-1296-5p molecular sponge, thereby conferring DDP resistance in NSCLC. The research provided the promoting effect of circ\_0017639 on DDP resistance in NSCLC.

## Highlights:

- (1) Circ\_0017639 was overexpressed in DDP-resistant NSCLC samples and cells.
- (2) Inhibition of circ\_0017639 decreased DDP resistance of DDP-resistant NSCLC cells.
- (3) Circ\_0017639 functioned as a miR-1296-5p molecular sponge.
- (4) SIX1 served as a miR-1296-5p target.

## Author's Contribution

Feiyun Chang conceived and designed the study, and drafted the first draft of the manuscript. All experiments were completed by all authors. Jiali Li, Quan Sun, Shuqing Wei, Yongming Song analyzed and collated the results. All authors reviewed and critiqued the manuscript, and agreed to the final submission of the manuscript. All authors read and approved the final manuscript.

## Disclosure statement

No potential conflict of interest was reported by the author(s).

## Funding

The author(s) reported that there is no funding associated with the work featured in this article.

## Data Availability Statement

No new data were generated or analyzed in support of this research.

## ORCID

Yongming Song  <http://orcid.org/0000-0003-1363-8444>

## References

- [1] Siegel RL, Miller KD, Jemal A. Cancer statistics, 2020. *CA Cancer J Clin.* 2020;70(1):7–30.
- [2] Bagcchi S. Lung cancer survival only increases by a small amount despite recent treatment advances. *Lancet Respir Med.* 2017;5(3):169.
- [3] Hirsch FR, Scagliotti GV, Mulshine JL, et al. Lung cancer: current therapies and new targeted treatments. *Lancet.* 2017;389(10066):299–311.
- [4] Sosa Iglesias V, Giuranno L, Dubois LJ, et al. Drug resistance in non-small cell lung cancer: a potential for NOTCH targeting? *Front Oncol.* 2018;8:267.
- [5] Siegel RL, Miller KD, Jemal A. Cancer statistics, 2018. *CA Cancer J Clin.* 2018;68(1):7–30.
- [6] Ferlay J, Soerjomataram I, Dikshit R, et al. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer.* 2015;136(5):E359–E386.
- [7] Song L, Cui Z, Guo X. Comprehensive analysis of circular RNA expression profiles in cisplatin-resistant non-small cell lung cancer cell lines. *Acta Biochim Biophys Sin (Shanghai).* 2020;52(9):944–953.
- [8] Kristensen LS, Andersen MS, Stagsted LVW, et al. The biogenesis, biology and characterization of circular RNAs. *Nat Rev Genet.* 2019;20(11):675–691.
- [9] Salzman J, Chen RE, Olsen MN, et al. Cell-type specific features of circular RNA expression. *PLoS Genet.* 2013;9(9):e1003777.
- [10] Wang Y, Mo Y, Gong Z, et al. Circular RNAs in human cancer. *Mol Cancer.* 2017;16(1):25.
- [11] Zhang P-F, Pei X, K-S L, et al. Circular RNA circFGFR1 promotes progression and anti-PD-1 resistance by sponging miR-381-3p in non-small cell lung cancer cells. *Mol Cancer.* 2019;18(1):179.
- [12] Huang M-S, Liu J-Y, Xia X-B, et al. Hsa\_circ\_0001946 Inhibits Lung Cancer Progression and Mediates Cisplatin Sensitivity in Non-small Cell Lung Cancer via the Nucleotide Excision Repair Signaling Pathway. *Front Oncol.* 2019;9:508.
- [13] Li B, Jin M, Cao F, et al. Hsa\_circ\_0017639 expression promotes gastric cancer proliferation and metastasis by sponging miR-224-5p and upregulating USP3. *Gene.* 2020;750:144753.
- [14] Hansen TB, Jensen TI, Clausen BH, et al. Natural RNA circles function as efficient microRNA sponges. *Nature.* 2013;495(7441):384–388.
- [15] Krol J, Loedige I, Filipowicz W. The widespread regulation of microRNA biogenesis, function and decay. *Nat Rev Genet.* 2010;11(9):597–610.
- [16] Wang L, Hu K, Chao Y, et al. MicroRNA-1296-5p suppresses the proliferation, migration, and invasion of human osteosarcoma cells by targeting NOTCH2. *Journal of Cellular Biochemistry.* 2020;121(2):2038–2046.
- [17] Chen G, He M, Yin Y, et al. miR-1296-5p decreases ERBB2 expression to inhibit the cell proliferation in ERBB2-positive breast cancer. *Cancer Cell Int.* 2017;17(1):95.
- [18] Zang H, Li Y, Zhang X, et al. Circ\_0000517 Contributes to Hepatocellular Carcinoma Progression by Upregulating TXNDC5 via Sponging miR-1296-5p. *Biosci Rep.* 2020;12:3457–3468.
- [19] Wu W, Ren Z, Li P, et al. Six1: a critical transcription factor in tumorigenesis. *Int J Cancer.* 2015;136(6):1245–1253.
- [20] Christensen KL, Patrick AN, McCoy EL, et al. The six family of homeobox genes in development and cancer. *Adv Cancer Res.* 2008; 101:93–126.
- [21] Armat M, Oghabi Bakhshaiesh T, Sabzichi M, et al. The role of Six1 signaling in paclitaxel-dependent apoptosis in MCF-7 cell line. *Bosn J Basic Med Sci.* 2016;16(1):28–34.
- [22] Chen G, Chen Z, Zhao H. MicroRNA-155-3p promotes glioma progression and temozolomide resistance by targeting Six1. *Journal of Cellular and Molecular Medicine.* 2020;24(9):5363–5374.
- [23] Li B, Zhao S, Geng R, et al. The Six1 Homeobox Homolog 1 (SIX1) Gene Regulates Paclitaxel Resistance by Affecting Reactive Oxygen Species and Autophagy in Human Hepatocellular Carcinoma Cell Line HepG2. *J Cell Mol Med.* 2018;24:2271–2279.
- [24] Li Z, Tian T, Hu X, et al. Six1 mediates resistance to paclitaxel in breast cancer cells. *Biochem Biophys Res Commun.* 2013;441(3):538–543.
- [25] Ge P, Cao L, Yao YJ, et al. lncRNA FOXD2-AS1 confers cisplatin resistance of non-small-cell lung cancer via regulation of miR185-5p-SIX1 axis. *Onco Targets Ther.* 2019;12:6105–6117.
- [26] Chen K, Wei H, Pan J, et al. Six1 is negatively correlated with poor prognosis and reduces 5-fluorouracil sensitivity via attenuating the stemness of hepatocellular carcinoma cells. *Eur J Pharmacol.* 2019;861:172599.
- [27] J-m L, Yang F, Li J, et al. Reelin Promotes Cisplatin Resistance by Induction of Epithelial-Mesenchymal



- Transition via p38/GSK3 $\beta$ /Snail Signaling in Non-Small Cell Lung Cancer. *Med Sci Monit.* **2020**;26:e925298.
- [28] Liu H, Bi J, Dong W, et al. Invasion-related circular RNA circFNDC3B inhibits bladder cancer progression through the miR-1178-3p/G3BP2/SRC/FAK axis. *Mol Cancer.* **2018**;17(1):161.
- [29] Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2<sup>-Delta Delta C(T)</sup> Method. *Methods.* **2001**;25(4):402–408.
- [30] Xiao H, Liu Y, Liang P, et al. TP53TG1 enhances cisplatin sensitivity of non-small cell lung cancer cells through regulating miR-18a/PTEN axis. *Cell & Bioscience.* **2018**;8:23.
- [31] Wu H, Mu X, Liu L, et al. Bone marrow mesenchymal stem cells-derived exosomal microRNA-193a reduces cisplatin resistance of non-small cell lung cancer cells via targeting LRRC1. *Cell Death Dis.* **2020**;11(9):801.
- [32] Zhang Y, Zhang Q, Chen H, et al. BCL9 promotes epithelial mesenchymal transition and invasion in cisplatin resistant NSCLC cells via  $\beta$ -catenin pathway. *Life Sci.* **2018**;208:284–294.
- [33] Chen X, Mao R, Su W, et al. Circular RNA circHIPK3 modulates autophagy via MIR124-3p-STAT3-PRKAA/AMPK $\alpha$  signaling in STK11 mutant lung cancer. *Autophagy.* **2020**;16(4):659–671.
- [34] Chen S, Wang L, Xu C, et al. Knockdown of REGY inhibits proliferation by inducing apoptosis and cell cycle arrest in prostate cancer. *Am J Transl Res.* **2017**;9(8):3787–3795.
- [35] Zhong Y, Du Y, Yang X, et al. Circular RNAs function as ceRNAs to regulate and control human cancer progression. *Mol Cancer.* **2018**;17(1):79.
- [36] Huang M, Li T, Wang Q, et al. Silencing circPVT1 enhances radiosensitivity in non-small cell lung cancer by sponging microRNA-1208. *Cancer Biomarkers.* **2021**;31(3):263–279.
- [37] Cai Y, Hao Y, Ren H, et al. miR-1305 Inhibits The Progression Of Non-Small Cell Lung Cancer By Regulating MDM2. *Cancer Management and Research.* **2019**;11:9529–9540.
- [38] Yao J, Xu G, Zhu L, et al. circGFRA1 Enhances NSCLC Progression by Sponging miR-188-3p. *Onco Targets Ther.* **2020**;13:549–558.
- [39] Liang YY, Huang JC, Tang RX, et al. Clinical value of miR-198-5p in lung squamous cell carcinoma assessed using microarray and RT-qPCR. *World Journal of Surgical Oncology.* **2018**;16(1):22.
- [40] Zhou J, Wang H, Sun Q, et al. miR-224-5p-enriched exosomes promote tumorigenesis by directly targeting androgen receptor in non-small cell lung cancer. *Mol Ther Nucleic Acids.* **2021**;23:1217–1228.
- [41] Cao J, Geng J, Chu X, et al. miRNA-885-3p inhibits docetaxel chemoresistance in lung adenocarcinoma by down-regulating Aurora A. *Oncol Rep.* **2019**;41(2):1218–1230.
- [42] Xu Q, Liu X, Liu Z, et al. MicroRNA-1296 inhibits metastasis and epithelial-mesenchymal transition of hepatocellular carcinoma by targeting SRPK1-mediated PI3K/AKT pathway. *Mol Cancer.* **2017**;16(1):103.
- [43] Medina PP, Castillo SD, Blanco S, et al. The SRY-HMG box gene, SOX4, is a target of gene amplification at chromosome 6p in lung cancer. *Hum Mol Genet.* **2009**;18(7):1343–1352.
- [44] Wu DM, Zhang T, Liu YB, et al. The PAX6-ZEB2 axis promotes metastasis and cisplatin resistance in non-small cell lung cancer through PI3K/AKT signaling. *Cell Death Dis.* **2019**;10(5):349.
- [45] Liang J, Sun T, Wang G, et al. Clinical significance and functions of miR-203a-3p/AVL9 axis in human non-small-cell lung cancer. *Personalized Medicine.* **2020**;17(4):271–282.
- [46] Du WW, Fang L, Li M, et al. MicroRNA miR-24 enhances tumor invasion and metastasis by targeting PTPN9 and PTPRF to promote EGF signaling. *J Cell Sci.* **2013**;126(Pt 6):1440–1453.
- [47] Zhou Y, Wu B, Li JH, et al. Rab22a enhances CD147 recycling and is required for lung cancer cell migration and invasion. *Exp Cell Res.* **2017**;357(1):9–16.
- [48] Wu P, Wang Y, Wu Y, et al. Expression and prognostic analyses of ITGA11, ITGB4 and ITGB8 in human non-small cell lung cancer. *PeerJ.* **2019**;7:e8299.
- [49] Liu X, Zhou X, Chen Y, et al. miR-186-5p targeting SIX1 inhibits cisplatin resistance in non-small-cell lung cancer cells (NSCLCs). *Neoplasma.* **2020**;67(1):147–157.
- [50] Cui C, Yang J, Li X, et al. Functions and mechanisms of circular RNAs in cancer radiotherapy and chemotherapy resistance. *Mol Cancer.* **2020**;19(1):58.
- [51] Zhang S, Qin W, Yang S, et al. Circular RNA SFMBT2 Inhibits the Proliferation and Metastasis of Glioma Cells Through Mir-182-5p/Mtss1 Pathway. *Technol Cancer Res Treat.* **2020**;19:1533033820945799.
- [52] Zhang HB, Qiu XM, Zhang YC, et al. Circ\_0017639 facilitates proliferative, migratory, and invasive potential of non-small cell lung cancer (NSCLC) cells via PI3K/AKT signaling pathway. *Bioengineered.* **2022**;13(1):1590–1601.
- [53] Jia Y, Zhao LM, Bai HY, et al. The tumor-suppressive function of miR-1296-5p by targeting EGFR and CDK6 in gastric cancer. *Biosci Rep.* **2019**;39(1):BSR20181556.
- [54] Deng H, Xie C, Ye Y, et al. MicroRNA-1296 expression is associated with prognosis and inhibits cell proliferation and invasion by Wnt signaling in non-small cell lung cancer. *Oncol Lett.* **2020**;19(1):623–630.
- [55] Wang CA, Jedlicka P, Patrick AN, et al. SIX1 induces lymphangiogenesis and metastasis via upregulation of VEGF-C in mouse models of breast cancer. *J Clin Invest.* **2012**;122(5):1895–1906.
- [56] Coletta RD, Christensen K, Reichenberger KJ, et al. The Six1 homeoprotein stimulates tumorigenesis by reactivation of cyclin A1. *Proc Natl Acad Sci U S A.* **2004**;101(17):6478–6483.
- [57] Li Z, Tian T, Lv F, et al. Six1 promotes proliferation of pancreatic cancer cells via upregulation of cyclin D1 expression. *PloS one.* **2013**;8(3):e59203.
- [58] Xia Y, Zhu Y, Ma T, et al. miR-204 functions as a tumor suppressor by regulating SIX1 in NSCLC. *FEBS Lett.* **2014**;588(20):3703–3712.