

## RESEARCH ARTICLE

# Prevalence of refractive errors and risk factors for myopia among schoolchildren of Almaty, Kazakhstan: A cross-sectional study

Ainagul Mukazhanova<sup>1,2</sup>, Neilya Aldasheva<sup>1</sup>, Juldyz Iskakbayeva<sup>1</sup>, Raushan Bakhytbek<sup>1</sup>, Aliya Ualiyeva<sup>2</sup>, Kaini Baigonova<sup>2</sup>, Damet Ongarbaeva<sup>2</sup>, Denis Vinnikov<sup>2,3\*</sup>

**1** Kazakh Eye Research Institute, Almaty, Kazakhstan, **2** Faculty of Medicine and Health Care, Al-Farabi Kazakh National University, Almaty, Kazakhstan, **3** Peoples' Friendship University of Russia (RUDN University), Moscow, Russian Federation

\* [denisvinnikov@mail.ru](mailto:denisvinnikov@mail.ru)



## Abstract

### Introduction

Very little is known about the prevalence of refractive errors among children in Kazakhstan. The aim of this study was to investigate the prevalence of refractive errors and risk factors of myopia among schoolchildren in Almaty, Kazakhstan.

### Methods

In the cross-sectional study of 2293 secondary school students (age 6–16), we examined cycloplegic autorefraction and offered a questionnaire in three age groups: 1<sup>st</sup> grade (N = 769), 5<sup>th</sup> grade (N = 768) and 9<sup>th</sup> grade (N = 756). The questionnaire covered main risk factors such as parental myopia, screen time, time outdoors, sports activities, near work, gender, grade, and school shift. Adjusted logistic regression analysis was applied to test the association of risk factors with myopia.

### Results

The mean spherical equivalent (SER) was  $-0.54 \pm 1.51$  diopters (D). The overall prevalence of refractive errors was 31.6% (95% confidence interval (CI) 29.7; 33.5); myopia 28.3% (95% CI 26.5; 30.1); hyperopia 3.4% (95% CI 2.7–4.1) and astigmatism 2.8% (95% CI 2.1; 3.5). In the multivariate adjusted regression analysis, higher class level (5<sup>th</sup> grade (odds ratio (OR) 1.78; 95% CI 1.26; 2.52) and 9<sup>th</sup> grade (OR 3.34; 95% CI 2.31; 4.82)) were associated with myopia, whereas outdoors activity more than 2 hours a day (OR 0.64; 95% CI 0.46; 0.89) and sports (OR 0.70; 95% CI 0.52; 0.93) were associated with a lower incidence of myopia.

### Conclusions

Myopia is a leading refractive error in schoolchildren in Almaty, Kazakhstan. Myopia prevention measures, including more time outdoors, should guide public health interventions in this population.

## OPEN ACCESS

**Citation:** Mukazhanova A, Aldasheva N, Iskakbayeva J, Bakhytbek R, Ualiyeva A, Baigonova K, et al. (2022) Prevalence of refractive errors and risk factors for myopia among schoolchildren of Almaty, Kazakhstan: A cross-sectional study. PLoS ONE 17(6): e0269474. <https://doi.org/10.1371/journal.pone.0269474>

**Editor:** Manbir Nagra, Anglia Ruskin University - Cambridge Campus, UNITED STATES

**Received:** October 6, 2021

**Accepted:** May 20, 2022

**Published:** June 3, 2022

**Peer Review History:** PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0269474>

**Copyright:** © 2022 Mukazhanova et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper.

**Funding:** The authors received no specific funding for this work.

**Competing interests:** The authors have declared that no competing interests exist.

## Introduction

Uncorrected refractive errors are the leading cause of moderate to severe visual impairment worldwide and the second most common cause of avoidable blindness [1, 2]. Studies on the prevalence of different types of refractive errors among children in different parts of the world are inconsistent and affected by age, gender, geography, and ethnicity [3–8]. Childhood myopia takes the lead in some countries of Southeast Asia with prevalence reaching 80% among adolescents, whereas hyperopia in children may be most prevalent in the Americas [9, 10]. At present, myopia, in particular childhood myopia, is a major public health issue, which in recent years has grown into an epidemic [4, 11, 12]. Many parts of the world have faced a dramatic increase in the number of people with myopia, especially among children and adolescents in Eastern Asian countries, in recent decades [10, 13–18].

The greatest burden of refractive error is myopia, with high prevalence rates in school-age children and adolescents, even greater in those with higher attained education [13, 19, 20]. According to Holden et al., half the world population (49.8%) will be myopic by the year 2050 and about 9.8% of people will have high myopia [11]. Childhood myopia, especially its early manifestation, increases the risk of complications, such as amblyopia, cataract, glaucoma, retinal detachment and myopic macular degeneration [21–23].

The studies of refractive errors from the countries of the former Soviet Union, especially those situated in Central Asia, are not very abundant, whereas the data from local studies are not available for the international audience. Sporadic reports from the countries of the former Soviet Union show a great variability in myopia prevalence. For example, the prevalence of myopia among urban children in Armenia was 23.3% in 2017 [24], but much greater in urban adolescents in Azerbaijan (34.7%) [25]. The prevalence of myopia in Russia also varies widely from 5.1% to 50.7%, depending on the region and age of the subjects [26–28]. Inconsistent data from the countries across the former Soviet Union may result from varying geographical and climatic conditions in addition to differences in ethnic and cultural composition of the peoples living in these countries [29, 30].

In the last 20 years, a number of studies reported the prevalence of visual impairments in school-age children in Kazakhstan [31–35]. In 2001, a pediatric examination in Almaty showed that 7.4% school-age children had any visual impairments, whereas the prevalence of refractive errors was not studied [34]. A cross-sectional study in the 2004 in Almaty, Kazakhstan revealed a 21% prevalence of refractive errors in schoolchildren, whereas myopia was confirmed in 14%, hyperopia in 3%, astigmatism in 1% and accommodation disorders in 3% of the studied population [31]. Another study (2010) reported that myopic refraction was found in 12% of rural school children and 22% among their urban counterparts [35]. Recently, there were only a few studies of the prevalence and structure of refractive errors among schoolchildren in Kazakhstan.

Therefore, the aim of this study was to ascertain the prevalence of refractive errors and predictors of myopia among school-age children and adolescents living in a metropolitan area of Almaty, Kazakhstan.

## Materials and methods

### Study population and sampling

It was a school-based cross-sectional study conducted from September to December 2019 in Almaty (Kazakhstan). From 1929 to 1997 Almaty was the capital of Kazakhstan. At present, Almaty remains the largest city in Kazakhstan with an area of 682 square kilometers and with a population of about 1.85 million inhabitants (about 10% of the country's total population). In 2018/2019, there were about 470,000 (25.3%) children 17 years old and younger living in Almaty [36, 37].

The educational system of Kazakhstan includes pre-school (pre-primary) education (kindergartens or nursery schools), secondary education, comprising schools for children 6 to 17 years old and higher education which includes universities, institutes, academies, etc. Public schools of Kazakhstan have a double-shift system, in which two separate groups of pupils attend schools during a school day. The first shift is in class from early morning until mid-day, whereas the second shift usually attends from mid-day to late afternoon.

This cross-sectional school-based study was performed according to the protocol of Refractive Error Study in Children (RESC). This protocol was designed to standardize the methodology used to obtain prevalence data on childhood refractive errors [38]. The sample size was calculated as  $N = \frac{Z^2 * (1-P) * P}{(B * P)^2}$ , where P is the anticipated prevalence of myopia, B is the desired error bound.

Prompted by previous research, the anticipated refractive error prevalence was 20% with a 15% error rate and a 95% confidence interval [35]. With a simple random sampling, 683 children were required for each class-level stratum. Assuming 5% non-participation, the required sample size increased to 751. The non-response rate was assumed to be 10%. Therefore, the calculated sample size was 751 for each of the three class-level strata, which corresponds to a total study sample of 2260 children. Considering that the average size of a class in the sampling frame was 36 (29–43) children, 63 randomly selected classes would be needed in total from three strata, 21 classes from each stratum.

This study was conducted between September to December 2019. We studied three age groups, including 1<sup>st</sup>, 5<sup>th</sup> and 9<sup>th</sup> grades from eight schools, which were randomly selected out of 201 Almaty schools. Sampling was performed maintaining the proportionality between the number of schools and 1<sup>st</sup>, 5<sup>th</sup> and 9<sup>th</sup> grades students. The number of studied classes in each school depended on the total number of students and varied from 2 to 4 grades of each link (on average, from 6 to 12 grades in each school). The study included students from two different types of secondary schools, such as general secondary schools and gymnasiums; however, all schools were state-owned. Both the general education school and the gymnasium were public secondary schools. Gymnasium was an advanced secondary education school with in-depth study of selected classes. These two school types differed in the in core academic hours and the overall structure of the academic workload. All children from the selected schools (N = 2442) were invited to participate in the study, but only 2293 (response rate 93.3%) accepted the invitation and took part in the study. Those children who had a history of any ocular surgery (including intraocular, refractive, trauma and strabismus surgeries), any inflammatory and infectious eye diseases, keratoconus, heterotropia, congenital cataract and pterygium were not included into the study. Students under orthokeratology treatment were excluded because their uncorrected visual acuity and uncorrected refractive error could not be obtained.

## Questionnaire

We also asked children's parents to complete a questionnaire, which included demographic information, history of parental myopia and participants' behavioral factors. Demographic information included age, gender and school grades. Behavioral factors included the average duration of daily gadgets use (computer, mobile phone, tab, games, etc.), near-work (extracurricular activities), and the outdoor activity estimated in hours per day. We used two questions to verify near work: (1) "How many hours does the child spend on the near work (reading, drawing, handicraft, homework, etc.) daily (time spent at school is not considered)?"; (2) "How many hours does the child spend with gadgets (computers, mobile phones, tabs, games, etc.) daily? In addition, outdoor and sport activity were assessed with two questions: (1) "How many hours does the child spend in the outdoor activities daily; (2) "Does your child attend

sports club or section?”. Finally, parental myopia was ascertained with a question “Does any parent have myopia?” with three response options (no; one of the parents; both parents).

### Examination protocol

Clinical examinations were conducted at temporary stations in the school premises. One team examined all children. The clinical team comprised one ophthalmologist and two ophthalmic assistants. The team was trained on the procedure for research in accordance with the RESC recommendations. All children underwent the following standard procedures: distance visual acuity testing, noncycloplegic autorefraction, slit-lamp examination followed by cycloplegia and cycloplegic autorefraction and ophthalmoscopy. The vision examinations and autorefractometry were performed by professional licensed ophthalmic nurses (ophthalmic assistants) in a well-lighted classroom. Visual acuity was measured at a distance of 5 meters, using a retro-illuminated Logarithm of the minimum angle of resolution (LogMAR) chart with Landolt C broken rings and was recorded as the smallest line read with one or no errors. Vision was recorded in decimal notation.

We measured cycloplegic refraction 30 minutes after the instillation of a drop of cyclopentolate 1% twice. Cycloplegia was considered complete if the pupils were dilated 6 mm or greater and the pupillary light reflex was absent. The pupil size was determined roughly with the help of a regular stationery ruler. If needed, a third drop was instilled. Children were asked to keep their eyes closed, for the duration of cycloplegia. We used RC-5000 (Tomey, Tokyo, Japan) for autorefraction with an average of three measurements. The autorefractor was calibrated at the beginning of each working day. Examination of the lens, vitreous, and fundus was performed by an ophthalmologist with a slit lamp and indirect ophthalmoscope.

### Definitions

Refractive errors are determined by the spherical equivalent refraction (SER) calculated as sphere plus the half of the cylindrical error. In accordance with international recommendations, myopia was defined as myopic refractive error when SER of  $\leq -0.50$  D [39]. Myopia was further divided into four refractive error groups: low myopia ( $-3.0$  D  $\leq$  SER  $\leq -0.50$  D), moderate myopia ( $-6.0$  D  $\leq$  SER  $< -3.0$  D) and high myopia (SER  $< -6.0$  D) [39]. Refractive errors of  $\geq +1.00$  D were classified as hyperopia [40]. Thus, emmetropia was defined as SER in the range  $-0.50$  D  $<$  SER  $< +1.00$  D. Clinically significant astigmatism (CSA) was defined as a cylindrical error as  $> 1.0$  D, regardless of sign [40]. Because the correlation of spherical equivalent and visual acuity between the eyes was high (Pearson correlation coefficient 0.91 (95% CI 0.90; 0.92;  $p < 0.001$ )), the spherical equivalent from the worse eye was used for analysis, similar to several previous cross-section studies in children [6, 7, 41, 42]. The worse eye was defined as the eye with the greater absolute value of the SER.

### Ethical issues

This study was conducted in accordance with the Declaration of Helsinki, and all procedures involving human subjects were approved by the Ethics committee of the Kazakh Eye Research Institute (No. 4 dated September 23, 2019) and the Ethics Committee of Al-Farabi Kazakh National University (No. 27–2019 dated May 21, 2019).

### Statistical analysis

We tested normality of all data and reported means with the standard deviation or medians with the corresponding interquartile range (IQR), prompted by the data normality. The main

outcome was the prevalence of refractive errors in different groups, which we reported as % with their 95% confidence interval (CI). Between-group comparisons of binary variables were tested using  $\chi^2$  test from contingency tables, whereas continuous variables were tested using t-test or Mann-Whitney U-test depending on the distribution. We used crude and multiple regression modelling to test the association of refraction errors with selected predictors, identified in the crude analyses. Because some categorical predictors of myopia could have been collinear, we used correlation matrix to test for correlations and found no significant correlations between them. The final adjusted model accounted for seven predictors, including gender, grade, parental myopia, near-work, time spent outdoors, full day of schooling and regular sports activities. The effect in both crude and adjusted analysis was quantified with the odds ratio (OR) with its 95% CI. The cut-off value of a probability of no effect was set to 0.05. All statistical analyses were performed using NCSS (NCSS, Utah, USA).

## Results

In the current cross-sectional study of 2293 analyzed subjects, 50.6% (n = 1161) were boys and 49.4% (n = 1132) were girls. The non-response rate in this study was 6.1%. The mean age of participants was  $11.2 \pm 3.6$  (range 6–16) years. There were no significant differences in gender between groups. Gender, grade and school type distribution is shown in Table 1.

Table 1 summarizes the prevalence of refractive errors among all examined children. The mean spherical equivalent in the total sample was  $-0.54 \pm 1.51$  D. There were no significant differences in refractive errors between boys and girls. The prevalence of refractive errors among all children was 31.6% (95% CI 29.7; 33.5) and increased with age and grade, from 22.2% (95% CI 19.30; 25.1) in 1<sup>st</sup> grade to 43.3% (95% CI 39.8; 46.8) in 9<sup>th</sup> grade. There were no differences in the prevalence of refractive errors between two types of school.

The overall prevalence of myopia was 28.3% (95% CI 26.5; 30.1). Of all participants with myopia, 79.2% had low myopia, 16.4% had moderate myopia, and 4.5% had high myopia. High myopia was more frequent among students of gymnasiums than among general education schools (6.5% vs 2.7%;  $p = 0.014$ ), while low myopia was more often observed in students of general education schools. Prevalence of hyperopia was 3.4%. Low hyperopia was found in 89.6% and moderate hyperopia was detected in 10.4% of all hyperopic participants. As demonstrated in Table 2, there were no significant differences in the prevalence of myopia and hyperopia among boys and girls.

**Table 1. The characteristics of sample and prevalence of refractive errors (N = 2293).**

Demographic variable	N (%)	Mean SER $\pm$ SD, D	Emmetropia, n (%)	Refractive error, n (%)	p
Gender					
Male	1161 (50.6)	-0.52 $\pm$ 1.54	802 (69.1)	359 (30.9)	0.468
Female	1132 (49.4)	-0.56 $\pm$ 1.48	766 (67.7)	366 (32.3)	
Grade					
1 <sup>st</sup>	769 (33.5)	-0.19 $\pm$ 1.07	598 (77.8)	171 (22.2)	<0.001
5 <sup>th</sup>	768 (33.5)	-0.48 $\pm$ 1.43	541 (70.4)	227 (29.6)	
9 <sup>th</sup>	756 (33.0)	-0.95 $\pm$ 1.84	429 (56.7)	327 (43.3)	
School type					
Gymnasiums	1062 (46.3)	-0.63 $\pm$ 1.61	722 (68.0)	340 (32.0)	0.705
General education schools	1231 (53.7)	-0.45 $\pm$ 1.42	846 (68.7)	385 (31.3)	
Total		-0.54 $\pm$ 1.51	1568 (68.4)	725 (31.6)	

Note: SER—spherical equivalent of refraction; SD—standard deviation; D—diopters

<https://doi.org/10.1371/journal.pone.0269474.t001>

**Table 2. The prevalence of different types of refractive errors by gender, school types and grades.**

Demographic variable	Myopia, n (%)	p	Hyperopia, n (%)	p	Astigmatism, n (%)	p
<b>Gender</b>						
Male	319 (27.5)	0.399	40 (3.4)	0.815	33 (2.8)	0.881
Female	329 (29.1)		37 (3.3)		31 (2.7)	
<b>Grade</b>						
1 <sup>st</sup>	135 (17.6)	<0.001	36 (4.7)	0.044	17 (2.2)	0.425
5 <sup>th</sup>	207 (27.0)		20 (2.6)		22 (2.9)	
9 <sup>th</sup>	306 (40.5)		21 (2.8)		25 (3.3)	
<b>School type</b>						
Gymnasiums	309 (29.1)	0.409	31 (2.9)	0.279	29 (2.7)	0.871
General education schools	339 (27.5)		46 (3.7)		35 (2.8)	
Total	648 (28.3)		77 (3.4)		64 (2.8)	

<https://doi.org/10.1371/journal.pone.0269474.t002>

As seen from Table 3, there was an increase in the prevalence of all refractive errors and myopia in particular in older grades in both school types ( $p < 0.05$ ). We did not observe an association between hyperopia and astigmatism with the grade of education. SER among 9<sup>th</sup> graders of general education schools was  $-0.84 \pm 1.70D$ , significantly higher among 9<sup>th</sup> graders from gymnasiums and equaled  $-1.08 \pm 1.99D$ .

When different types of schools were compared, there were no significant differences in the prevalence of refractive errors (myopia, hyperopia and astigmatism) between the corresponding grades of gymnasiums and secondary schools ( $p > 0.05$  for all grades).

The odds of myopia were higher among the children in 5<sup>th</sup> grades (OR 1.73; 95% CI 1.36; 2.21) and 9<sup>th</sup> grade (OR 3.19; 95% CI 2.52; 4.04) as compared to first graders. We also observed dramatic increase of moderate myopia in 5<sup>th</sup> grade group (OR 2.26; 95% CI 1.22; 4.19) and in 9<sup>th</sup> grades group (OR 4.18; 95% CI 2.35; 7.44), when comparing with 1<sup>st</sup> grades children. The risk of high myopia significantly increased in the 9<sup>th</sup> grade students (OR 10.42; 95% CI 2.43; 44.73). Hyperopia among students in 5<sup>th</sup> grade (2.6%) was twice less prevalent compared to 1<sup>st</sup> grade (OR 0.54; 95% CI 0.31; 0.95).

Response rate to the questionnaires filled out by the parents on behavioral risk factors and the role of parental myopia was 50.9% (1166 subjects). As seen from Table 4, the prevalence of myopia among students whose parents were not myopic was 25.2%. Compared with having parents without myopia, having at least one parent with myopia was a risk factor for myopia (OR 1.50; 95% CI 1.08; 2.09) in the crude regression analysis, whereas the presence of myopia

**Table 3. The prevalence of refractive errors in students of different grades.**

Grade	Mean SER $\pm$ SD, D	Emmetropia, n (%)	Refractive errors, n (%)	Myopia, n (%)	High myopia, n (%)	Hyperopia, n (%)	Astigmatism, n (%)
<b>General education schools</b>							
1 <sup>st</sup>	-0.11 $\pm$ 1.11	316 (76.5)	97 (23.5)	72 (17.4)	1 (1.4)	25 (6.1)	10 (2.4)
5 <sup>th</sup>	-0.40 $\pm$ 1.26	288 (71.6)	114 (28.4)	105 (26.1)*	1 (1.0)	9 (2.2)	9 (2.2)
9 <sup>th</sup>	-0.84 $\pm$ 1.70	242 (58.2)	174 (41.8) *	162 (38.9) *	7 (4.3)*	12 (2.9)	16 (3.8)
<b>Gymnasium</b>							
1 <sup>st</sup>	-0.27 $\pm$ 1.01	282 (79.2)	74 (20.8)	63 (17.7)	1 (1.6)	11 (3.1)	7 (2.0)
5 <sup>th</sup>	-0.57 $\pm$ 1.60	253 (69.1)	113 (30.9)*	102 (27.9)*	6 (5.9)	11 (3.0)	13 (3.6)
9 <sup>th</sup>	-1.08 $\pm$ 1.99	187 (55.0)	153 (45.0)*	144 (42.4)*	13 (9.0)	9 (2.6)	9 (2.6)

Note: SER—spherical equivalent of refraction; SD—standard deviation; D—diopters. p-value was calculated for two identical grades of different types of schools:

\*p < 0.05

<https://doi.org/10.1371/journal.pone.0269474.t003>



Table 4. Risk factors of myopia from regression analysis.

Risk factor	Crude			Adjusted <sup>a</sup>		
	OR	95% CI	p	OR	95% CI	p
Gender						
Male	Ref			Ref		
Female	1.09	0.91–1.30	0.399	1.06	0.80–1.41	0.668
Grade						
1 <sup>st</sup> grade	Ref			Ref		
5 <sup>th</sup> grade	1.73	1.36–2.21	<0.001	1.78	1.26–2.52	<0.001
9 <sup>th</sup> grade	3.19	2.52–4.04	<0.001	3.34	2.31–4.82	<0.001
Parental myopia						
neither	Ref			Ref		
either	1.50	1.08–2.09	0.015	1.38	0.98–1.94	0.063
both	1.63	0.88–3.00	0.117	Ref		
School type						
General education school	Ref			-		
Gymnasium	1.07	0.90–1.29	0.409	-		
Uses smartphone (h per day)						
no	Ref					
<1 hour per day	1.06	0.64–1.75	0.836	-		
1.01–2.00	1.23	0.75–2.02	0.405	-		
> 2.01	1.6	0.95–2.67	0.074	-		
Near work (h per day)						
<1 hour per day	Ref			Ref		
1.01–2.00	1.08	0.74–1.59	0.696	Ref		
> 2.01	1.54	1.05–2.24	0.026	1.16	0.87–1.55	0.317
Outdoors (h per day)						
<1 hour per day	Ref			Ref		
1.01–2.00	0.75	0.55–1.03	0.074	Ref		
> 2.01	0.61	0.43–0.86	0.005	0.64	0.46–0.89	0.009
School shift						
morning	Ref			Ref		
afternoon	1.36	0.98–1.89	0.068	Ref		
full day	1.56	1.13–2.17	0.008	1.31	0.95–1.80	0.096
Sports activity						
no	Ref			Ref		
yes	0.71	0.54–0.94	0.016	0.70	0.52–0.93	0.015

Note: Ref—reference; OR—odds ratio; CI—confidence interval; h—hour; <sup>a</sup>adjusted models are adjusted for all the significant predictors in the crude models (school grade, myopia of one of the parents, near-work more than 2 hours a day, outdoors more than 2 hours a day, full school day and sports activity).

<https://doi.org/10.1371/journal.pone.0269474.t004>

in both parents was not associated with the presence of myopia in a student (OR 1.63; 95% CI 0.88; 3.00).

Hours spent per day performing various activities in groups are shown in Table 4. Myopic children spent significantly more screen time, than did non-myopic children (OR 2.29; 95% CI 1.54; 3.42). Extracurricular near work over 2 hours per day increased the odds of myopia (OR 2.29; 95% CI 1.54; 3.42). The prevalence of myopia among students who spent more than 2 hours daily on outdoor activities was 22.2%, whereas that among students who spent less than 1 hours a day outdoors was 32.0%. Non-myopic children spent more total hours outdoors (OR 0.61; 95% CI 0.43; 0.86) than did myopic children ( $p < 0.01$ ). More sports activities were

also a protective factor (OR 0.65; 95% CI 0.49; 0.85). A number of selected predictors were associated with myopia in crude models (Table 4). In the multivariate adjusted regression analysis, significant positive predictors of myopia were the higher class-level: 5<sup>th</sup> grade (OR 1.78; 95% CI 1.26; 2.52) and 9<sup>th</sup> grade (OR 3.34; 95% CI 2.31; 4.82). Outdoor activity more than 2 hours a day (OR 0.64; 95% CI 0.46; 0.89) and sports activity (OR 0.70; 95% CI 0.52; 0.93) were protective against myopia in such multivariate adjusted analyses.

## Discussion

This study was designed to estimate the prevalence of refractive errors in school children in Almaty, Kazakhstan. The prevalence of refractive errors was 31.6%, and the most common refractive error was myopia, which accounted for 89.4% of all refractive errors.

In general, the prevalence of myopia among schoolchildren was 28.3% (95% CI 26.5; 30.1), indicative of the growing trend compared to the previous study in 2010 with 22.1% prevalence in the same age group [35]. We believe that constant increase in educational workloads, leading to more time spent for reading and studying with concurrent reduction in time spent outdoors can explain that [43–46]. Differences in examination methods or discrepancies in myopia definition and thresholds [39] can also explain inconsistent results of studies elsewhere. For example, refractive errors in previous studies were only determined by a spherical refraction component and equivalent refraction, but not SER. In addition, refraction was measured using retinoscopy as opposed to autorefraction [31–35]. Although both methods are reliable [47, 48], direct comparison between studies may not be easy. As expected, our results demonstrated an increase of the prevalence and severity of myopia throughout each successive grades: from 17.6% myopic and 0.3% high myopic students in the 1st grades to 40.5% and 2.5% in 9<sup>th</sup> grades, respectively. Moreover, 5<sup>th</sup> grade (OR 1.78, 95% CI 1.26; 2.52) and 9<sup>th</sup> grade (OR 3.34, 95% CI 2.32; 4.83) were significant factors for myopia progression in an adjusted analysis.

Furthermore, our findings were consistent with other studies in children confirming growing prevalence of myopic refraction and a decrease in the rate of hyperopia with age, especially in the second decade of life [8, 49–57]. Our study did not show the association of myopia with gender, as in some previous studies [58], but data were not consistent across other reports [59]. Such disagreements may indicate both cultural and behavioral characteristics of certain regions, for example, greater sports activity among boys or a more diligent attitude towards school among girls.

Another major change in lifestyle over the past 10 years has been the increase in extracurricular activities [60], decrease of the time spent on homework activities, outdoor activities and sports [61, 62]. This is confirmed by the significantly lower prevalence of myopia in rural areas and economically undeveloped areas, where time spent outside and for household chores is greater compared to the city [63–65]. However, recent research does not find a clear relationship between myopia and screen time [66].

This is the first study of myopia risk factors in schoolchildren in Kazakhstan, in which the effect of independent associated risk factors on myopia was assessed using multiple logistic regression. Contrary to the some available data on the association of myopia with near work range and using smartphones, there was no such relationship in our study [55]. Myopia prevalence, which increased with class-level (age), was almost 30% lower among children attending sports clubs and spending more than 2 hours a day outdoor. These results underscore the significance of environmental factors for myopia and confirm the need for a balance between workload and physical activity. Increased time outdoors are the interventions that are proven to reduce the onset of myopia and are simple interventions to implement in public schools and



at home [67]. Of note, outdoor activities for myopia prevention were implemented in several countries [68–72].

Our study did not confirm the association of myopia in a child with having both myopic parents. Indeed, the OR for both parents was higher than for one (1.63 versus 1.50), but it was statistically non-significant. However, a more likely explanation of such finding was the subjectivity when filling out the questionnaires, resulting in the exposure misclassification. Besides, there might have been a lack of parental awareness; some respondents could have mistaken for myopia any refractive errors requiring spectacle correction, such as presbyopia or hyperopia. In addition, it was likely that not all respondents knew the term 'myopia', instead using non-scientific 'nearsightedness' or 'short-sightedness', which could also affect the results of the study. Therefore, subsequent studies would need a greater sample size and provide an explanation in the myopia questionnaire for more accurate and reliable results.

The advantages of the study were large sample, strict adherence to the study protocol, including clear randomization, and determination of cycloplegic refraction. In general, the protocol of our study was close to RESC with some exceptions, including the use of optotypes in the form of Landolt C rings to determine visual acuity; the sample in a continuous array, and the inclusion of all students of the corresponding grades [38].

The study had several limitations. The study was conducted in governmental schools only. Because a number of studies have shown the association of socioeconomic status with myopia [73–75], private schools with children from families with higher socioeconomic status would likely have more myopia. Our study was cross-sectional and, therefore, we were unable to establish causal relationship between the associations we observed in the risk factor analyses. Moreover, our measurements were taken once at the beginning of the school year, in autumn. It may have biased the effect reducing myopia prevalence, because children may spend more time outdoors in the warm season, particularly during summer holidays with more exposure to natural daylight [76]. With regard to classification of near work in the questionnaire we used, we only asked about extracurricular part of it, which may entail some classification bias and thus is another limitation of our presentation. Furthermore, some selection bias may be present because we looked at the worst eye to diagnose disorders. Finally, we assessed only one urban region of Kazakhstan. The prevalence of myopia was likely higher in cities than in rural areas [77]. Thus, data from this study cannot be extrapolated to the rural or smaller cities of Kazakhstan, and further research is needed to ascertain the prevalence and risk factors in other regions. In addition, this study did not consider the association of myopia with ethnicity. Thus, to get a complete picture of the vision in schoolchildren in the country, research is needed in other regions of Kazakhstan.

## Conclusion

In summary, our study demonstrated that more than a third of school-age children in Almaty had refractive errors, whereas myopia was most common. Compared to previous local studies in this age group, myopia prevalence among urban children likely increased over the past 10 years. We also found that time spent outdoors may be associated with reduced odds of myopia; however, the effects we identified should be further studied in rural population of Kazakhstan.

## Acknowledgments

We would like to thank all survey staff of the Kazakh Eye Research Institute and Faculty of medicine and health care of al-Farabi Kazakh National University. We thank the education department of Almaty and schools' staff for their help in organizing the study. We are grateful to the reviewers and editor for their careful review and insightful comments.

## Author Contributions

**Conceptualization:** Ainagul Mukazhanova, Denis Vinnikov.

**Data curation:** Ainagul Mukazhanova, Neilya Aldasheva.

**Formal analysis:** Aliya Ualiyeva, Kaini Baigonova.

**Methodology:** Ainagul Mukazhanova, Neilya Aldasheva, Juldyz Iskakbayeva, Raushan Bakhytbek, Aliya Ualiyeva, Damet Ongarbaeva, Denis Vinnikov.

**Project administration:** Juldyz Iskakbayeva.

**Resources:** Kaini Baigonova.

**Software:** Kaini Baigonova.

**Supervision:** Ainagul Mukazhanova, Neilya Aldasheva, Raushan Bakhytbek, Damet Ongarbaeva.

**Validation:** Juldyz Iskakbayeva, Aliya Ualiyeva, Kaini Baigonova, Damet Ongarbaeva.

**Writing – original draft:** Ainagul Mukazhanova, Denis Vinnikov.

**Writing – review & editing:** Neilya Aldasheva, Juldyz Iskakbayeva, Raushan Bakhytbek, Aliya Ualiyeva, Kaini Baigonova, Damet Ongarbaeva.

## References

1. Flaxman SR, Bourne RRA, Resnikoff S, Ackland P, Braithwaite T, Cicinelli MV, et al. Global causes of blindness and distance vision impairment 1990–2020: a systematic review and meta-analysis. *Lancet Glob Health*. 2017; 5: e1221–e1234. [https://doi.org/10.1016/S2214-109X\(17\)30393-5](https://doi.org/10.1016/S2214-109X(17)30393-5) PMID: 29032195
2. Lou L, Yao C, Jin Y, Perez V, Ye J. Global Patterns in Health Burden of Uncorrected Refractive Error. *Invest Ophthalmol Vis Sci*. 2016; 57: 6271–6277. <https://doi.org/10.1167/iovs.16-20242> PMID: 27893092
3. Recko M, Stahl ED. Childhood myopia: epidemiology, risk factors, and prevention. *Mo Med*. 2015; 112: 116–121. PMID: 25958656
4. Foster PJ, Jiang Y. Epidemiology of myopia. *Eye*. 2014; 28: 202–208. <https://doi.org/10.1038/eye.2013.280> PMID: 24406412
5. Harrington SC, Stack J, O'Dwyer V. Risk factors associated with myopia in schoolchildren in Ireland. *Br J Ophthalmol*. 2019; 103: 1803–1809. <https://doi.org/10.1136/bjophthalmol-2018-313325> PMID: 30745305
6. Prevalence of Myopia and Hyperopia in 6 to 72 Months Old African American and Hispanic Children: The Multi-Ethnic Pediatric Eye Disease Study. *Ophthalmology*. 2010; 117: 140. <https://doi.org/10.1016/j.ophtha.2009.06.009> PMID: 19926137
7. Wen G, Tarczy-Hornoch K, McKean-Cowdin R, Cotter SA, Borchert M, Lin J, et al. Prevalence of Myopia, Hyperopia and Astigmatism in Non-Hispanic White and Asian Children: Multi-Ethnic Pediatric Eye Disease Study. *Ophthalmology*. 2013; 120: 2109–2116. <https://doi.org/10.1016/j.ophtha.2013.06.039> PMID: 23953098
8. Rudnicka AR, Kapetanakis VV, Wathern AK, Logan NS, Gilmartin B, Whincup PH, et al. Global variations and time trends in the prevalence of childhood myopia, a systematic review and quantitative meta-analysis: implications for aetiology and early prevention. *Br J Ophthalmol*. 2016; 100: 882–890. <https://doi.org/10.1136/bjophthalmol-2015-307724> PMID: 26802174
9. Hashemi H, Fotouhi A, Yekta A, Pakzad R, Ostadimoghaddam H, Khabazkhoob M. Global and regional estimates of prevalence of refractive errors: Systematic review and meta-analysis. *J Curr Ophthalmol*. 2018; 30: 3–22. <https://doi.org/10.1016/j.joco.2017.08.009> PMID: 29564404
10. Rim TH, Kim S-H, Lim KH, Choi M, Kim HY, Baek S-H, et al. Refractive Errors in Koreans: The Korea National Health and Nutrition Examination Survey 2008–2012. *Korean J Ophthalmol KJO*. 2016; 30: 214–224. <https://doi.org/10.3341/kjo.2016.30.3.214> PMID: 27247521
11. Holden BA, Fricke TR, Wilson DA, Jong M, Naidoo KS, Sankaridurg P, et al. Global Prevalence of Myopia and High Myopia and Temporal Trends from 2000 through 2050. *Ophthalmology*. 2016; 123: 1036–1042. <https://doi.org/10.1016/j.ophtha.2016.01.006> PMID: 26875007

12. Morgan IG, French AN, Ashby RS, Guo X, Ding X, He M, et al. The epidemics of myopia: Aetiology and prevention. *Prog Retin Eye Res.* 2018; 62: 134–149. <https://doi.org/10.1016/j.preteyeres.2017.09.004> PMID: 28951126
13. Yam JC, Tang SM, Kam KW, Chen LJ, Yu M, Law AK, et al. High prevalence of myopia in children and their parents in Hong Kong Chinese Population: the Hong Kong Children Eye Study. *Acta Ophthalmol (Copenh).* 2020. <https://doi.org/10.1111/aos.14350> PMID: 31981300
14. Lam CS-Y, Lam C-H, Cheng SC-K, Chan LY-L. Prevalence of myopia among Hong Kong Chinese schoolchildren: changes over two decades. *Ophthalmic Physiol Opt.* 2012; 32: 17–24. <https://doi.org/10.1111/j.1475-1313.2011.00886.x> PMID: 22150587
15. Tsai T-H, Liu Y-L, Ma I-H, Su C-C, Lin C-W, Lin LL-K, et al. Evolution of the Prevalence of Myopia among Taiwanese Schoolchildren: A Review of Survey Data from 1983 through 2017. *Ophthalmology.* 2020. <https://doi.org/10.1016/j.ophtha.2020.07.017> PMID: 32679159
16. Kang M-T, Jan C, Li S, Yusufu M, Liang X, Cao K, et al. Prevalence and risk factors of pseudomyopia in a Chinese children population: the Anyang Childhood Eye Study. *Br J Ophthalmol.* 2020. <https://doi.org/10.1136/bjophthalmol-2020-316341> PMID: 32859718
17. Jung S-K, Lee JH, Kakizaki H, Jee D. Prevalence of Myopia and its Association with Body Stature and Educational Level in 19-Year-Old Male Conscripts in Seoul, South Korea. *Invest Ophthalmol Vis Sci.* 2012; 53: 5579–5583. <https://doi.org/10.1167/iovs.12-10106> PMID: 22836765
18. Karupiah V, Wong L, Tay V, Ge X, Kang LL. School-based programme to address childhood myopia in Singapore. *Singapore Med J.* 2019. <https://doi.org/10.11622/smedj.2019144> PMID: 31680176
19. Plotnikov D, Williams C, Atan D, Davies NM, Ghorbani Mojarrad N, Guggenheim JA, et al. Effect of Education on Myopia: Evidence from the United Kingdom ROSLA 1972 Reform. *Invest Ophthalmol Vis Sci.* 2020; 61: 7. <https://doi.org/10.1167/iovs.61.11.7> PMID: 32886096
20. Pärssinen O. The increased prevalence of myopia in Finland. *Acta Ophthalmol (Copenh).* 2012; 90: 497–502. <https://doi.org/10.1111/j.1755-3768.2011.02210.x> PMID: 21902818
21. Negrel AD, Maul E, Pokharel GP, Zhao J, Ellwein LB. Refractive Error Study in Children: sampling and measurement methods for a multi-country survey. *Am J Ophthalmol.* 2000; 129: 421–426. [https://doi.org/10.1016/s0002-9394\(99\)00455-9](https://doi.org/10.1016/s0002-9394(99)00455-9) PMID: 10764848
22. Gilbert C, Foster A. Childhood blindness in the context of VISION 2020—the right to sight. *Bull World Health Organ.* 2001; 79: 227–232. PMID: 11285667
23. Fricke TR, Jong M, Naidoo KS, Sankaridurg P, Naduvilath TJ, Ho SM, et al. Global prevalence of visual impairment associated with myopic macular degeneration and temporal trends from 2000 through 2050: systematic review, meta-analysis and modelling. *Br J Ophthalmol.* 2018; 102: 855–862. <https://doi.org/10.1136/bjophthalmol-2017-311266> PMID: 29699985
24. Giloyan A, Harutyunyan T, Petrosyan V. Risk Factors for Developing Myopia among Schoolchildren in Yerevan and Gegharkunik Province, Armenia. *Ophthalmic Epidemiol.* 2017; 24: 97–103. <https://doi.org/10.1080/09286586.2016.1257028> PMID: 28032802
25. Abdieva YJ. The prevalence of refractive anomalies among children and adolescents of Azerbaijan depending on the place of residence. In: *Russian Ophthalmological Journal [Internet].* 26 May 2020 [cited 4 Oct 2020] pp. 7–11–11. <https://doi.org/10.21516/2072-0076-2020-13-2-7-11>
26. Bikbov MM, Kazakbaeva GM, Gilmanshin TR, Zainullin RM, Arslangareeva II, Salavatova VF, et al. Axial length and its associations in a Russian population: The Ural Eye and Medical Study. *PloS One.* 2019; 14: e0211186. <https://doi.org/10.1371/journal.pone.0211186> PMID: 30707718
27. Mamatkhuzhaeva GN. [Prevalence of refractive anomalies among school children]. *Vestn Oftalmol.* 2002; 118: 47–49. PMID: 11898364
28. Proskurina OP, Markova EYu, Brzheskij VV, Efimova EL, Efimova MN, Chvatova NN, et al. [The Prevalence of Myopia in Schoolchildren in Some Regions of Russia]. *Ophthalmol Russ.* 2018; 15: 348–353. <https://doi.org/10.18008/1816-5095-2018-3-348-353>
29. Bhandary SK, Dhakal R, Sanghavi V, Verkicharla PK. Ambient light level varies with different locations and environmental conditions: Potential to impact myopia. *PloS One.* 2021; 16: e0254027. <https://doi.org/10.1371/journal.pone.0254027> PMID: 34234353
30. Jones-Jordan LA, Sinnott LT, Chu RH, Cotter SA, Kleinstein RN, Manny RE, et al. Myopia Progression as a Function of Sex, Age, and Ethnicity. *Invest Ophthalmol Vis Sci.* 2021; 62: 36. <https://doi.org/10.1167/iovs.62.10.36> PMID: 34463720
31. Iskakbayeva D. The study of myopia prevalence in the city of Almaty and working out its treatment options. Thesis of the Candidate of Medical Sciences, 2004 (in Russian).
32. Aubakirova A, Kulzhanov M, Toksambaeva G, Dzhanabaev C. The reasons for vision worsening in children 12 years old as of the Republican 10-year data. Abstracts of the 5th International conference, 2006; 19–22 (in Russian).

33. Aubakirova A, Toksambaeva G, Auzhanova R. The efficacy of myopia prevention and treatment in children at school. *Kazakhstan Oftalmologialyk Zhurnaly* 2010; 12–14 (in Russian).
34. Aubakirova A, Kenzhebaeva K, Iskakbayeva D. Clinical and statistical characteristics of myopia and Almaty schoolchildren and the peculiarities of its treatment. *Kazakhstan Oftalmologialyk Zhurnaly* 2001; 8–10 (in Russian).
35. Kenzhebaeva K. Clinical and pathogenetic aspects of myopia in children and methods of its correction. Thesis of the Candidate of Medical Sciences. 2010 (in Russian).
36. National Research Center for Health Development. [cited 9 Jun 2018]. Available: [http://rcrz.kz/index.php/ru/?option=com\\_content&view=article&id=591](http://rcrz.kz/index.php/ru/?option=com_content&view=article&id=591)
37. Bureau of National statistics. [cited 11 Feb 2022]. Available: <https://stat.gov.kz/>
38. Assessment of the prevalence of visual impairment attributable to refractive error or other causes in school children. Protocol and manual Ginebra. World Health Organization; 2007.
39. Flitcroft DI, He M, Jonas JB, Jong M, Naidoo K, Ohno-Matsui K, et al. IMI—Defining and Classifying Myopia: A Proposed Set of Standards for Clinical and Epidemiologic Studies. *Invest Ophthalmol Vis Sci.* 2019; 60: M20–M30. <https://doi.org/10.1167/iovs.18-25957> PMID: 30817826
40. Williams KM, Verhoeven VJM, Cumberland P, Bertelsen G, Wolfram C, Buitendijk GHS, et al. Prevalence of refractive error in Europe: the European Eye Epidemiology (E(3)) Consortium. *Eur J Epidemiol.* 2015; 30: 305–315. <https://doi.org/10.1007/s10654-015-0010-0> PMID: 25784363
41. Wang J, Ying G, Fu X, Zhang R, Meng J, Gu F, et al. Prevalence of myopia and vision impairment in school students in Eastern China. *BMC Ophthalmol.* 2020; 20: 2. <https://doi.org/10.1186/s12886-019-1281-0> PMID: 31898504
42. Chen M, Wu A, Zhang L, Wang W, Chen X, Yu X, et al. The increasing prevalence of myopia and high myopia among high school students in Fenghua city, eastern China: a 15-year population-based survey. *BMC Ophthalmol.* 2018; 18: 159. <https://doi.org/10.1186/s12886-018-0829-8> PMID: 29970057
43. Oecd. Does Homework Perpetuate Inequities in Education? OECD Publishing; 2014 Dec. Report No.: 46. Available: <https://ideas.repec.org/p/oec/eduddd/46-en.html>
44. Jiang Y, Tian B. Understanding Modifiable Risk Factors for the Development of Myopia. *Ophthalmology.* 2019; 126: 221–222. <https://doi.org/10.1016/j.ophtha.2018.09.001> PMID: 30196067
45. Ku P-W, Steptoe A, Lai Y-J, Hu H-Y, Chu D, Yen Y-F, et al. The Associations between Near Visual Activity and Incident Myopia in Children: A Nationwide 4-Year Follow-up Study. *Ophthalmology.* 2019; 126: 214–220. <https://doi.org/10.1016/j.ophtha.2018.05.010> PMID: 29934268
46. French AN, Morgan IG, Mitchell P, Rose KA. Patterns of myopic activities with age, gender and ethnicity in Sydney schoolchildren. *Ophthalmic Physiol Opt J Br Coll Ophthalmic Opt Optom.* 2013; 33: 318–328. <https://doi.org/10.1111/opo.12045> PMID: 23452023
47. Jorge J, Queirós A, Almeida JB, Parafita MA. Retinoscopy/autorefraction: which is the best starting point for a noncycloplegic refraction? *Optom Vis Sci Off Publ Am Acad Optom.* 2005; 82: 64–68. PMID: 15630406
48. Adyanthaya S. A comparison between retinoscopy and autorefraction in acceptance of subjective correction in school age children. *Indian J Clin Exp Ophthalmol.* 2020; 6: 418–421. <https://doi.org/10.18231/j.ijceo.2020.090>
49. Hongjie Yu Yueqin Shao, HongYuan BoYan. Age-determined referral criteria of myopia for large-scale vision screening. *眼科学报.* 2015; 30: 151–155.
50. Lin LLK, Shih YF, Hsiao CK, Chen CJ. Prevalence of myopia in Taiwanese schoolchildren: 1983 to 2000. *Ann Acad Med Singapore.* 2004; 33: 27–33. PMID: 15008558
51. Fotouhi A, Hashemi H, Khabazkhoob M, Mohammad K. The prevalence of refractive errors among schoolchildren in Dezful, Iran. *Br J Ophthalmol.* 2007; 91: 287–292. <https://doi.org/10.1136/bjo.2006.099937> PMID: 17035280
52. Czepita D, Mojsa A, Ustianowska M, Czepita M, Lachowicz E. Prevalence of refractive errors in schoolchildren ranging from 6 to 18 years of age. *Ann Acad Med Stetin.* 2007; 53: 53–56. PMID: 18561610
53. Goh P-P, Abqariyah Y, Pokharel GP, Ellwein LB. Refractive error and visual impairment in school-age children in Gombak District, Malaysia. *Ophthalmology.* 2005; 112: 678–685. <https://doi.org/10.1016/j.ophtha.2004.10.048> PMID: 15808262
54. Wang SK, Guo Y, Liao C, Chen Y, Su G, Zhang G, et al. Incidence of and Factors Associated With Myopia and High Myopia in Chinese Children, Based on Refraction Without Cycloplegia. *JAMA Ophthalmol.* 2018; 136: 1017–1024. <https://doi.org/10.1001/jamaophthalmol.2018.2658> PMID: 29978185
55. Xie Z, Long Y, Wang J, Li Q, Zhang Q. Prevalence of myopia and associated risk factors among primary students in Chongqing: multilevel modeling. *BMC Ophthalmol.* 2020; 20: 146. <https://doi.org/10.1186/s12886-020-01410-3> PMID: 32295555

56. Truckenbrod C, Meigen C, Brandt M, Vogel M, Wahl S, Jurkutat A, et al. Reference curves for refraction in a German cohort of healthy children and adolescents. *PLoS One*. 2020; 15: e0230291. <https://doi.org/10.1371/journal.pone.0230291> PMID: 32160245
57. Theophanous C, Modjtahedi BS, Batech M, Marlin DS, Luong TQ, Fong DS. Myopia prevalence and risk factors in children. *Clin Ophthalmol Auckl NZ*. 2018; 12: 1581–1587. <https://doi.org/10.2147/OPHT.S164641> PMID: 30214142
58. Theophanous C, Modjtahedi BS, Batech M, Marlin DS, Luong TQ, Fong DS. Myopia prevalence and risk factors in children. *Clin Ophthalmol Auckl NZ*. 2018; 12: 1581–1587. <https://doi.org/10.2147/OPHT.S164641> PMID: 30214142
59. Qian X, Liu B, Wang J, Wei N, Qi X, Li X, et al. Prevalence of refractive errors in Tibetan adolescents. *BMC Ophthalmol*. 2018; 18: 118. <https://doi.org/10.1186/s12886-018-0780-8> PMID: 29747615
60. LeBlanc AG, Katzmarzyk PT, Barreira TV, Broyles ST, Chaput J-P, Church TS, et al. Correlates of Total Sedentary Time and Screen Time in 9–11 Year-Old Children around the World: The International Study of Childhood Obesity, Lifestyle and the Environment. *PLoS One*. 2015; 10: e0129622. <https://doi.org/10.1371/journal.pone.0129622> PMID: 26068231
61. Zhang Y, Zhang X, Li J, Zhong H, Pan C-W. Associations of outdoor activity and screen time with adiposity: findings from rural Chinese adolescents with relatively low adiposity risks. *BMC Public Health*. 2020; 20: 1769. <https://doi.org/10.1186/s12889-020-09897-7> PMID: 33228624
62. Cleland V, Timperio A, Salmon J, Hume C, Baur LA, Crawford D. Predictors of time spent outdoors among children: 5-year longitudinal findings. *J Epidemiol Community Health*. 2010; 64: 400–406. <https://doi.org/10.1136/jech.2009.087460> PMID: 19778909
63. Mahayana IT, Indrawati SG, Pawiroranu S. The prevalence of uncorrected refractive error in urban, suburban, exurban and rural primary school children in Indonesian population. *Int J Ophthalmol*. 2017; 10: 1771–1776. <https://doi.org/10.18240/ijo.2017.11.21> PMID: 29181324
64. Tomaz SA, Hinkley T, Jones RA, Watson ED, Twine R, Kahn K, et al. Screen Time and Sleep of Rural and Urban South African Preschool Children. *Int J Environ Res Public Health*. 2020; 17. <https://doi.org/10.3390/ijerph17155449> PMID: 32751089
65. Muthuri SK, Wachira L-JM, Leblanc AG, Francis CE, Sampson M, Onyvera VO, et al. Temporal trends and correlates of physical activity, sedentary behaviour, and physical fitness among school-aged children in Sub-Saharan Africa: a systematic review. *Int J Environ Res Public Health*. 2014; 11: 3327–3359. <https://doi.org/10.3390/ijerph110303327> PMID: 24658411
66. Wang J, Li M, Zhu D, Cao Y. Smartphone Overuse and Visual Impairment in Children and Young Adults: Systematic Review and Meta-Analysis. *J Med Internet Res*. 2020; 22: e21923. <https://doi.org/10.2196/21923> PMID: 33289673
67. Zhang L, Wang W, Dong X, Zhao L, Peng J, Wang R. Association between time spent outdoors and myopia among junior high school students: A 3-wave panel study in China. *Medicine (Baltimore)*. 2020; 99: e23462. <https://doi.org/10.1097/MD.00000000000023462> PMID: 33327278
68. Wu P-C, Chen C-T, Chang L-C, Niu Y-Z, Chen M-L, Liao L-L, et al. Increased Time Outdoors Is Followed by Reversal of the Long-Term Trend to Reduced Visual Acuity in Taiwan Primary School Students. *Ophthalmology*. 2020; 127: 1462–1469. <https://doi.org/10.1016/j.ophtha.2020.01.054> PMID: 32197911
69. Rose KA, Morgan IG, Smith W, Burlutsky G, Mitchell P, Saw S-M. Myopia, lifestyle, and schooling in students of Chinese ethnicity in Singapore and Sydney. *Arch Ophthalmol Chic Ill 1960*. 2008; 126: 527–530. <https://doi.org/10.1001/archophth.126.4.527> PMID: 18413523
70. Wu P-C, Chen C-T, Lin K-K, Sun C-C, Kuo C-N, Huang H-M, et al. Myopia Prevention and Outdoor Light Intensity in a School-Based Cluster Randomized Trial. *Ophthalmology*. 2018; 125: 1239–1250. <https://doi.org/10.1016/j.ophtha.2017.12.011> PMID: 29371008
71. Dirani M, Tong L, Gazzard G, Zhang X, Chia A, Young TL, et al. Outdoor activity and myopia in Singapore teenage children. *Br J Ophthalmol*. 2009; 93: 997–1000. <https://doi.org/10.1136/bjo.2008.150979> PMID: 19211608
72. Rose KA, Morgan IG, Ip J, Kifley A, Huynh S, Smith W, et al. Outdoor activity reduces the prevalence of myopia in children. *Ophthalmology*. 2008; 115: 1279–1285. <https://doi.org/10.1016/j.ophtha.2007.12.019> PMID: 18294691
73. Tideman JWL, Polling JR, Hofman A, Jaddoe VW, Mackenbach JP, Klaver CC. Environmental factors explain socioeconomic prevalence differences in myopia in 6-year-old children. *Br J Ophthalmol*. 2018; 102: 243–247. <https://doi.org/10.1136/bjophthalmol-2017-310292> PMID: 28607175
74. Grzybowski A, Kanclerz P, Tsubota K, Lanca C, Saw S-M. A review on the epidemiology of myopia in school children worldwide. *BMC Ophthalmol*. 2020; 20: 27. <https://doi.org/10.1186/s12886-019-1220-0> PMID: 31937276

75. Saxena R, Vashist P, Tandon R, Pandey RM, Bhardawaj A, Menon V, et al. Prevalence of Myopia and Its Risk Factors in Urban School Children in Delhi: The North India Myopia Study (NIM Study). *PLoS One*. 2015; 10: e0117349. <https://doi.org/10.1371/journal.pone.0117349> PMID: 25719391
76. Rusnak S, Salcman V, Hecova L, Kasl Z. Myopia Progression Risk: Seasonal and Lifestyle Variations in Axial Length Growth in Czech Children. *J Ophthalmol*. 2018; 2018: 5076454. <https://doi.org/10.1155/2018/5076454> PMID: 29692929
77. Ip JM, Rose KA, Morgan IG, Burlutsky G, Mitchell P. Myopia and the urban environment: findings in a sample of 12-year-old Australian school children. *Invest Ophthalmol Vis Sci*. 2008; 49: 3858–3863. <https://doi.org/10.1167/iovs.07-1451> PMID: 18469186