

## Research Article

# Spatiotemporal Study of COVID-19 in Fars Province, Iran, October–November 2020: Establishment of Early Warning System

Ali Semati <sup>1</sup>, Azimeh Zare <sup>2</sup>, Marjan Zare <sup>3</sup>, Alireza Mirahmadizadeh <sup>1</sup>,  
and Mostafa Ebrahimi <sup>4</sup>

<sup>1</sup>Non-Communicable Diseases Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

<sup>2</sup>Department of Anesthesiology, Gerash University of Medical Sciences, Gerash, Fars, Iran

<sup>3</sup>Maternal-Fetal Medicine Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

<sup>4</sup>Communicable Disease Control Center, Shiraz University of Medical Sciences, Shiraz, Iran

Correspondence should be addressed to Marjan Zare; [marjan.zare@gmail.com](mailto:marjan.zare@gmail.com)

Received 11 January 2022; Revised 8 May 2022; Accepted 12 May 2022; Published 30 May 2022

Academic Editor: Marcos Christian Bautista

Copyright © 2022 Ali Semati et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Background.** Using time series and spatiotemporal analyses, this study aimed to establish an Early Warning System (EWS) for COVID-19 in Fars province Iran. **Methods.** A EWS was conducted on (i) daily basis city-level time series data including 53 554 cases recorded during 18 February–30 September 2020, which were applied to forecast COVID-19 cases during 1 October–14 November 2020, and (ii) the spatiotemporal analysis, which was conducted on the forecasted cases to predict spatiotemporal outbreaks of COVID-19. **Results.** A total of 55 369 cases were forecasted during 1 October–14 November 2020, most of which (26.9%) occurred in Shiraz. In addition, 65.80% and 34.20% of the cases occurred in October and November, respectively. Four significant spatiotemporal outbreaks were predicted, with the Most Likely Cluster (MLC) occurring in ten cities during 2–22 October ( $P < 0.001$  for all). Moreover, subgroup analysis demonstrated that Zarrindasht was the canon of the epidemic on 6 October ( $P = 0.04$ ). As a part of EWS, the epidemic was triggered from Jahrom, involving the MLC districts in the center, west, and south parts of the province. Then, it showed a tendency to move towards Zarrindasht in the south and progress to Lar in the southernmost part. Afterwards, it simultaneously progressed to Fasa and Sepidan in the central and northwestern parts of the province, respectively. **Conclusion.** EWS, which was established based on the current protocol, alarmed policymakers and health managers on the progression of the epidemic and on where and when to implement medical facilities. These findings can be used to tailor province-level policies to servile the ongoing epidemic in the area; however, governmental level effort is needed to control the epidemic at a larger scale in the future.

## 1. Introduction

The surveillance systems for infectious diseases are encouraging well-developed techniques that can warn them by detecting the exact time and place of disease occurrence and, subsequently, targeting interventions and resources to more risky areas [1]. Some infections follow space and time trends for which Early Warning Systems (EWS) can be useful tools to find the potential outbreaks [2]. EWS are alert systems to predict epidemic outbreaks in the region. In the 1910s, Captain S.R Christophers from the British army set a system to predict malaria in India. Since then, it has been used as a

useful tool to aid infected nations, make health policies, and manage services to the infected area [1]. The epidemic statue of SARS-CoV-2 disease (COVID-19) was officially declared by the World Health Organization in January 2020 [3, 4]. What makes it important to set a EWS for ongoing COVID-19 epidemic is that it could early notify the policy makers and health managers on the time and location of the outbreaks, so that they could focus on more needy areas and dedicate more medical facilities. Finally, people are helped with less COVID-19 threat, infections, and death.

The question is that, using COVID-19 time series data, if the current study could establish a EWS for future COVID-19

outbreaks, the surveillance system could act timely, prioritize times and locations for targeting interventions, allocate resources more efficiently, and improve decision-making in the area.

Time series data refer to a sequence of successive naturally temporal data, which are equally spaced in points of time. A time series analysis consists of methods for analyzing time series data and forecasting the future values based on the previously observed values [5]. Spatial and Temporal Scan (SaTScan) modeling discovers more about the space and time of an event occurrence. Space-time permutation scan statistic, which is derived from SaTScan, was first introduced by Martin Kulldorff in 2015 [6]. This model aims at determining whether the cases are distributed uniformly over time and space. The statistic utilizes scanning windows moving systematically across the study area to identify the clusters of cases. This modeling helps surveillance systems decide timely on more at risk areas [7]. Using one model output as another mode input, this study could develop a protocol of EWS in the area by (i) forecasting the COVID-19 cases, (ii) discovering the future space-time outbreaks of COVID-19, (iii) predicting the progression of epidemic, and (iv) having more control on the epidemic.

## 2. Methods

**2.1. Study Design, Data Source, and Study Variable.** Daily basis city-level time series data including 53 554 cases recorded in 26 cities of Fars province, Iran, during 18 February–30 September 2020 (225 days) were applied to forecast COVID-19 cases during 1 October–14 November 2020 (45 days). The cases were obtained from 44 COVID-19 sampling centers affiliated to Shiraz University of Medical Sciences (SUMS). All data were reported to SUMS either online, <https://www.coronalab.sums.ac.ir>, or by manual registration. There were no exclusion criteria, meaning that all cases were included in the study. The COVID-19 cases were the positive cases confirmed by real-time Reverse Transcription Polymerase Chain Reaction (RT-PCR). Additionally, the geographical coordinates of the locations were obtained through Google-Earth (US Dept. of State Geographer 2021) based on the latitude-longitude coordinate system.

**2.2. Case Study and Area.** Fars province, the fourth most populated (4 852 274 dwellers) and widespread (122 608 km<sup>2</sup>) province amongst 31 provinces in Iran, is located at 27°3' and 31°40' northern latitude and 50°36' and 55°35' eastern longitude in south of Iran. It includes 26 cities, with Shiraz, the fifth most populated city in Iran, being the capital. Marvdasht, Jahrom, Fasa, Kazerun, Darab, Firoozabad, Lar, and Abade, also, are the most populated areas (more than 100 000 dwellers per capita). 67.6% of the province are urban dwellers (urban areas and suburbs), 31% are villagers (small towns and rural areas), and the rest are nomads (no fixed habitation who regularly move to and from the same areas breeding livestock) [8].

**2.3. Statistical Analysis.** Mean difference and Std. Error of difference were used to describe quantitative variables, and frequency (relative frequency) was used to describe qualitative variables.

The distribution of COVID-19 cases across cities and times are tested using independent-sample one way ANOVA and pair wise comparisons (LSD test). Additionally, time series analysis including (i) Akaike Information Criterion (AIC)/Bayesian Information Criterion (BIC), which are the estimators of the relative quality of time series models comparative to other models for a given set of data, (ii) Auto Correlation Function (ACF)/Partial Auto Correlation Function (PACF) sample/model, which assess the conformity of the observed and fitted pattern in the data, and (iii) residual analysis, which evaluates the presumptions of white noise parameter, was used to forecast the COVID-19 cases during 1 October–14 November 2020 (45 days). A set of time series models were applied, and the best ones were selected based on lower AIC/BIC scores, ACF/PACF conformity, and residual analysis.

Descriptions on time series models:

A simple additive decomposed time series model is

$$x_t = m_t + s_t + e_t, \quad (1)$$

$m_t$  is the trend,  $s_t$  is the seasonality, and  $e_t$  is the error or random white noise.

To be used to predict well founded values, a time series must be stationary, meaning that it should have a constant mean and variance across the time series. So, in order to allow forecasting, several time series models can be created including Autoregressive Model (AR), Moving Average Model (MA), and Autoregressive Moving Average (ARMA) model, which are used on stationary time series. Seasonal Autoregressive Moving Average (SARMA) model, however, is used for non-stationary time series.

Autoregressive Model: AR ( $p$ )

AR model uses observations from previous time steps as input to a regression equation to predict the value at the next step. It takes one argument,  $p$ , which determines how many previous time steps will be input. The order,  $p$ , can be determined by looking at the PACF. The PACF gives the partial correlation of a time series with its own lagged value, regressed of the time series at all shorter lags.

Moving Average Model: MA ( $q$ )

MA model is a time series model that accounts for very short-run autocorrelation. Basically, it states that the next observation is the mean of every past observation. The order of MA model,  $q$ , can usually be estimated by looking at ACF plot.

Autoregressive Moving Average: ARMA ( $p, q$ )

ARMA and ARIMA models are simply a combination of an AR model and MA model.

TABLE 1: Distribution of 53 554 COVID-19 cases in 26 cities of Fars province, Iran, February 18–September 30, 2020.

City	Shiraz	Lar	Jahrom	Fasa	Kazerun	Marvdasht	Darab	Abade
COVID-19 (%)	53.49	8.64	6.25	5.95	5.24	3.32	2.07	1.63
City	Zarindasht	Firoozabad	Lamerd	Mamasani	Ghirkarzin	Safashahr	Sepidan	Stahban
COVID-19 (%)	1.39	1.18	1.16	1.10	1.08	1.04	0.99	0.85
City	Eghlid	Farashband	Pasargad	Neireez	Arsenjan	Rostam	Khonj	Kharama
COVID-19 (%)	0.78	0.73	0.67	0.57	0.47	0.38	0.29	0.29
City	Bavanat	Sarvestan						
COVID-19 (%)	0.24	0.20						

TABLE 2: Distribution of 53 554 COVID-19 cases in Fars province, Iran, February 18–September 30, 2020.

Month	February	March	April	May	June	July	August	September
COVID-19 (%)	0.30	3.50	6.50	4.70	19.70	24	18.70	22.60

Seasonal Autoregressive Moving Average model: SARMA<sub>s</sub>(*p*, *q*)

SARMA model is an extension of ARMA model. The only difference is that a seasonal component is added now. *s* is the number of periods in the season. SARMA can adjust a nonstationary time series by removing trend and seasonality. [9].

Applying time series forecasted cases in spatiotemporal analysis, high-rate clusters were detected considering the minimum and maximum temporal sizes that were equal to one day and 50% of the study period, respectively. The minimum and maximum spatial sizes were also equal to two cases and 50% of at risk population, respectively. Temporal data were checked to ensure that all the cases were within the specified temporal study period. In addition, geographical data check was done to ensure that all the observations were within the specified geographical area. No geographical overlap and subgroup analysis were used.

R v.3.6.3, ITSM 2002, ArcGIS v. 10, and SaTScan software were applied for data analysis. Significance level was considered 0.05 for all tests.

**2.4. Validation Study.** A validation study using real data from 1 October–14 November 2020 was done to assess the accuracy of the results.

**2.5. Ethical Statement.** All ethical steps including data collection and analysis as well as reporting the results were in accordance with the standards approved by the Ethic Committee of the Ministry of Health, Treatment, and Medical Education (IR.SUMS.REC.1399.574). The work processes were anonymous, and the results were reported to the study participants.

### 3. Results

**3.1. Distributions of 53 554 COVID-19 Cases in Fars Province by City and Month, February 18–September 30, 2020.** From 53 554 COVID-19 cases, the maximum and minimum cases happened in Shiraz (53.49%) and Sarvestan (0.2%), respectively. In addition, the maximum and minimum cases happened in July (24%) and February (0.3%), respectively.

The distribution of 53 554 COVID-19 cases by city and month is shown in Tables 1 and 2.

COVID-19 cases were not distributed uniformly across cities (test statistics = 170.83, *df* = 25, *P* < 0.001) and months (test statistics = 21.3, *df* = 7, *P* < 0.001) in Fars province during February 18–September 30, 2020. Pairwise comparisons of 53 554 COVID-19 cases by city and month have been shown in Tables 3 and 4.

COVID-19 cases were significantly higher in Shiraz than those in 25 other cities (*P* < 0.05 for all). Jahrom had more cases than Farashband, Khonj, Darab, Rostam, Kharama, Zarindasht, Sarvestan, Firoozabad, Ghirkarzin, Marvdasht, Lamerd, Mamasani, and Sepidan (*P* < 0.05 for all); however, fewer cases were seen in Jahrom compared with Lar (*P* < 0.05). In addition, Safashahr has lower cases compared with Fasa, Jahrom, Kazerun, Marvdasht, and Lar (*P* < 0.05 for all). Also, there were more cases in Fasa than those in Neireez, Abade, Arsenjan, Stahban, Eghlid, Bavanat, Pasargad, Kharama, Farashband, khonj, Darab, Rostam, Zarindasht, Sarvestan, Firoozabad, Ghirkarzin, Marvdasht, Lamerd, Sepidan, and Mamasani (*P* < 0.05 for all); there, however, were fewer cases in Fasa compared with Lar (*P* < 0.05). Moreover, fewer cases were seen in Neireez, Arsenjan, Stahban, Eghlid, Pasargad, and Bavanat than those in Jahrom, Kazerun, Marvdasht, and Lar (*P* < 0.05 for all, respectively). Furthermore, the cases were lower in Kharama, Khonj, Rostam, Ghirkarzin, Sarvestan, and Farashband than those in Kazerun, Marvdasht, and Lar (*P* < 0.05 for all, respectively). Additionally, the cases were lower in Abade than those in Jahrom, Kazerun, and Lar (*P* < 0.05 for all). Likewise, the cases were lower in Darab, Firoozabad, and Zarrindasht than those in Kazerun, and Lar (*P* < 0.05 for all, respectively). Besides, the cases were more in Kazerun than those in Lamerd, Mamasani, and Sepidan (*P* < 0.05 for all); however, less cases were seen in Kazerun compared with Lar (*P* < 0.05). Also, the cases were higher in Marvdasht than those in Mamasani, and Sepidan (*P* < 0.05 for both); however, fewer cases were seen compared with Lar (*P* < 0.05). Moreover, there were fewer cases in Lamerd and Mamasani than those in Lar (*P* < 0.05 for both). The cases were higher in Lar than those in Sepidan (*P* < 0.05).

COVID-19 cases in February, March, April, and May were significantly lower than those in June, July, August, and

TABLE 3: Pairwise comparisons of 53 554 COVID-19 cases across 26 cities of Fars province, Iran, during February 18–September 30, 2020.

City	Mean difference	<i>P</i>	City	Mean difference	<i>P</i>	City	Mean difference	<i>P</i>	City	Mean difference	<i>P</i>
Shiraz-Safashahr	124.28	<0.001	Safashahr-Fasa	-11.62	<0.001	Neireez-Jahrom	-13.46	<0.001	Jahrom-Kharame	14.13	<0.001
Shiraz-Fasa	112.66	<0.001	Safashahr-Jahrom	-12.33	<0.001	Neireez-Kazerun	-11.07	<0.001	Jahrom-Farashband	13.10	<0.001
Shiraz-Neireez	125.41	<0.001	Safashahr-Kazerun	-9.95	<0.001	Neireez-Marvdasht	-6.52	0.01	Jahrom-Khonj	14.11	<0.001
Shiraz-Abade	122.9	<0.001	Safashahr-Marvdasht	-5.39	0.04	Neireez-Lar	-19.13	<0.001	Jahrom-Darab	9.91	<0.001
Shiraz-Arsenjan	125.63	<0.001	Safashahr-Lar	-18	<0.001	Abade-Jahrom	-10.95	<0.001	Jahrom-Rostam	13.90	<0.001
Shiraz-Stahban	124.73	<0.001	Fasa-Neireez	12.7	<0.001	Abade-Kazerun	-8.56	0.001	Jahrom-Zarindasht	11.1	<0.001
Shiraz-Eghlid	124.92	<0.001	Fasa-Abade	10.24	<0.001	Abade-Lar	-16.62	<0.001	Jahrom-Sarvestan	14.32	<0.001
Shiraz-Bavanat	126.18	<0.001	Fasa-Arsenjan	12.98	<0.001	Arsenjan-Jahrom	-13.69	<0.001	Jahrom-Firoozabad	12	<0.001
Shiraz-Pasargad	125.17	<0.001	Fasa-Stahban	12.08	<0.001	Arsenjan-Kazerun	-11.3	<0.001	Jahrom-Ghirkarzin	12.26	<0.001
Shiraz-Jahrom	111.95	<0.001	Fasa-Eghlid	12.25	<0.001	Arsenjan-Marvdasht	-6.7	0.01	Jahrom-Marvdasht	6.94	0.009
Shiraz-Kharame	126.08	<0.001	Fasa-Bavanat	13.52	<0.001	Arsenjan-Lar	-19.36	<0.001	Jahrom-Kharame	1.13	<0.001
Shiraz-Farashband	125.03	<0.001	Fasa-Pasargad	12.51	0.01	Stahban-Jahrom	-12.78	<0.001	Jahrom-Lamerd	12.07	<0.001
Shiraz-khonj	126.03	<0.001	Fasa-Kharame	13.42	<0.001	Stahban-Kazerun	-10.4	<0.001	Jahrom-Mamasani	12.04	<0.001
Shiraz-Darab	121.85	<0.001	Fasa-Farashband	12.37	<0.001	Stahban-Marvdasht	-5.84	0.03	Jahrom-lar	-5.67	0.03
Shiraz-Rostam	125.85	<0.001	Fasa-khonj	13.4	0.01	Stahban-Lar	-18.46	<0.001	Jahrom-Sepidan	12.46	<0.001
Shiraz-Zarindasht	123.46	<0.001	Fasa-Darab	9.19	0.001	Eghlid-Jahrom	-12.96	<0.001	Kharame-Kazerun	-11.74	<0.001
Shiraz-Sarvestan	126.27	<0.001	Fasa-Rostam	13.19	<0.001	Eghlid-Kazerun	-10.57	<0.001	Kharame-Marvdasht	-7.19	0.007
Shiraz-Firoozabad	123.96	<0.001	Fasa-Zarindasht	10.8	<0.001	Eghlid-Marvdasht	-6.02	0.02	Kharame-Lar	-19.8	<0.001
Shiraz-Ghirkarzin	124.21	<0.001	Fasa-Sarvestan	13.61	<0.001	Eghlid-Lar	-18.63	<0.001	Farashband-Kazerun	-10.69	<0.001
Shiraz-Kazerun	114.34	<0.001	Fasa-Firoozabad	11.30	<0.001	Bavanat-Jahrom	-14.23	<0.001	Farashband-Marvdasht	-6.14	0.02
Shiraz-Marvdasht	118.89	<0.001	Fasa-Ghirkarzin	11.55	<0.001	Bavanat-Kazerun	-11.8	<0.001	Farashband-Lar	-18.75	<0.001
Shiraz-Lamerd	124.02	<0.001	Fasa-Marvdasht	6.23	0.01	Bavanat-Marvdasht	-7.29	0.006	Khonj-Kazerun	-11.73	<0.001
Shiraz-Mamasani	124.15	<0.001	Fasa-Lamerd	11.36	<0.001	Bavanat-Lar	-19.9	<0.001	Khonj-Marvdasht	-7.17	0.007
Shiraz-Lar	106.28	<0.001	Fasa-Mamasani	11.50	<0.001	Pasargad-Jahrom	-13.22	<0.001	Khonj-Lar	-19.78	<0.001
Shiraz-Sepidan	124.41	<0.001	Fasa-Lar	-6.38	0.02	Pasargad-Kazerun	-10.84	<0.001	Marvdasht-Mamasani	5.26	0.04
Darab-Kazerun	-7.52	0.005	Fasa-Sepidan	11.75	<0.001	Pasargad-Marvdasht	-6.28	0.02	Marvdasht-Lar	-12.61	<0.001
Darab-Lar	-15.57	<0.001	Sarvestan-Kazerun	-11.94	<0.001	Pasargad-Lar	-19	<0.001	Marvdasht-Sepidan	5.52	0.03
Rostam-Kazerun	-11.52	<0.001	Sarvestan-Marvdasht	-7.38	0.005	Ghirkarzin-Kazerun	-9.87	<0.001	Kazerun-Lamerd	9.68	<0.001
Rostam-Marvdasht	-6.95	0.009	Sarvestan-Lar	-19.99	<0.001	Ghirkarzin-Marvdasht	-5.32	0.04	Kazerun-Mamasani	9.82	<0.001

TABLE 3: Continued.

City	Mean difference	<i>P</i>	City	Mean difference	<i>P</i>	City	Mean difference	<i>P</i>	City	Mean difference	<i>P</i>
Rostam-Lar	-19.57	<0.001	Firoozabad-Kazerun	-9.62	<0.001	Ghirkarzin-Lar	-18	<0.001	Kazerun-Lar	-8.06	0.002
Zarindasht-Kazerun	-9.13	0.001	Firoozabad-Lar	-17.68	<0.001	Lamerd-Lar	-17.74	<0.001	Kazerun-Sepidan	10.08	<0.001
Zarindasht-Lar	-17.19	<0.001	Lar-Sepidan	18.13	<0.001	Mamasani-Lar	-17.88	<0.001			

Note: std. error for all difference *s* is 2.66; just statistically significant differences were reported.

TABLE 4: Pairwise comparisons of 53 554 COVID-19 cases by month in Fars province, Iran, during 18 February–30 September 2020.

Sammonth	Mean difference	Std. error	Adj. <i>P</i>	Sample median	Mean difference	Std. error	Adj. <i>P</i>
February–June	-13.02	2.45	<0.001	April–June	-9.05	1.85	<0.001
February–July	-16	2.45	<0.001	April–July	-12.04	1.85	<0.001
February–August	-12	2.42	<0.001	April–August	-7.23	1.81	<0.001
February–September	-15.08	2.45	<0.001	April–September	-11.2	1.85	<0.001
March–June	-11.15	1.84	<0.001	May–June	-10.30	1.85	<0.001
March–July	-1.14	1.84	<0.001	May–July	-13.30	1.85	<0.001
March–August	-9.33	1.80	<0.001	May–August	-8.48	1.81	<0.001
March–September	-13.21	1.84	<0.001	May–September	-12.36	1.85	<0.001
July–August	4.81	1.81	0.008	August–September	-3.88	1.81	0.03

Note: just statistically significant differences were reported.

September ( $P < 0.05$  for all, respectively); COVID-19 cases in July were significantly higher than those in August ( $P < 0.05$  for all); COVID-19 cases in August were significantly lower than those in September ( $P < 0.05$  for all).

**3.2. Forecasting COVID-19 Cases in Fars Province during 1 October–14 November 2020.** From the 53 554 observed cases during 18 February–30 September 2020, 55 369 cases were forecasted during 1 October–14 November 2020 using variety of time series models. The frequency of 55 369 forecasted cases alongside the results of time series analysis for the 26 cities has been presented in Table 5.

From the 55 369 forecasted cases during 1 October–14 November 2020, the highest proportion was related to Shiraz (26.9%). Additionally, 65.80% and 34.20% of the cases were forecasted to occur in October and November, respectively. The time series results for Jahrom and Shiraz have been presented in Figures 1 and 2, respectively. The interpretation of the results was similar for the 24 other cities.

**3.2.1. COVID-19 Time Series Analysis in Jahrom during 1 October–14 November 2020.** To reach the stability of variance, Box-Cox transformation was set zero. Besides, SARMA<sub>12</sub>(3, 4) was fitted with the observed quadratic trend and seasonality of 12. The estimated COVID-19 cases during 18 February–30 September 2020, the predicted COVID-19 cases during 1 October–14 November 2020 (45 days), and ACF/PACF sample/model for Jahrom have been presented in Figure 1.

SARMA<sub>12</sub>(3, 4) model was shown to be a good fit that can predict COVID-19 cases in Jahrom during October<sup>1st</sup>–November<sup>14th</sup> 2020.

**3.2.2. COVID-19 Time Series Analysis in Shiraz during 1 October–14 November 2020.** To reach the stability of variance, Box-Cox transformation was set zero. Besides, MA(8) was fitted with the observed quadratic trend. The estimated COVID-19 cases during 18 February–30 September 2020, the predicted COVID-19 cases during 1 October–14 November 2020 (45 days), and ACF/PACF sample/model for Shiraz have been presented in Figure 2.

**3.3. Spatiotemporal Analysis.** The results of the spatiotemporal analysis including 55 369 COVID-19 forecasted cases at the city level on a daily basis together with the results of subgroup analysis have been presented in Table 6.

Overall, four outbreaks occurred in Fars province during 1 October–14 November 2020, the largest and longest of which occurred in ten cities during 2–22 October, with Jahrom being the canon (MLC). Outbreaks 2 and 3 occurred in Sepidan and Fasa, respectively, and lasted from 3 to 13 November. Besides, outbreak 4 occurred in Lar and Zarindasht on 21 October. The results of subgroup analysis demonstrated that Zarrindasht was the canon of the epidemic on 6 October.

As a part of EWS, the progression of COVID-19 epidemic resulting from spatiotemporal outbreaks of the 55 369 COVID-19 cases in Fars province, Iran, during 1 October–14 November 2020 has been shown in Figure 3.

COVID-19 epidemic inclined to trigger from Jahrom on 2 October, involving the MLC districts in the center, west, and south parts of the province; then, it showed a tendency to move towards the south in Zarrindasht on 6 October. Afterwards, it progressed to the southernmost part in Lar on 21 October. Afterwards, it simultaneously progressed to Fasa and Sepidan in the central and northwestern parts of the

TABLE 5: The frequency 55 369 forecasted cases during 1 October–14 November 2020 accompanied by the results of time series analysis in the 26 cities of Fars province, Iran.

City	Forecasted cases		Model	Time series analysis results							Order of min AICC YW	AIC	BIC
	n	%		Residual analysis tests (P)									
				Ljung-box	Diff sign points	Rank test statistic	McLeod-Li	Turning points	Jarque-bera				
Shiraz	14895	26.9	MA (8)	0.230	0.30	0.69	0.030	0.100	<0.001	0	8280	8303	
Lar	2762	4.99	SARMA <sub>12</sub> (2, 2)	0.400	0.9	0.49	0.040	0.560	<0.001	0	1098	1101	
Jahrom	1716	3.10	SARMA <sub>12</sub> (3, 4)	0.770	0.73	0.37	<0.001	0.010	<0.001	0	1221	1225	
Fasa	1772	3.20	ARMA (1, 1)	0.460	0.73	0.12	<0.001	0.090	<0.001	0	1371	1371	
Kazerun	1661	3	AR (1)	0.440	0.30	0.04	0.020	0.220	<0.001	0	1267	1266	
Marvdasht	2215	4	SARMA <sub>12</sub> (1, 1)	0.610	0.9	0.04	0.010	0.560	<0.001	0	1247	1247	
Darab	1440	2.60	ARMA (1, 1)	0.960	0.02	0.80	0.040	0.180	<0.001	0	1360	1361	
Abade	1661	3	SARMA <sub>12</sub> (1, 1)	0.960	0.42	0.60	0.040	0.180	<0.001	0	1360	1361	
Zarindasht	1883	3.40	AR (1)	0.860	0.03	0.85	0.010	0.950	0.030	0	1357	1356	
Firoozabad	1495	2.70	SARMA <sub>12</sub> (1, 2)	0.060	0.08	0.21	0.030	0.010	<0.001	0	1391	1389	
Lamerd	1440	2.60	SARMA <sub>12</sub> (2, 2)	0.190	0.21	0.34	<0.001	0.0002	0.030	0	1364	1368	
Mamasani	1495	2.70	SARMA <sub>12</sub> (1, 2)	0.580	0.04	0.32	0.040	0.490	<0.001	0	1381	1380	
Ghirkarzin	1440	2.60	ARIMA (3, 1)	0.980	0.73	0.51	0.020	0.450	0.010	0	1364	1366	
Safashahr	1440	2.60	ARIMA (1, 1)	0.500	0.45	0.76	0.0010	0.350	0.020	0	1348	1348	
Sepidan	1993	3.60	SARMA <sub>12</sub> (3, 1)	0.130	0.30	0.92	0.030	0.950	0.040	0	1344	1342	
Stahban	1440	2.60	ARMA (1, 1)	0.980	0.2	0.81	<0.001	0.180	<0.001	0	1391	1392	
Eghlid	1440	2.60	MA (1)	0.140	0.73	0.17	0.020	0.670	<0.001	0	1391	1388	
Farashband	1440	2.60	SARMA <sub>12</sub> (2, 2)	0.920	0.73	0.40	<0.001	0.040	0.001	0	1369	1370	
Pasargad	1661	3	MA (1)	0.060	0.04	0.47	0.010	0.490	0.050	0	1334	1332	
Neireez	1440	2.60	AR (1)	0.230	0.45	0.34	0.002	0.450	0.010	0	1364	1363	
Arsenjan	1440	2.60	ARMA (2, 1)	0.080	0.42	0.14	0.030	0.830	<0.001	0	1372	1367	
Rostam	1440	2.60	AR (1)	0.090	0.07	0.20	<0.001	<0.001	<0.001	0	1321	1321	
Khonj	1440	2.60	SARMA <sub>12</sub> (2, 1)	0.680	0.06	0.05	<0.001	0.830	<0.001	0	1295	1291	
Kharama	1440	2.60	SARMA <sub>12</sub> (1, 3)	0.450	0.43	0.23	0.001	0.001	0.010	0	1356	1357	
Bavanat	1440	2.60	MA (1)	0.220	0.91	0.02	<0.001	0.310	<0.001	0	1308	1306	
Sarvestan	1440	2.60	SARMA <sub>12</sub> (2, 2)	0.540	0.9	0.96	0.010	0.590	0.130	0	1260	1258	

AR, autoregressive; MA, moving average; ARMA, autoregressive moving average; ARIMA, autoregressive integrated moving average; SARMA, seasonal autoregressive moving average. It is noteworthy to mention that if four out of the six residual analysis test statistics are statistically significant, it is enough to say that the model is a good fit; based on residual analysis McLeod- Li test statistic needs to be significant and the other five tests should be greater than or equal to 0.05; in addition, the order of min AICC YW, which assesses the mean of white noise residual, needs to be zero. The lower the AIC/BIC scores, the better the model fits.

province, respectively on 3 November. This progression alarmed the policy makers and health managers on the potential starting point of ongoing epidemic (Jahrom city on 2 October, 2020) and, also, on when and where to prioritize the medical facilities and staffs; moreover, it could intervene with the transmission chain of COVID-19 in the area.

**3.4. Validation Study.** 59 980 cases were recorded during 1 October to 14 November 2020, from which 64.2%, 5.7%, 5.5%, 2.2%, 2.2%, 2.1%, 2%, 2%, 1.8%, 1.4%, 1.2%, 1.2%, 1.1%, 1%, 1%, 0.9%, 0.8%, 0.7%, 0.7%, 0.4%, 0.4%, 0.4%, 0.4%, 0.3%, 0.3%, and 0.2% occurred in Shiraz, Jahrom, Fasa, Kazerun, Lar, Abade, Firoozabad, Neireez, Sepidan, Eghlid, Stahban, Mamasani, Darab, Kharama, Pasargad, Safashahr, Bavanat, Arsenjan, Rostam, Sarvestan, Zarindasht, Farashband, Marvdasht, Ghirkazin, Lamerd, and Khonj, respectively. In addition, 51.2% and 48.8% of the cases occurred in October and September, respectively. The space-time outbreaks of the 59 980 COVID-19 incidence cases accompanied by subcluster analysis in Fars province, Iran, during 1 October to 14 November 2020 are shown in Table 7.

#### Cluster 1:

Real data: an outbreak occurred in Zarindasht (canon), Darab, and Lar during 2020/11/8–2020/11/14. Forecasted in EWS: Zarindasht and Lar (canons) were predicted to get involved in an outbreak on 2020/10/21.

#### Cluster 2:

Real data: an outbreak occurred in Fasa during 2020/11/4–2020/11/14. Forecasted in EWS: Fasa was predicted to get involved in an outbreak on 2020/11/3.

#### Cluster 3:

Real data: an outbreak occurred in Sepidan during 2020/11/13–2020/11/14. Forecasted in EWS: Sepidan was predicted to get involved in an outbreak during 2020/11/3–2020/11/13.

#### Cluster 4:

Real data: an outbreak occurred in Firoozabad (canon), Farashband, Ghirkarzin, and Sarvestan during 2020/11/8–2020/11/14. Forecasted in EWS: Firoozabad, Farashband, Ghirkarzin, and Sarvestan were predicted




FIGURE 1: (a) SARMA<sub>12</sub> (3, 4) estimating the number of COVID-19 cases in Jahrom during February<sup>18th</sup>–September<sup>30th</sup> 2020; the green dashed line represents the number of observed COVID-19 cases during February<sup>18th</sup>–September<sup>30th</sup> 2020; the simple red line indicates SARMA<sub>12</sub> (3, 4) model estimating COVID-19 cases during February<sup>18th</sup>–September<sup>30th</sup> 2020. (b) SARMA<sub>12</sub> (3, 4) model predicting COVID-19 cases in Jahrom during October<sup>1st</sup>–November<sup>14th</sup> 2020; the green line represents the estimated COVID-19 cases during February<sup>18th</sup>–September<sup>30th</sup> 2020; the red line represents the predicted COVID-19 cases during October<sup>1st</sup>–November<sup>14th</sup> 2020 with its confidence interval. (c) Sample/model SARMA<sub>12</sub> (3, 4) auto correlation function (ACF) and (d) partial auto correlation function (PACF) for Jahrom during October<sup>1st</sup>–November<sup>14th</sup> 2020; the green correlations are derived from the sample and the red correlations are taken from the fitted model; two dashed horizontal lines are confidence bands; a model is considered good in case the sample and model correlations out of the bands overlap.

to get involved in an outbreak during 2020/10/2–2020/10/22.

However, Bavanat (2020/11/11–2020/11/14), Rostam (2020/11/1–2020/11/14), and Darab (2020/11/14), which were detected in validation study, were not predicted in EWS outbreaks.

The prediction of the epidemic consideration: real data showed that the epidemic was started from Jahrom on 2020/10/27 and progressed to Sarvestan, Rostam, Fasa, Zarindasht, Ghirkarzin, Farashband, Bavanat, Sepidan, Fir-o-zabad and Darab; moreover, in EWS, the epidemic was triggered from Jahrom on 2020/10/2 and progressed to MLC districts, Zarindasht, Lar, Fasa, and Sepidan.

A total of 12 cities got involved in the outbreaks during 1 October–14 November 2020 using real data, nine of which were predicted in the EWS. The EWS predicted time periods were either the same as the observed periods or occurred earlier. Based on the validation study findings, the results of EWS were shown to be valid enough for our purpose.

#### 4. Discussion

Covid-19 cases did not distributed uniformly across cities and times during 18 February–30 September 2020 in Fars

province, Iran. In this protocol presented as EWS, 55 369 COVID-19 cases were forecasted from the 53 554 daily basis city-level time series data recorded during 18 February–30 September 2020. The forecasted cases resulted in the prediction of four significant spatiotemporal outbreaks of COVID-19 in the area including the MLC and three other clusters. The MLC, which was also the largest and the longest cluster, included ten cities during 2–22 October, with Jahrom being the canon. This finding, schematically presented as the progression of the epidemic during 1 October–14 November 2020, revealed that the epidemic inclined to trigger from Jahrom on 2 October, involving the MLC districts in the center, west, and south parts of the province; then, it showed a tendency to move towards the south in Zarrindasht on 6 October. Afterwards, it progressed to the southernmost part in Lar on 21 October. Later, it simultaneously progressed to Fasa and Sepidan in the central and northwestern parts of the province, respectively on 3 November. To the best of our knowledge, this protocol alarmed the surveillance system to focus on the progression of the epidemic and especially on the starting point, Jahrom, so that proper interventions could be done on the transmission chain of SARS-CoV-2 resulting in lower prevalence of the disease in the area. Moreover, the surveillance system can




FIGURE 2: (a) MA (8) estimating the number of COVID-19 cases in Shiraz during February<sup>18th</sup>–September<sup>30th</sup> 2020; the green dashed line represents the number of observed COVID-19 cases during February<sup>18th</sup>–September<sup>30th</sup> 2020; the simple red line indicates MA (8) model estimating COVID-19 cases during February<sup>18th</sup>–September<sup>30th</sup> 2020. (b) MA (8) model predicting COVID-19 cases in Shiraz during October<sup>1st</sup>–November<sup>14th</sup> 2020; the green line represents the estimated COVID-19 cases during February<sup>18th</sup>–September<sup>30th</sup> 2020; the red line represents the predicted COVID-19 cases during October<sup>1st</sup>–November<sup>14th</sup> 2020 with its confidence interval. (c) Sample/model MA (8) auto correlation function (ACF) and (d) partial auto correlation function (PACF) for Shiraz during October<sup>1st</sup>–November<sup>14th</sup> 2020; the green correlations are derived from the sample and the red correlations are taken from the fitted model; two dashed horizontal lines are confidence bands; a model is considered good in case the sample and model correlations out of the bands overlap.

TABLE 6: The spatiotemporal outbreaks of the 55 369 COVID-19 cases in Fars province, Iran during 1 October–14 November 2020.

Cluster	Location	Radius (km)	Canon	Start date	End date	Test statistic	Critical value <sup>†</sup>	P*
Cluster 1 (MLC)	Ghirkarzin, Jahrom, Firoozabad, Khonj, Fasa, Farashband, Sarvestan, Lamerd, Kharam, Stahban	135.53	Jahrom	2020/10/02	2020/10/22	223.31	8.70	<0.001
Cluster 2	Sepidan	0	Sepidan	2020/11/03	2020/11/13	185.98	8.70	<0.001
Cluster 3	Fasa	0	Fasa	2020/11/03	2020/11/13	117.80	8.70	<0.001
Cluster 4	Lar, Zarindasht	76.03	Lar	2020/10/21	2020/10/21	32.28	8.70	<0.001
<i>Sub-cluster results</i>								
Cluster 4	Zarrindasht	0	Zarrindasht	2020/10/06	2020/10/09	3.23	3.08	0.04
Cluster 1 (MLC)	Fasa	0	Fasa	2020/10/06	2020/10/09	6.30	6.90	0.09
	Ghirkarzin	0	Ghirkarzin	2020/11/08	2020/11/09	3.44	6.90	0.90

\*Statistical significance was evaluated using Monte Carlo hypothesis testing at 0.05 significance level. <sup>†</sup>Standard Monte Carlo critical value. MLC, most likely cluster.

dedicate more test sites, medical resources, community quarantine, vaccination coverage, and travel bans on more at risk locations at specified detected times.

Despite being the most important transmission line of the disease, it should be explained why the most populated city, Shiraz, with the maximum number of COVID-19 cases





FIGURE 3: The progression of COVID-19 epidemic in Fars province, Iran during 1 October–14 November 2020.

was not included in the predicted outbreaks. Shiraz was ignored in permutation scan statistics analysis, in which clusters are ranked by their likelihood ratio scores rather than by the population density or the area size. This can be attributed to the fact that, with the capital city of Fars province being with advanced medical services, many patients from nearby cities travel to Shiraz to receive health

services, thereby transmitting the infection to other cities of the province.

Time series models based on statistical methods were revealed to fit well for forecasting time dependent data. They presented well-fit results in case of the COVID-19 epidemic as well. For instance, the AR model could forecast the confirmed COVID-19 cases in Iran [10]. Additionally,

TABLE 7: The space-time outbreaks of the 59 980 COVID-19 incidence cases accompanied by subcluster analysis in Fars province, Iran, during 1 October–14 November 2020.

Cluster	Location	Radius (km)	Canon	Start date	End date	Test statistic	Critical value <sup>†</sup>	P value*
Cluster1	Zarindasht, Darab, Lar	76.03	Zarindasht	2020/11/8	2020/11/14	189.3	5.6	<0.001
Cluster2	Fasa	0	Fasa	2020/11/4	2020/11/14	138.18	5.6	<0.001
Cluster3	Sepidan	0	Sepidan	2020/11/13	2020/11/14	55.54	5.6	<0.001
Cluster4	Firoozabad, Farashband, Ghirkarzin, Sarvestan	79.70	Firoozabad	2020/11/8	2020/11/14	36.89	5.6	<0.001
Cluster5	Bavanat	0	Bavanat	2020/11/11	2020/11/14	21.47	5.6	<0.001
Cluster6	Rostam	0	Rostam	2020/11/1	2020/11/14	17.86	5.6	<0.001
Cluster7	Jahrom	0	Jahrom	2020/10/27	2020/11/14	6.72	5.6	0.02
<i>Sub-cluster results</i>								
Cluster1	Darab, Zarindasht	46.57	Darab	2020/11/14	2020/11/14	6.08	2.4	<0.001
	Zarrindasht	0	Zarindasht	2020/11/5	2020/11/14	2.1	1.53	0.01
	Ghirkarzin	0	Ghirkarzin	2020/11/7	2020/11/14	18.23	2.71	<0.001
Cluster4	Sarvestan	0	Sarvestan	2020/10/31	2020/11/14	14.58	2.71	<0.001
	Farashband	0	Farashband	2020/11/8	2020/11/14	14.25	2.71	<0.001
	Firoozabad	0	Firoozabad	2020/11/14	2020/11/14	7.61	2.71	<0.001

\*Statistical significance was evaluated using Monte Carlo hypothesis testing at 0.05 significance level. <sup>†</sup>Standard Monte Carlo critical value. MLC, most likely cluster.

ARIMA-based time series analysis forecasting near future could assist the government to be prepared for the upcoming emergencies of the current epidemic in India [11–14]. In another study, ARIMA models were fitted to forecast short-term COVID-19 cases in the five most affected countries of the world including Italy, Spain, France, the United Kingdom, and the United States of America [15].

Space-time permutation scan statistics could well manage space-time dependent variables through informing the policymakers on the progression of the epidemic in China [16–18]. In the United States, the space-time outbreaks of COVID-19 were detected during two periods of time (22 January to 9 March 2020 and 18 January to 27 March 2020) using space-time statistics [19]. Space-time scan statistics also resulted in the detection of space-time outbreaks in Hong Kong during 23 January–14 April 2020 [20]. Furthermore, space-time permutation scan statistics worked better for predicting highly frequent infectious diseases with high transmission rates [8, 21, 22]. Another study employed almost the same protocol as the one used to establish the EWS in the current study in order to detect the vulnerable districts for COVID-19 in India. Using ARIMA time series model for forecasting short-term COVID-19 cases and performing spatial analysis, they indicated that the western and southern parts of the country were highly vulnerable for COVID-19 infection [23, 24].

EWS establishment could effectively warn the surveillance system about the progression of the epidemic in Fars province, Iran. However, it had several limitations, which might have affected the results. Firstly, the surveillance system of infectious diseases in Iran is a passive system, and many SARS-CoV-2 infected people with mild to moderate symptoms did not refer to health centers and were not diagnosed. Therefore, the forecasted cases of COVID-19 might have been underestimated. This could affect estimating the required health services and medical facilities. Secondly, permutation scan statistics discover

more than one cluster in a given area and time. Hence, interpretation of the results could be somewhat difficult, especially when incorporating main-cluster and sub-cluster results. Regarding clinical and statistical significance, prioritizing the clusters can also be difficult. Another limitation of scan statistics is that it uses a circular scan window. Thus, the precise limits of the detected clusters remain uncertain. In other words, either the site within the detected clusters may not have a high risk, or a high risk site may be out of the cluster. Yet, using subgroup analysis for detecting significant clusters within large clusters could solve the problem. Further investigations are suggested to assess the genetic variation and meteorological risk factors of COVID-19 in the long run.

## 5. Conclusion

The EWS presented here would alarm the policy makers and health managers on the potential starting point of ongoing epidemic and, also, on when and where to prioritize the medical facilities and staffs. The MLC, which was also the largest and the longest cluster, included ten cities during 2–22 October, with Jahrom being the canon. This finding revealed that the epidemic inclined to trigger from Jahrom on 2 October, involving the MLC districts in the center, west, and south parts of the province; then, it showed a tendency to move towards the south in Zarrindasht on 6 October. Afterwards, it progressed to the southernmost part in Lar on 21 October. Afterwards, it simultaneously progressed to Fasa and Sepidan in the central and northwestern parts of the province, respectively, on 3 November. To the best of our knowledge, this protocol alarmed the surveillance system to focus on the starting point and progression of the epidemic as well as to dedicate more test sites, medical resources, stricter quarantine, vaccination coverage, and travel bans on more at risk locations at specified times. These findings can

be used to tailor province-level policies to servile the on-going epidemic in the area; however, governmental level effort is needed to control the epidemic at a larger scale in the future.

## Data Availability

The health service data used to support the findings of this study are restricted by the Ethics Committee of the Ministry of Health, Treatment, and Medical Education affiliated to the Shiraz University of Medical Sciences, Shiraz, Iran, in order to protect patient privacy. Data are available from Marjan Zare, Central Building of Shiraz University of Medical Sciences, Vice Chancellor for Health, fourth floor, Zand St, Shiraz, Iran, telephone number: 07132122368, E-mail: marjan.zare@gmail.com, for researchers who meet the criteria for access to confidential data.

## Ethical Approval

All ethical steps including data collection and analysis as well as reporting the results were in accordance with the standards approved by the Ethic Committee of the Ministry of Health, Treatment, and Medical Education (IR.SUMS.-REC.1399.574). The work processes were anonymous, and the results were reported to the study participants.

## Conflicts of Interest

No conflicts of interest were declared by the authors.

## Acknowledgments

The authors would like to thank Ms. A. Keivanshekouh at the Research Consultation Center (RCC) of the Shiraz University of Medical Sciences for improving the use of English in the manuscript. As a purely academic work, this study did not receive any funds and is scientifically supported by research deputy of Shiraz University of Medical Sciences, Shiraz, Iran.

## References

- [1] M. Zare, A. Rezaianzadeh, H. Tabatabaee, H. Faramarzi, M. Aliakbarpour, and M. Ebrahimi, "Establishment of an early warning system for cutaneous leishmaniasis in Fars province, Iran," *Asian Pacific Journal of Tropical Biomedicine*, vol. 9, no. 6, pp. 232–239, 2019.
- [2] L. F. Chaves and M. Pascual, "Comparing models for early warning systems of neglected tropical diseases," *PLoS Neglected Tropical Diseases*, vol. 1, no. 1, p. e33, 2007.
- [3] M. Kasraeian, M. Zare, H. Vafaei et al., "COVID-19 pneumonia and pregnancy; a systematic review and meta-analysis," *Journal of Maternal-Fetal and Neonatal Medicine*, vol. 35, pp. 1–8, 2020.
- [4] A. Semati, M. Zare, A. Mirahmadizadeh, A. Hemmati, and M. Ebrahimi, "Epidemiological study of infection and death due to COVID-19 in Fars province, Iran, from February to September 2020," *Iranian Journal of Medical Sciences*, vol. 47, 2021.
- [5] M. Laine, "Introduction to dynamic linear models for time series analysis," *Geodetic Time Series Analysis in Earth Sciences*, pp. 139–156, Springer, Berlin, Germany, 2020.
- [6] M. Kulldorff, *SaTScan—Software for the Spatial, Temporal, and Space-Time Scan Statistics*, Harvard Medical School and Harvard Pilgrim Health Care, Boston, MA, USA, 2015.
- [7] M. Zare, A. Rezaianzadeh, H. Tabatabaee, H. Faramarzi, M. Aliakbarpour, and M. Ebrahimi, "Determining endemic values of cutaneous leishmaniasis in Iranian Fars province by retrospectively detected clusters and receiver operating characteristic curve analysis," *Asian Pacific Journal of Tropical Biomedicine*, vol. 9, no. 9, pp. 359–364, 2019.
- [8] M. Zare, A. Rezaianzadeh, H. Tabatabaee, M. Aliakbarpour, H. Faramarzi, and M. Ebrahimi, "Spatiotemporal clustering of cutaneous leishmaniasis in Fars province, Iran," *Asian Pacific Journal of Tropical Biomedicine*, vol. 7, no. 10, pp. 862–869, 2017.
- [9] A. A. Jake Esprabens and J. Kim, *A Short Guide to Time Series Analysis*, <https://bookdown.org/JakeEsprabens/431-Time-Series/introduction-to-time-series.html>, 2020.
- [10] M. Maleki, M. R. Mahmoudi, D. Wraith, and K.-H. Pho, "Time series modelling to forecast the confirmed and recovered cases of COVID-19," *Travel Medicine and Infectious Disease*, vol. 37, Article ID 101742, 2020.
- [11] H. Tandon, P. Ranjan, T. Chakraborty, and V. Suhag, "Coronavirus (COVID-19): ARIMA based time-series analysis to forecast near future," 2020, <https://arxiv.org/abs/200407859>.
- [12] F. M. Khan and R. Gupta, "ARIMA and NAR based prediction model for time series analysis of COVID-19 cases in India," *Journal of Safety Science and Resilience*, vol. 1, no. 1, pp. 12–18, 2020.
- [13] M. Mousavi, R. Salgotra, D. Holloway, and A. H. Gandomi, "COVID-19 time series forecast using transmission rate and meteorological parameters as features," *IEEE Computational Intelligence Magazine*, vol. 15, no. 4, pp. 34–50, 2020.
- [14] L. Fang, D. Wang, and G. Pan, "Analysis and estimation of COVID-19 spreading in Russia based on ARIMA model," *SN Comprehensive Clinical Medicine*, vol. 2, no. 12, pp. 2521–2527, 2020.
- [15] S. Singh, K. S. Parmar, S. J. S. Makkhan, J. Kaur, S. Peshoria, and J. Kumar, "Study of ARIMA and least square support vector machine (LS-SVM) models for the prediction of SARS-CoV-2 confirmed cases in the most affected countries," *Chaos, Solitons & Fractals*, vol. 139, Article ID 110086, 2020.
- [16] C. Mo, D. Tan, T. Mai et al., "An analysis of spatiotemporal pattern for COVID-19 in China based on space-time cube," *Journal of Medical Virology*, vol. 92, no. 9, pp. 1587–1595, 2020.
- [17] W. Yang, M. Deng, C. Li, and J. Huang, "Spatio-temporal patterns of the 2019-nCoV epidemic at the county level in Hubei province, China," *International Journal of Environmental Research and Public Health*, vol. 17, no. 7, p. 2563, 2020.
- [18] L. Ye and L. Hu, "Spatiotemporal distribution and trend of COVID-19 in the Yangtze river delta region of the people's Republic of China," *Geospatial Health*, vol. 15, no. 1, 2020.
- [19] M. Desjardins, A. Hohl, and E. Delmelle, "Rapid surveillance of COVID-19 in the United States using a prospective space-time scan statistic: detecting and evaluating emerging clusters," *Applied Geography*, vol. 118, Article ID 102202, 2020.
- [20] Z. Kan, M.-P. Kwan, M. S. Wong, J. Huang, and D. Liu, "Identifying the space-time patterns of COVID-19 risk and their associations with different built environment features in

- Hong Kong." *The Science of the Total Environment*, vol. 772, Article ID 145379, 2021.
- [21] A. Rezaianzadeh, M. Zare, M. Aliakbarpoor, H. Faramarzi, and M. Ebrahimi, "Space-time cluster analysis of malaria in Fars province-Iran," *International Journal of Infection*, vol. 7, no. 3, Article ID e107238, 2020.
- [22] A. Rezaianzadeh, M. Zare, H. Tabatabaee, M. Ali-Akbarpour, H. Faramarzi, and M. Ebrahimi, "Does prospective permutation scan statistics work well with cutaneous leishmaniasis as a high-frequency or malaria as a low-frequency infection in Fars province, Iran?" *Asian Pacific Journal of Tropical Biomedicine*, vol. 8, no. 10, pp. 478–484, 2018.
- [23] S. Roy, G. S. Bhunia, and P. K. Shit, "Spatial prediction of COVID-19 epidemic using ARIMA techniques in India," *Modeling Earth Systems and Environment*, vol. 7, no. 2, pp. 1385–1391, 2021.
- [24] M. R. Rahman, A. H. M. H. Islam, and M. N. Islam, "Geospatial modelling on the spread and dynamics of 154 day outbreak of the novel coronavirus (COVID-19) pandemic in Bangladesh towards vulnerability zoning and management approaches," *Modeling Earth Systems and Environment*, vol. 7, no. 3, pp. 2059–2087, 2020.