


REVIEW

Open Access



# Multidimensional insights into the repeated electromagnetic field stimulation and biosystems interaction in aging and age-related diseases

Felipe P. Perez<sup>1,2\*</sup> , Joseph P. Bandeira<sup>1,2</sup>, Cristina N. Perez Chumbiauca<sup>1,3</sup>, Debomoy K. Lahiri<sup>1,4,5</sup>, Jorge Morisaki<sup>6</sup> and Maher Rizkalla<sup>7</sup>

## Abstract

We provide a multidimensional sequence of events that describe the electromagnetic field (EMF) stimulation and biological system interaction. We describe this process from the quantum to the molecular, cellular, and organismal levels. We hypothesized that the sequence of events of these interactions starts with the oscillatory effect of the repeated electromagnetic stimulation (REMFS). These oscillations affect the interfacial water of an RNA causing changes at the quantum and molecular levels that release protons by quantum tunneling. Then protonation of RNA produces conformational changes that allow it to bind and activate Heat Shock Transcription Factor 1 (HSF1). Activated HSF1 binds to the DNA expressing chaperones that help regulate autophagy and degradation of abnormal proteins. This action helps to prevent and treat diseases such as Alzheimer's and Parkinson's disease (PD) by increasing clearance of pathologic proteins. This framework is based on multiple mathematical models, computer simulations, biophysical experiments, and cellular and animal studies. Results of the literature review and our research point towards the capacity of REMFS to manipulate various networks altered in aging (Reale et al. *PLoS one* 9, e104973, 2014), including delay of cellular senescence (Perez et al. 2008, *Exp Gerontol* 43, 307-316) and reduction in levels of amyloid- $\beta$  peptides (A $\beta$ ) (Perez et al. 2021, *Sci Rep* 11, 621). Results of these experiments using REMFS at low frequencies can be applied to the treatment of patients with age-related diseases. The use of EMF as a non-invasive therapeutic modality for Alzheimer's disease, specifically, holds promise. It is also necessary to consider the complicated and interconnected genetic and epigenetic effects of the REMFS-biological system's interaction while avoiding any possible adverse effects.

**Keywords:** Repeated electromagnetic field stimulation, Autophagy, Chaperones, HSF1, Amyloid beta, Alzheimer's disease, Aging, Age-related disease

## Introduction

### REMFS in current literature

The massive proliferation of EMF devices has awakened great curiosity to understand the mechanism of

their interaction with biological systems. Recently, numerous researchers have evaluated this interaction [1–4]. Their results suggest specific conditions of experimental and clinical RF exposure may lead to multi-target effects [5] through activation of several biological pathways [5]. Relevant effects of EMF exposures on the pathways known to be involved in the aging process have been identified by *in vitro* studies

\*Correspondence: fpperez@iu.edu

<sup>1</sup> Indiana University School of Medicine, Indianapolis, IN, USA  
Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

**Table 1** In vitro experiments

Biological effects	EMF	Bio-system	Field strength	Exposure time	References #
HSF1 activation	60 Hz	Human HL60 cells	0.8 $\mu$ T, 80 $\mu$ T	20 min	[6]
Mitochondrial activity	50 Hz	Human SH-SY5Y	100- $\mu$ T	24 h	[7]
ROS production	900 MHz	Rat astroglia	10 V/m	5, 10, 20 min	[8]
Blastogenesis	3 Hz	Human lymphocytes	60 G	72 h	[9]
<sup>45</sup> Ca incorporation	30 Hz	Human lymphocytes	0.15 mT		[10]
Channel-activity	42.25 GHz	Kidney cells	2 mW/cm <sup>2</sup> continuous	20–30 min	[11]
Voltage-gated calcium Growth-related	ELF and Microwave 60 Hz	Human and animal cells Human lymphoma	Pulsed and continuous 10 mV/cm continuous	Seconds to min 1 h	[12] [13]
RNA synthesis	60 Hz	HL-60 cells	5.7 $\mu$ T continuous	20 min	[14]
DNA synthesis	15 Hz to 4 kHz	Human fibroblasts	0.023–5.6 G, sinusoidal	1,2,24 h	[15]
DNA synthesis	75 Hz	Human chondrocytes	2.3 mT pulsed	6 to 30 h	[16]
Expression of microRNA	75 Hz	Mononuclear cell from AD	3 mT pulsed	15, 30, 60 min	[17]
Expression of microRNA	50 Hz	Mouse GC–2 cells	1 mT, 2 mT and 3 mT pulsed	72 h (5 min on/10 min off)	[18]
reduction of oxidative stress	75 Hz	Human SH-SY5Y	2 mT pulsed	10 min, 4 times a week	[19]
reduction of oxidative stress	60 Hz	Human SH-SY5Y	4 to 10 mT continuous	20 min	[20]
expression of hsp70	60 Hz	Human breast cells	3 mT continuous	1 to 3 h	[21]
Cytoprotection	60 Hz	Rodent cardiomyocytes	8 $\mu$ T	30 min	[22]
Ubiquitin–proteasome system	50 Hz	Caco 2 cells	1 mT	24–72 h	[23]
Ubiquitin–proteasome system	1.95 MHz	KB cells	3 mW/g	1, 2, 3 h	[24]
Ubiquitin–proteasome system	100 mT	Rat hippocampal cells	100 mT continuous	15 min	[25]
Autophagy-lysosome systems	75 Hz	SH-SY5Y cells	2 mT pulsed	1 h	[26]
Inflammation	27.12 MHz	Human dermal fibroblasts	591 V/m pulsed	30 min	[27]
Cellular senescence	50 MHz	Mouse fibroblasts	0.5 W/Kg continuous	30 min/d	
$\beta$ -amyloid (A $\beta$ ) deposition	64 MHz	Human neurons	0.4-to 0.9 continuous	1 h/d $\times$ 21 days	

(Table 1) and in vivo studies (Table 2). These studies have looked at specific techniques involving various field strengths and exposure time (dosimetry).

The results listed in Tables 1 and 2 demonstrate that the effects of biological mechanisms influenced by REMFS are likely extensive and may act in multiple distinct pathways. These data support possible therapeutic implications of REMFS on the aging process and age-related diseases, such as Late Onset Alzheimer's disease (LOAD), which is also supported by the results of our prior experiments and theories [39]. In later sections of this paper, we will specify these potential mechanisms.

### The consistencies and inconsistencies between the in vivo and in vitro data

As DNA damage is frequently a prerequisite for cancerous diseases, reviews on this topic provide an experimental body of evidence on the effect of EMF on genetic material. Diab's studies on the matter showed contradictory data in in vivo and in vitro experiments [40]. Several studies of both in in vivo and in vitro samples showed a detrimental effect on DNA when exposed to EMF. Some reports showed no injury to DNA in either in in vivo or in vitro experiments. Results from other experiments were inconclusive [40]. These conflicting findings were probably caused by variability in the EMF generators,

**Table 2** In vivo experiments

Biological effects	EMF	Bio-system	Field strength or SAR	Exposure	References #
Oxidative stress	15 $\mu$ T	Vicia faba L	15 $\mu$ T	8 h/d $\times$ 8 days	[28]
Expression of microRNA	2.4 GHz	Rat brain	2420 $\mu$ W/kg	24 h/d $\times$ 12 months	[29]
Reduction of oxidative stress	10 kHz	Maize	3mT	6 h/d $\times$ 4 days	[30]
Reduction of oxidative stress	900 MHz	Rat	0.18 W/kg	1 h/d $\times$ 21 days	[31]
Reduction of oxidative stress	60 Hz	Chick embryos	8 $\mu$ T	20 min	[32]
Inflammation	1–100 Hz	Mice	1–100 Gauss pulsed	30–45 min	[33]
Inflammation	27.12 MHz	Human after breast surgery	591 V/m pulsed	30 min	[34]
Mitochondrial enhancement	918 MHz	AD mice	0.25–1.05 W/kg, pulsed	2 h/d $\times$ 1 month	[35]
Neuronal activity	918 MHz	AD mice	0.25–1.05 W/kg, pulsed	2 h/d $\times$ 2 months	[36]
$\beta$ -amyloid (A $\beta$ ) deposition	918 MHz	AD mice	0.25–1.05 W/kg, pulsed	2 h/d $\times$ 7 to 9 months	[37]
Osteoarthritis	37 and 75 Hz	Guinea pigs	3mT pulsed	6 h/day $\times$ 6 months, pulsed	[38]

SAR specific absorption rate

different experimental methods including time of exposure, and characteristics of the specific samples (age, genetic differences, size, tissue penetration, anatomical differences [41], etc.).

It is important to understand that the in vitro results cannot easily be extrapolated to in vivo results. This is because the energy absorbed by an object is dependent on the way the EMF is able to penetrate the object [42]. The physiological characteristics of in vivo specimens differ significantly from that of an in vitro cell, and exposure to the same external field would result in an entirely different internal field. Hence, it becomes important to determine what external fields would produce similar internal fields inside both in vitro and in vivo specimens before we reach any conclusions about the biological effects of a specific EMF frequency or field intensity.

Another concern is tissue penetration, which is inversely proportional to the frequency of the EMF. With in vitro specimens there is no need to calculate tissue penetration, but in vivo samples have complex calculations for tissue penetration due to the presence of multiple tissue layers, the geometry of the tissues, and their specific dielectric properties (conductivity and permittivity). As the frequency is increased, the penetration depths of the tissue layers change such that the largest part of the incident energy may be transmitted at one frequency and absorbed at another. Sufficiently high frequencies should result in a small penetration depth, resulting in superficial penetration.

## REMFS in cell death and senescence

### REMFS delays cell death

In our REMFS initial study, we exposed T-cells and lymphoblasts cultures to a frequency of 50 MHz and a

power of 0.5W [43]. We found a 20% reduction ( $p < 0.05$ ) in LDH over 3 weeks of REMFS treatments. The 30-min REMFS exposures were the most stable between REMFS treatment times, so this was selected as the minimal optimal treatment regimen. We demonstrated a 34% reduction in LDH release in REMFS-treated quiescent T-cells compared to control cells following treatment with 30 min of REMFS for 7 consecutive days ( $p < 0.01$ ), suggesting a significant protective effect from REMFS. We corroborated the protective effects of REMFS with a Trypan blue exclusion study. Results showed that REMFS decreased T-cell death for 7 days, with maximal benefit using 30 min of daily treatments. These data demonstrated that REMFS of low energy and intensity (50 MHz / 0.5 W) can play a cytoprotective role [43].

### REMFS delays cellular senescence

REMFS treatments produced effects associated with cellular senescence [43]. We treated knockout (KO) and control mouse fibroblasts at 100% of lifespan completed or 23 cell population doubling (CPDL) with REMFS at 50 MHz / 0.5 W every 3 days for 14 days. We found that when cells passed from CPDL 3 to 23, they became larger, vacuolated cells with more diverse morphotypes than cells at earlier CPDL. Interestingly, REMFS reversed and delayed senescent morphology, enlargement, and variation of HSF1 +/+ mouse fibroblasts, but not HSF1 -/- mouse fibroblasts. These results emphasize the importance of REMFS effects on HSF1. REMFS treated HSF1 +/+ mouse fibroblasts remained smaller in size and more spindle-shaped with more parallel positioning of the cells. In addition, there were less multinucleated cells. We also observed that REMFS prolonged the replicative lifespan to 29 CPDL of murine fibroblast HSF

1 +/+ compared to 23 CPDL in non-REFMS treated fibroblasts [43].

We also compared the CPDL tables of HSF1 +/+ and HSF1 knockout mouse fibroblasts in treated vs non-treated cultures [43]. Both groups grew at similar rates until CPDL 18, after which REFMS-exposed fibroblasts CPDL appeared to prevent the decline in cell proliferations rates observed in untreated cells. REMFS-treated fibroblasts demonstrated 138 days of proliferative lifespan, compared to 118 days in non-treated cultures. This represented an increase of 17% in lifespan. Similar to prior experiments, HSF1- knockout cultures did not show response to the treatments, achieving only 23 CPDLs with 100 days of replicative life span.

## REMFS at the organismal level

### REMFS in aging

Several short-term exposure studies have shown that REMFS increases lifespan in mice, worms, and flies. In a recent study, mice had an increased average lifespan when exposed to REMFS with an alternating magnetic field of 100 nT and 60  $\mu$ T [44]. In another study with rotating Magnetic Field (0.2 T, 4 Hz), REMFS exposure slowed the aging process and prolonged the lifespan of *C. elegans* and of Human Umbilical Vein Endothelial Cells (HUVECs). REMFs also improved activity, reduced pigment accumulation, and delayed paralysis induced by A $\beta$ , as well as increased heat tolerance and oxidative stress resistance [45]. A higher frequency study (10 GHz) extended the life span of *Basc* females fruit fly *Drosophila melanogaster* [46]. In another high frequency (1–10 THz) study, there was no increase in survival in early life but increased survival in later life [47].

An interesting study of *Drosophila melanogaster* lifespan showed that population was decreased or increased depending on parameters of the REMFS exposures [48]. There are, however, many other studies that show that prolonged EMF exposures do not increase lifespan in multiple organisms. This is most likely due to the use of different frequencies, powers, times or cell types [49–52].

### REMFS in Alzheimer's disease

AD and Lewy body dementia (LBD) usually emerge during aging, when the proteostasis quality control is unable to prevent the aggregation of misfolded proteins. AD is characterized by A $\beta$  peptides, an APP fragment of 39–43-amino acids [53]. Efficacy and safety of REMFS have been demonstrated in Transgenic (Tg) AD mouse models in vivo. An initial REMFS study prevented or reversed memory loss in Tg AD mouse model (A $\beta$ PPsw) when a pulsed and modulated RF-EMF at 918 MHz with a SAR of 0.25–1.05 W/kg was applied over a 7 to 9 month period [54]. REMFS exposed Tg mice preserved good

cognitive function, whereas control Tg mice showed cognitive decline. Tg mice of advanced age (21–27 months) with daily REMFS exposure for 2 months showed improved memory in the Y-maze task, although not in more complex tasks [36]. These older Tg controls showed high levels of A $\beta$  aggregates with treated mice showing a 24–30% decrease of A $\beta$  deposits. These data suggest a degradation of A $\beta$  deposits with REMFS exposure. In addition, these long-term treatments were found to be safe (daily for up to 9 months) without any toxic effects on multiple health parameters, including oxidative stress, brain histology, brain heating, damage to DNA, or cancer in peripheral tissues [55].

A higher frequency study (1950 MHz) showed decreased AD pathology in Tg-5xFAD transgenic mice, which overexpress APP, and wild type (WT) mice treated with REMFS at 1950 MHz with SAR 5 W/kg for 2 h per day, 5 days per week [56]. This long-term exposure to REMFS decreased A $\beta$  plaques, APP, and APP carboxyl-terminal fragments in the brain. REMFS also decreases the expression of  $\beta$  Beta secretase 1 (BACE1) to prevent inflammation.

Additionally, REMFS reverses cognitive decline in AD mice. REMFS treatment showed that when compared to WT mice, 5 genes that are all implicated in A $\beta$  processing (Tshz2, Gm12695, St3gal1, Isx and Tll1), are affected in Tg-5xFAD mice treated with REMFS. Specifically, WT showed the same genetic profile to non-REFMS-treated Tg mice, while REMFS-treated Tg mice demonstrated different patterns. Therefore, these data suggest that chronic REMFS treatment influence A $\beta$  processing in AD mice, but not in wild or Tg controls [56].

Altogether, AD mouse studies and human brain cell studies revealed that REMFS exposures reduce A $\beta$ . It also prevents and decreases brain A $\beta$  aggregation without causing any inflammatory reaction as seen in passive immunity treatment trials [57, 58]. This represents a potential therapeutic strategy in the treatment of AD patients who already have large amounts of A $\beta$  deposits. Other investigators have demonstrated improved cognitive function that accompanied reduction of A $\beta$  in AD mouse models [36, 54–56]. Taken together, these data suggest a potential therapeutic role of REMFS in human diseases, such as LOAD.

Other studies suggested that EMF exposures enhance pathways involved in A $\beta$  degradation through upregulation of the HSF1 pathway [43], the autophagy-lysosome system [26], the ubiquitin–proteasome system [23, 25], and a reduction in  $\beta$ -secretase activity following REMFS thus producing a protective effect through reduction of A $\beta$  [56]. Furthermore, REMFS also targets multiple aging and cell defense pathways that are involved in AD, including oxidative stress [19], cytoprotection [20],

inflammation [27], mitochondrial enhancement, and neuronal activity [55], thereby making REMFS a potential multi-target therapeutic strategy for AD and other age related diseases.

## Proposed mechanisms of REMFS and biosystems interactions

### Low energy challenges

There are two challenges that underly an explanation of the REMFS and bio-systems interaction: (1) the low energy in these EMFs is several orders of magnitude lower than  $k_B T$  ( $k_B$ : the Boltzmann constant,  $T$ : room temperature). This leads to information being transmitted via a phenomenon known as the  $k_B T$  paradox in which information is hidden in thermal noise [59]. (2) Classical models of molecular dynamics would hold that the excited state produced by the EMF would promptly dissolve due to the thermal excitations that restart when the EMF exposure is eliminated, which is not seen [60].

REMFS exposure transmits a very low energy that is insufficient to excite electrons and is thereby considered non-ionizing. The photon energy of REMFS at 50 MHz is  $2.0678 \text{ eV}^{-7}$ , at 64 MHz it is  $2.64 \text{ eV}^{-7}$ , and at 915 MHz it is  $3.7841 \times 10^{-6} \text{ eV}$ . Additionally, protein conformational changes cannot occur under direct electric field magnitudes lower than  $10^8 \text{ V/m}$  [61] and REMFS only produce  $16.22 \text{ V/m}$  [62]. REMFS energies are incapable of directly causing the dissociation of chemical bonds such as the H–O–H covalent bond of a water molecule ( $\text{H}_2\text{O}$ ) because this type of reaction would require  $493.4 \text{ kJ/mol}$  or  $5.1138 \text{ eV}$  [63, 64], an exponentially higher amount of energy.

Thus, classical physics is unable to explain the biological responses to REMFS. Nevertheless, quantum physics provides an explanation of how this reaction occurs. Here we consider low energy EMF with frequencies below the THz wavelength. Interestingly, high energy EMF are not able to produce the biological effects of the low energy EMF [65]. In addition, Panagopoulos found that oscillating EMF with frequencies lower than  $1.6 \times 10^4 \text{ Hz}$  produce biological effects, even at very low intensities. Conversely, as the frequency of the EMF increases to more than  $1.6 \times 10^4 \text{ Hz}$  a higher field intensity is required to produce biological effects [66].

One plausible explanation why high EMF frequency is less likely to produce biological effects could be the reduced hydrogen bonding seen in higher temperatures which is directly correlated with higher energy [67]. In an interesting study, THz-exposed cells exhibited some biological responses such as increase in heat shock protein expression. Results suggest that the biological effects imposed by THz radiation appear to be primarily thermal in nature [68]. Conversely, the effects of the RF and

microwave are primarily non-thermal, therefore suggesting a different mechanism. This difference could be due to the effects that RF and microwave range correlates to the rotation of polyatomic molecules and higher frequency correlates to the vibrations of flexible bonds [68].

Another possible explanation is the effect of the REMFS oscillation on the H-bond at the quantum level. It may be produced by the fact that the REMFS frequency is much slower (Hz to GHz) than the H-bond frequency (74 THz), thus inducing the H-bond to act as a driven quantum harmonic oscillator under REMFS exposures [69, 70] (Eq. 1) in a time-dependent adiabatic perturbation [71]. REMFS is a continuous field exposure (perturbation) which acts slowly enough to allow the quantum system sufficient time for the functional form to adapt [71] (adiabatic process), and consequently become able to cause changes in the probability density and amplitude. Under faster excitatory frequencies the driven harmonic oscillator has no time to follow the excitatory frequency like in the classical solution [72].

### Changes in water depend on EMF properties

EMF produce their effects by the electric field [67], rather than the magnetic field. Depending on the frequency and intensity of the EMF, it can change water structure through different influences. Consequently, there are some inconsistencies in the effect of EMF on the H-bond network (HBN) [73] and on the biological response. De Nino demonstrated a decreasing coherent population accompanied by the increase of the intermediate population under high amplitude of field (0.15 T) [74]. Conversely, Shen found stronger polarization and a higher degree of association in exposed water to low frequency EMF and low amplitude of field (0.15 T) [75]. These data suggest that the effects of EMF on H-bond and its biological effects are determined by the strength of the external field. The values and distribution of the internal fields depend on the frequency, polarization, field strength, field distribution of the external fields, the configuration of the tissue, and its dielectric properties [76].

### Potential mechanisms

There is no generally accepted mechanism to explain the role of low frequency EMF in biological systems, though multiple mechanisms have been proposed. There is inconsistency amongst these theories, which can be explained by a lack of consideration for varying energy levels, tissues, or the quantum effects of these fields on water molecules. Below are examples of these proposed mechanisms:

- 1) RF-EMF alters the structure of the water surrounding some biomolecules, which allows water to store

and release a greater amount of energy under EMF [77]. The theory is that RF-EMF exposure induces water auto-ionization to produce hydronium, which in turn protonates biomolecules to activate biological pathways. The energy that would be generated by this proposed mechanism is so high that it would cause an increase in temperature that has not been observed in REMFS exposures.

- 2) Protein and protein complexes (HSF1-Hsp90) [43], as well as elements of RNA and DNA [78], are EMF sensitive and can behave as EMF-sensors that operate by disruption of their conformation to form secondary structures in response to EMF variations. Structural transitions can uncover or obscure important regions of RNA, such as binding sites, or lead to dissociation of protein complexes which can release active transcriptional factors. These changes, then, affect the translation rate of nearby protein-coding genes to activate biological pathways. It is an unlikely mechanism because we observed that the initial event in our experiments is not DNA activation; heat stress is required to initiate translation.
- 3) Cells (e.g., neurons) possess the ability to yield constructive interference effects that enhance their intensities at several points [79–81], so that an applied EMF could be amplified to produce conformational changes in some proteins, transcriptional factors, and RNA. This hypothesis also requires very high energy to break bonds and would cause a thermal response.
- 4) Second-harmonic generation increases energy of photons whereby water molecules align during EMF exposures [82]. In this mechanism two photons with the same frequency interact with a nonlinear material to create a new photon with twice the energy of the initial photons. This hypothesis is unlikely to be the mechanism because even this doubled energy of the new photon would be still too low to break any chemical bonds.
- 5) EMF forces affect electrons in a way that weakens H bonds. This destabilization can act on H bonds holding DNA strands together, thereby affecting transcription. The low electron affinity of the bases, which has been previously identified in electromagnetic response elements (EMREs), are needed for EM field interaction with DNA. This theory is also less likely because we demonstrated that DNA activation is not the initial event in our experiments.
- 6) Another hypothesis is that of a resonant frequency as the mechanism of this interaction [83]. However, the exposures are billions of folds different [78, 84–87], thereby creating a wide range of frequencies capable of causing the same biological effects. This makes the

hypothesis of a resonant frequency very unlikely and difficult to substantiate.

- 7) High energy vibrations of ions have are less likely the cause of the RF and biological systems interaction because the mobility of ions is low among these exposures [88].


### Summary of our hypothesis

In this paper, we do not intend to produce a systematic review of all the EMF bio-effects; instead, we are trying to develop a theoretical framework for our experimental results. The mechanism postulated here explains the activation of the Heat Shock Factor1 (HSF1) via REMFS. We believe that different EMF frequencies produce different mechanisms of action on biological systems. For example, ionizing or thermal mechanisms produce the high energy required to remove electrons and break covalent bonds. Also, high (THz) frequencies produce their effects mainly by a thermal mechanism [68].

Here, we concentrate on studies performed on the low energy EMF spectrum (Hz to GHz) to describe mechanisms by which non-ionizing, non-thermal, non-modulated, continuous EM waves induce biological effects. We use results collected through our research on human cell cultures and other researchers' recent results on mouse AD models to support these theories.

Our initial REMFS experiment used an EMF frequency of 50 MHz and a specific absorption rate (SAR) of 0.5 W/Kg. We determined that REMFS activated HSF1 in cell cultures of in lymphocytes and fibroblasts [43], increasing 70-kDa heat shock proteins (Hsp70) chaperone levels and ultimately postponing aging and death in cell cultures. Recently, we demonstrated that REMFS at 64 MHz with a SAR of 0.6 W/Kg for 14 days reduced potentially toxic amyloid- $\beta$  peptide ( $A\beta$ ) levels by 46% in cultures of primary human brain (PHB) when compared to non-exposed controls. A decrease of  $A\beta$  levels in PHB cultures also appeared with different duration and power protocols. Of note,  $A\beta$  precursor protein (APP) levels and non-APP processing pathway products were not altered by the treatments, suggesting enhancement of  $A\beta$  degradation as the possible mechanism of  $A\beta$  reduction.

We hypothesized that a multidimensional sequence of events explains the REMFS and biological system interaction (Fig. 1) from the quantum to the molecular, cellular, and organismal levels. The REMFS mechanism is a combination of the oscillatory quantum [89] and molecular [67] effects on the interfacial water HBN surrounding biomolecules; specifically in REMFS, those H-bond's confined to the first layer of the interfacial water in the vicinity of the non-coding RNA Heat Shock RNA-1 (HSR1) [90]. This EMF oscillation causes the H-bond to



behave like a driven quantum harmonic oscillator [91], thereby increasing the amplitude of the H-bond vibration [92] within the interfacial water that naturally surrounds nucleic acids. This shortens the length of the H-bond, increasing the probability of proton tunneling [93] and protonation of the nucleic acids [94]. This leads to the formation of tautomers [95] that produce conformational changes in HSR1 [90] to allow binding and activation of HSF1. Subsequently, HSF1 binds to DNA to express chaperones that initiate chaperone autophagy and degradation of abnormal proteins such as Aβ with consequent clinical improvement in Alzheimer’s disease (AD).

The main question in the EMG and biological systems interactions is what the target of these fields is

and how the target is affected to produce a response. We hypothesized REMFS oscillations on the interfacial water cause a combination of quantum and molecular vibrations responsible for the biological effects under these fields. The mechanism of the EMF and biological systems interaction could be thermal or non-thermal. Previous studies showed that REMFS acts by a non-thermal mechanism [43, 96]. A temperature-dependent (i.e. thermal) mechanism produces changes in the rates of biochemical reactions as a result of heat energy transfer to the target receptor. In contrast, nonthermal mechanisms are not associated with a change in temperature, but rather with oscillations of the RF which cause vibrational energy transfer, and ultimately a

change in the REMFS receptor [97]. We will discuss the changes at the quantum and molecular levels in more detail in the following subsections.

### Exposure times and regimens

Exposure time is a very important factor to achieve an effective dose that produces biological effects. To find out possible advantages and mechanisms of REMFS, it would be much more valuable to perform experimental studies to determine the effects accumulated with time and profiles of repetition. In comparison, computational studies are limited by very short exposure times. The time exposure and repetition regimen parameters differ regarding the case under study and depend on the physical and biological conditions of the exposed target.

In the particular case of REMFS and biological pathways associated with protein degradation systems in humans and mice, we observed that the minimum exposure time to produce biological effects was between 15 and 30 min of exposure with a peak response of one hour. We also observed that the minimum regimen was 3 times per week. Marchesi found that briefer EMF exposures of 15 and 30 min did not show significant differences compared to the untreated control when measuring miR-30a expression levels. However, lengthier EMF exposures of 1 h, and to a lesser extent 3–24 h, produced biological effects.

This minimum time exposure is necessary to activate and recruit enough molecules of the HSR1 to initiate protein degradation. A repetition exposure regimen leads to maintaining high chaperone levels and degradation of abnormal proteins; otherwise, cells would continue to accumulate abnormal proteins that form during cellular metabolism [26]. There should also be a balanced process of degradation and protein synthesis, which is obtained by an intermittent regimen. There is evidence that effective exposure times can vary according to the type of cell or organism, biological pathway affected, and the physical conditions of the exposure. For example, a study showed that longer time exposures are needed to obtain the maximal biological effect. They found a minimal effect in mammalian stem cells after 2 h of EMF exposure and maximum effect after 9 h of radiation [98].

Therefore, studies that focus on the minimum exposure times (MET) and minimum exposure regimens (MER) to produce biological effects can shed light on thresholds of cell capabilities. In addition, it is more likely to reduce the complexity of the EMF interaction targets in cell cultures and organisms by lowering the exposure times and regimens, which at least reduces the overall rise in temperature.

### REMFS mechanism at the quantum level

#### REMFS exposures perturbs intracellular quantum systems

The REMFS interaction with biological systems allocated to multiple layers, which themselves also radiate different EM frequencies, represents a complex subject due to the scattering problem and the many-body perturbation theory (MBPT) from layered structures [55, 99]. The challenge owes its complexity to the mathematical difficulties in describing electromagnetic interaction within intracellular structures, which can be exceedingly complex due to the extremely heterogeneous nature of physical and biological processes. The interactions between multiple biomolecules, atoms, ions, rough interfaces scattering, and the applied EMF exposures poses strong limitations to the development of comprehensive models for the effects of REMFS on biological system. Herein, we will describe the REMFS receptor and biological systems interaction as a consequence of quantum confinement of the interfacial water H<sub>2</sub>O by electronegative biomolecules and the electropositive nucleus [100, 101].

#### REMFS receptor

The main reason we hypothesized that there is one receptor for REMFS instead of multiple receptors for each frequency is the fact that a wide range of frequencies (Hz to GHz) can induce similar biological responses. One well recognized example of this phenomenon is observed in the heat shock response [24, 49–52]. The fact that billions of frequencies (from Hz to GHz) produce the same biological effect makes the possibility that there is a receptor for each type of electromagnetic frequency improbable. Rather, it is more likely that some common receptor mediates this effect [102]. This common receptor would have to be capable of responding to the oscillating energy of the EMF exposures within a wide range of frequencies to induce the activation of biological pathways [102]. If so, this receptor must be able to induce conformational changes, specifically biomolecules that result in a secondary structure formation that regulates transcriptional or post-transcriptional mechanisms.

Interestingly several studies have found that the interfacial water is an integral partner of biomolecules and the modulator of their activity [103]. In addition, interfacial water plays an important role in the structure of DNA and RNA, forming bridges between bases within the same strand or between two strands, and in proteins, stabilizing  $\beta$ -sheets and  $\alpha$ -helices [104]. In RNA, the first layer of the interfacial water is considered a part of nucleic acid structure because it defines structure, folding, and intra-molecular interactions [105, 106]. The explanation of the interfacial water as a modulator of biological activity is also proposed by Mentré. He suggested that the key properties of the interfacial water in the cell are



that H-bonds are cooperative, currents of protons, osmosis, hydrostatic pressure, density variations, and selective exclusions of ions. These changes make stronger and shorter H-bonds in the interfacial water with higher heat capacity than bulk water because more energy is necessary to break its H-bonds [103].

Experimental studies have demonstrated EMF interfacial effects can produce biological effects [107–109]. Beruto et al. grew *Chlorella vulgaris* with and without the application of a low intensity, low frequency EMF of about 3 mT for 30 days. These exposures produce a significant effect on the exothermic clusterization step. The investigators proposed that this effect is produced by the interfacial water of glycocalyx of the external region of microalgal membrane because it can interact with chemical species present in the environment [109].

The organization of the interfacial water depends on the type of bio-surface electrostatic forces [103]. The first layer of the interfacial water is in special quantum confinement, since the polar and charged groups of the nucleic acid interact with the surrounding H bonds [110] from the interfacial water. This interaction actively affects the structure, function, and H bond of biomolecules [111]. Such polar and charged groups are sources of electric fields [112], which are very important for proton transfer [113]. This proton transfer is based on the electropositive applied electric field (EF) from REMFS (16.22 V/m) on the interfacial water and the intracellular EF rearrangements from RNA HSR1 electronegative attraction of EF (– 30 to 100 kT/e) and the electropositive EF from the cytoplasm ( $38 \times 10^6$  V/m).

Data indicates that intracellular EF rearrangements cause an electrostatic confinement of the first layer of the interfacial water, which gives it quantum properties [114]. These interfacial water molecules are 'pseudo-immobilized' and, therefore, confined to sub-diffraction volume. Of special note, this electrokinetic confined or trapped water exhibits a quantum tunneling behavior [115]. These data suggest that the interfacial water is a potential target for the effects of REMFS, which is confirmed by other investigators that proposed that water is a sensor for low energy EMF [116].

### H bond under REMFS

The pico- to sub-picosecond lifetimes of H-bonds are too short for experimental techniques such as nuclear magnetic resonance (NMR) and dielectric spectroscopy time window, so it is hard to perform experiments with water under EMF [117]. However, in a recent study applying an intense THz pulse (peak electric field strength of 14.9 MV/cm) to liquid water led to increasing H-bond stretching and bending vibrations. [118]

Similar evidence that polarized REMFS radiation causes its biological effects comes from the fact that it induces the dissociation of water into its constituent elements [119]. Rao examined distilled water under a polarized 2.45 GHz exposure. The Raman spectra of the treated water showed significant changes in the O–H stretch bond, which predisposes to proton tunneling and protonation of the surrounding molecules. Interestingly, despite different experimental conditions such as frequency (THz versus Hz), most of the conclusions are consistent with the fact that very different REMFS can produce water dissociation [120].

Furthermore, EMF exposures strengthen and shorten the H-bonds on the surface of biomolecules. For example, when hemoglobin and bovine serum albumin in water solutions were exposed to 50 Hz, samples revealed a significant increase in the absorbance signal of the Amide II band and an up-shift toward the high energies after exposure. These results suggested that EMF exposures strengthened the H-bonds of the secondary structures of these proteins [121].

Although REMFS increases the rotational kinetic energy of bulk water, the effects on the water of the first layer surrounding biomolecules are most prominent. The water of the first layer in the vicinity of biomolecules has a forced orientation and cannot rotate easily. However, under REMFS, it can undergo large-amplitude librational motions [122, 123]. REMFS oscillations at the molecular level produce rotation of water molecules within the first layer of the interfacial water as they try to "flip" their polar directions to match the polarity of the radio wave radiation. As a result, the oscillating electric field from the REMFS forces the water dipole moments to reorient themselves [67], which affects the H-bond that connects the first layer of the interfacial water.

This effect can be observed in biological tissues, where all polar molecules, such as water, are forced to oscillate in phase with the field and on planes parallel to its polarization [66, 124]. This is one of the most important factors for the quantum effects of REMFS: the man-made polarization of the excitatory oscillation of REMFS on contact with the interfacial water of the bio-system. These oscillations have a lower frequency relative to the exposed quantum system. They will change the frequency of the system to the excitatory frequency [65] like a driven harmonic oscillator [91, 125]. The system is driven by energy imparted upon the harmonic oscillator continuously by an external force [126]. If the excitatory frequencies are slower, the oscillator frequency is pulled towards the excitatory frequency [126].

### REMFS produces proton tunneling

Proton tunneling is a type of quantum tunneling that causes the transfer of a proton in one site to the closest site isolated by a potential barrier. Proton tunneling is commonly related to H-bonds. The hydrogen atoms are linked to two non-hydrogen atoms via an H-bond at one end and a covalent bond at the other.

H-bonds are classified based on energy or on geometry [127]. Several studies have identified the transitions from weak to moderate to strong H-bonds and the physical bases of the main geometry-based H-bond strength classifications. In this study, we use the geometric classification where the hydrogen bond is very strong when the distance between donor and acceptor atoms is in the range 2.2–2.5 Å, strong if it is in the range 2.5–3.2 Å, and weak if it is in the range 3.2–4.0 Å [128].

As mentioned above EMF decrease the water H-bond length and increase the H-bond angles as a function of the large amplitude motions [129] (Eq. 2). These large amplitude vibrations decrease the H-bond length of the first layer of the interfacial water to the oxygen of the nucleic acid (O–H...O) to values below 1.85 Å [130], which are ideal lengths for rising the probability of proton tunneling [131, 132]. This effect depends on the spatial location of the molecule with respect to the field [133], affecting the HBN [134] in an anisotropic manner. Thus, as the front water molecule is rotated [135] and hydrogen pairs with net dipole moment, the new configuration changes the hydrogen-bonding energy and distance. This water reorientation is of importance in multiple quantum processes, including proton transfer [136], proton transport [137, 138], and hydration of RNA or proteins for their function [139].

The amplitude of the H-bond varied between 0.18 and 0.22 Å for liquid water at 25 °C [129, 140]. In ab initio calculations large-amplitude motions caused by EMF exposures affect the bond distances and decrease the barrier distance from the H of the interfacial water to the O from the carbonyl group of the nucleic acid, consequently increasing the probability of proton tunneling [141] (Eq. 3).

Evidence that EMF cause tunneling is shown in a combined experiment and computer simulation demonstrated that the Hydrogen–Oxygen (H...O) distance is critical for tunneling and the rotation of the hydrogen from the confined water toward the O modulates the H...O distance [142]. Transitions arising from both pure rotation and rotation-tunneling can occur [143]. These data indicate thermodynamically balanced motions that control the donor–acceptor distance and site of active electrostatics, developing conformations apt for proton tunneling [144].

Additionally, a mathematical model found that changing the H-bond length by the radius of a hydrogen atom (0.05 nm) changes the transmission coefficient or tunneling current by 210%, suggesting an extreme sensitivity of tunneling to distance changes on the scale of atomic dimensions [145]. Furthermore, dynamical complexity increases with the exposure to the REMFS frequency during barrier penetration of the tunneling process [141]. For example, studies have found evidence for Coherent Proton Tunneling in a HBN at a tunneling frequency of 35 MHz, a frequency somewhat close to the REMFS exposures (64 MHz) used in our experiments [146].

Further evidence that supports tunneling in REMFS is a quantum calculation study that found that the flipping processes of water under quasi-one-dimensional (1D) confinement produces quantum tunneling effects [147]. The confinement in this study is very similar to the electrostatic confinement of the interfacial water of the HSR1 mentioned above during REMFS exposures.

In addition, several studies have found that EMF cause proton tunneling to produce tautomers in nucleic acids. Cerón-Carrasco found that electric fields induce proton transfer which produce tautomers in nucleic acids using Quantum Mechanical (QM) calculations [148]. The electric field decreased the potential barrier leading to the tautomer by 20–55 kJ mol<sup>-1</sup>. The study concluded that in the presence of EFs, only Guanine-Cytosine fit the necessary kinetic criteria to be considered a viable route to formation of tautomers.

In a more recent study Cerón-Carrasco investigated a more accurate DNA fragment in a simulation. The study found that at higher electric fields, tautomers are more stable than canonical bases [93]. In a classical molecular dynamics study, Cerón-Carrasco found that a continuous electric field exposure produces conformational changes in nucleic acids in 10 picoseconds [149].

Furthermore, Gheorghiu found that EM effects occur when the electric field is parallel to the H-bond axis [150]. Parallel electric fields were found to have a great influence on the energetics of the Guanine-Cytosine proton transfer tautomerism. It is important to consider that these effects occur under high electric fields like the found in the intracellular interfacial water of the HSR1 as mentioned above.

All these data suggest that REMFS promotes proton tunneling by oscillations that increase the amplitude of the H-bond vibrations of the interfacial water and modulate proton-acceptor distance, which increases the probability of tunneling proportional to the amplitude and the proton-acceptor distance [141, 151]. This protonation creates tautomers in RNA and DNA that affect biological changes.

**Mathematical model**

Our challenge was how to explain why a low energy wave causes biological effects. Therefore, we hypothesized the quantum effects of REMFS. We established a numerical model for the interaction between the REMFS exposures and biological systems at the quantum level [152]. For simplicity, we divided them into three stages, each with its equation (see Fig. 2):

*Stage 1.* REMFS vibrating energy produces a time-dependent adiabatic perturbation on the H-bond of the first layer of the interfacial water (FLIFW) [152]. During the REMFS exposures that increase the amplitude of the H-bond vibrations, the H-bond of the FLIFW changes state to a driven quantum harmonic oscillator. REMFS affects the H-bond situated near the oxygen (O) of the guanine of the RNA (GRNA). The following formula (Eq. 1) estimates the amplitude increment of H-bond oscillation as a driven quantum harmonic oscillator (see Fig. 3) system under REMFS [70].

*Equation 1:*

First, we obtain the time dependent periodic force:

$$F(t) = \frac{A \cos(\omega t + \varphi)}{\sqrt{2m\hbar\omega_0}} \tag{1.1}$$

Then we obtain the amplitude:

$$A = \frac{F(t)\sqrt{2m\hbar\omega_0}}{\cos(\omega t + \varphi)} \tag{1.2}$$

where: A=amplitude,  $\omega$ =vibration frequency of the periodic force,  $\omega_0$  = frequency of the oscillator, m=mass of the oscillator,  $\phi$  = phase of the driving field, t=time of

the exposure, F (t) = time dependent periodic force, and  $\hbar$  = reduced Planck's constant. [70]

In our experiments,  $F = q (E + v \times B)$ ,  $q = 1.062 \times 10^{-19}$  C (proton charge),  $v = 3 \times 10^8$  m/sec,  $E = 17$  V/m = 17 N/C,  $B = 3.89 \times 10^2$  A/m =  $146.2 \times 10^3$  V/m =  $146.2 \times 10^3$  N/C,  $m = 1.673 \times 10^{-27}$  k (proton mass),  $\hbar = 1.054 \times 10^{-34}$  J/s,  $t = 3,600$  s,  $\varphi = 0$ ,  $\omega = 64$  MHz, and  $\omega_0 = 74$  THz.

*Stage 2.* The value of the calculated H–O distance is very short (1.85 Å) [153]. REMFS shortens the distance between H of the FLIFW to O of the GRNA by increasing amplitude of the H-bond vibration. The change in the H-bond distance as a function of the amplitude of the oscillation is calculated in Eq. 2 [129].

*Equation 2:*


First, we find the average over the inter-nuclear configurations of the interfacial water and RNA nucleic acids:

$$r_a^{exp} = \langle r \rangle = \int F_r(q)P(q)dq \tag{2.1}$$


where:  $r_a^{exp}$  = the average over the inter-nuclear configurations of the interfacial water and RNA nucleic acids,  $F_r(q)$  = Variation of Bond distance,  $P(q)$  = Probability function.

The equation of P (q) in the classical Boltzmann approximation is:

$$P(q) = 1/N \int \exp\left(-\frac{V(q)}{RT}\right) dq \tag{2.2}$$



**Fig. 2** REMFS quantum effects on the first layer of the interfacial water of RNA. REMFS (repeated electromagnetic field stimulation)



where:  $V(q)$  = Potential function of floppy motion of the molecule,  $N$  = Normalization constant,  $q$  = amplitude coordinate of the oscillations.

Then, the dynamical correction term  $\delta_{dyn}$  is:

$$\delta_{dyn} = F_r(q_{min}) - r_a^{exp} \tag{2.3}$$

where:  $r_a^{exp}$  = operational parameter determined from the least squares fit to the experimental electron diffraction intensity curves.  $F_r(q_{min})$  = bond distance at the minimum of the potential function.

This equation predicts the shortening of the H-bond of the interfacial water under the time dependent perturbation caused by REMFS.

*Stage 3.* REMFS oscillation increases the amplitude of the H bond and decreases distance of the H from the FLIFW to the O from the GRNA, predisposing for H tunneling [70]. The probability of tunneling is proportional to the square of the amplitude. The barrier thickness or the decreased distance estimates the quantum tunneling probability by Eq. (3) [141].

*Equation 3:*

$$-\frac{\hbar^2}{2m} \frac{d^2\Psi(x)}{dx^2} = (E - U_0)\Psi(x) \tag{3.1}$$

where:  $E < U_0$ .

Hydrogen energy

Energy of the incoming particle/ Hydrogen energy  $\epsilon = 0.306$  eV, mass ( $m$ ) =  $1.673 \times 10^{-27}$  kg, height of barrier ( $U_0$ ) =  $5.099$  eV, thickness ( $d$ ) =  $2.6$  nm, attenuation factor ( $\alpha$ ) =  $0.475678 \times 10^{12}$  1/m, ratio of the exit and incident amplitudes =  $0.999952433311635$

$$\Psi = Ae^{-\alpha x}, \text{ where } \alpha = \sqrt{\frac{2m(U_0 - E)}{\hbar^2}} \tag{3.2}$$

where:  $E < U_0$ .

Energy of the incoming particle/Hydrogen energy ( $E$ ) =  $0.306$  eV, mass ( $m$ ) =  $1.673 \times 10^{-27}$  kg, height of barrier ( $U_0$ ) =  $5.099$  eV, thickness ( $d$ ) =  $2.6$  nm, attenuation factor ( $\alpha$ ) =  $0.475678 \times 10^{12}$  1/m, ratio of the exit and incident amplitudes =  $0.999952433311635$ , This equation finds the probability of proton tunneling.

These three stages and their respective equations have allowed us to develop a numerical model that can predict why the time dependent perturbation produced by REMFS alter the H-bond of the water of first layer that surrounds biomolecules. This H-bond acts as a quantum damped harmonic oscillator to increase the probability of tunneling and allows for the protonation of the nucleic acids of the surrounding RNA that in turn activates biological pathways.

## REMFS mechanism at the molecular level

### Nucleic acids tautomers cause conformational changes in RNA and DNA

In RNA, nucleic acid bases occur in several tautomeric forms due to protonation of the nucleobases [95]. Tautomers [154] are used by multiple RNA to produce their functions [95, 155].

Furthermore, it is well known that tautomeric equilibria are affected by several chemical and physical factors such as metals, temperature [156], pH [156] and recently EMF exposures [93, 157]. Also, tautomer inter-conversions can adopt various secondary structures responsible for a variety of functions during biological processes like RNA conformational changes, DNA replication, packaging, and transcription [158, 159]. Often, such conformational changes promote binding to activating factors that in turn affect transcription and translation of proteins.

Tautomerism causes conformational changes in the catalysis of self-cleaving RNA [95] to produce a wide array of biological functions [160–162]. The importance of tautomerism in the function of a biomolecule is highlighted in all crystal structure changes of TPP dependent riboswitches. TPP is in an extended conformation, with the orientation of the thiazolium and pyrimidine rings within hydrogen-bonding distance [163]. Here, a glutamic acid protonates and stabilizes the imino tautomer of the pyrimidine ring to lead to the ylide conformation bound in either the TPP-dependent riboswitch or a TPP-dependent enzyme [164].

Investigators using quantum chemical computations detected all possible protonated base pairs in RNA crystal structures. Data showed 18 different protonated base pair combinations from RNA and proposed a theoretical model for base pair combination [165].

Guanine and cytosine protonation affects RNA structure and function. An interesting QM analyses suggested that the guanine protonation can be a crucial factor in structure and function of RNAs [166]. The different RNA structures come from the changes of single and double bonds in the ring systems of purines and pyrimidines [167]. In a theoretical study, Chaudry calculates the quantum source of the tautomerism in DNA [168]. This tautomerism shows enhancement under EMF exposures.

Evidence suggests that REMFS protonates biomolecules [60], resulting in important tautomeric interconversions and conformational changes [95, 169]. Experimental and theoretical studies show that externally applied electric and EMF produce biologically relevant tautomers and conformational changes in RNA. An experimental study showed the effects of electric fields on RNA conformational changes and orientation by ultraviolet absorbance and electric dichroism [170]. In another study they found long-lived conformational changes in RNA by electrical

impulses. The researchers applied an electric field of about 20 kV/cm to induce large dipole moments by shifting the ionic atmosphere, which caused strand repulsion and conformational changes in the RNA. Interestingly, the fields used were of the same intensity as those found in nerve transmission.

Furthermore, other biomolecule structures are affected by REMFS. De Ninno observed structural changes in Glutamic acid induced by exposure of REMFS at 50 Hz. The IR spectra contained stretching and bending bands of the protonated COOH which is attributable to the coupled C-O stretch and O-H bend of the COOH group [171]. Another study determined the tautomeric protonation of N-methyl piperazine. They performed theoretical calculations and practical experiments. These results suggest that proton relocation occurs by solvent assistance in water or proton jump. They found that predicted activation free energy was about 10 kcal/mol based on variable temperature nuclear magnetic resonance experiments [172].

All these data suggest that REMFS can cause tautomerism and conformational changes in RNA. This mechanism is similar to the regulation of HSR by RNA thermometers [173] in bacteria [174].

### REMFS activates HSF1 and chaperone expression

REMFS, heat, alcohol, hypoxia, metal ions, peroxide, amino acid analogs, and other stressors activate HSF1 and the HSR [175]. Most of these factors cause denaturation and accumulation of abnormal proteins, which induce the HSR [176, 177]. However, REMFS exposures are not likely to produce protein denaturation, so the mechanism must be related to an EMF-sensitive biomolecule such as HSR1. EMF exposure also increases HSF1-heat shock element binding activity, thereby directly contributing to the activation of HSF1 and the stress-induced Hsp70 [175] transcription and translation in cells exposed to REMFS [178, 179].

HSF1 is a transcriptional factor that is a master regulator of stress gene expression (molecular chaperones) [180]. Recently, in addition to chaperone expression, accumulating evidence indicates multiple additional functions for HSF1 beyond chaperone production. HSF1 acts in diverse stress-induced cellular processes and molecular mechanisms, including the endoplasmic reticulum, unfolded protein response and ubiquitin-proteasome system, multidrug resistance, autophagy, apoptosis, immune response, cell growth arrest, differentiation underlying developmental diapause, chromatin remodeling, cancer development, and aging [181].

REMFS produces biological effects through HSR1 [182] which activates HSF1. HSR1 employs a similar mechanism as that of bacterial RNA thermometers to sense

temperature and energy changes in the cell and ultimately regulate the translational machinery [183]. HSR1 is a long non-coding RNA that undergoes conformational changes from a close to an open structure under thermal radiation exposure (THz to GHz frequencies). These conformational changes in HSR1 are required for the binding and activation of HSF1 [90]. Computer simulations reveal that HSR1 is composed of an extensive secondary structure that changes predictably within a physiological range of temperatures [90] and EMF exposures without heating [184].

Another important co-factor in the activation of HSF1 is the translation elongation factor (eEF1A), which is a key component regulating the actin cytoskeleton architecture in the cell [185]. A full HSF1 activation requires a combination of purified HSR1 and eEF1A in-vitro at physiological concentrations [186]. Under normal conditions HSR1 is present in an inactive “closed” conformation. During heat shock or EMF exposures, HSR1 “switches” to the “open” conformation that activates HSF1 and releases it from its repressor Hsp90, while after a stress a massive release of eEF1A from cytoskeleton collapse (from misfolded proteins) can then fully activate the newly freed HSF1 [90]. In contrast, under REMFS exposure alone, there is no cytoskeleton collapse [43]. The role of REMFS in this process is to promote binding of HSR1 to HSF1, with a subsequent release of HSF1 from its repressor Hsp90 [43].

Triggering the HSR by stressors after REMFS treatment produces a fast and vigorous expression of Heat shock proteins (Hsps) [187]. Protein aggregation is an important factor in the progression of aging and age-related diseases such as AD [188]. Several pathways are associated with abnormal protein clearance, including molecular chaperones, the ubiquitin–proteasome system, and autophagy pathways [189]. The production of these chaperones depends on the activation of HSF1, an event attenuated by the aging process [190]. HSF1 is repressed by the Hsp90 complex and released to get activated under several cellular stresses [191].

Similarly, REMFS exposure also releases HSF1 from the Hsp90 complex [43]. Once released from the Hsp90 complex, it trimerizes spontaneously to bind DNA, an event that produces increased amounts of Heat shock proteins such as Hsp70, Hsp90, etc. Once chaperones are produced, they bind abnormal proteins. Excess Hsp90 binds to HSF1 trimers and causes them to dissociate and revert once again to the inactive, monomeric state [192].

The HSF1 effects on Hsps play a role in aging [193] and protein accumulation diseases [188]. The role of HSF1 in the aging process and age-related diseases such as AD suggests a deeper relationship between the molecular mechanisms of these two processes. HSF1 activation

prevents the decline in proteostasis, the primary contributor to aging, thus delaying the aging process [39, 194, 195]. This suggests a potential role for HSF1-based therapeutic tools, such as REMFS, in the treatment of a wide array of age-related diseases [196].

Hence, it is here where molecular chaperones such as Hsp70 take on an essential role by deterring protein aggregation [197]. Two protein degradation pathways, macroautophagy and chaperone-mediated autophagy (CMA) [198], undergo age-dependent decline probably subsequent to the age-related attenuation of the HSF1 [199, 200], which is an early molecular event in the aging process [201].

### REMFS upregulates HSF1 to promote A $\beta$ autophagy

REMFS studies have demonstrated that repeated exposures increase Hsp70 levels by activation of HSF1 [6, 43] and autophagy [26]. Additionally, HSF1 upregulates ATG7 and RIPK1 to promote autophagy [202, 203]. Several studies have suggested that heat shock proteins prevent protein accumulation such as amyloid deposition [204–207]. Genetic overexpression of HSF1, reduces A $\beta$  levels, induces autophagy, and up-regulates production of chaperones. Increased expression of HSF1 ameliorates AD-like cognitive deficits in PDAPP transgenic mice, which produce excess levels of human APP [208]. Furthermore, Hsp70 promotes degradation and inhibits accumulation of amyloid [204, 209–211]. Hsp70 decreased A $\beta$  levels when given to microglia from rats [212]. Hsps binds to APP and decreases the levels of A $\beta$ 40 and A $\beta$ 42 [213]. 78-kDa glucose-regulated protein (GRP78) is another member of the HSP70 family with a role in AD. In a HEK cell model co-transfected with APP and GRP78, this Hsp70 binds to APP in the ER, preventing the  $\beta$ / $\gamma$ -secretase cleavage necessary to produce A $\beta$ , thereby decreasing A $\beta$  intracellular toxicity [214]. In addition, the overexpression of GRP78 decreases the level of A $\beta$ 40 and A $\beta$ 42 in mutant APP (APPsw) cells [214].

Hsp70 binds to APP by the KFERQ motif (see Fig. 1). HSP70 transports APP to lysosomes for CMA or endosomal microautophagy (eMI) for degradation to reduce A $\beta$  oligomers levels [215]. Additionally, many pathogenic proteins including tau,  $\alpha$ -synuclein, and huntingtin are degraded by CMA [216–218]. Hsps binds to these proteins and degrades them through the CMA or proteasome system [219]. In an interesting study, they modified A $\beta$  as a substrate for CMA and eMI (termed as Hsc70-based autophagy) by tagging its oligomers with multiple CMA motifs. This method significantly reduced A $\beta$  oligomers in induced pluripotent stem cell (iPSC), which are cortical neurons derived from AD patient fibroblasts [220].

Hsp70 also suppresses oligomerization of A $\beta$  by binding to the hydrophobic region to modify their conformation [219]. Structural changes in oligomers occurred when Hsps interacted with oligomers and fibrils. However, Hsps did not cause any direct effect on fibrils, suggesting that Hsps suppress the early stages of self-assembly [221–224]. Taken together these studies confirm that the activation of HSF1 and subsequent increase in chaperone levels, especially Hsp70, by EMF exposures influence A $\beta$  degradation pathways [225–227] by autophagy [228, 229] to lower A $\beta$  levels.

### REMFS decreases A $\beta$ levels in primary human brain cultures

We recently utilized REMFS to lower A $\beta$  levels in cell cultures of primary human mixed brain (PHB). REMFS treatment decreased A $\beta$ 40 and A $\beta$ 42 levels without evidence of toxicity. Treatment started on day 7 in vitro (DIV 7). After 14 days of REMFS, we measured levels of A $\beta$ 40 peptide in exposed and non-exposed cells. The REMFS parameters were frequency of 64 MHz with a SAR of 0.6 W/Kg for 1 h daily; this treatment achieved a 46% reduction in A $\beta$ 40 levels ( $p=0.001$ ,  $g=0.798$ ), compared to the non-treated cultures. The same REMFS parameters achieved a 36% decrease in A $\beta$ 42 levels. Subsequently, we demonstrated that REMFS at 64 or 100 MHz with a lower SAR of 0.4 W/kg for 14 days achieved a comparable reduction in A $\beta$ 40 and A $\beta$ 42 levels. Furthermore, when we increased the exposure time from 1 to 2 h, there was a similar reduction in the A $\beta$  levels. Additionally, when we increased the frequency from 64 to 100 MHz, we found a comparable difference in A $\beta$  levels. The results of our experiments suggest that REMFS at 64 MHz with a SAR of 0.4 W/kg for 1 h (typical of that already utilized in clinical MRI contexts) would be the minimum energy needed to produce bio-effects in human neurons, specifically a reduction in levels of toxic A $\beta$  peptides.

### REMFS safety

In contrast to our experiments some studies have shown possible harmful effects of REMFS on biological systems which resulted from longer exposure times (minutes vs days), higher power (0.5 vs 5 W/m) and higher SAR (0.5 vs.10 W/kg) [230]. The energy produced by REMFS is extremely low, making it unlikely that our studied REMFS exposures would lead to adverse health effects, especially as we did not observe any evidence of cellular toxicity or morphological changes in our cell culture experiments [43]. Furthermore, long-term mouse AD experiments (daily for up to 9 months) and a recent phase 1 clinical trial with REMFS were safe, with no toxic effects observed on multiple safety factors evaluated.

Specifically, there were no toxic effects on brain oxidative stress, brain histology, brain heating, DNA in circulating blood cells, and changes in peripheral tissues [35, 36, 55]. Additionally, our experiments use the same REMFS frequency that has been used by MRI machines for decades. Since their implementation for clinical imaging, MRI exposures have had no demonstrable negative health impacts [231]. Lastly, a recent phase one clinical trial in AD did not find any behavioral side-effect, pain, tumor growth, hemorrhage, or abnormal physiological responses after 2 months of treatment with REMFS.

### Future perspectives

Any organ that shows functional decline, including the brain, kidneys, joints, liver, or heart, may benefit from engineered REMFS exposures to induce protein disaggregation by activation of the HSF1 pathway and autophagy. Therefore, we will initiate human head exposure to treat the protein aggregation caused by AD. The major technical difficulties for developing an exposure system are the human head geometry, the multiple tissue layers of the head, and development of an antenna that produce a homogeneous SAR on the whole brain.

Therefore, before clinical trials are considered we must determine the best electromagnetic settings for human exposures such as power output, power deposition, far field, antenna type, distance from antenna, electric field, magnetic field, etc. that will produce homogeneous internal fields when applied to a human brain with a target SAR of 0.4–0.9 W/kg. We will start with determining by mathematical and computer modeling the REMFS exposures in our biological studies that deliver safe thermal and SAR measurements to the human head [62].

Using these results, we will develop a virtual exposure system by numerical model and computer simulation. We will design a virtual antenna that delivers a SAR of around 0.6 W/kg to a simulated phantom of a human brain. With these simulations we will find the REMFS parameters that will deliver a homogeneous radiation to a human head in clinical trials [232]. In the near future, we will experimentally confirm these results using an appropriate practical antenna to expose a Specific Anthropomorphic Mannequin (SAM) human head phantom [233] with internal and external probes oriented vertically to determine the EMF parameters that will provide an effective and safe SAR for future AD treatment. Data suggest that the ideal environment for these treatments should be an anechoic chamber to prevent RF wave reflections and provide a uniform exposure to the subjects [234]. The final step will be to initiate phase 1 clinical trials in patient with early AD to determine safety and efficacy of this new potential therapeutic strategy.

## Conclusion

The current study proposes a multidimensional mechanism at the quantum, molecular, cellular, and organismal level inside a theoretical framework that may explain the results of our experiments, and those of other investigators. The proposed quantum tunneling mechanism here is the first to provide an explanation of how low energy radio-frequency radiation may induce a biological response. Quantum tunneling allows for an understanding of events occurring between single photons and biomolecules that would otherwise be extremely difficult to visualize in experimental studies. Hence, it is by way of quantum tunneling that we finally understand the intimate relationship between REMFS and the HBN of the interfacial water of biomolecules. The process is a time dependent adiabatic perturbation of the HBN that is set into motion as a photon carried along an EM wave (with a frequency lower than the H bond frequency) that forces the H bond to change its frequency to that of the EM wave, thereby increasing the amplitude of the H bond vibrations in a process similar to a driven quantum oscillator [42]. The increased amplitude will decrease the H bond donor–acceptor distance and result in an increased probability of proton tunneling [235]. Consequently, interfacial water will donate its hydrogen toward protonation of nucleic acids, and the tautomeric interconversions that ensue result in structural changes in biomolecules and RNA, namely HSR1. The secondary structure produced will then bind to HSF1 and cause its dissociation from the multi-chaperone complex, freeing it from inhibition. Once activated, the HSF1 monomer undergoes trimerization and accumulation, inducing the expression of Hsp70 and thereby activating A $\beta$ -clearance pathways to delay cellular senescence.

Data suggests HSR attenuation goes hand-in-hand with aging, and may even be the initial event in the aging process [201], presumably due to a decrease in HSF1-DNA binding [199]. Hence, the failure of proteostasis associated with aging may be the initial event in the development of AD [39]. Consequently, it is plausible that a treatment that enhances or restores HSF1-DNA binding would improve the loss of the proteostasis observed with aging. Therapeutic implications could also be expanded to involve other age-related diseases associated with protein accumulation, such as AD, PD, LBD, and frontotemporal diseases.

The theory herein comprises an important framework that lays the foundation for understanding the interactions between EMF and organisms and provides a valuable contribution to the foundational principles that should underlie any discussion on the biological effects of EMF stimulation. Importantly, the results of our literature review and research also point to the capacity of

REMFS to influence various networks within known biological systems dysregulated in AD. The potential implications of REMFS as a therapeutic modality are likely to be far in the future, but the ability of RF-EMF to significantly reduce A $\beta$ 40 and A $\beta$ 42 levels in human neurons, coupled with animal model results, indicate a pathway worth further exploration. These results in cell and animal systems are likely achieved through a combination of efficient A $\beta$  degradation, autophagy-lysosome system [26], and proteasome system activation [23], as well as the reduction of  $\beta$ -secretase activity [56]. Yet, we must note that quantum tunneling-based model could explain the conformational changes of molecules involved in other biological pathways not mentioned here.

The use of REMFS as a non-invasive therapy for the management of AD holds promise and the results of a recent phase 1 clinical trial confirm its safety in humans [236]. Nevertheless, it remains necessary to take into account the complex network of genetic [237–239] and epigenetic [240] effects occurring under REMFS. As regulation pathways triggered by REMFS have yet to be clearly elucidated, current knowledge of EMF-biological systems interaction and possible adverse effects remain limited. Quantum and classical molecular computer simulations, complemented by in vitro and in vivo laboratory studies as well as clinical trials, are needed to investigate the initial and late effects of REMFS. These studies will help develop the conditions useful for its therapeutic use while avoiding any possible adverse effects. Finally, there is a need to perform mathematical modeling and computer simulation that elucidates the appropriate EMF settings for human treatments [62, 232]. Regardless, the theories that we have proposed provide the framework for observed outcomes in several cellular and animal studies that prove the potential therapeutic implications of REMFS on age-related diseases in humans.

### Acknowledgements

The authors thank Wesley Fry for his assistance with the graphics.

### Author contributions

FP, JB, CP, DL, JM, and MR established review idea, information collection and were major contributors in writing the manuscript. All authors read and approved the final manuscript.

### Funding

This work was supported by Grant NIHMS (831498).

### Availability of data and materials

Not applicable.

### Declarations

### Ethics approval and consent to participate

Not applicable.

### Consent for publication

Not applicable.



**Competing interests**

The authors declare that they have no competing interests.

**Author details**

<sup>1</sup>Indiana University School of Medicine, Indianapolis, IN, USA. <sup>2</sup>Division of General Internal Medicine and Geriatrics, Department of Medicine, Indiana University School of Medicine, Indianapolis, IN, USA. <sup>3</sup>Division of Rheumatology, Department of Medicine, Indiana University School of Medicine, Indianapolis, IN, USA. <sup>4</sup>Department of Psychiatry, Institute of Psychiatric Research, Neuroscience Research Center, Indiana University School of Medicine, Indianapolis, IN, USA. <sup>5</sup>Department of Medical and Molecular Genetics, Indiana University School of Medicine, Indianapolis, IN, USA. <sup>6</sup>Department of Bioengineering, University of Illinois at Chicago, Chicago, IL, USA. <sup>7</sup>Department of Electrical and Computer Engineering, Indiana University-Purdue University, Indianapolis, IN, USA.

Received: 20 December 2021 Accepted: 7 June 2022

Published online: 13 June 2022

**References**

- Rosen A, Stuchly MA, Vorst AV. Applications of RF/microwaves in medicine. *IEEE Trans Microw Theory Tech*. 2002;50:963–74.
- Michaelson SM. Health implications of exposure to radiofrequency/microwave energies. *Br J Ind Med*. 1982;39:105–19.
- Singh S, Kapoor N. Health implications of electromagnetic fields, mechanisms of action, and research needs. *Adv Biol*. 2014;2014:24.
- Hardell L, Sage C. Biological effects from electromagnetic field exposure and public exposure standards. *Biomed Pharmacother*. 2008;62:104–9.
- Gherardini L, Ciuti G, Tognarelli S, Cinti C. Searching for the perfect wave: the effect of radiofrequency electromagnetic fields on cells. *Int J Mol Sci*. 2014;15:5366–87.
- Lin H, Opler M, Head M, Blank M, Goodman R. Electromagnetic field exposure induces rapid, transitory heat shock factor activation in human cells. *J Cell Biochem*. 1997;66:482–8.
- Luukkonen J, Liimatainen A, Juutilainen J, Naarala J. Induction of genomic instability, oxidative processes, and mitochondrial activity by 50Hz magnetic fields in human SH-SY5Y neuroblastoma cells. *Mutat Res*. 2014;760:33–41.
- Campisi A, Gulino M, Acquaviva R, Bellia P, Raciti G, Grasso R, Musumeci F, Vanella A, Triglia A. Reactive oxygen species levels and DNA fragmentation on astrocytes in primary culture after acute exposure to low intensity microwave electromagnetic field. *Neurosci Lett*. 2010;473:52–5.
- Conti P, Gigante GE, Alesse E, Cifone MG, Fieschi C, Reale M, Angeletti PU. A role for Ca<sup>2+</sup> in the effect of very low frequency electromagnetic field on the blastogenesis of human lymphocytes. *FEBS Lett*. 1985;181:28–32.
- Rozek RJ, Sherman ML, Liboff AR, McLeod BR, Smith SD. Nifedipine is an antagonist to cyclotron resonance enhancement of <sup>45</sup>Ca incorporation in human lymphocytes. *Cell Calcium*. 1987;8:413–27.
- Fesenko EE, Geletyuk VI, Kazachenko VN, Chemeris NK. Preliminary microwave irradiation of water solutions changes their channel-modifying activity. *FEBS Lett*. 1995;366:49–52.
- Pall ML. Electromagnetic fields act via activation of voltage-gated calcium channels to produce beneficial or adverse effects. *J Cell Mol Med*. 2013;17:958–65.
- Byus CV, Pieper SE, Adey WR. The effects of low-energy 60-Hz environmental electromagnetic fields upon the growth-related enzyme ornithine decarboxylase. *Carcinogenesis*. 1987;8:1385–9.
- Goodman R, Henderson AS. Mechanistic approaches to interactions of electric and electromagnetic fields with living systems. New York: Plenum Press; 1987.
- Liboff AR, Williams T Jr, Strong DM, Wistar R Jr. Time-varying magnetic fields: effect on DNA synthesis. *Science*. 1984;223:818–20.
- Pezzetti F, De Mattei M, Caruso A, Cadossi R, Zucchini P, Carinci F, Traina G, Sollazzo V. Effects of pulsed electromagnetic fields on human chondrocytes: an in vitro study. *Calcif Tissue Int*. 1999;65:396–401.
- Capelli E, Torrisi F, Venturini L, Granato M, Fassina L, Lupo GFD, Ricevuti G. Low-frequency pulsed electromagnetic field is able to modulate miRNAs in an experimental cell model of Alzheimer's Disease. *J Healthc Eng*. 2017;2017:10.
- Liu Y, Liu WB, Liu KJ, Ao L, Cao J, Zhong JL, Liu JY. Extremely low-frequency electromagnetic fields affect the miRNA-mediated regulation of signaling pathways in the GC-2 Cell Line. *PLoS ONE*. 2015;10:e0139949.
- Osera C, Amadio M, Falone S, Fassina L, Magenes G, Amicarelli F, Ricevuti G, Govoni S, Pascale A. Pre-exposure of neuroblastoma cell line to pulsed electromagnetic field prevents H<sub>2</sub>O<sub>2</sub>-induced ROS production by increasing MnSOD activity. *Bioelectromagnetics*. 2015;36:219–32.
- Osera C, Fassina L, Amadio M, Venturini L, Buoso E, Magenes G, Govoni S, Ricevuti G, Pascale A. Cytoprotective response induced by electromagnetic stimulation on SH-SY5Y human neuroblastoma cell line. *Tissue Eng Part A*. 2011;17:2573–82.
- Han L, Lin H, Head M, Jin M, Blank M, Goodman R. Application of magnetic field-induced heat shock protein 70 for presurgical cytoprotection. *J Cell Biochem*. 1998;71:577–83.
- Kato T. On the adiabatic theorem of quantum mechanics. *J Phys Soc Jpn*. 1950;5:435–9.
- Eleuteri AM, Amici M, Bonfili L, Cecarini V, Cuccioloni M, Grimaldi S, Giuliani L, Angeletti M, Fioretti E. 50 Hz extremely low frequency electromagnetic fields enhance protein carbonyl groups content in cancer cells: effects on proteasomal systems. *J Biomed Biotechnol*. 2009;2009:834239.
- Caraglia M, Marra M, Mancinelli F, D'Ambrosio G, Massa R, Giordano A, Budillon A, Abbruzzese A, Bismuto E. Electromagnetic fields at mobile phone frequency induce apoptosis and inactivation of the multi-chaperone complex in human epidermoid cancer cells. *J Cell Physiol*. 2005;204:539–48.
- Hirai T, Taniura H, Goto Y, Ogura M, Sng JC, Yoneda Y. Stimulation of ubiquitin-proteasome pathway through the expression of amidohydrolase for N-terminal asparagine (Ntan1) in cultured rat hippocampal neurons exposed to static magnetism. *J Neurochem*. 2006;96:1519–30.
- Marchesi N, Osera C, Fassina L, Amadio M, Angeletti F, Morini M, Magenes G, Venturini L, Biggiogera M, Ricevuti G, Govoni S, Caorsi S, Pascale A, Comincini S. Autophagy is modulated in human neuroblastoma cells through direct exposition to low frequency electromagnetic fields. *J Cell Physiol*. 2014;229:1776–86.
- Kubat NJ, Moffett J, Fray LM. Effect of pulsed electromagnetic field treatment on programmed resolution of inflammation pathway markers in human cells in culture. *J Inflamm Res*. 2015;8:59–69.
- Jouni FJ, Abdolmaleki P, Ghanati F. Oxidative stress in broad bean (*Vicia faba* L.) induced by static magnetic field under natural radioactivity. *Mutat Res*. 2012;741:116–21.
- Dasdag S, Akdag MZ, Erdal ME, Erdal N, Ay OI, Ay ME, Yilmaz SG, Tasdelen B, Yegin K. Effects of 2.4 GHz radiofrequency radiation emitted from Wi-Fi equipment on microRNA expression in brain tissue. *Int J Radiat Biol*. 2015;91:555–61.
- Hajnorouzi A, Vaezzadeh M, Ghanati F, Jamnezhad H, Nahidian B. Growth promotion and a decrease of oxidative stress in maize seedlings by a combination of geomagnetic and weak electromagnetic fields. *J Plant Physiol*. 2011;168:1123–8.
- Maaroufi K, Had-Aissouni L, Melon C, Sakly M, Abdelmelek H, Poucet B, Save E. Spatial learning, monoamines and oxidative stress in rats exposed to 900 MHz electromagnetic field in combination with iron overload. *Behav Brain Res*. 2014;258:80–9.
- Di Carlo AL, White NC, Litovitz TA. Mechanical and electromagnetic induction of protection against oxidative stress. *Bioelectrochemistry*. 2001;53:87–95.
- Pena-Philippides JC, Yang Y, Bragina O, Hagberg S, Nemoto E, Roitbak T. Effect of pulsed electromagnetic field (PEMF) on infarct size and inflammation after cerebral ischemia in mice. *Transl Stroke Res*. 2014;5:491–500.
- Rohde CH, Taylor EM, Alonso A, Ascherman JA, Hardy KL, Pilla AA. Pulsed electromagnetic fields reduce postoperative interleukin-1 $\beta$ , pain, and inflammation: a double-blind, placebo-controlled study in TRAM flap breast reconstruction patients. *Plast Reconstr Surg*. 2015;135:808e–17e.

35. Dragicevic N, Bradshaw P, Mamcarz M, Lin X, Wang L, Cao C, Arendash G. Long-term electromagnetic field treatment enhances brain mitochondrial function of both Alzheimer's transgenic mice and normal mice: a mechanism for electromagnetic field-induced cognitive benefit? *Neuroscience*. 2011;185:135–49.
36. Arendash GW, Mori T, Dorsey M, Gonzalez R, Tajiri N, Borlongan C. Electromagnetic treatment to old Alzheimer's mice reverses beta-amyloid deposition, modifies cerebral blood flow, and provides selected cognitive benefit. *PLoS ONE*. 2012;7: e35751.
37. Arendash GW, Sanchez-Ramos J, Mori T, Mamcarz M, Lin X, Runfeldt M, Wang L, Zhang G, Sava V, Tan J, Cao C. Electromagnetic field treatment protects against and reverses cognitive impairment in Alzheimer's disease mice. *J Alzheimers Dis*. 2010;19:191–210.
38. Veronesi F, Torricelli P, Giavaresi G, Sartori M, Cavani F, Setti S, Cadossi M, Ongaro A, Fini M. In vivo effect of two different pulsed electromagnetic field frequencies on osteoarthritis. *J Orthop Res*. 2014;32:677–85.
39. Perez FP, Bose D, Maloney B, Nho K, Shah K, Lahiri DK. Late-onset Alzheimer's disease, heating up and foxed by several proteins: pathomolecular effects of the aging process. *J Alzheimer's Dis*. 2014;40:1–17.
40. Diab KA. The impact of the low frequency of the electromagnetic field on human. *Cell Biol Transl Med*. 2019;7:135–49.
41. Alon L, Deniz CM, Carluccio G, Brown R, Sodickson DK, Collins CM. Effects of anatomical differences on electromagnetic fields, SAR, and temperature change. *Concepts Magn Reson Part B: Magn Reson Eng*. 2016;46:8–18.
42. Perez FP, Morisaki JJ, Bandeira JP. Repeated electromagnetic field stimulation in aging and health. In *The science of hormesis in health and longevity*. Amsterdam: Elsevier; 2019, pp 189–197.
43. Novoselova EG, Novikov VV, Lunin SM, Glushkova OV, Novoselova TV, Parfenyuk SB, Novoselov SV, Khrenov MO, Fesenko EE. Effects of low-level combined static and weak low-frequency alternating magnetic fields on cytokine production and tumor development in mice. *Electromagn Biol Med*. 2019;38:74–83.
44. Xu J, Liu K, Chen T, Zhan T, Ouyang Z, Wang Y, Liu W, Zhang X, Sun Y, Xu G. Rotating magnetic field delays human umbilical vein endothelial cell aging and prolongs the lifespan of *Caenorhabditis elegans*. *Aging (Albany NY)*. 2019;11:10385.
45. Memmi BK, Ünlü H. The effects of short duration microwave exposure on the life span and the induction of sex-linked recessive lethal mutations in *Drosophila melanogaster*. *Hacettepe J Biol Chem*. 2007;35:173–9.
46. Weisman NY, Fedorov V, Nemova E, Nikolaev N. Survival and life span of *Drosophila melanogaster* in response to terahertz radiation. *Adv Gerontol*. 2014;4:187–92.
47. Makarov VI, Khmelinskii I. External control of the *Drosophila melanogaster* lifespan by combination of 3D oscillating low-frequency electric and magnetic fields. *Electromagn Biol Med*. 2014;33:276–81.
48. Bua L, Tibaldi E, Falcioni L, Lauriola M, De Angelis L, Gnudi F, Manservigi M, Manservigi F, Manzoli I, Menghetti I. Results of lifespan exposure to continuous and intermittent extremely low frequency electromagnetic fields (ELFEMF) administered alone to Sprague Dawley rats. *Environ Res*. 2018;164:271–9.
49. Shi Z, Yu H, Sun Y, Yang C, Lian H, Cai P. The energy metabolism in *Caenorhabditis elegans* under the extremely low-frequency electromagnetic field exposure. *Sci Rep*. 2015;5:1–11.
50. Gao Y, Lu Y, Yi J, Li Z, Gao D, Yu Z, Wu T, Zhang C. A genome-wide mRNA expression profile in *Caenorhabditis elegans* under prolonged exposure to 1750MHz radiofrequency fields. *PLoS ONE*. 2016;11: e0147273.
51. Hung Y-C, Lee J-H, Chen H-M, Huang GS. Effects of static magnetic fields on the development and aging of *Caenorhabditis elegans*. *J Exp Biol*. 2010;213:2079–85.
52. Mattson MP. Pathways towards and away from Alzheimer's disease. *Nature*. 2004;430:631–9.
53. Arendash GW, Sanchez-Ramos J, Mori T, Mamcarz M, Lin X, Runfeldt M, Wang L, Zhang G, Sava V, Tan J, Cao C. Electromagnetic field treatment protects against and reverses cognitive impairment in Alzheimer's disease mice. *J Alzheimer's Dis JAD*. 2010;19:191–210.
54. Arendash GW. Transcranial electromagnetic treatment against Alzheimer's disease: why it has the potential to trump Alzheimer's disease drug development. *J Alzheimer's Dis JAD*. 2012;32:243–66.
55. Jeong YJ, Kang GY, Kwon JH, Choi HD, Pack JK, Kim N, Lee YS, Lee HJ. 1950 MHz electromagnetic fields ameliorate abeta pathology in Alzheimer's disease mice. *Curr Alzheimer Res*. 2015;12:481–92.
56. Ahmad RH, Fakhoury M, Lawand N. Electromagnetic field in Alzheimer's disease: a literature review of recent preclinical and clinical studies. *Curr Alzheimer Res*. 2020;17:1001–12.
57. Giacobini E, Gold G. Alzheimer disease therapy—moving from amyloid- $\beta$  to tau. *Nat Rev Neurol*. 2013;9:677–86.
58. Binhi VN, Rubin AB. Magnetobiology: the kT paradox and possible solutions. *Electromagn Biol Med*. 2007;26:45–62.
59. De Ninno A, Congiu Castellano A. Deprotonation of glutamic acid induced by weak magnetic field: an FTIR-ATR study. *Bioelectromagnetics*. 2011;32:218–25.
60. Pompa P, Bramanti A, Maruccio G, Cingolani R, De Rienzo F, Corni S, Di Felice R, Rinaldi R. Retention of native-like conformation by proteins embedded in high external electric fields. *AIP*. 2005.
61. Perez F, Millholland G, Peddinti SV, Thella AK, Rizkalla J, Salama P, Rizkalla M, Morisaki J, Rizkalla ME. Electromagnetic and thermal simulations of human neurons for SAR applications. *J Biomed Sci Eng*. 2016;9:437–44.
62. Nelson DL, Cox MM, Lehninger AL. *Principles of biochemistry*, fourth edition edition 1. 4th ed. New York: WH Freeman and Company; 2005. p. 2.
63. Goss DJ, Petrucci RH. *General chemistry principles and modern applications*, Petrucci, Harwood, Herring, Madura: Study Guide, Pearson/Prenice Hall. 2007.
64. Panagopoulos DJ, Johansson O, Carlo GL. Polarization: a key difference between man-made and natural electromagnetic fields, in regard to *Biological Activity*, 5, 14914.
65. Panagopoulos DJ, Messini N, Karabarbounis A, Philippetis AL, Margaritis LH. A mechanism for action of oscillating electric fields on cells. *Biochem Biophys Res Commun*. 2000;272:634–40.
66. English NJ, MacElroy J. Molecular dynamics simulations of microwave heating of water. *J Chem Phys*. 2003;118:1589–92.
67. Wilmink GJ, Rivest BD, Ibey BL, Roth CL, Bernhard J, Roach WP. Quantitative investigation of the bioeffects associated with terahertz radiation, In *Optical interactions with tissues and cells XXI*, pp 139–148, SPIE. 2010.
68. Hänggi P. Driven quantum systems, Quantum transport and dissipation, 250. 1998.
69. Piilo J, Maniscalco S. Driven harmonic oscillator as a quantum simulator for open systems. *Phys Rev A*. 2006;74: 032303.
70. Teufel S. *Adiabatic perturbation theory in quantum dynamics*. Berlin: Springer; 2003.
71. Castañón L, Zuñiga-Segundo A. The forced harmonic oscillator: coherent states and the RWA. *Am J Phys*. 2019;87:815–23.
72. Zhao L, Ma K, Yang Z. Changes of water hydrogen bond network with different externalities. *Int J Mol Sci*. 2015;16:8454–89.
73. De Ninno A, Castellano AC. On the effect of weak magnetic field on solutions of glutamic acid: the function of water. *J Phys Conf Ser*. 2011. p 012025, IOP Publishing.
74. Shen X. Increased dielectric constant in the water treated by extremely low frequency electromagnetic field and its possible biological implication, *J Phys Conf Ser*. 2011. 012019, IOP Publishing.
75. Protection ICoNIR. Guidelines for limiting exposure to time-varying electric, magnetic, and electromagnetic fields (up to 300 GHz). *Health Phys*. 1998;74:494–522.
76. Del Emilio G. Old and new views on the structure of matter and the special case of living matter. *J Phys: Conf Ser*. 2007;67: 012006.
77. Blank M, Goodman R. DNA is a fractal antenna in electromagnetic fields. *Int J Radiat Biol*. 2011;87:409–15.
78. Gluckman BJ, Netoff TI, Neel EJ, Ditto WL, Spano ML, Schiff SJ. Stochastic resonance in a neuronal network from mammalian brain. *Phys Rev Lett*. 1996;77:4098.
79. Radman T, Parra L, Bikson M. Amplification of small electric fields by neurons; implications for spike timing, In: *Engineering in Medicine and Biology Society*, 2006. EMBS'06. 28th Annual International Conference of the IEEE. IEEE; 2006. pp 4949–4952.
80. Buzsáki G, Anastassiou CA, Koch C. The origin of extracellular fields and currents—EEG, ECoG, LFP and spikes. *Nat Rev Neurosci*. 2012;13:407–20.

81. Cisek R, Tokarz D, Krouglov S, Steup M, Emes MJ, Tetlow IJ, Barzda V. Second harmonic generation mediated by aligned water in starch granules. *J Phys Chem*. 2014;B118:14785–94.
82. Balcavage WX, Alvager T, Swez J, Goff CW, Fox MT, Abdullyava S, King MW. A mechanism for action of extremely low frequency electromagnetic fields on biological systems. *Biochem Biophys Res Commun*. 1996;222:374–8.
83. Nikolova T, Czyz J, Rolletschek A, Blyszczuk P, Fuchs J, Jovtchev G, Schuderer J, Kuster N, Wobus AM. Electromagnetic fields affect transcript levels of apoptosis-related genes in embryonic stem cell-derived neural progenitor cells. *FASEB J*. 2005;19:1686–8.
84. Remondini D, Nylund R, Reivinen J, Poullietier de Gannes F, Veyret B, Lagroye I, Haro E, Trillo MA, Capri M, Franceschi C, Schlatterer K, Gminski R, Fitzner R, Tauber R, Schuderer J, Kuster N, Leszczynski D, Bersani F, Maercker C. Gene expression changes in human cells after exposure to mobile phone microwaves. *Proteomics*. 2006;6:4745–54.
85. Zhou J, Ming L-G, Ge B-F, Wang J-Q, Zhu R-Q, Wei Z, Ma H-P, Xian CJ, Chen K-M. Effects of 50Hz sinusoidal electromagnetic fields of different intensities on proliferation, differentiation and mineralization potentials of rat osteoblasts. *Bone*. 2011;49:753–61.
86. Blank M, Goodman R. Electromagnetic fields stress living cells. *Pathophysiology*. 2009;16:71–8.
87. Kucera O, Cifra M. Radiofrequency and microwave interactions between biomolecular systems. *J Biol Phys*. 2016;42:1–8.
88. Weinberg P, Bukov M, D'Alessio L, Polkovnikov A, Vajna S, Kolodrubetz M. Adiabatic perturbation theory and geometry of periodically-driven systems. *Phys Rep*. 2017;688:1–35.
89. Shamovsky I, Nudler E. New insights into the mechanism of heat shock response activation. *Cell Mol Life Sci CMLS*. 2008;65:855–61.
90. Philbin TG. Quantum dynamics of the damped harmonic oscillator. *New J Phys*. 2012;14: 083043.
91. Cassone G. Ab initio molecular dynamics simulations of H-bonded systems under an electric field, Université Pierre et Marie Curie-Paris VI. 2016.
92. Cerón-Carrasco JP, Jacquemin D. Electric field induced DNA damage: an open door for selective mutations. *Chem Commun*. 2013;49:7578–80.
93. Goryainov S. A model of phase transitions in double-well Morse potential: application to hydrogen bond. *Physica B*. 2012;407:4233–7.
94. Singh V, Fedeles BI, Essigmann JM. Role of tautomerism in RNA biochemistry. *RNA*. 2015;21:1–13.
95. Malíková I, Janoušek L. Non-thermal effects of low-frequency electromagnetic field on biological cells, in 2014 ELEKTRO, pp 599–602, IEEE.
96. Hundt PM, Jiang B, van Reijzen ME, Guo H, Beck RD. Vibrationally promoted dissociation of water on Ni (111). *Science*. 2014;344:504–7.
97. Bock J, Fukuyo Y, Kang S, Phipps ML, Alexandrov LB, Rasmussen KØ, Bishop AR, Rosen ED, Martinez JS, Chen H-T. Mammalian stem cells reprogramming in response to terahertz radiation. *PLoS ONE*. 2010;5: e15806.
98. Shavitt I, Bartlett RJ. Many-body methods in chemistry and physics: MBPT and coupled-cluster theory. Cambridge: Cambridge University Press; 2009.
99. Dos Santos C, Drigo Filho E, Ricotta R. Quantum confinement in hydrogen bond of DNA and RNA. *J Phys Conf Ser*, p 012033, IOP Publishing. 2015.
100. Milet A, Moszynski R, Wormer PE, van der Avoird A. Hydrogen bonding in water clusters: pair and many-body interactions from symmetry-adapted perturbation theory. *J Phys Chem A*. 1999;103:6811–9.
101. Fesenko EE, Gluvstein AY. Changes in the state of water, induced by radiofrequency electromagnetic fields. *FEBS Lett*. 1995;367:53–5.
102. Mentré P. Interfacial water: a modulator of biological activity. *J Biol Phys Chem*. 2004;4:115–23.
103. Wahl MC, Sundaralingam M. C–H... O hydrogen bonding in biology. *Trends Biochem Sci*. 1997;22:97–102.
104. Westhof E. Water: an integral part of nucleic acid structure. *Annu Rev Biophys Biophys Chem*. 1988;17:125–44.
105. Otting G, Liepinsh E, Wuthrich K. Protein hydration in aqueous solution. *Science*. 1991;254:974–80.
106. Beruto D, Botter R, Perfumo F, Scaglione S. Interfacial effect of extremely low frequency electromagnetic fields (EM-ELF) on the vaporization step of carbon dioxide from aqueous solutions of body simulated fluid (SBF). *Bioelectromagnetics*. 2003;24:251–61.
107. Ravera S, Repaci E, Morelli A, Pepe I, Botter R, Beruto D. Effects of extremely low frequency electromagnetic fields on the adenylate kinase activity of rod outer segment of bovine retina. *Bioelectromagn J Bioelectromagn Soc*. 2004;25:545–51.
108. Beruto DT, Lagazzo A, Frumento D, Converti A. Kinetic model of *Chlorella vulgaris* growth with and without extremely low frequency-electromagnetic fields (EM-ELF). *J Biotechnol*. 2014;169:9–14.
109. Arunan E, Desiraju GR, Klein RA, Sadlej J, Scheiner S, Alkorta I, Clary DC, Crabtree RH, Dannenberg JJ, Hobza P. Defining the hydrogen bond: an account (IUPAC Technical Report). *Pure Appl Chem*. 2011;83:1619–36.
110. Ball P. Water as an active constituent in cell biology. *Chem Rev*. 2008;108:74–108.
111. Pal SK, Zewail AH. Dynamics of water in biological recognition. *Chem Rev*. 2004;104:2099–124.
112. Nguyen TH, Zhang C, Weichselbaum E, Knyazev DG, Pohl P, Carloni P. Interfacial water molecules at biological membranes: structural features and role for lateral proton diffusion. *PLoS ONE*. 2018;13: e0193454.
113. Kounovsky-Shafer KL, Hernandez-Ortiz JP, Potamouis K, Tsvid G, Place M, Ravindran P, Jo K, Zhou S, Odijk T, De Pablo JJ. Electrostatic confinement and manipulation of DNA molecules for genome analysis. *Proc Natl Acad Sci*. 2017;114:13400–5.
114. Kolesnikov AI, Reiter GF, Choudhury N, Prisk TR, Mamontov E, Podlesnyak A, Ehlers G, Seel AG, Wesolowski DJ, Anovitz LM. Quantum tunneling of water in beryl: a new state of the water molecule. *Phys Rev Lett*. 2016;116: 167802.
115. Lobyshev V. Water is a sensor to weak forces including electromagnetic fields of low intensity. *Electromagn Biol Med*. 2005;24:449–61.
116. Elgabarty H, Kaliannan NK, Kühne TD. Enhancement of the local asymmetry in the hydrogen bond network of liquid water by an ultrafast electric field pulse. *Sci Rep*. 2019;9:1–8.
117. Zhao H, Tan Y, Zhang L, Zhang R, Shalaby M, Zhang C, Zhao Y, Zhang X-C. Ultrafast hydrogen bond dynamics of liquid water revealed by terahertz-induced transient birefringence. *Light Sci Appl*. 2020;9:1–10.
118. Rao ML, Sedlmayr SR, Roy R, Kanzius J. Polarized microwave and RF radiation effects on the structure and stability of liquid water. *Curr Sci*. 2010;98:1500–4.
119. Vallée P, Lafait J, Legrand L, Mentré P, Monod M-O, Thomas Y. Effects of pulsed low-frequency electromagnetic fields on water characterized by light scattering techniques: role of bubbles. *Langmuir*. 2005;21:2293–9.
120. Calabrò E, Magazù S. Response of hydrogen bonding to low-intensity 50 Hz electromagnetic field in typical proteins in bi-distilled water solution. *Spectrosc Lett*. 2017;50:330–5.
121. Mundy JN. Solid state: nuclear methods, vol. 21. Cambridge: Academic Press; 1983.
122. James DW, Armishaw RF. Structure of aqueous solutions: infrared spectra of the water librational mode in solutions of monovalent halides. *Aust J Chem*. 1975;28:1179–86.
123. Panagopoulos DJ, Margaritis LH. Theoretical considerations for the biological effects of electromagnetic fields. Biological effects of electromagnetic fields. Springer; 2003. pp 5–33. 2003.
124. Chew W, Liu A, Salazar-Lazaro C, Sha W. Quantum electromagnetics: a new look—part I. *IEEE J Multiscale Multiphys Comput Tech*. 2016;1:73–84.
125. Pikovsky A, Rosenblum M, Kurths J. Synchronization—a unified approach to nonlinear science. Cambridge: Cambridge University Press; 2001.
126. Mariam YH, Musin RN. Transition from moderate to strong hydrogen bonds: its identification and physical bases in the case of O—H... O intramolecular hydrogen bonds. *J Phys Chem A*. 2008;112:134–45.
127. Steiner T. Weak hydrogen bonds. In: Implications of molecular and materials structure for new technologies. Springer; 1999. pp 185–196.
128. Samdal S. The effect of large amplitude motion on the comparison of bond distances from ab initio calculations and experimentally determined bond distances, and on root-mean-square amplitudes of vibration, shrinkage, asymmetry constants, symmetry constraints, and inclusion of rotational constants using the electron diffraction method. *J Mol Struct*. 1994;318:133–41.
129. Kirillova S, Carugo O. Hydration sites of unpaired RNA bases: a statistical analysis of the PDB structures. *BMC Struct Biol*. 2011;11:41.
130. Grifoni M, Hänggi P. Driven quantum tunneling. *Phys Rep*. 1998;304:229–354.

131. Besancon R. The encyclopedia of physics. Berlin: Springer Science & Business Media; 2013.
132. Shun-Ping S, Quan Z, Li Z, Rong W, Zheng-He Z, Gang J, Yi-Bei F. Geometrical structures, vibrational frequencies, force constants and dissociation energies of isotopic water molecules (H<sub>2</sub>O, HDO, D<sub>2</sub>O, HTO, DTO, and T<sub>2</sub>O) under dipole electric field. *Chin Phys B*. 2011;20: 063102.
133. Zong D, Hu H, Duan Y, Sun Y. Viscosity of water under electric field: anisotropy induced by redistribution of hydrogen bonds. *J Phys Chem B*. 2016;120:4818–27.
134. Baranyai A, Bartók A, Chialvo AA. Computer simulation of the 13 crystal-line phases of ice. *J Chem Phys*. 2005;123: 054502.
135. Laage D, Hynes JT. A molecular jump mechanism of water reorientation. *Science*. 2006;311:832–5.
136. Agmon N. The Grotthuss mechanism. *Chem Phys Lett*. 1995;244:456–62.
137. Marx D, Tuckerman ME, Hutter J, Parrinello M. The nature of the hydrated excess proton in water. *Nature*. 1999;397:601.
138. Bagchi B. Water dynamics in the hydration layer around proteins and micelles. *Chem Rev*. 2005;105:3197–219.
139. Iijima T. The mean amplitude of the OH...O hydrogen bond in ice and liquid water. *J Mol Struct*. 1996;376:525–9.
140. Ankerhold J. Introduction. In: *Quantum tunneling in complex systems*. Springer; 2007. pp 1–5.
141. Masgrau L, Roujeinikova A, Johannissen LO, Hothi P, Basran J, Ranaghan KE, Mulholland AJ, Sutcliffe MJ, Scrutton NS, Leys D. Atomic description of an enzyme reaction dominated by proton tunneling. *Science*. 2006;312:237–41.
142. Dyke TR, Mack KM, Muentner JS. The structure of water dimer from molecular beam electric resonance spectroscopy. *J Chem Phys*. 1977;66:498–510.
143. Klinman JP, Kohen A. Hydrogen tunneling links protein dynamics to enzyme catalysis. *Annu Rev Biochem*. 2013;82:471–96.
144. Razavy M, Muga JG. Quantum theory of tunneling. *Phys Today*. 2004;57:66–7.
145. Horsewill AJ, Jones NH, Caciuffo R. Evidence for coherent proton tunneling in a hydrogen bond network. *Science*. 2001;291:100–3.
146. Jin L, Yang X, Zhu Y, Zhang Z, Liu R, Wang Z. Quantum resonant tunneling enhances hydrogen bond rotation of confined water, arXiv preprint [arXiv:2107.02619](https://arxiv.org/abs/2107.02619). 2021.
147. Cerón-Carrasco JP, Jacquemin D. Electric-field induced mutation of DNA: a theoretical investigation of the GC base pair. *Phys Chem Chem Phys*. 2013;15:4548–53.
148. Cerón-Carrasco JP, Cerezo J, Jacquemin D. How DNA is damaged by external electric fields: selective mutation vs. random degradation. *Phys Chem Chem Phys*. 2014;16:8243–6.
149. Gheorghiu A, Coveney PV, Arabi AA. The influence of external electric fields on proton transfer tautomerism in the guanine–cytosine base pair. *Phys Chem Chem Phys*. 2021;23:6252–65.
150. Roston D, Cheatum CM, Kohen A. Hydrogen donor–acceptor fluctuations from kinetic isotope effects: a phenomenological model. *Biochemistry*. 2012;51:6860–70.
151. Perez FP, Rizkalla J, Jeffers M, Salama P, Chumbiaua CNP, Rizkalla M. The effect of repeated electromagnetic fields stimulation in biological systems. In: *Ionizing and non-ionizing radiation*. IntechOpen; 2019.
152. Chandra AK, Nguyen MT, Uchimaru T, Zeegers-Huyskens T. Protonation and deprotonation enthalpies of guanine and adenine and implications for the structure and energy of their complexes with water: comparison with uracil, thymine, and cytosine. *J Phys Chem A*. 1999;103:8853–60.
153. Antonov L. *Tautomerism: concepts and applications in science and technology*. New Jersey: Wiley; 2016.
154. Cochrane JC, Strobel SA. Riboswitch effectors as protein enzyme cofactors. *RNA*. 2008;14:993–1002.
155. Singh V, Peng CS, Li D, Mitra K, Silvestre KJ, Tokmakoff A, Essigmann JM. Direct observation of multiple tautomers of oxythiamine and their recognition by the thiamine pyrophosphate riboswitch. *ACS Chem Biol*. 2013;9:227–36.
156. Felicissimo VC, da Rocha Martins J, Boldt IS, Chacham H. Molecular switches activated by electromagnetic pulses: first-principles calculations. *Phys Rev A*. 2009;80: 063410.
157. Peng CS, Jones KC, Tokmakoff A. Anharmonic vibrational modes of nucleic acid bases revealed by 2D IR spectroscopy. *J Am Chem Soc*. 2011;133:15650–60.
158. Abou-Zied OK, Jimenez R, Romesberg FE. Tautomerization dynamics of a model base pair in DNA. *J Am Chem Soc*. 2001;123:4613–4.
159. Klein DJ, Been MD, Ferré-D'Amaré AR. Essential role of an active-site guanine in glmS ribozyme catalysis. *J Am Chem Soc*. 2007;129:14858–9.
160. Cochrane JC, Strobel SA. Catalytic strategies of self-cleaving ribozymes. *Acc Chem Res*. 2008;41:1027–35.
161. Cochrane JC, Lipchock SV, Strobel SA. Structural investigation of the GlmS ribozyme bound to its catalytic cofactor. *Chem Biol*. 2007;14:97–105.
162. Frank R, Leeper F, Luisi B. Structure, mechanism and catalytic duality of thiamine-dependent enzymes. *Cell Mol Life Sci*. 2007;64:892–905.
163. Malandrinis G, Louloudi M, Hadjilias N. Thiamine models and perspectives on the mechanism of action of thiamine-dependent enzymes. *Chem Soc Rev*. 2006;35:684–92.
164. Chawla M, Sharma P, Halder S, Bhattacharyya D, Mitra A. Protonation of base pairs in RNA: context analysis and quantum chemical investigations of their geometries and stabilities. *J Phys Chem B*. 2011;115:1469–84.
165. Halder A, Bhattacharya S, Datta A, Bhattacharyya D, Mitra A. The role of N7 protonation of guanine in determining the structure, stability and function of RNA base pairs. *Phys Chem Chem Phys*. 2015;17:26249–63.
166. Goh GB, Knight JL, Brooks CL. Constant pH molecular dynamics simulations of nucleic acids in explicit solvent. *J Chem Theory Comput*. 2011;8:36–46.
167. Ranjan C. Tautomeric mutation in DNA. A theoretical comparative study and synthesis between various proposed quantum models. *Наноструктуры Математическая физика и моделирование*. 2015;12:85–98.
168. Yerkin A, Valentina Y, TussupbayevNessipbay ST, Zhadyra Y, Maxat B, Evgeniy B. Explanation of tautomerism and isomerization in terms of the magnetic field. *J Chem*. 2016;10:96–8.
169. Revzin A, Neumann E. Conformation changes in rRNA induced by electric impulses. *Biophys Chem*. 1974;2:144–50.
170. Salzberg CA, Cooper-Vastola SA, Perez F, Viehbeck MG, Byrne DW. The effects of non-thermal pulsed electromagnetic energy on wound healing of pressure ulcers in spinal cord-injured patients: a randomized, double-blind study. *Ostomy Wound Manage*. 1995; 41, 42–44, 46, 48 passim.
171. Nagy PI, Maheshwari A, Kim Y-W, Messer WS Jr. Theoretical and experimental studies of the isomeric protonation in solution for a prototype aliphatic ring containing two nitrogens. *J Phys Chem B*. 2010;114:349–60.
172. Giudice ED, Tedeschi A. Water and autocatalysis in living matter. *Electromagn Biol Med*. 2009;28:46–52.
173. Cimdins A, Klinkert B, Aschke-Sonnenborn U, Kaiser FM, Kortmann J, Narberhaus F. Translational control of small heat shock genes in mesophilic and thermophilic cyanobacteria by RNA thermometers. *RNA Biol*. 2014;11:594–608.
174. Goodman R, Blank M, Lin H, Dai R, Khorkova O, Soo L, Weisbrot D, Henderson A. Increased levels of hsp70 transcripts induced when cells are exposed to low frequency electromagnetic fields. *Bioelectrochem Bioenerg*. 1994;33:115–20.
175. Morimoto RI, Kroeger P, Cotto J. The transcriptional regulation of heat shock genes: a plethora of heat shock factors and regulatory conditions. In: *Stress-inducible cellular responses*. Springer; 1996. pp 139–163.
176. Morimoto RI. Regulation of the heat shock transcriptional response: cross talk between a family of heat shock factors, molecular chaperones, and negative regulators. *Genes Dev*. 1998;12:3788–96.
177. Mannerling A-C, Simkó M, Mild KH, Mattsson M-O. Effects of 50-Hz magnetic field exposure on superoxide radical anion formation and HSP70 induction in human K562 cells. *Radiat Environ Biophys*. 2010;49:731–41.
178. Goodman R, Henderson AS. Exposure of salivary gland cells to low-frequency electromagnetic fields alters polypeptide synthesis. *Proc Natl Acad Sci USA*. 1988;85:3928–32.
179. Lang BJ, Guerrero ME, Prince TL, Okusha Y, Bonorino C, Calderwood SK. The functions and regulation of heat shock proteins; key

- orchestrators of proteostasis and the heat shock response. *Arch Toxicol.* 2021;95:1943–70.
180. Barna J, Csermely P, Vellai T. Roles of heat shock factor 1 beyond the heat shock response. *Cell Mol Life Sci.* 2018;75:2897–916.
  181. Shamovsky I, Nudler E. New insights into the mechanism of heat shock response activation. *Cell Mol Life Sci.* 2008;65:855–61.
  182. Narberhaus F, Waldminghaus T, Chowdhury S. RNA thermometers. *FEMS Microbiol Rev.* 2005;30:3–16.
  183. de Pomerai DI, Smith B, Dawe A, North K, Smith T, Archer DB, Duce IR, Jones D, Candido EPM. Microwave radiation can alter protein conformation without bulk heating. *FEBS Lett.* 2003;543:93–7.
  184. Gross SR, Kinzy TG. Translation elongation factor 1A is essential for regulation of the actin cytoskeleton and cell morphology. *Nat Struct Mol Biol.* 2005;12:772–8.
  185. Morita MT, Tanaka Y, Kodama TS, Kyogoku Y, Yanagi H, Yura T. Translational induction of heat shock transcription factor  $\zeta$ 32: evidence for a built-in RNA thermosensor. *Genes Dev.* 1999;13:655–65.
  186. Lindquist S, Craig E. The heat-shock proteins. *Annu Rev Genet.* 1988;22:631–77.
  187. Morley JF, Morimoto RI. Regulation of longevity in *Caenorhabditis elegans* by heat shock factor and molecular chaperones. *Mol Biol Cell.* 2004;15:657–64.
  188. Nixon RA. The role of autophagy in neurodegenerative disease. *Nat Med.* 2013;19:983–97.
  189. Anckar J, Sistonen L. Regulation of HSF1 function in the heat stress response: implications in aging and disease. *Annu Rev Biochem.* 2011;80:1089–115.
  190. Zou J, Guo Y, Guettouche T, Smith DF, Voellmy R. Repression of heat shock transcription factor HSF1 activation by HSP90 (HSP90 complex) that forms a stress-sensitive complex with HSF1. *Cell.* 1998;94:471–80.
  191. Shi Y, Mosser DD, Morimoto RI. Molecular chaperones as HSF1-specific transcriptional repressors. *Genes Dev.* 1998;12:654–66.
  192. Shemesh N, Ben-Zvi A. HSF1 regulation in aging and its role in longevity. In: *Heat shock factor*. Springer; 2016. pp 93–113.
  193. Alavez S, Vantipalli MC, Zucker DJ, Klang JM, Lithgow GJ. Amyloid-binding compounds maintain protein homeostasis during ageing and extend lifespan. *Nature.* 2011;472:226–9.
  194. Neef DW, Jaeger AM, Thiele DJ. Heat shock transcription factor 1 as a therapeutic target in neurodegenerative diseases. *Nat Rev Drug Discov.* 2011;10:930–44.
  195. Perez FP, Moinuddin SS, ul ain Shamim Q, Joseph DJ, Morisaki J, Zhou X. Longevity pathways: HSF1 and FoxO pathways, a new therapeutic target to prevent age-related diseases. *Curr Aging Sci.* 2012;5:87–95.
  196. Hut HM, Kampinga HH, Sibon OC. Hsp70 protects mitotic cells against heat-induced centrosome damage and division abnormalities. *Mol Biol Cell.* 2005;16:3776–85.
  197. Dice JF. Chaperone-mediated autophagy. *Autophagy.* 2007;3:295–9.
  198. Heydari AR, You S, Takahashi R, Gutschmann-Conrad A, Sarge KD, Richardson A. Age-related alterations in the activation of heat shock transcription factor 1 in rat hepatocytes. *Exp Cell Res.* 2000;256:83–93.
  199. Gutschmann-Conrad A, Heydari AR, You S, Richardson A. The expression of heat shock protein 70 decreases with cellular senescence in vitro and in cells derived from young and old human subjects. *Exp Cell Res.* 1998;241:404–13.
  200. Ben-Zvi A, Miller EA, Morimoto RI. Collapse of proteostasis represents an early molecular event in *Caenorhabditis elegans* aging. *Proc Natl Acad Sci USA.* 2009;106:14914–9.
  201. Luan Q, Jin L, Jiang CC, Tay KH, Lai F, Liu XY, Liu YL, Guo ST, Li CY, Yan XG. RIPK1 regulates survival of human melanoma cells upon endoplasmic reticulum stress through autophagy. *Autophagy.* 2015;11:975–94.
  202. Desai S, Liu Z, Yao J, Patel N, Chen J, Wu Y, Ahn EE-Y, Fodstad O, Tan M. Heat shock factor 1 (HSF1) controls chemoresistance and autophagy through transcriptional regulation of autophagy-related protein 7 (ATG7). *J Biol Chem.* 2013;288:9165–76.
  203. Dul JL, Davis DP, Williamson EK, Stevens FJ, Argon Y. Hsp70 and antifibrillogenic peptides promote degradation and inhibit intracellular aggregation of amyloidogenic light chains. *J Cell Biol.* 2001;152:705–16.
  204. Diamant S, Ben-Zvi AP, Bukau B, Goloubinoff P. Size-dependent disassembly of stable protein aggregates by the DnaK chaperone machinery. *J Biol Chem.* 2000;275:21107–13.
  205. Raman B, Ban T, Sakai M, Pasta SY, Ramakrishna T, Naiki H, Goto Y, Rao CM.  $\alpha$ B-crystallin, a small heat-shock protein, prevents the amyloid fibril growth of an amyloid  $\beta$ -peptide and  $\beta$ 2-microglobulin. *Biochem J.* 2005;392:573–81.
  206. Barral JM, Broadley SA, Schaffar G, Hartl FU. Roles of molecular chaperones in protein misfolding diseases. In: *Seminars in cell and developmental biology*. Elsevier; 2004. pp 17–29.
  207. Pierce A, Podlutskaya N, Halloran JJ, Hussong SA, Lin PY, Burbank R, Hart MJ, Galvan V. Over-expression of heat shock factor 1 phenocopies the effect of chronic inhibition of TOR by rapamycin and is sufficient to ameliorate Alzheimer's-like deficits in mice modeling the disease. *J Neurochem.* 2013;124:880–93.
  208. Craig EA, Weissman JS, Horwich AL. Heat shock proteins and molecular chaperones: mediators of protein conformation and turnover in the cell. *Cell.* 1994;78:365–72.
  209. Georgopoulos C, Welch W. Role of the major heat shock proteins as molecular chaperones. *Annu Rev Cell Biol.* 1993;9:601–34.
  210. Hartl FU. Molecular chaperones in cellular protein folding. *Nature.* 1996;381:571–9.
  211. Kakimura J-I, Kitamura Y, Takata K, Umeki M, Suzuki S, Shibagaki K, Taniguchi T, Nomura Y, Gebicke-Haerter PJ, Smith MA. Microglial activation and amyloid- $\beta$  clearance induced by exogenous heat-shock proteins. *FASEB J.* 2002;16:601–3.
  212. Frautschy S, Hu W, Kim P, Miller S, Chu T, Harris-White M, Cole G. Phenolic anti-inflammatory antioxidant reversal of A $\beta$ -induced cognitive deficits and neuropathology. *Neurobiol Aging.* 2001;22:993–1005.
  213. Yang Y, Turner RS, Gaut JR. The chaperone BiP/GRP78 binds to amyloid precursor protein and decreases A $\beta$ 40 and A $\beta$ 42 secretion. *J Biol Chem.* 1998;273:25552–5.
  214. Park J-S, Kim D-H, Yoon S-Y. Regulation of amyloid precursor protein processing by its KFERQ motif. *BMB Rep.* 2016;49:337.
  215. Wang Y, Martinez-Vicente M, Krüger U, Kaushik S, Wong E, Mandelkow E-M, Cuervo AM, Mandelkow E. Tau fragmentation, aggregation and clearance: the dual role of lysosomal processing. *Hum Mol Genet.* 2009;18:4153–70.
  216. Vogiatzi T, Xilouri M, Vekrellis K, Stefanis L. Wild type  $\alpha$ -synuclein is degraded by chaperone-mediated autophagy and macroautophagy in neuronal cells. *J Biol Chem.* 2008;283:23542–56.
  217. Bauer PO, Goswami A, Wong HK, Okuno M, Kurosawa M, Yamada M, Miyazaki H, Matsumoto G, Kino Y, Nagai Y. Harnessing chaperone-mediated autophagy for the selective degradation of mutant huntingtin protein. *Nat Biotechnol.* 2010;28:256–63.
  218. Dickey C, Kraft C, Jinwal U, Koren J, Johnson A, Anderson L, Lebson L, Lee D, Dickson D, De Silva R. Aging analysis reveals slowed tau turnover and enhanced stress response in a mouse model of tauopathy. *Am J Pathol.* 2009;174:228–38.
  219. Dou J, Su P, Xu C, Wen Z, Mao Z, Li W. Targeting Hsc70-based autophagy to eliminate amyloid  $\beta$  oligomers. *Biochem Biophys Res Commun.* 2020;524:923–8.
  220. Evans CG, Wisen S, Gestwicki JE. Heat shock proteins 70 and 90 inhibit early stages of amyloid beta-(1–42) aggregation in vitro. *J Biol Chem.* 2006;281:33182–91.
  221. Leissring MA, Farris W, Chang AY, Walsh DM, Wu X, Sun X, Frosch MP, Selkoe DJ. Enhanced proteolysis of  $\beta$ -amyloid in APP transgenic mice prevents plaque formation, secondary pathology, and premature death. *Neuron.* 2003;40:1087–93.
  222. Leissring MA. Proteolytic degradation of the amyloid beta-protein: the forgotten side of Alzheimer's disease. *Curr Alzheimer Res.* 2006;3:431–5.
  223. Bonini NM. Chaperoning brain degeneration. *Proc Natl Acad Sci.* 2002;99:16407–11.
  224. Miyakoshi J, Takemasa K, Takashima Y, Ding GR, Hirose H, Koyama S. Effects of exposure to a 1950 MHz radio frequency field on expression of Hsp70 and Hsp27 in human glioma cells. *Bioelectromagnetics.* 2005;26:251–7.
  225. Wang J, Koyama S, Komatsubara Y, Suzuki Y, Taki M, Miyakoshi J. Effects of a 2450 MHz high-frequency electromagnetic field with a wide range of SARs on the induction of heat-shock proteins in A172 cells. *Bioelectromagnetics.* 2006;27:479–86.

226. Tokalov SV, Gutzzeit HO. Weak electromagnetic fields (50 Hz) elicit a stress response in human cells. *Environ Res.* 2004;94:145–51.
227. Westerheide SD, Morimoto RI. Heat shock response modulators as therapeutic tools for diseases of protein conformation. *J Biol Chem.* 2005;280:33097–100.
228. Morelli L, Llovera R, Ibendahl S, Castaño EM. The degradation of amyloid  $\beta$  as a therapeutic strategy in Alzheimer's disease and cerebrovascular amyloidoses. *Neurochem Res.* 2002;27:1387–99.
229. Goodman EM, Greenebaum B, Marron MT. Effects of electromagnetic fields on molecules and cells. *Int Rev Cytol.* 1995;158:279–338.
230. Shellock F, Crues J. MRI: potential adverse effects and safety considerations. *MRI Decis.* 1988;2:25–30.
231. Perez FP, Bandeira JP, Morisaki JJ, Peddinti SVK, Salama P, Rizkalla J, Rizkalla ME. Antenna design and SAR analysis on human head phantom simulation for future clinical applications. *J Biomed Sci Eng.* 2017;10:421–30.
232. Christ A, Chavannes N, Nikoloski N, Gerber HU, Poković K, Kuster N. A numerical and experimental comparison of human head phantoms for compliance testing of mobile telephone equipment. *Bioelectromagn Soc.* 2005;26:125–37.
233. Chung B, Chuah H. Modeling of RF absorber for application in the design of anechoic chamber. *Prog Electromagn Res.* 2003;43:273–85.
234. Perez FP, Rizkalla J, Jeffers M, Salama P, Chumbiauca CNP, Rizkalla M. The effect of repeated electromagnetic fields stimulation in biological systems. In: *The effect of electromagnetic fields on living beings*, IntechOpen; 2019.
235. Arendash G, Cao C, Abulaban H, Baranowski R, Wisniewski G, Becerra L, Andel R, Lin X, Zhang X, Wittwer D. A clinical trial of transcranial electromagnetic treatment in Alzheimer's disease: cognitive enhancement and associated changes in cerebrospinal fluid, blood, and brain imaging. *J Alzheimer's Dis.* 2019; 1–26.
236. Phillips JL. Effects of electromagnetic field exposure on gene transcription. *J Cell Biochem.* 1993;51:381–6.
237. Lupke M, Frahm J, Lantow M, Maercker C, Remondini D, Bersani F, Simkó M. Gene expression analysis of ELF-MF exposed human monocytes indicating the involvement of the alternative activation pathway. *Biochim Biophys Acta BBA Mol Cell Res.* 2006;1763:402–12.
238. Frahm J, Mattsson M-O, Simkó M. Exposure to ELF magnetic fields modulate redox related protein expression in mouse macrophages. *Toxicol Lett.* 2010;192:330–6.
239. Fassina L, Saino E, Visai L, Silvani G, Cusella De Angelis MG, Mazzini G, Benazzo F, Magenes G. Electromagnetic enhancement of a culture of human SAOS-2 osteoblasts seeded onto titanium fiber-mesh scaffolds. *J Biomed Mater Res, Part A.* 2008;87:750–9.
240. Perez FP, Zhou X, Morisaki J, Ilie J, James T, Jurivich DA. Engineered repeated electromagnetic field shock therapy for cellular senescence and age-related diseases. *Rejuvenation Res.* 2008;11:1049–57.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

