

Synthetic Cooling Agents in US-marketed E-cigarette Refill Liquids and Popular Disposable E-cigarettes: Chemical Analysis and Risk Assessment

Sairam V. Jabba, DVM, PhD^{*1,2}, Hanno C. Erythropel, PhD^{*2,3} , Deyri Garcia Torres⁴,
Lauren A. Delgado⁴, Jackson G. Woodrow⁴, Paul T. Anastas⁵, Julie B. Zimmerman PhD^{2,3},
Sven-Eric Jordt, PhD^{1,2,6} 

¹Department of Anesthesiology, Duke University School of Medicine, Durham, NC, USA

²Yale Center for the Study of Tobacco Products (YCSTP), Department of Psychiatry, Yale School of Medicine, New Haven, CT, USA

³Department of Chemical and Environmental Engineering, Yale University, New Haven, CT, USA

⁴Yale College, New Haven, CT, USA

⁵Environmental Health Sciences Department, School of Public Health, Yale University, New Haven, CT, USA

⁶Integrated Toxicology & Environmental Health Program (ITEHP), Duke University, Durham, NC, USA

*These authors contributed equally.

Corresponding Author: Sven-Eric Jordt, PhD, Department of Anesthesiology, Duke University School of Medicine, 3 Genome Ct., Durham, NC 27710-3094, USA.

E-mail: sven.jordt@duke.edu

Abstract

Introduction: Menthol, through its cooling and pleasant sensory effects, facilitates smoking and tobacco product initiation, resulting in the high popularity of mint/menthol-flavored E-cigarettes. More recently, E-cigarette vendors started marketing synthetic cooling agents as additives that impart a cooling effect but lack a characteristic minty odor. Knowledge about content of synthetic coolants in US-marketed E-cigarette products and associated health risks is limited.

Aims and Methods: E-liquid vendor sites were searched with the terms “koolada”, “kool/cool”, “ice”, or WS-3/WS-23, denoting individual cooling agents, and relevant refill E-liquids were purchased. “Ice” flavor varieties of Puffbar, the most popular disposable E-cigarette brand, were compared with non-“ice” varieties. E-liquids were characterized, and synthetic coolants quantified using GC/MS. Margin of exposure (MOE), a risk assessment parameter, was calculated to assess the risk associated with synthetic coolant exposure from E-cigarette use.

Results: WS-3 was detected in 24/25 refill E-liquids analyzed. All Puffbar flavor varieties contained either WS-23 (13/14) or WS-3 (5/14), in both “ice”- and non-“ice” flavors. Modeling consumption of WS-3 from vaped E-liquids, resulted in MOEs below the safe margin of 100 for most daily use scenarios. MOEs for WS-23 were <100 for 10/13 Puffbar flavors in all use scenarios. Puffbar power specifications are identical to Juul devices.

Conclusions: Synthetic cooling agents (WS-3/WS-23) were present in US-marketed E-cigarettes, at levels that may result in consumer exposures exceeding safety thresholds set by regulatory agencies. Synthetic coolants are not only found in mint- or menthol-flavored products but also in fruit- and candy-flavored products, including popular disposable E-cigarette products such as Puffbar.

Implications: Synthetic cooling agents are widely used in “kool/cool”- and “ice”-flavored E-liquids and in E-liquids without these labels, both as a potential replacement for menthol or to add cooling “notes” to nonmenthol flavors. These agents may be used to bypass current and future regulatory limits on menthol content in tobacco products, and not just E-cigarettes. Because synthetic cooling agents are odorless, they may not fall under the category of “characterizing flavor”, potentially circumventing regulatory measures based on this concept. Regulators need to consider the additional health risks associated with exposure to synthetic cooling agents.

Introduction

Additives and flavors are added to tobacco products to counteract the harshness, sensory irritation and bitter tastes of tobacco and nicotine. In the United States, the Family Smoking Prevention and Tobacco Control Act (FSPTCA) of 2009 authorized the Food and Drug Administration (FDA) to regulate constituents, additives, and flavors in tobacco products.¹ The FSPTCA banned the sales of flavored combustible cigarettes, with the exception of menthol cigarettes.¹ Menthol in tobacco products is well understood to reduce harshness, impart a pleasant minty and cooling flavor and increase palatability.² Menthol affects tobacco and nicotine use behaviors such as

initiation, dependence, abuse liability and reduces the ability to quit, especially among the youth and young adults.² The use of mentholated tobacco products is associated with tobacco use disparities, with 85% of African American smokers using menthol cigarettes.^{2,3} The US FDA announced in May 2021 that it will issue a rule to ban menthol cigarettes.⁴ Menthol is also a popular flavor additive in E-cigarettes, especially after the FDA banned all flavored closed pod systems (eg, Juul) except for menthol and tobacco.⁵ Sales of menthol Juul pods increased 6-fold from August 2019 to May 2020 (10.7%–61.8%) when all other Juul flavors (besides tobacco) were removed from the marketplace.^{6,7}

Received: June 17, 2021. Revised: December 6, 2021. Accepted: February 12, 2022

© The Author(s) 2022. Published by Oxford University Press on behalf of the Society for Research on Nicotine and Tobacco. All rights reserved.

For permissions, please e-mail: journals.permissions@oup.com.

Synthetic cooling agents share the sensory cooling effects of menthol, but lack menthol's strong minty odor, especially at the concentrations at which they are added to E-liquids.^{8,9} These agents were developed beginning in the 1970s, to replace menthol in body care and food products. In the 1970s and 80s, the tobacco industry including R.J. Reynolds and Phillip Morris, carried out consumer tests with cigarettes containing synthetic cooling agents, to impart a "cooling without menthol" experience.^{8–11} A 2018 analysis of combustible cigarettes (both menthol and non-menthol) marketed in Germany detected WS-3, indicating that coolants other than menthol are added to tobacco products.^{12,13} In a recent study, we demonstrated that European-marketed Juul menthol pods contained the synthetic cooling agent WS-3, whereas US and Canadian pods did not.¹⁴ More recently, Omaiye et al. demonstrated the presence of synthetic coolants in US-marketed disposables.¹⁵ Although synthetic cooling agents are sold to (Do-It-Yourself) DIY users in the USA, often under the label "Koolada," the identity and concentrations of these agents in US-marketed E-cigarettes and E-cigarette refill liquids ("E-liquids") remains unknown.¹⁶

While natural and synthetic mint- and cooling compound are widely used in food and other consumer products, there are rising concerns about their toxicity, especially in poorly regulated products such as E-cigarettes. For example, we detected concerning levels of pulegone, a carcinogenic mint flavorant banned by FDA in food, in a variety of mint- and menthol-flavored E-cigarette and smokeless tobacco products.^{17,18} Synthetic cooling agents such as WS-3 are classified by the Flavor Extracts Manufacturers Association (FEMA) as GRAS (Generally Recognized As Safe). However, this applies only to their intended use in food products, but not to E-cigarette use resulting in inhalational exposure.¹⁹ The Joint Expert Committee on Food Additives (JECFA) of the World Health Organization (WHO) established a threshold of concern of 90 µg/day/person for intake of WS-3 and related compounds, a level that is often exceeded by consumers.²⁰ Based on oral toxicity studies, an oral No-Observed-Adverse-Effect-Level (NOAEL) of 8 mg/kg-bw for WS-3^{20–22} was established, and 5 mg/kg-bw for WS-23, another widely used synthetic cooling agent. Chronic or higher doses were observed to cause kidney and liver lesions.^{20,23} In vitro genotoxicity studies in mammalian cells demonstrated WS-23 to be clastogenic (ie, inducing disruption or breakages of chromosomes), leading to a call by JECFA for additional toxicological studies to further evaluate the safety of WS-23.²⁰ Exposure of airway-epithelial cells in vitro to various concentrations of synthetic coolants and WS-23 containing e-liquids also demonstrated a dose-dependent cellular toxicity.¹⁵ A recent inhalation exposure study in rodents by an E-liquid manufacturer for WS-23 determined a NOAEL of 29 mg/kg-bw (342 mg/m³) in rodents.²⁴ Due to their evident toxicity and increased use in E-cigarettes, it is critical to evaluate the potential health risks associated with synthetic coolant exposure from E-cigarette use.

In the present study, we determined the levels of synthetic cooling agents (WS-3, WS-23) as well as important peppermint (menthol and menthone) and spearmint (carvone) flavor chemicals, in US-marketed refill E-liquids and in disposable E-cigarettes of the brand Puffbar. The rationale to include menthol, menthone and carvone in chemical analysis is that, they are the most frequent mint flavors encountered in E-liquids. Several recent E-cigarette prevalence studies indicated high use of disposables, especially "ice"-flavored, among youth and young adults.^{6,7,25–29} Disposable E-cigarette

brand, Puffbar, was selected because, according to the 2021 US National Youth Tobacco Survey it was determined as the most popular E-cigarette device among middle and high school students.^{6,7,28} In addition, the power specifications for Puffbar devices were determined. We then assessed the health risk associated with synthetic coolant exposure via E-cigarette use by calculating the Margin of Exposure (MOE), a risk assessment parameter, for a range of product use scenarios modeling low, moderate and frequent E-cigarette use.

Materials and Methods

Analysis of Synthetic Cooling Agents and Other Flavor Content in US-Marketed E-cigarette Liquids and Their Aerosols

E-liquid selection

E-cigarette vendor websites keyword searches for "koolada," "kool," "WS-3," and "WS-23" were carried out to identify E-liquids potentially containing synthetic coolants. Identified products, including a concentrate ("Agent Cool") for do-it-yourself (DIY) users, were purchased directly from vendors in 2019 and 2020. In addition, previously purchased menthol/mint/spearmint/wintergreen E-liquids (2019) were re-analyzed for synthetic coolant presence. In early 2020, all available flavors of "Puffbar" were purchased from puffcig.com due to the frequent occurrence of the words "cool/ice" in Puffbar flavor names (see Tables 1 and 2 for an overview). Purchased E-liquids were stored in dark at room temperature.

Power specifications measurement

Puffbar devices were opened manually, and the heating coil was carefully pulled toward the front of the opened device by the connecting wires. Resistance was measured in three devices by clamping a digital multimeter (Fluke 116) to the connecting wires. Battery specifications were printed onto the battery pack and reported as such.

E-liquid chemical analysis

E-liquids were analyzed using an established GC-MS method and selected compounds (WS-3, WS-23, menthol, menthone, carvone, nicotine) were quantified in triplicate using an established GC-FID method.³⁰ To do so, commercially available standards of *N*-ethyl-*p*-menthane-3-carboxamide (WS-3; 99%; CAS# 39711-79-0), (*S*)-nicotine (>99%; CAS# 54-11-5), (*dl*)-menthol (99%; CAS# 89-78-1), *d*-carvone (>96%; CAS# 2244-16-8; all Sigma-Aldrich, Saint Louis, MO), 2-isopropyl-*N*-2,3-trimethylbutyramide, (WS-23; >98%; CAS# 51115-67-4, TCI, Portland, OR), *l*-menthone (97%; CAS# 14073-97-3; Alfa Aesar, Haverhill, MA), propylene glycol (>99.5%; CAS# 57-55-6), and glycerol (99.6%; CAS 56-81-5; both Fisher Scientific, Hampton, NH) were used to construct calibration curves in the relevant concentration range. Samples were diluted in 1 mL of methanol [JT Baker, Center Valley, PA] containing 1 g/L of 1,4-dioxane [Alfa Aesar] as internal standard (IS) and injected; the detailed analytical methods have been described previously.³⁰ PG/VG ratios for Puffbars were rounded to the nearest 5 for simplicity.

Coolant carryover analysis

To assess the carryover of the synthetic coolants WS-3 and WS-23 from refill E-liquids to E-cigarette-generated aerosol, the Suorin iShare refillable pod E-cigarette device was used as its

Table 1. Concentrations of synthetic cooling agents (WS-3 and WS-23), menthol, menthone, carvone, nicotine and solvent chemicals in US-marketed E-cigarette refill liquids

	E-cigarette Flavor (Manufacturer)	Flavor Concentration ± s.d. (µg/g)				Menthone	Carvone	Nicotine concentration ± s.d. (mg/g)	PG/VG ratio(%/%)	Synthetic coolant carryover ± s.d. (%)
		WS-3	WS-23	Menthol	Menthone					
1	Arctic Rush (Central Vapors)	14,650 ± 230					9.2 ± 0.09	50/50	106 ± 4	
2	Frzn Mango (BLVK)	2,450 ± 00		3,640 ± 10			5.08 ± 0.03	30/70		
3	Nuclear White Menthol (Vapor HQ)	2,440 ± 10		23,490 ± 130	310 ± 20	190 ± 70	9.88 ± 0.02	50/50	104 ± 4	
4	Rich Girl (Stunna)	2,260 ± 30					3.31 ± 0.06	70/30	107 ± 11	
5	Frozen Berries (Shinjii Vapor)	2,220 ± 20		2,320 ± 20			3.36 ± 0.05	30/70		
6	Frozen Limeade (Shinjii Vapor)	1,540 ± 20		3,470 ± 50			3.38 ± 0.04	30/70		
7	Cool Lime (Chill)	2,110 ± 60					1.63 ± 0.05	30/70		
8	Tropicool (Chill)	2,040 ± 40					3.02 ± 0.07	30/70	91*	
9	Kool Cali (Chill)	1,690 ± 20					4.20 ± 0.06	40/60		
10	Sakura 3 mg Nic (Sugoi)	1,090 ± 20		910 ± 10			2.32 ± 0.03	30/70	105 ± 2	
11	Coconut Ice (EC Blend)	2,240 ± 10					1.41 ± 0.02	60/40		
12	Clove + Cool Hit (EC Blend)	1,600 ± 20					17.91 ± 0.23	50/50		
13	Popsicle USA (EC Blend)	150 ± 10				10 ± 1		60/40		
14	Frozen (FJ's)	1,550 ± 50						30/70		
15	Wintergreen Menthol (VaporHQ)	870 ± 20		4,850 ± 30			9.72 ± 0.08	50/50		
16	Icy Menthol (Vista Vapors)	560 ± 10		9,240 ± 90	1,320 ± 50	2,890 ± 10	9.80 ± 0.02	50/50		
17	Ice Dragon (ELD Flavors)	410 ± 30					2.75 ± 0.01	20/80		
18	Wintermint (Gothic Vapor)	330 ± 50		1,110 ± 190		510 ± 10	9.03 ± 0.84	50/50		
19	Climate Change (What Juice)	290 ± 10		330 ± 10			9.48 ± 0.12	30/70		
20	Arctic (Vista Select)	280 ± 20		3,470 ± 10	30 ± 10	930 ± 70	6.59 ± 0.06	30/70		
21	Dillinger (Nasty Juice X Kilo)		1,860 ± 40		280 ± 10		4.29 ± 0.07	30/70	99 ± 10	
22	Kool Peach (Salty Man)	990 ± 20					46.4 ± 1.09	40/60	109 ± 1	
23	Sakura Nic Salts 35 mg (Sugoi)	1,170 ± 30		1,170 ± 20			28.74 ± 0.43	50/50		
24	Sakura Nic Salts 50 mg (Sugoi)	1,140 ± 60		1,140 ± 90	14,110 ± 670		39.19 ± 2.20	50/50	97*	
25	Unicorn Nicotine Salt Spearmint (BLVK)	2,950 ± 20		10,920 ± 160	25,450 ± 600	6,830 ± 40	41.36 ± 0.33	40/60	107 ± 8	
26	Agent Cool Super Concentrate (TBD Liquids)	8,840 ± 130	124,060 ± 720							

Carryover rates for synthetic coolants are provided for select refill liquids. *single experiment.

Table 2. Chemical concentrations of synthetic cooling agents (WS-3 and WS-23), menthol and nicotine in flavors of US-marketed Puffbar disposable E-cigarettes

Puffbar flavor	Flavor Concentration± s.d. (µg/g)					Nicotine concentration± s.d. (mg/g)	PG/VG ratio (%/%)
	WS-3	WS-23	Menthol	Menthone	Carvone		
1 Cool Mint	10,770 ± 210	36,660 ± 530	620 ± 50			32.79 ± 0.38	55/45
2 Café Latte		2300 ± 30	1500 ± 50			36.56 ± 0.30	55/45
3 Orange Ice	1,750 ± 20	22,950 ± 450				37.61 ± 0.76	30/70
4 Pomegranate						29.37 ± 0.47	50/50
5 Lush Ice		25,830 ± 450	2,250 ± 190			32.26 ± 0.53	55/45
6 Melon Ice		8,810 ± 60				32.81 ± 0.29	65/35
7 Cucumber	3,770 ± 20	3,940 ± 10				31.67 ± 0.15	60/40
8 Lychee		26,240 ± 240				33.22 ± 0.27	55/45
9 Menthol	3,540 ± 30	1,610 ± 1,950	3,680 ± 550			33.25 ± 0.22	45/55
10 Blueberry Ice	1,880 ± 20	14,120 ± 210	2,300 ± 40			32.71 ± 0.45	50/50
11 Strawberry Banana		5,420 ± 90				32.12 ± 0.52	50/50
12 OMG		7,390 ± 330	1,330 ± 40			34.01 ± 1.35	50/50
13 Peach Ice		11,540 ± 270				33.11 ± 0.63	55/45
14 Banana Ice		13,110 ± 50	2,040 ± 110			40.66 ± 0.22	55/45

No menthone or carvone was found in any of the analyzed Puffbar e-liquids. PG/VG ratio measured in triplicate and rounded to next 5. Empty cells represent “not detected”. Yellow highlights indicate WS-3; green shade represent WS-23; and blue indicates menthol.

power specifications are very similar to the Puffbar and Juul device (Suorin iShare: 3.7V, 1.8 Ω, resulting max. power 7.6 W; Puffbar and Juul: 3.7 V, 1.7 Ω, resulting max. power 8.1 W). Moreover, Suorin remains highly popular among vapers, and especially high school age vapers, as a favorite refillable device.²⁹ The Sourin device was attached via a custom-3D printed connector to a previously described house-built vaping machine.³¹ In brief, the device was controlled by a programmable Arduino board that activates a micro-diaphragm pump at an operator-defined puff length, inter-puff break, and number of puffs. A microneedle valve in combination with a flow meter (Omega, Norwalk, CT) was used to control the flow rate. Generated aerosol was trapped by a series of liquid nitrogen-chilled cold-finger traps. The carryover puffing regime was 20 puffs, 2 L/min flow rate, 2.4 s puff length, 80 mL puff volume, 30 s inter-puff break and carryover was determined in triplicate (three separate cartridges). The chosen puffing regime is at the lower spectrum of observed user behavior^{31,32} and has been utilized previously to determine carryover in popular Juul E-cigarettes.³³ After 20 puffs were collected for each experiment, the traps were allowed to thaw and the captured material was taken up in 1 mL of methanol containing IS and injected into the GC/FID. Percent carryover was calculated as coolant content in trapped aerosol over coolant content of consumed liquid, as determined by weighing the e-liquid-containing cartridge before and after the carryover experiment using an analytical balance (Sartorius Praxium, Göttingen, Germany). E-liquids for carryover experiments were chosen based on type of synthetic coolant present (WS-3, WS-23), and based on high, average, or low synthetic coolant and/or nicotine content (Table 1).

Margin of exposure (MOE) calculation for synthetic cooling agents

MOE, a preferred risk assessment parameter for toxic compounds by the JECFA,^{34,35} was calculated for WS-3 and WS-23 containing E-liquids using the following formula, as described previously¹⁷:

$$\text{MOE} = \frac{\text{NOAEL for synthetic coolant}}{\text{Estimated exposure dose (EED)}}$$

where NOAEL is the No-Observed-Adverse-Effect-Level, a dose determined as the threshold producing a toxic effect in animal oral toxicity studies. Though benchmark dose (BMD) is the preferred toxicological threshold for MOE calculations, NOAEL-values are applicable when available toxicity data are not amenable for statistical modeling of BMD,³⁶ as was the case for WS-3 and WS-23. JECFA determined NOAELs for WS-3 and WS-23 are 8 and 5 mg/kg-bw/day, respectively.^{21–23} Daily Estimated Exposure Doses were determined from WS-3 and WS-23 concentrations in the E-liquids and their estimated average daily amount consumed. Nicotine concentrations in the majority of refill E-liquids ranged between 0.1% and 1.0%, and user surveys found consumption volumes for such low nicotine E-cigarette liquids to be 9 mL/day, ranging from 1.7 to 17.7 mL/day).^{18,37,38} To cover a wide range of user behaviors, MOEs for these refill E-liquids were calculated for consumption volumes of 1, 3, 5, 10 and 15 mL/day. For nicotine salt-containing refill liquids (nicotine content >3%), MOEs were calculated for 1, 3, 5 mL/day to account for the higher nicotine content. Puffbar E-liquid MOEs were calculated for consumption amounts of half-, 1- and 2-Puffbar(s) to model volumes consumed by light, moderate, and heavy users of 3%–5% nicotine salt-containing disposable E-cigarettes (Puffbar volume is 1.3 mL). For the calculations, a user body weight of 60 kg was assumed; to convert E-cigarette liquid consumption from volume in mL/day to g/day, the densities of propylene glycol and glycerol (1.04 and 1.26 g/cm³, respectively) were taken into consideration.

Results

Chemical Analysis of Synthetic Cooling Agents, Nicotine and Other Flavorants in US-Marketed E-cigarette Refill Liquids, Concentrates and Their Aerosols

Using our search criteria, 25 E-cigarette refill liquids and 1 concentrate from 19 different brands were identified and their con-

tents of synthetic coolants, mint-related compounds and nicotine identified by GC/MS and quantified by GC/FID. Of the analyzed liquids, 21 contained free-base nicotine and 4 nicotine salts. WS-3 was present in 24/25 E-liquids, with amounts varying from 150 to 14,650 $\mu\text{g/g}$ (Table 1). One E-liquid contained WS-23 exclusively (Dillinger; 1,860 $\mu\text{g/g}$) (Table 1). The concentrate contained both WS-3 (8,840 $\mu\text{g/g}$) and WS-23 (124,060 $\mu\text{g/g}$) (Table 1).

Carryover rates for WS-3 from 8-different E-liquids into aerosol ranged between 91% and 109% (vs. neat E-liquid content) (average: $105 \pm 6.4\%$; $n = 20$; Table 1). For the only refill E-liquid it was detected in, the carryover rate for WS-23 was $99 \pm 10\%$ (vs. neat E-liquid content; $n = 3$; Table 1).

In addition to WS-3 and WS-23, menthol, menthone, carvone, and nicotine were also quantified, which were detected in 13/25 E-liquids for menthol (330–23,490 $\mu\text{g/g}$), 6/25 for menthone (30–25,450 $\mu\text{g/g}$) and carvone (10–6,830 $\mu\text{g/g}$), and 4/25 for menthol, menthone, carvone, WS-3 combined (4,940–46,150 $\mu\text{g/g}$; see Table 1 for details). Nicotine concentrations in 20/21 free-base nicotine containing E-liquids were below 1% (1.41–9.97 mg/g), except for Clove + Cool Hit (17.91 mg/g). Nicotine contents in nicotine-salts containing E-liquids ranged from 28.74 to 44.6 mg/g (Table 1). Carryover rates for menthol, menthone and carvone were calculated from vaping of select refill E-liquids and ranged from 103%–107%, 85%–106% and 97%–105%, respectively (Supplementary Table 1). Taken together, these results suggest that synthetic coolants are added to US-marketed E-liquids of a wide range of flavors (Table 1), including to those already containing the “natural” coolant menthol. Indicated PG/VG ratios for refill E-liquids in Table 1 represent label information.

Chemical Analysis of Synthetic Cooling Agents and Menthol in US-Marketed Disposable Puffbar E-cigarettes

Several Puffbar flavors are marketed as “Ice” flavors, indicating the presence of a cooling agent. WS-23 was detected in 13/14 Puffbar flavors (1,610–36,660 $\mu\text{g/g}$), WS-3 in 5/14 flavors (1,880–10,770 $\mu\text{g/g}$) and menthol in 7/14 analyzed flavors (620–3,680 $\mu\text{g/g}$; Table 2). 3/14 Puffbar flavors contained all three compounds, with a combined concentrations of 8,830–48,050 $\mu\text{g/g}$ (Table 2). The nicotine concentrations in the Puffbar E-liquids ranged from 29.4 to 40.7 mg/g. As no PG/VG ratios were indicated on the product label, ratios were measured by GC/FID (Table 2).

Puffbar Power Specifications

Contrary to pod-based E-cigarettes, the disposable puffbar device does not contain a separate “pod” or cartridge that can be exchanged, but the e-liquid is housed within the metal frame of the device, where it is stored in an absorbent material much like in the first generation “cig-a-likes”.³¹ The coil resistance was measured at 1.7 Ω and the battery indicated a maximum power output of 3.7 V, which means the device has the exact power specifications as the Juul device,¹⁴ in addition to its look that also seem to mimic the Juul device. Maximum power output is thus calculated as 8.05 W.

Margin of Exposure Calculations for Synthetic Coolants in US-Marketed Refill E-liquids and Disposable E-cigarettes

To assess the risk associated with daily intake of WS-3 and WS-23 from E-cigarette use, we calculated the MOEs for different use scenarios (see Methods Section for details). For

organ toxicity, a safety margin MOE of >100 is considered low risk, while MOEs <100 require prioritization by regulatory agencies for risk mitigation for a given additive^{36,39}

For the E-liquid with the highest WS-3 concentration (Arctic Rush), MOEs ranged from 28 (1 mL/day) to 2 (15 mL/day; Supplementary Table S2 and Figure 1), and are thereby significantly lower than the recommended safety margin of 100. For consumption volumes of 3 mL/day and 5 mL/day, 13/24 and 18/24, respectively, of WS-3 containing refill E-liquids had a MOE of <100 . For 10 and 15 mL/day, MOEs were calculated only for the 20 free-base nicotine containing E-liquids: For 10 mL/day and 15 mL/day consumption volumes, 16/20 and 19/20 WS-3 containing E-liquids, respectively, yielded a MOE <100 (Supplementary Table S2 and Figure 1). MOEs for the one WS-23-containing refill E-liquid was <100 in all assessed consumption scenarios, except for 1 mL/day consumption (Supplementary Table S2 and Figure 1). The tested concentrate (“Agent Cool”) contained both WS-3 and WS-23. Based on vendor recommendations to add 1–2 drops of concentrate per pod, we conservatively estimated MOEs for a resulting dilution range of 1%–3%. Resulting MOEs for WS-23 when added at 1%–3% to E-liquids were <100 at all consumption rates, except for the lowest consumption rate (1-mL/day) at 1% dilution. Taken together, these results indicate that users of synthetic coolants flavored E-liquids are exposed to WS-3 and WS-23 levels that can potentially pose a health risk upon long-term use.




Figure 1. Margins of Exposure (MOE) for synthetic cooling agents from E-Cigarette refill liquids and coolant concentrate. MOEs are plotted for representative WS-3 (left) and WS-23 (right) -containing E-liquids and diluents with high, intermediate and low coolant concentrations, for daily consumption of 1, 3, 5, 10 and 15 mL E-liquid. No-Observed-Adverse-Effect-Levels (NOAEL) determined by JECFA in animal studies of organ toxicity upon oral administration were used for MOE calculations. A MOE of <100 signals a safety concern and requires regulatory mitigation. For WS-23 concentrate (Agent Cool, TBD Liquids), MOEs were calculated for dilutions within the range of manufacturer-recommendations to 1% and 3%. MOEs of the oral average daily intake of US consumers (US-OADI) determined by JECFA are displayed for comparison.

Next, we calculated the MOEs for synthetic coolants for daily consumption of Puffbar devices. For the Puffbar flavor with the highest content of both WS-3 and WS-23 (Cool Mint), the MOEs for consumption of 0.5–2 Puffbars/day ranged from 11 to 2 for WS-23 and 57–14 for WS-3, values that are significantly lower than the safety margin threshold of 100 (Supplementary Table S3 and Figure 2). MOEs for WS-3 were lower than the safety threshold of 100 at the highest estimated consumption rate (2-Puffbars/day) for all five WS-3 containing Puffbar flavors. MOEs for WS-3 were also <100 for Cucumber and Menthol flavors at 1-Puffbar/day consumption rate. For all other flavors, WS-23 MOEs were <100 for all use scenarios, with the exception of Café Latte (0.5 Puffbar/day), Cucumber (0.5 Puffbar/day), and Menthol (0.5 and 1-Puffbar/day). These results indicate that users of Puffbar disposable E-cigarettes are potentially exposed to levels of synthetic coolants that may pose organ toxicity risks, especially from WS-23 exposure.

Discussion

We detected the presence of synthetic coolants (WS-3 and WS-23) in US-marketed E-cigarette refill liquids and almost all tested flavors of the popular disposable E-cigarette brand, Puffbar, which was found to be the most popular device among US middle- and high school students in 2021.²⁸ Using a relatively low-power E-cigarette that resembles the power specifications of Puffbar and the popular Juul device, we demonstrate that these coolants can efficiently be aerosolized and carried over to reach user airways. Margins of

Exposure (MOE) were calculated to assess the health risks associated with exposure to these coolants from E-cigarette use, a parameter used by the US FDA and other global regulatory agencies to regulate the safe use of food additives with carcinogenic and organotoxic effects. These calculations suggest that daily intake of WS-3 or WS-23 by E-cigarette users exceed levels considered acceptable for consumption by FDA and the European Flavour Association (EFFA).

Natural cooling agents such as menthol provide a cooling sensation by activating cold-sensitive trigeminal nerve fibers innervating the upper respiratory tract and oral cavity. These sensory neurons express TRPM8 ion channels, the receptor for menthol that is also activated by cooling.^{40,41} Menthol also has counterirritant, antitussive and analgesic properties and inhibits respiratory irritation responses to tobacco smoke and its various irritant constituents through a TRPM8-dependent mechanism in rodent models.^{42,43} Menthol, through its actions on TRPM8, attenuated the aversive oral effects of nicotine in mice and increased nicotine intake.⁴⁴ The half-maximal concentrations of menthol to activate human TRPM8 is 4 μM .^{45,46} This concentration is known to be exceeded after inhalation of menthol cigarette smoke.⁴⁷ A menthol concentration of 0.65 μM (16 ppm), was sufficient to reduce sensory irritation by cigarette smoke exposure in mice, illustrating the strong counterirritant properties of menthol even at low concentrations.⁴² In E-cigarette users, menthol is transferred to the airways with high efficiency, where menthol concentrations per puff are comparable to concentrations in puffs generated from menthol cigarettes.^{14,33} Previous analytical studies demonstrated that menthol concentrations in marketed E-liquids range from 2 to 68,000 $\mu\text{g}/\text{mL}$ (13 μM –435 mM).⁴⁸ In the present study, menthol levels in E-cigarette refill liquids and Puffbar devices ranged from 330–23,490 $\mu\text{g}/\text{g}$ (~2.4–173 mM) and 620–3,680 $\mu\text{g}/\text{g}$ (~4.6–27 mM), respectively. These levels are sufficient to reduce sensory perception of irritation and harshness elicited by nicotine in humans.⁴⁹ Similar to menthol, synthetic cooling agents (including WS-3 and WS-23) also activate TRPM8 with increased efficacy (WS-3) and with comparable half-maximal concentrations (WS-3: 2–4 μM ; WS-23: 44 μM).^{45,50,51} Several of the refill E-liquids analyzed in this study contained WS-3 at levels comparable to menthol levels in mint- and menthol-flavored E-liquids (150–14,650 $\mu\text{g}/\text{g}$; ~816 μM –80 mM), with some containing synthetic coolants in addition to “natural” cooling agents. The concentrations of WS-3 in several Puffbar E-liquids, up to 1% (1,880–10,770 $\mu\text{g}/\text{g}$; 10–59 mM), are similar to menthol concentrations in liquids of Juul, the market-leading pod E-cigarette system in the United States.^{14,33} In addition, the majority of the Puffbar E-liquids analyzed in this study contained WS-23 ranging between 1% and 3.7% (up to 250 mM). Carryover of synthetic cooling agents from E-liquids into the vapor was highly efficient, approximating 100%, as previously demonstrated for menthol (Table 1, Supplementary Table S1).³³ Based on the puffing regimen in our study, the aerosol concentrations of WS-3 for majority of refill E-liquids (18/24 E-liquids) was calculated to be higher than 5 ppm (range: 5.8–98 ppm). Similarly, WS-3 and WS-23 concentrations for all of disposable Puffbars tested exceeded 10 ppm (WS-3: 12–72 ppm; WS-23: 13–303 ppm), which is sufficient to mediate TRPM8-dependent counterirritant effects. Since these synthetic coolants are either more potent or comparable TRPM8 agonists to menthol, it is likely they facilitate inhalation in vapers using the products analyzed by




Figure 2. Margins of Exposure (MOE) for synthetic cooling agents from Puffbar-branded disposable E-cigarettes. MOEs are plotted for exposures from Puffbar devices with flavored E-liquids containing both WS-3 and WS-23, at high (Cool mint), intermediate (Blueberry ice) or low (Menthol) concentrations, separated for each coolant (left: WS-3, right: WS-23), calculated for daily consumption of 1/2 Puffbar (0.65 mL), 1 Puffbar (1.3 mL), and 2 bars (2.6 mL). JECFA-determined NOAEL values were used for MOE calculations. MOEs of the oral average daily intake of US consumers (US-OADI) determined by JECFA are displayed for comparison.

us. One recent study also quantified menthol and the synthetic coolants (WS-3 and WS-23) in Puffbar devices.¹⁵ This study (Omaiye et al.) found similar quantities of synthetic coolants and a higher quantity of menthol in the Puffbar “Cool Mint” product, and higher quantities of all 3 compounds in the “Menthol” product.¹⁵ Some of these differences in amounts could be that, the tested “Cool Mint” device in this study was smaller in size (1.3 mL) than what was analyzed in Omaiye et al. study (3.2 mL). It is thus possible that new products were introduced and that e-liquid contents might have been adjusted simultaneously. Another observation to explain this difference could be the degree of e-liquid content variability (i.e., production inconsistency and manufacturing sites) of the Puffbar devices, such as evident in WS-23 contents in the “Menthol” product in this study and menthol content in the “Cool Mint” product reported earlier.¹⁵ This clearly illustrates the variability and inconsistencies of the manufacturing practices in the market place.

Our risk assessment analysis suggests that users of E-cigarettes containing synthetic cooling agents are exposed to amounts of WS-3 and WS-23 that exceed the amounts considered acceptable by regulators for intake from food. Detected amounts were much higher than the thresholds set by JECFA for human intake of flavor additives of this particular class of chemicals (aliphatic and aromatic amines and amides, FEMA Nos. 1594-1601, 1606 and 1613) at 90 µg/person/day, either added as individual flavoring or combined total.²⁰ Even consumption of just 1-mL of WS-3 containing E-cigarette liquid (150–14,650 µg/g) results in exposures exceeding JECFA’s safety threshold by ~2–180-fold. Vaping of 1 disposable Puffbar E-cigarette containing synthetic cooling agents would expose a user to amounts of WS-3 (1,880–10,770 µg/g) that exceed the threshold of concern by ~29–176-fold, or ~26–598-fold for WS-23 (3,940–36,660 µg/g). There were 5 Puffbar varieties that contained both WS-3 and WS-23, and when added together, would expose a user vaping 1 Puffbar device to ~126–774-fold the recommended maximal daily amount for amides, based on JECFA recommendations. These results strongly suggest that E-cigarette users are exposed to synthetic coolant amounts which would be considered unsafe when consumed in food, raising concerns for health risks upon chronic exposure.

The available chemical toxicity margins for WS-3 and WS-23, such as provided by JECFA, are derived from toxicological studies in animals in which coolants were administered orally.²⁰ Calculating MOE for inhalation of a compound based on past oral toxicity studies requires some caution. For human risk assessment and to calculate risk assessment parameters from exposure to these chemicals through various routes, it is routine practice for US and International regulatory agencies (US Environmental Protection Agency, EPA; European Chemical Agency, ECHA; UK’s Interdepartmental Group on Health Risks from Chemicals, IGHRC) to employ route-to-route (R2R) extrapolation.^{52,53} Regulatory agencies agree that the absorption efficiency for chemicals by the respiratory system is higher than the efficiency of the digestive tract and usually, an oral-to-inhalation ratio of 2 is applied for route-to-route extrapolation, reflecting the higher absorption of toxicants following inhalation.^{52–54} In this study, R2R extrapolation was not used, however, if applied, the MOE values would be even lower, reflecting an even higher risk associated with inhalation of these two synthetic cooling agents at various daily consumption volumes. This would suggest

even greater urgency for regulatory measures mitigating the risk from WS-3 and WS-23 exposures in E-cigarette users.

Recently, an e-vapor product company (RLX technology, China) has conducted a nose-only acute and subacute inhalation study for WS-23 (28-day exposure at doses of 19 and 29 mg/kg-bw/day).²⁴ Whole-body plethysmography was conducted after 28 days of repeated exposure to measure various parameters of pulmonary function. Compared to control and solvent groups, mice exposed to WS-23 demonstrated significant changes in various parameters, such as time of expiratory (Te), peak expiratory flow (PEF), relaxation time (RT), minute volume (MV), respiratory rate (F), end-inspiration pause (EIP), and end-expiratory pause (EEP). These results indicate that WS-23 may alter pulmonary function of users of WS-23 containing E-cigarettes. However, the study reports no remarkable histopathological changes in either the respiratory organs (nose, throat, trachea, and lungs) or other visceral organs (liver, kidney, heart, etc.). It has to be noted that the histopathological analysis was qualitative, and not based on a quantitative histopathologic scoring. The absence of histopathological evidence for organ toxicity contradicts prior WS-23 oral toxicity studies in rats that reported kidney lesions and hepatic toxicity starting at 10 mg/kg-bw/day (NOAEL of 5 mg/kg-bw/day).^{20,21} The discrepancy between these observations could be due to the different routes of exposure (oral vs. inhalation), or more importantly, may have resulted from the different exposure durations (14-weeks oral dosing vs. 4-weeks inhalational exposure).

In vitro genotoxicity studies of WS-23 in mammalian cells showed evidence for clastogenicity.²⁰ Clastogenicity was observed only in the presence of metabolic activation, indicating that the effect is mediated by the formation of a reactive metabolite. Due to the increased exposure to WS-23 from use of E-cigarettes, it is imperative that more studies are conducted to understand the mechanisms of their metabolism and further identify WS-23’s reactive clastogenic metabolites.

Previous studies have examined the toxicological effects of synthetic cooling agents, including WS-3 and WS-23, in isolation, and not in mixtures. As demonstrated in this study, refill E-cigarette liquids and popular disposable E-cigarettes contain mixtures of natural and synthetic cooling agents. Inhalation toxicity studies are required to examine the effects of acute, subacute and chronic exposures to mixtures of these chemicals and marketed E-cigarettes that contain such mixtures. Another limitation in assessing the risk associated from use of these E-cigarettes is the rapidly evolving product variety and the dynamic nature of their composition. The products analyzed in this study were purchased in the last 1–2 years. More recent marketed products may contain different compositions. Hence, continuous monitoring of new E-cigarette products by determining their chemical compositions for presence of potentially toxic flavorant levels is crucial for both regulatory and risk assessment purposes.

Following the withdrawal of flavored Juul cartridges from the US market, disposable E-cigarette products of the brand “Puffbar” brand have become increasingly popular among US youth and young adults.^{6,7} In Puffbar products, synthetic cooling agents were not only found in mint/menthol-flavored, or “cool”- or “ice”-labelled E-liquids but also in almost all products tested (13/14), suggesting that a cooling effect is preferred by consumers. Cooling agents were not only added to fruit-flavored E-liquids named “Orange Ice” and “Blueberry Ice” but also “Strawberry Banana” and “Lychee”, without the

“Ice” label, and in “Cucumber” and “OMG” flavors. This illustrates the utility of odorless synthetic cooling agents to add a cooling effect to flavors that likely would not be favored by users when combined with menthol, due to the incompatible minty odor the “natural” coolant menthol would add. Synthetic cooling agents may allow tolerance of more intense flavor combinations that would otherwise be irritating.

The fact that synthetic coolants are odorless also raises another important question: Legislation such as the FSPTCA or the European Tobacco Product Directive (TPD)^{1,55} banned “characterizing” flavorants from combustible tobacco products (with exemption for menthol in the United States), yet is unclear how synthetic coolants lacking a characterizing flavor would be treated. It has to be noted that TPD does not allow additives in tobacco products that facilitate inhalation, but no specific compounds that facilitate inhalation or method to measure facilitation are currently listed in the TPD. So, a replacement of the cooling compound menthol (with a minty characterizing flavor) with a synthetic coolants (lacking characterizing flavor) could be a strategy to circumvent a menthol ban. Policy makers should consider this when implementing menthol bans, such as both Canada and Germany have done in respective regulations that either maintain a positive list of compounds permitted as additives in tobacco products (Canada), or specifically ban a whole class of compounds that facilitates inhalation, including all natural and synthetic derivatives of menthol and other cooling agents (Germany).^{56,57}

Conclusions

Synthetic cooling agents are added to US-marketed E-cigarettes in a wide range of amounts that are comparable and relative to menthol amounts in flavored E-cigarettes. Synthetic coolants were found not only in mint- and menthol-flavored products but also in fruit-, dessert- and sweet-flavored E-cigarettes. These coolants are added to refill liquids, E-cigarette liquid diluents and popular disposable E-cigarettes such as Puffbar, at levels that pose potential health risks to users, thereby requiring for these products to be prioritized for risk mitigation measures.

Supplementary Material

A Contributorship Form detailing each author’s specific involvement with this content, as well as any supplementary data, are available online at <https://academic.oup.com/ntr>.

Funding

This work was supported by the National Institute on Drug Abuse (NIDA) of the National Institutes of Health (NIH) and the Center for Tobacco Products of the US Food and Drug Administration (FDA) (P50DA036151 and U54DA036151; Yale Tobacco Center of Regulatory Science, TCORS); and National Institute of Environmental Health Sciences (NIEHS) (R01ES029435 to Dr. Sven-Eric Jordt).

Declaration of Interests

The funding organization had no role in the design and conduct of the study; the collection, management, analysis, and interpretation of the data; the preparation, review, or approval of the manuscript; nor in the decision to submit the

manuscript for publication. The content is solely the responsibility of the authors and does not necessarily represent the views of National Institutes of Health (NIH) or the Food and Drug Administration (FDA). Dr. Jordt reports receiving personal fees from Hydra Biosciences LLC and Sanofi S.A. and nonfinancial support from GlaxoSmithKline Pharmaceuticals outside the submitted work. No other financial disclosures were reported by the authors of this paper.

Author Contributions

SVJ, HCE and SEJ conceptualized and designed the study; DGT, LD, and HCE acquired and analyzed the chemical analytical data; SVJ analyzed and interpreted the data for risk analysis calculations; SVJ, SEJ and HCE drafted the manuscript; SVJ, HCE, PTA, JBZ, and SEJ provided supervision; JBZ critically revised the manuscript for important intellectual content.

Data Availability

The data underlying this article are available in the article and in its online supplementary material.

References

1. FDA Center for Tobacco Products. Family Smoking Prevention and Tobacco Control Act - An Overview. *FDA*. Published online June 3, 2020. <https://www.fda.gov/tobacco-products/rules-regulations-and-guidance/family-smoking-prevention-and-tobacco-control-act-overview>. Accessed June 7, 2021.
2. FDA Center for Tobacco Products. Menthol and Other Flavors in Tobacco Products. *FDA*. Published online April 28, 2021. <https://www.fda.gov/tobacco-products/products-ingredients-components/menthol-and-other-flavors-tobacco-products>. Accessed May 14, 2021.
3. Villanti AC, Mowery PD, Delnevo CD, *et al*. Changes in the prevalence and correlates of menthol cigarette use in the USA, 2004–2014. *Tob Control*. 2016;25(Suppl 2):ii14–ii20.
4. Food and Drug Administration. FDA Commits to Evidence-Based Actions Aimed at Saving Lives and Preventing Future Generations of Smokers. *FDA*. Published April 30, 2021. <https://www.fda.gov/news-events/press-announcements/fda-commits-evidence-based-actions-aimed-saving-lives-and-preventing-future-generations-smokers>. Accessed May 10, 2021.
5. Office of the Commissioner. FDA finalizes enforcement policy on unauthorized flavored cartridge-based e-cigarettes that appeal to children, including fruit and mint. *FDA*. Published March 24, 2020. <https://www.fda.gov/news-events/press-announcements/fda-finalizes-enforcement-policy-unauthorized-flavored-cartridge-based-e-cigarettes-appeal-children>. Accessed May 14, 2021.
6. Wang TW, Neff LJ, Park-Lee E, *et al*. E-cigarette use among middle and high school students — United States, 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69:1310–1312. doi:10.15585/mmwr.mm6937e1.
7. Ali FRM, Diaz MC, Vallone D, *et al*. E-cigarette unit sales, by product and flavor type — United States, 2014–2020. *MMWR Morb Mortal Wkly Rep*. 2020;69:1313–1318. doi:10.15585/mmwr.mm6937e2.
8. Leffingwell JC. Cool without Menthol & Cooler than Menthol and Cooling Compounds as Insect Repellents. *Cool without Menthol & Cooler than Menthol*. Published June 8, 2018. http://www.leffingwell.com/cooler_than_menthol.htm. Accessed May 14, 2021.
9. Leffingwell JC, Rowsell D. Wilkinson sword cooling compounds: from the beginning to now. *Perfumer Flavorist*. 2014;39(3):34–43.
10. Leffingwell JC. WILKINSON-SWORD “CWM”. Published online March 5, 1975. <https://www.industrydocuments.ucsf.edu/docs/tkjin0089>

11. Newman F. Memorandum From Philip Morris In-House Counsel (Newman) To Philip Morris In-House Counsel (Holtzman) Providing Legal Advice And Analysis Regarding Content Of Draft Press Release For New Product Introduction; Northwind Comments On 3rd Tier Inquires. March 14, 1981. <https://www.industrydocuments.ucsf.edu/docs/kgxy0101>. Accessed May 24, 2021.
12. Reger L, Julia M, Hahn H, Hahn J. Analysis of menthol, menthol-like, and other tobacco flavoring compounds in cigarettes and in electrically heated tobacco products. *Beiträge Tabakforsch Int Contrib Tobacco Res.* 2018;28(2):93–102.
13. Jabba SV, Jordt SE. Turbocharged Juul device challenges European tobacco regulators. *Eur Respir J.* 2020;56(2):2002430.
14. Erythropel HC, Anastas PT, Krishnan-Sarin S, et al. Differences in flavourant levels and synthetic coolant use between USA, EU and Canadian Juul products. *Tob Control.* 2021;30:453–455. doi:10.1136/tobaccocontrol-2019-055500.
15. Omaiye EE, Luo W, McWhirter KJ, Pankow JF, Talbot P. Flavour chemicals, synthetic coolants and pulegone in popular mint-flavoured and menthol-flavoured e-cigarettes. *Tob Control.* 2021. doi:10.1136/tobaccocontrol-2021-056582.
16. Liquid Nicotine Wholesalers. Koolada 10% PG (TFA). *Liquid Nicotine Wholesalers.* <https://liquidnicotinewholesalers.com/flavor-apprentice-koolada-10-pg.html>. Accessed June 8, 2021.
17. Jabba SV, Jordt SE. Risk analysis for the carcinogen pulegone in mint- and menthol-flavored e-cigarettes and smokeless tobacco products. *JAMA Intern Med.* 2019. doi:10.1001/jamainternmed.2019.3649.
18. Jabba SV, Jordt SE. Estimating fluid consumption volumes in electronic cigarette use-reply. *JAMA Intern Med* 2020;180(3):468–469.
19. Hallagan J. The Safety Assessment and Regulatory Authority to Use Flavors – Focus on E-Cigarettes. 2015. https://www.femaflavor.org/sites/default/files/FEMAGRAS_Ecig_March_3_2015.pdf. Accessed June 8, 2021.
20. Joint Expert Committee on Food Additives, ed. *Evaluation of Certain Food Additives: Sixty-Ninth Report of the Joint FAO/WHO Expert Committee on Food Additives; [Rome, 17-26 June 2008]; 2009.* <http://www.inchem.org/documents/jecfa/jecmono/v952je01.pdf>
21. Cheng M. *A Subchronic Oral Toxicity Study of T0332.04 in Rats. Unpublished Report to the Flavor Manufacturers Association. Submitted to WHO by the Flavor and Extract Manufacturers Association of the United States, Washington DC, USA.* Madison, Wisconsin, USA: Hazleton Laboratories; 1982.
22. Miyata K. *Summary of 28-Day Repeated Dose Oral Toxicity Study of WS-3. Unpublished Report to the Flavor Manufacturers Association.* Washington DC, USA: WHO by the Flavor and Extract Manufacturers Association of the United States; 1995 (Submitted).
23. Joint FAO/WHO Expert Committee on Food Additives. Meeting (65th: 2005: Geneva Switzerland), International Program on Chemical Safety. *Safety Evaluation of Certain Food Additives.* World Health Organization, International Programme on Chemical Safety; 2006.
24. Wu ZH, Liu YS, Li XD, et al. Acute and subacute inhalation toxicity assessment of WS-23 in Sprague-Dawley rats. *J Appl Toxicol.* 2021. doi:10.1002/jat.4166.
25. Leventhal A, Dai H, Barrington-Trimis J, Sussman S. “Ice” flavoured e-cigarette use among young adults. *Tob Control.* 2021. doi:10.1136/tobaccocontrol-2020-056416.
26. Galimov A, Leventhal A, Meza L, et al. Prevalence of disposable pod use and consumer preference for e-cigarette product characteristics among vape shop customers in Southern California: a cross-sectional study. *BMJ Open* 2021;11(10):e049604.
27. Davis DR, Morean ME, Bold KW, et al. Cooling e-cigarette flavors and the association with e-cigarette use among a sample of high school students. *PLoS One.* 2021;16(9):e0256844.
28. Park-Lee E. Notes from the Field: E-Cigarette use among middle and high school students—National Youth Tobacco Survey, United States, 2021. *MMWR Morb Mortal Wkly Rep.* 2021;70. doi:10.15585/mmwr.mm7039a4.
29. Gaiha SM, Lempert LK, McKelvey K, Halpern-Felsher B. E-cigarette devices, brands, and flavors attract youth: Informing FDA’s policies and priorities to close critical gaps. *Addict Behav.* 2022;126:107179. doi:10.1016/j.addbeh.2021.107179.
30. Erythropel HC, Jabba SV, DeWinter TM, et al. Formation of flavorant-propylene glycol adducts with novel toxicological properties in chemically unstable e-cigarette liquids. *Nicotine Tob Res.* 2019;21(9):1248–1258.
31. Rosbrook K, Erythropel HC, DeWinter TM, et al. The effect of sucralose on flavor sweetness in electronic cigarettes varies between delivery devices. *PLoS One.* 2017;12(10):e0185334.
32. Talih S, Balhas Z, Eissenberg T, et al. Effects of user puff topography, device voltage, and liquid nicotine concentration on electronic cigarette nicotine yield: measurements and model predictions. *Nicot Tobacco Res.* 2015;17(2):150.
33. Erythropel HC, Davis LM, de Winter TM, et al. Flavorant-solvent reaction products and menthol in JUUL E-cigarettes and aerosol. *Am J Prev Med.* 2019;57(3):425–427.
34. WHO/IPC. *Environmental Health Criteria 239: Principles for Modelling Dose-Response for the Risk Assessment of Chemicals.* World Health Organization https://apps.who.int/iris/bitstream/handle/10665/44065/WHO_EHC_240_8_eng_Chapter5.pdf?sequence=8
35. Benford D, Bolger PM, Carthew P, et al. Application of the Margin of Exposure (MOE) approach to substances in food that are genotoxic and carcinogenic. *Food Chem Toxicol.* 2010;48:S2–S24. doi:10.1016/j.fct.2009.11.003.
36. Baumung C, Rehm J, Franke H, Lachenmeier DW. Comparative risk assessment of tobacco smoke constituents using the margin of exposure approach: the neglected contribution of nicotine. *Sci Rep.* 2016;6(1):35577.
37. Chaumont M, de Becker B, Zaher W, et al. Differential effects of e-cigarette on microvascular endothelial function, arterial stiffness and oxidative stress: a randomized crossover trial. *Sci Rep.* 2018;8(1):10378.
38. Smets J, Baeyens F, Chaumont M, Adriaens K, Van Gucht D. When less is more: vaping low-nicotine vs. high-nicotine E-Liquid is compensated by increased wattage and higher liquid consumption. *Int J Environ Res Public Health.* 2019;16(5). doi:10.3390/ijerph16050723.
39. National Research Council (U.S.), ed. *Risk Assessment in the Federal Government: Managing the Process.* National Academy Press; 1983.
40. McKemy DD, Neuhauser WM, Julius D. Identification of a cold receptor reveals a general role for TRP channels in thermosensation. *Nature* 2002;416(6876):52–58.
41. Peier AM, Moqrich A, Hergarden AC, et al. A TRP channel that senses cold stimuli and menthol. *Cell* 2002;108(5):705–715.
42. Willis DN, Liu B, Ha MA, Jordt SE, Morris JB. Menthol attenuates respiratory irritation responses to multiple cigarette smoke irritants. *FASEB J.* 2011;25(12):4434–4444.
43. Ha MA, Smith GJ, Cichocki JA, et al. Menthol attenuates respiratory irritation and elevates blood cotinine in cigarette smoke exposed mice. *PLoS One.* 2015;10(2):e0117128.
44. Fan L, Balakrishna S, Jabba SV, et al. Menthol decreases oral nicotine aversion in C57BL/6 mice through a TRPM8-dependent mechanism. *Tob Control.* 2016;25(Suppl 2):ii50–ii54.
45. Behrendt HJ, Germann T, Gillen C, Hatt H, Jostock R. Characterization of the mouse cold-menthol receptor TRPM8 and vanilloid receptor type-1 VR1 using a fluorometric imaging plate reader (FLIPR) assay. *Br J Pharmacol.* 2004;141(4):737–745.
46. Klein AH, Iodi Carstens M, McCluskey TS, et al. Novel menthol-derived cooling compounds activate primary and second-order trigeminal sensory neurons and modulate lingual thermosensitivity. *Chem Senses.* 2011;36(7):649–658.
47. Tobacco Products Scientific Advisory Committee. *Menthol Cigarettes and Public Health: Review of the Scientific Evidence and Recommendations.* US Department of Health and Human Services,

- Food and Drug Administration; 2011:1–252. <https://wayback.archive-it.org/7993/20170405201731/https://www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/TobaccoProductsScientificAdvisoryCommittee/UCM269697.pdf>. Accessed April 27, 2021.
48. Omaiye EE, McWhirter KJ, Luo W, *et al.* High concentrations of flavor chemicals are present in electronic cigarette refill fluids. *Sci Rep.* 2019;9(1):2468.
49. Rosbrook K, Green BG. Sensory effects of menthol and nicotine in an e-cigarette. *Nicot Tobacco Res.* 2016;18(7):1588–1595.
50. Sherkheli MA, Vogt-Eisele AK, Bura D, *et al.* Characterization of selective TRPM8 ligands and their structure activity response (S.A.R) relationship. *J Pharm Pharm Sci.* 2010;13(2):242–253.
51. Johnson S, Tian M, Sheldon G, Dowd E. Trigeminal receptor study of high-intensity cooling agents. *J Agric Food Chem.* 2018;66(10):2319–2323.
52. European Chemicals Agency (ECHA). *Guidance on Information Requirements and Chemical Safety Assessment Chapter R.8: Characterisation of Dose [Concentration]-Response for Human Health.* European Chemicals Agency; 2012. https://echa.europa.eu/documents/10162/13632/information_requirements_r8_en.pdf
53. The Interdepartmental Group on Health Risks from Chemicals (IGHRC). *Guidelines on Route-to-Route Extrapolation of Toxicity Data When Assessing Health Risks of Chemicals.* Cranfield University; 2006. http://www.iehconsulting.co.uk/IEH_Consulting/IEHCPubs/IGHRC/cr12.pdf
54. Schröder K, Escher SE, Hoffmann-Dörr S, *et al.* Evaluation of route-to-route extrapolation factors based on assessment of repeated dose toxicity studies compiled in the database RepDose®. *Toxicol Lett.* 2016;261:32–40. doi:10.1016/j.toxlet.2016.08.013.
55. European Union. Revision of the Tobacco Products Directive. Public Health - European Commission. Published November 25, 2016. https://ec.europa.eu/health/tobacco/products/revision_en. Accessed June 8, 2021.
56. Tabakerz V - Verordnung über Tabakerzeugnisse und verwandte Erzeugnisse. <https://www.gesetze-im-internet.de/tabakerzv/BJNR098010016.html>. Accessed June 8, 2021.
57. Health Canada. Order Amending the Schedule to the Tobacco Act (Menthol). Published August 2, 2017. <https://www.canada.ca/en/health-canada/programs/consultation-changes-tobacco-act-address-menthol/order-amending-schedule-tobacco-act-menthol.html>. Accessed June 8, 2021.