

Design and Application of a Biosensor for Monitoring Toxicity of Compounds to Eukaryotes

R. P. HOLLIS,¹ K. KILLHAM,² AND L. A. GLOVER^{1*}

Department of Molecular and Cell Biology, University of Aberdeen, Institute of Medical Sciences, Aberdeen AB25 2ZD,¹ and Department of Plant and Soil Science, University of Aberdeen, Aberdeen AB24 3UU,² United Kingdom

Received 21 September 1999/Accepted 31 December 1999

Here we describe an alternative approach to currently used cytotoxicity analyses through applying eukaryotic microbial biosensors. The yeast *Saccharomyces cerevisiae* was genetically modified to express firefly luciferase, generating a bioluminescent yeast strain. The presence of any toxic chemical that interfered with the cells' metabolism resulted in a quantitative decrease in bioluminescence. In this study, it was demonstrated that the luminescent yeast strain senses chemicals known to be toxic to eukaryotes in samples assessed as nontoxic by prokaryotic biosensors. As the cell wall and adaptive mechanisms of *S. cerevisiae* cells enhance stability and protect from extremes of pH, solvent exposure, and osmotic shock, these inherent properties were exploited to generate a biosensor that should detect a wide range of both organic and inorganic toxins under extreme conditions.

Many luminescent bacterial biosensors have been produced which detect a wide range of pollutants while simultaneously assessing bioavailability in environmental samples. *Saccharomyces cerevisiae* has been used previously to assess toxicity through the use of an amperometric gas diffusion (oxygen) electrode, which quantifies changes in culture respiration (1, 7, 8, 9, 17). Alternatively, the effect of a compound on *S. cerevisiae* cultures was measured directly through inhibition of maximum growth rates (2, 10). However, such toxicity assays are time-consuming and expensive in comparison to luminometry analysis. Walmsley et al. (18) created an *S. cerevisiae* biosensor that induces green fluorescent protein expression on exposure to genotoxic agents. The luminescent biosensor designed in the present study works on a different principal (reduction in reporter gene product activity) and complements the biosensor designed by Walmsley et al. (18) by detecting a wide range of toxins, not just genotoxic agents.

For the novel *S. cerevisiae* biosensor described here, a luminescence detection system was constructed using firefly luciferase (*luc*) from *Photinus pyralis*. The firefly luciferase light reaction relies on ATP being supplied by actively metabolizing cells. This dependence on endogenous energy supplies enables a luciferase assay system to report directly on cell health upon exposure to toxins. The luciferin substrate for *P. pyralis* luciferase is an amphipathic molecule with a charged carboxyl group at physiological pH. This prevents easy passage of luciferin across cell membranes, leading to problems during the exogenous addition for in vivo assays. This was overcome through the development of a novel assay system where the biosensor preparation was acidified after exposure to the toxicant and before luminescence quantification.

MATERIALS AND METHODS

Strains, media, and chemicals. *S. cerevisiae* strain W303-1B (*MAT α leu2-3,112 his3-11,15 trp1-1 can1-100 ade2-1 ura3-1*) (16) was the host for the chromosomal

luciferase-expressing construct. Chromosomal insertion was carried out using a published method (5). *S. cerevisiae* was grown in synthetic complete medium (15) in a shaking incubator at 200 rpm and 30°C. Diuron [3-(3,4-dichlorophenyl)-1,1-dimethylurea] and mecoprop [(+/-)-2-(4-chloro-*o*-tolylxy)propionic acid] were obtained from Greyhound Chemicals. D-Luciferin (potassium salt) was obtained from Molecular Probes.

Construction of the chromosomal luciferase expression system. The pBlue-script-based yeast centromeric plasmid pRS316 (14), was modified to create pLUC Δ P. A *PGK* terminator was amplified by PCR from YCpPLP (4). Primer 5SR (5'-GTGTTGCTTCTTATCCGCGGAGAAAATAAATTGAAT-3') introduced a *SacII* site at the 5' end of the terminator region, and a *SacI* site was inserted by the 3SR (5'-TTTTTCGAAACGCAGAGCTCTCGAGTTATTTAACTT-3') oligonucleotide at the 3' end. The PCR to amplify *luc* used the forward primer 5LEADL (5'-ACAGATCACCGGATCCATCAAGACACCAATCAAAACAATAAAAACATCATCACAATGGAAGACGCCAAAACATAAAGA AAGGCCG-3') and reverse primer N3RA (5'-TCTAGAGCGGCCGCTGAA TACAGTTACATTTTACTTTCCGCCCTTCTTGGCCTTT-3') for the inclusion of a *NoI* site 3' to the luciferase gene. The template for this PCR was the pGL2 vector from Promega. The *luc Δ* and *URA3* genes from the pLUC Δ P plasmid vector were amplified separately using PCR. Primers rLUC (5'-AGCC TCATAAATAAAGGTAGATAGTAAAGTATACAAGAGAAGAATCCCA AGATGGAAGACGCCAAAACATAAAGAAGGCCG-3') and S3R amplify a promoterless version of the *luc Δ* gene (includes the *PGK* terminator). Primers rURA (5'-ATCAAACATCATTCTGCAGAACTGAAAACATACTT GAACACTTGGGACAGCTGACCTGATGCGGTATTTTCTCCTTACGCA TCT-3') and ST1K1 (5'-GAGCTCTGCGTTTCGAAAACCGGAGACGGTC ACAGCTT-3') amplify the *URA3* gene, including control regions. Homologous sequence for directing genomic integration at *rps16a* were included in the rLUC and rURA primers. Homology to the luciferase amplification product (present in ST1K1) allowed the luciferase and *URA3* genes to fuse and amplify through this homologous region, resulting in a 3.5-kb product.

Bioassay procedure. Stock concentrations of toxicants were prepared in deionized water, and the pH was adjusted to 5.5 using HCl or NaOH. *S. cerevisiae* biosensor cells were harvested at peak luminescence (optical density at 600 nm of around 3.7), centrifuged at 700 \times g, and washed twice in 0.1 M KCl (twice the original volume). A potential toxicant (450 μ l) was added to each 1-ml cuvette (Clinicon, catalog no. 2174 701), and the final volume was made up to 500 μ l by adding 50 μ l of diluted *S. cerevisiae* cells (approximately 2×10^7 cells per assay). Following a 10-min exposure to the sample, 500 μ l of pH 2.5 citrate phosphate buffer (containing luciferin to make the final concentration 0.1 mM per cuvette) was added. Bioluminescence was then monitored in a BioOrbit 1251 luminometer using a Multiuse software package (version 1.01, April 1991, JN). The units of luminescence were expressed as relative light units, which equated to 10 mV s⁻¹ ml⁻¹. For comparative bacterial assays, lyophilized *Escherichia coli* HB101 cultures containing the plasmid pUCD607 were prepared as described by Weitz (19). These lyophilized cells were resuscitated for 1 h in 10 ml of 0.1 M KCl at 25°C and 200 rpm. For the assay, 100 μ l of resuscitated cells was added to 900 μ l of toxicant standard in each cuvette. In order to interpret the bioassay data, the relative light units were translated into a percentage of maximum luminescence (the 100% value was determined by the blank cuvette). For toxicity detection in solvents, the standards were prepared using 10% solvent as a diluent. The maximum luminescence for these assays was determined using the relevant

* Corresponding author. Mailing address: Department of Molecular and Cell Biology, University of Aberdeen, Institute of Medical Sciences, Foresterhill, Aberdeen AB25 2ZD, United Kingdom. Phone: 44 (0)1224 273099. Fax: 44 (0)1224 273144. E-mail: l.a.glover@abdn.ac.uk.




FIG. 1. Differential response of *E. coli* HB101(pUCD607) and *S. cerevisiae* Luc Δ to diuron. *S. cerevisiae* and *E. coli* cells were exposed to identical concentrations of diuron. *S. cerevisiae* (●) exhibits a toxic response at levels of toxin where the *E. coli* (○) cells do not sense any toxicity. The experiment was carried out in triplicate at 25°C, and the error bars represent standard errors of the mean triplicate values. For comparative assays the same biosensor sample was used; however, for the assays presented here, they were repeated on at least three occasions and the trends were found to be identical.

solvent diluent. Samples were run alongside standards prepared in H₂O for comparative analysis. To determine the effects of external pH on light output, a pH range of 1 to 12 was prepared using HCl and NaOH to adjust the pH of H₂O. To assay toxicity at pH extremes, the standards were prepared in the normal fashion except that the H₂O for dilution had its pH adjusted accordingly.

RESULTS

The herbicide diuron, a known eukaryotic toxin, was toxic to *S. cerevisiae* but not to a *lux*-marked luminescent bacterial strain (Fig. 1). Investigations performed to examine the effects of external pH on *S. cerevisiae* demonstrated that external pH had minimal effects on the metabolism of the cells if the cells were acidified prior to luminescence quantification (data not shown). To discover if altering the pH affected toxicity detection, two test compound solutions (copper and the organic toxin mecoprop) were pH adjusted. Copper was no longer found to be bioavailable at extremes of pH, presumably due to speciation effects (data not shown). However, the sensitivity of the *S. cerevisiae* biosensor to mecoprop was greatest at pH 1 and was reduced at pH 12 (Fig. 2), with a similar dose response over a pH range of pH 3 to 10. *S. cerevisiae* was found, in acute exposure, to be relatively tolerant to fairly high concentrations of solvents, with a reduction in light output of between 10 to 40% in the presence of 9% solvent. Two compounds were assayed to determine how the presence of solvent affected toxicity, and results were obtained for the heavy metal copper and the organic toxin diuron (data for copper are shown in Fig. 3). The presence of methanol severely reduced the biosensor response to both organic and inorganic toxins. Conversely, the presence of ethanol considerably enhanced the biosensor sensitivity to both toxins.

DISCUSSION

During screening of known toxic herbicides, the *S. cerevisiae* biosensor sensed toxicity in samples that were not toxic to existing *lux*-based bacterial biosensors (Fig. 1). However, the bacterial biosensors did not utilize a *luc*-based luminescence system; therefore, the results demonstrated in Fig. 1 are not directly comparable. The effects of external pH on mecoprop toxicity, as detected by the luminescent *S. cerevisiae* biosensor, are shown in Fig. 2. The most dramatic increases in toxicity sensing were seen in toxic responses at pH 1 and 2. A possible explanation for this is that weak acids have been demonstrated to dissipate the proton motive force across plasma membranes in *S. cerevisiae* (6). To counteract this effect, the plasma membrane pumps out protons using the membrane ATPase at the expense of ATP production (12). Many organic toxins disrupt cell plasma membranes (13) which are required for defense against extreme pH, and therefore, the increase in toxicity at low pH may be caused by mecoprop disrupting the cell membrane, which in turn disrupts *S. cerevisiae* pH defense mechanisms, resulting in an amplified toxic effect.

S. cerevisiae has been applied in xenobiotic testing in other studies where its solvent tolerance was advantageous (2). However, possible synergistic effects of these solvents and xenobiotics were never discussed. The differences in toxicity detection in ethanol and methanol are quite distinct. Methanol did not act synergistically with either the organic or inorganic toxins (results for copper are shown in Fig. 3). This is unlikely to be explained by slightly reduced membrane disruption, as *S. cerevisiae* has many mechanisms that will readily import ethanol into the cell (3). Therefore, during ethanol transport into cells, it is possible that toxins simultaneously enter through similar




FIG. 2. Effects of pH on mecoprop toxicity in *S. cerevisiae* Luc Δ . The external pH was found to have minimal effects on dose-response curves for pH 3 (Δ), pH 5.5 (\square , solid lines), and pH 10 (\blacksquare , solid lines). External pHs of 11 (\bullet) and pH 12 (\circ) resulted in a loss of toxin sensitivity. When the external pH was lowered (dashed lines) to pH 1 (\square) and pH 2 (\blacksquare), there were dramatic increases in sensitivity to the toxin. The experiment was carried out in triplicate at 25°C, and the error bars represent standard errors of the mean triplicate values. For comparative assays the same biosensor sample was used; however, the assays were repeated on at least three separate occasions and the trends were found to be identical.




FIG. 3. Effects of solvents on copper toxicity sensing in *S. cerevisiae* Luc Δ . The presence of 9% ethanol (\blacksquare) resulted in an increase in the toxic response, and a loss in sensitivity was observed for 9% methanol (\square), compared to the dose-response curve for double-distilled H₂O (Δ). The presence of 9% acetone (\bullet) resulted in a dose-response curve similar to that for double-distilled H₂O. These experiments were carried out in triplicate at 25°C, and the error bars represent standard errors of the mean triplicate value. For comparative assays the same biosensor sample was used; however, the assays were repeated on at least three separate occasions and the trends were found to be identical.

uptake mechanisms. Comparable findings have been observed for higher eukaryotes; for example, the presence of ethanol increases the toxicity of paraquat in rabbits (11).

Toxic responses from these cytotoxicity analyses are indicative rather than definitive. This is because *S. cerevisiae* responds to some toxic compounds which are not deleterious to higher eukaryotes, for example, cyclohexane (17) and antifungal agents. However, *luc*-marked *S. cerevisiae* may be a very useful reporter of toxicity to other fungi, such as mycorrhizal fungi, which have major ecological importance.

ACKNOWLEDGMENTS

We acknowledge the support of the Natural Environment Research Council (NERC) and Yorkshire Water Services Ltd. for funding a studentship for R. P. Hollis.

REFERENCES

1. Campanella, L., G. Favero, D. Mastrofino, and M. Tomassetti. 1996. Toxicity order of cholanic acids using an immobilised cell biosensor. *J. Pharm. Biomed. Anal.* **14**:1007–1013.
2. Cascorbi, L., H. Bittrich, J. Ricklinkat, W. Voss, A. Seyfarth, and M. Forêt. 1993. Effects of a heterogeneous set of xenobiotics on growth and plasma membranes of mammalian and fungal cell cultures. *Ecotoxicol. Environ. Saf.* **26**:113–126.
3. D'Amore, T., C. J. Panchal, I. Russell, and G. G. Stewart. 1990. A study of ethanol tolerance in yeast. *Crit. Rev. Biotechnol.* **9**:287–304.
4. Dickson, L. M., and A. J. P. Brown. 1998. mRNA translation in yeast during entry into stationary phase. *Mol. Gen. Genet.* **259**:282–293.
5. Gietz, D., A. St. Jean, R. A. Woods, and R. H. Schiestl. 1992. Improved method for high efficiency transformation of intact yeast cells. *Nucleic Acids Res.* **20**:1425.
6. Guldfeldt, L. U., and N. Arneborg. 1998. Measurement of the effects of acetic acid and extracellular pH on intracellular pH of nonfermenting, individual *Saccharomyces cerevisiae* cells by fluorescence microscopy. *Appl. Environ. Microbiol.* **64**:530–534.
7. Haubenstricker, M. E., P. G. Meier, K. H. Mancy, and M. J. Brabec. 1990. Rapid toxicity testing based on yeast respiratory activity. *Bull. Environ. Contam. Toxicol.* **44**:669–674.
8. Henderson, G. 1989. A comparison of effects of chromate, molybdate and cadmium oxide on respiration in the yeast *Saccharomyces cerevisiae*. *Biol. Metals* **2**:83–88.
9. Ikebukuro, K., A. Miyata, S. J. Cho, Y. Nomura, S. M. Chang, Y. Yamauchi, Y. Hasebe, S. Uchiyama, and I. Karube. 1996. Microbial cyanide sensor for monitoring river water. *J. Biotechnol.* **48**:73–80.
10. Koch, H. P. 1993. The yeast test: an alternative method for the testing of acute toxicity of drug substances and environmental chemicals. *Methods Find. Exp. Clin. Pharmacol.* **15**:141–152.
11. Kuo, T. L., and R. Nanikawa. 1990. Effect of ethanol on acute paraquat toxicity in rabbits. *Nippon Hoigaku Zasshi.* **44**:12–17.
12. Pampulha, M. E., and M. C. Loureiro-Dias. 1989. Combined effect of acetic acid, pH and ethanol on intracellular pH of fermenting yeast. *Appl. Microbiol. Biotechnol.* **31**:547–550.
13. Sikkema, J., J. A. M. de Bont, and B. Poolman. 1995. Mechanisms of membrane toxicity of hydrocarbons. *Microbiol. Rev.* **59**:201–222.
14. Sikorski, R. S., and P. Hieter. 1989. A system of shuttle vectors and yeast host strains designed for efficient manipulation of DNA in *Saccharomyces cerevisiae*. *Genetics* **122**:19–27.
15. Strathern, J. N. 1994. Ty Insertional mutagenesis, p. 118. *In* J. R. Johnston (ed.), *Molecular genetics of yeast: a practical approach*. Oxford University Press, Oxford, United Kingdom.
16. Thomas, B. J., and R. Rothstein. 1989. Elevated recombination rates in transcriptionally active DNA. *Cell* **56**:619–630.
17. Uribe, S., P. Rangel, G. Espinola, and G. Aguirre. 1990. Effects of cyclohexane, an industrial solvent, on the yeast *Saccharomyces cerevisiae* and on isolated yeast mitochondria. *Appl. Environ. Microbiol.* **56**:2114–2119.
18. Walmsley, R. M., N. Billinton, and W.-D. Heyer. 1997. Green fluorescent protein as a reporter for the DNA damage-induced gene *RAD54* in *Saccharomyces cerevisiae*. *Yeast* **13**:535–545.
19. Weitz, H. J. 1999. Ph.D. thesis. University of Aberdeen, Aberdeen, United Kingdom.