

Inpatient glycemic control and community-acquired pneumonia outcomes in the pre-COVID-19 era: reviewing the evidence to pave the road for future studies

Paras B Mehta ,¹ Aidar R Gosmanov ^{2,3}

To cite: Mehta PB, Gosmanov AR. Inpatient glycemic control and community-acquired pneumonia outcomes in the pre-COVID-19 era: reviewing the evidence to pave the road for future studies. *BMJ Open Diab Res Care* 2022;10:e003011. doi:10.1136/bmjdr-2022-003011

Received 24 June 2022
Accepted 27 June 2022



► <http://dx.doi.org/10.1136/bmjdr-2022-002880>



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Division of Endocrinology and Metabolism, University of California San Francisco, San Francisco, California, USA

²Division of Endocrinology, Department of Medicine, Albany Medical College, Albany, New York, USA

³Endocrinology Section, Stratton VAMC, Albany, New York, USA

Correspondence to

Dr Paras B Mehta;
paras.mehta@ucsf.edu

Over the past several decades, our understanding of inpatient glycemic control has grown, and paradigms of management have shifted with results of key studies such as the Leuven Surgical Trial in 2001 and NICE-SUGAR trial in 2009.^{1 2} Inpatient glycemic targets in society guidelines, including the 2012 Endocrine Society recommendations and the Surviving Sepsis Campaign, significantly changed to a more liberal glycemic range as a result of the NICE-SUGAR results.^{3 4} Despite these advances, gaps exist in our knowledge, including more precise evaluations of inpatient glycemic targets and whether hyperglycemia is associated with or potentially causes unfavorable clinical outcomes in hospitalized patients.

In this issue of the journal, Barmanray *et al* report the findings of a thorough and methodical systematic review and meta-analysis of observational studies on associations of inpatient glycemic control and community-acquired pneumonia (CAP) prior to the COVID-19 pandemic.⁵ They note that inpatient hyperglycemia was associated with in-hospital mortality and intensive care unit (ICU) admission, while the presence of diabetes (irrespective of glycemic control) was not. This systematic review provides key insights into the likely relationship between inpatient hyperglycemia and CAP outcomes, and also highlights several important gaps in our knowledge of inpatient glycemic management.

This study importantly adds to the literature on the association between hyperglycemia and hospital outcomes, as such analyses are limited. A review of literature by an Endocrine Society task force noted that intensive glycemic control might be linked

to decreased risk of infection in hospitalized persons with diabetes.⁶ Of note, studies that included patients with stress hyperglycemia or hyperglycemia from previously unrecognized diabetes were not part of this analysis. Additionally, a retrospective study of the Veterans Affairs database found an association between increasing mean glucose and mortality in critically ill patients hospitalized with sepsis or pneumonia.⁷

Delving further into the impact of hyperglycemia, Barmanray *et al* demonstrate that an acutely elevated blood glucose portends a poor inpatient outcome more so than having underlying diabetes.⁵ The authors describe the pathophysiologic mechanism between hyperglycemia and infection, and suggest that chronic non-glycemic effects of diabetes may not play a role in the acute setting. Prior research has similarly suggested that hyperglycemia in patients without diabetes leads to increased mortality and poorer functional status than in those with diabetes.^{8 9} The underlying reason for this connection is not clearly understood, though we hypothesize that hyperglycemia in individuals without a known diagnosis of diabetes hospitalized for reasons such as CAP may be considered a stress response, initially raising the threshold for therapeutic antihyperglycemic interventions. With essential time to pharmacologically lower hyperglycemia potentially being lost, known detrimental effects of hyperglycemia on the immune system may prevent effective treatment of infection including CAP. Alternatively, inpatient hyperglycemia could be just a marker of infection severity. Though only future prospective interventional trials will be able to delineate whether non-diabetic hyperglycemia impacts inpatient

outcomes, we believe that proactive and early management of stress hyperglycemia should be implemented in patients without diabetes, similarly to the innate attention devoted to glycemic control in patients with diabetes.

The publication by Barmanray *et al* also reminds us of the challenge in defining evidence-based inpatient glycemic targets. While 17 studies examining the relationship between hyperglycemia and CAP outcomes were identified by the authors, significant heterogeneity existed in definitions of hyperglycemia. For example, a glucose of 126–180 mg/dL was considered ‘mild hyperglycemia’ in one study but would qualify as ‘severe hyperglycemia’ in another. Similar heterogeneity also applies to the Leuven Surgical Trial and NICE-SUGAR trial, two landmark studies on critically ill patients that shaped inpatient guidelines. The glycemic target comparisons are slightly different, particularly in the ‘conventional’ glycemic groups, with the Leuven patients having a target of 180–200 mg/dL and NICE-SUGAR having a target of <180 mg/dL. Additionally, no large randomized controlled trials have examined non-critically ill patients (which comprise the vast majority of hospitalized patients) nor an inpatient glycemic target of 110–140 mg/dL.

Lastly, as this systematic review and meta-analysis evaluated CAP outcomes prior to COVID-19, we must examine the newest data derived from the CAP studies conducted during the pandemic. Though not a universal finding, it appears that the majority of studies reporting on COVID-19 pneumonia and glycemic control noted a significant association between hyperglycemia and poor hospital outcomes,^{10–12} similar to the findings by Barmanray *et al*. A potential factor introducing bias is that patients with more severe COVID-19 pneumonia likely received glucocorticoids, which may in turn have worsened glycemic control. With regard to the studies looking at the impact of underlying diabetes, there seems to be a consistent association with the presence of diabetes, independent of glycemic control, portending an increased risk of severe COVID-19 pneumonia and death.^{13–15} This is in contrast to the findings of Barmanray *et al* that diabetes alone was not associated with CAP mortality or ICU admission in the pre-COVID-19 era.

We have gained new insights on diabetes, glycemic control, and outcomes in hospitalized patients with pneumonia during the COVID-19 pandemic. The systematic review and meta-analysis on pre-COVID-19 CAP and glycemic control highlights both the extent and the limitations of our knowledge. Furthermore, prospective studies on alternate glycemic targets, such as 110–140 mg/dL, can improve the precision and confidence of our consensus guidelines. We hope the manuscript by Barmanray *et al*, as well as the scientific community’s recent experience with the COVID-19 pandemic, further catalyzes clinical research in inpatient diabetes management.

Contributors PBM drafted and revised the manuscript. ARG revised and edited the manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Disclaimer Opinions expressed in this paper are those of the authors and do not represent the views of the Department of Veterans Affairs or the US Government.

Competing interests ARG is an employee of the US Department of Veterans Affairs. ARG serves as an Associate Editor of the *BMJ Open Diabetes Research & Care*.

Patient consent for publication Not required.

Ethics approval Not applicable.

Provenance and peer review Commissioned; internally peer reviewed.

Data availability statement Data sharing not applicable as no datasets generated and/or analyzed for this study.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Paras B Mehta <http://orcid.org/0000-0001-5434-5424>

Aidar R Gosmanov <http://orcid.org/0000-0002-4197-5344>

REFERENCES

- 1 NICE-SUGAR Study Investigators, Finfer S, Chittock DR, *et al*. Intensive versus conventional glucose control in critically ill patients. *N Engl J Med* 2009;360:1283–97.
- 2 van den Berghe G, Wouters P, Weekers F, *et al*. Intensive insulin therapy in critically ill patients. *N Engl J Med* 2001;345:1359–67.
- 3 Rhodes A, Evans LE, Alhazzani W, *et al*. Surviving sepsis campaign: international guidelines for management of sepsis and septic shock: 2016. *Intensive Care Med* 2017;43:304–77.
- 4 Umpierrez GE, Hellman R, Korytkowski MT, *et al*. Management of hyperglycemia in hospitalized patients in non-critical care setting: an endocrine Society clinical practice guideline. *J Clin Endocrinol Metab* 2012;97:16–38.
- 5 Barmanray RD, Cheuk N, Fourlanos S. In-Hospital hyperglycemia but not diabetes mellitus alone is associated with increased in-hospital mortality in community-acquired pneumonia (CAP): a systematic review and meta-analysis of observational studies prior to COVID-19. *BMJ Open Diabetes Res Care* 2022;10.
- 6 Murad MH, Coburn JA, Coto-Yglesias F, *et al*. Glycemic control in non-critically ill hospitalized patients: a systematic review and meta-analysis. *J Clin Endocrinol Metab* 2012;97:49–58.
- 7 Falciglia M, Freyberg RW, Almenoff PL, *et al*. Hyperglycemia-related mortality in critically ill patients varies with admission diagnosis. *Crit Care Med* 2009;37:3001–9.
- 8 Farokhi F, Smiley D, Umpierrez GE. Glycemic control in non-diabetic critically ill patients. *Best Pract Res Clin Endocrinol Metab* 2011;25:813–24.
- 9 Umpierrez GE, Isaacs SD, Bazargan N, *et al*. Hyperglycemia: an independent marker of in-hospital mortality in patients with undiagnosed diabetes. *J Clin Endocrinol Metab* 2002;87:978–82.
- 10 Klonoff DC, Messler JC, Umpierrez GE, *et al*. Association between achieving inpatient glycemic control and clinical outcomes in hospitalized patients with COVID-19: a multicenter, retrospective hospital-based analysis. *Diabetes Care* 2021;44:578–85.
- 11 Mehta PB, Kohn MA, Koliwad SK, *et al*. Lack of association between either outpatient or inpatient glycemic control and COVID-19 illness severity or mortality in patients with diabetes. *BMJ Open Diab Res Care* 2021;9:e002203.
- 12 Saand AR, Flores M, Kewan T, *et al*. Does inpatient hyperglycemia predict a worse outcome in COVID-19 intensive care unit patients? *J Diabetes* 2021;13:253–60.
- 13 Barron E, Bakhai C, Kar P, *et al*. Associations of type 1 and type 2 diabetes with COVID-19-related mortality in England: a whole-population study. *Lancet Diabetes Endocrinol* 2020;8:813–22.
- 14 Izz-Engbeaya C, Distaso W, Amin A, *et al*. Adverse outcomes in COVID-19 and diabetes: a retrospective cohort study from three London teaching hospitals. *BMJ Open Diabetes Res Care* 2021;9:e001858.
- 15 Shang J, Wang Q, Zhang H, *et al*. The relationship between diabetes mellitus and COVID-19 prognosis: a retrospective cohort study in Wuhan, China. *Am J Med* 2021;134:e6–14.