



HHS Public Access

Author manuscript

Evid Based Pract Child Adolesc Ment Health. Author manuscript; available in PMC 2023 March 23.

Published in final edited form as:

Evid Based Pract Child Adolesc Ment Health. 2022 ; 7(2): 176–197.

doi:10.1080/23794925.2022.2042874.

Equity in Mental Health Services for Youth at Clinical High Risk for Psychosis: Considering Marginalized Identities and Stressors

Joseph S. DeLuca¹, Derek M. Novacek^{2,3}, Laura H. Adery³, Shaynna N. Herrera¹, Yulia Landa^{1,4}, Cheryl M. Corcoran^{1,4}, Elaine F. Walker^{5,6}

¹Department of Psychiatry, Icahn School of Medicine at Mount Sinai, , New York, NY, USA

²Desert Pacific Mental Illness Research, Education, and Clinical Center, Veterans Affairs Greater Los Angeles Healthcare System, , Los Angeles, CA, USA

³Department of Psychiatry and Biobehavioral Sciences, University of California, Los Angeles, CA, USA

⁴New York Mental Illness Research, Education and Clinical Center, James J. Peters Veterans Affairs Medical Center, Bronx, NY, USA

⁵Department of Psychology, Emory University, Atlanta, GA, USA

⁶Department of Psychiatry and Behavioral Sciences, Emory University School of Medicine, Atlanta, GA, USA

Abstract

Prevention and early intervention programs have been initiated worldwide to serve youth at Clinical High Risk for Psychosis (CHR-P), who are adolescents and young adults experiencing subclinical psychosis and functional impairment. The primary goals of these efforts are to prevent or mitigate the onset of clinical psychosis, while also treating comorbid issues. It is important to consider issues of diversity, equity, and inclusion in CHR-P work, especially as these programs continue to proliferate around the world. Further, there is a long history in psychiatry of misdiagnosing and mistreating psychosis in individuals from racial and ethnic minority groups. Although there have been significant developments in early intervention psychosis work, there is evidence that marginalized groups are underserved by current CHR-P screening and intervention efforts. These issues are compounded by the contexts of continued social marginalization and significant mental health disparities in general child/adolescent services. Within this narrative review and call to action, we use an intersectional and minority stress lens to review and discuss current issues related to equity in CHR-P services, offer evidence-based recommendations, and propose next steps. In particular, our intersectional and minority stress lenses incorporate perspectives for a range of marginalized and underserved identities related to race, ethnicity, and culture; faith; immigration status; geography/residence; gender identity; sexual orientation; socioeconomic status/class; and ability status.

Correspondence concerning this article should be addressed to Joseph S. DeLuca, Department of Psychiatry, 1399 Park Ave, 3rd Floor, New York NY, 10029. joseph.deluca@mssm.edu.

We have no known conflict of interest to disclose.

Keywords

prodrome; equity; inclusion; intersectionality; stigma

There have been worldwide efforts to identify and treat youth at Clinical High Risk for Psychosis (CHR-P), a heterogeneous group of youth who experience distressing attenuated psychosis symptoms (e.g., subthreshold hallucinations, delusions, disorganization) (Catalan, Salazar de Pablo, et al., 2020; Fusar-Poli, Salazar de Pablo, et al., 2020; Kotlicka-Antczak et al., 2020). Compared to other youth clinical groups, individuals at CHR-P are at a heightened risk for developing a psychotic disorder in the future and often experience distressing comorbid psychopathology (Addington et al., 2017; Lim et al., 2015), as well as functional impairment (Fusar-Poli et al., 2013). Current estimates suggest that approximately 25% of youth at CHR-P go on to develop threshold psychosis within three years (Catalan, Salazar de Pablo, et al., 2020; Fusar-Poli, Salazar de Pablo, et al., 2020), with non-converters still demonstrating clinically significant difficulties that warrant treatment (Addington et al., 2011; Beck et al., 2019). CHR-P programs typically provide transdiagnostic treatment to meet the needs of these youth and their families, and the goals of these programs generally include preventing or delaying the transition to threshold psychosis by reducing distress and building coping skills (Thompson et al., 2015). To fully realize the goals of programs such as these, it is imperative to have equitable and culturally sensitive research and services that identify and treat diverse youth and families. Given the association between psychosis and social determinants (Anglin et al., 2020), as well as historical abuse and racial disparities related to psychosis conceptualizations and diagnoses (Loring & Powell, 1988; Metzl, 2010; Nazroo et al., 2020; Schwartz & Blankenship, 2014), it is especially important to consider issues of diversity, equity, and inclusion in CHR-P work.

Contexts

Research and Policy Context

CHR-P programs are proliferating worldwide (Catalan, Salazar de Pablo, et al., 2020; Fusar-Poli et al., 2017; Kotlicka-Antczak et al., 2020). Just in the United States (US), the Substance Abuse and Mental Health Services Administration (SAMHSA, 2018) recently funded more than 20 CHR-P low-cost treatment programs in the past three years, and the National Institute of Mental Health (NIMH, 2020) recently partnered with multiple worldwide agencies to support a collaborative effort to advance CHR-P research. This large-scale NIMH project, the *Accelerating Medicines Partnership for those at risk of developing Schizophrenia*, will include research on biological and psychosocial factors related to disease progression, with aims of better understanding CHR-P trajectories and informing treatments. Additionally, worldwide and cross-cultural CHR-P work is ongoing elsewhere, including work focused on improving detection, identifying biological correlates, and implementing services (Zhang et al., 2020).

Broader Social Context

These promising services and research endeavors come at a time when social inequities have become amplified for many marginalized groups due to the global COVID-19 pandemic and rising racially and ethnically motivated terrorism (e.g., white supremacist terrorism) (Laster Pirtle, 2020; State, 2019). Residual health impacts are expected for racial and ethnic minority (REM) groups (Novacek et al., 2020), and the effects of the current social context have also already negatively impacted—or have been predicted to impact—people at CHR-P or those experiencing threshold psychosis or other serious mental illness (Cowan, 2020; DeLuca et al., 2020; Nemani et al., 2021). These issues are compounded by the context of significant mental health disparities in general child/adolescent health services (Merikangas et al., 2011; Whitney & Peterson, 2019), which may prevent or delay the identification of young people at CHR-P since many CHR-P referrals come from general outpatient and inpatient psychiatric departments (Cornblatt, 2002; Fusar-Poli et al., 2016).

Social Inequities and CHR-P Experiences

Research has documented that social inequities and CHR-P experiences are intertwined (Anglin et al., 2020). For example, high neighborhood crime has been related to high suspiciousness among individuals at CHR-P (Vargas et al., 2020; Wilson et al., 2016). Psychotic experiences, which conceptually overlap with the CHR-P state, have also been associated with police victimization (DeVylder et al., 2017), racial discrimination (Anglin et al., 2014; Oh et al., 2016), other trauma exposure (Mayo et al., 2017; Redman et al., 2017), and urbanicity and air pollution exposure (Newbury et al., 2019). In general, REM individuals have been found to endorse higher levels of subclinical psychosis experiences (Calkins et al., 2014; Morgan et al., 2009; Wigman et al., 2011), as well as potentially unique symptom presentations that intersect with discrimination and trauma (Anglin et al., 2015; Thompson et al., 2009). More broadly, components of inequality (e.g., income inequality) have been associated with a higher schizophrenia incidence worldwide (Boydell et al., 2004; Burns et al., 2014). Further, lack of involvement of REM youth in early intervention CHR-P programs may have negative long-term, downstream effects, such as delayed mental health diagnosis and disproportionate criminal justice system involvement (Al-Rousan et al., 2017; Youman et al., 2010).

Successes and Limitations of Current CHR-P Programs

These findings suggest that there is likely an even greater need for psychosis prevention work and specialized CHR-P research programs and services among marginalized groups. Notwithstanding the major successes of CHR-P programs (i.e., worldwide hubs of research and clinical experts who improve the lives of youth and families) (McGorry et al., 2021), youth may be hesitant to seek and engage in CHR-P services (Ben-David et al., 2019a; Ben-David et al., 2019b; Leanza et al., 2020) and there is evidence that marginalized groups are underserved by current early-stage psychosis prevention screening, intervention, and research efforts (Anglin et al., 2020; Deriu et al., 2018; Oluwoye et al., 2021). This raises major questions regarding outreach to, and identification of, marginalized groups for psychosis prevention. In particular, although the reliability and validity of semi-structured

assessments have been found to be generally consistent across countries, more research is needed overall (Woods et al., 2019), including research on cultural group differences in the same country. To this end, mixed results have been found regarding the precision of common CHR-P screening tools for REM youth in the US (Cicero et al., 2019; Karcher et al., 2018; Lång et al., 2020; Millman et al., 2019; Rakhshan Rouhakhtar et al., 2019), which may contribute to inequity since screening tools are generally the first line of assessment to determine potential psychosis risk and further inclusion in research studies and services.

Research has also demonstrated that REM youth are less likely to visit specialized psychosis services (compared to non-Hispanic White youth) before their first presentation of psychosis (Coleman et al., 2019), suggesting that culturally sensitive outreach to these groups may be particularly warranted (Lynch et al., 2016). Additionally, research on first-episode psychosis (FEP) service use—which the CHR-P state may be a precursor to—has also demonstrated racial disparities, namely that Black youth (compared to White youth) are more likely to experience homelessness and housing insecurity, to have a significantly poorer quality of life, exhibit more severe symptoms, and are less likely to have private insurance (Nagendra et al., 2018). Non-Hispanic Black youth in FEP programs have also been found to be less likely to receive important early intervention services, such as family psychoeducation (Oluwoye et al., 2018). Overall, results from FEP studies on disparities strengthen the premise that increasing equity in CHR-P services may be one upstream method to reduce disparities in FEP programs.

Our Framework

The framework of this narrative review and call to action is based on minority stress models (Brooks, 1981; Meyer, 2003) and intersectional models (Crenshaw, 1989; Seng et al., 2012) of mental health, and thus acknowledges that understanding and focusing on stressors affecting marginalized groups and those of oppressed identities are paramount to increasing equity in CHR-P services. For presentation clarity, we describe various identities separately below while acknowledging that intersectionality is a process by which multiple oppressed identities *interact* and *combine* to increase inequity and stress. We note some of these interactions and nuances within each section, as well as in our Discussion section. The intersectional identities below were chosen based on foundational past minority stress and intersectional research (Crenshaw, 1990; Hays, 1996), including emerging research with individuals at CHR-P (Ruiz et al., 2021; Saleem et al., 2014). These identities include developmental stage and associated factors; race, ethnicity, and culture; faith; immigration status; geography/residence; gender identity; sexual orientation; socioeconomic status/class; and ability status. We tie together these identities in a concluding section on minority stress and trauma, acknowledging the high prevalence and confluence of various stressors that further perpetuate disparities.

Method

We adopted a narrative review and call to action approach for this work, similar to other recently published early-stage psychosis articles focused on exploring distinct literatures and generating empirically-based recommendations (Brooke et al., 2019; Harrop et al.,

2015). Building on our interest in reviewing a broad range of CHR-P equity issues using an intersectional and minority stress lens, we believed that a narrative review and call to action approach would be most appropriate. Some authors have referred to this approach as a critical review that allows one to “take stock” of and synthesize broad work in the field, while fostering “conceptual innovation” to help move the field forward (Brooke et al., 2019; Grant & Booth, 2009). Our search terms included CHR-P and related terms, terms related to the various identities discussed above (Crenshaw, 1990; Hays, 1996), and terms including and related to diversity, equity, and inclusion. Snowballing techniques were used by searching the reference lists of relevant articles. All searches were conducted in February and March 2021.

Below, we provide the results of our narrative review organized by intersectional identity, with a concluding section on the overlap of trauma and stressful life experiences across these identities. We then provide our recommendations and “call to action” based on this review, followed by a discussion and conclusion.

Results

Developmental Stage and Associated Factors

As noted, CHR-P experiences typically first occur during adolescence and young adulthood, which are key developmental periods. During this time, youth are contending with significant psychological, social, and biological/physical changes. Youth at CHR-P present to different clinics (e.g., child/pediatric v. adult) and with different presenting concerns and current life problems, depending on their age and developmental stage. For example, research has found that younger youth (~age 13 and younger) endorse more items on psychosis-risk screening tools compared to older youth (Rouhakhtar et al., 2019), which may increase the false-positive rate for this group. Nonetheless, early identification of true psychotic experiences among younger teens (~age 11-12) may signal vulnerability for future psychopathology (Rimvall et al., 2020). In general, researchers have called for a careful evaluation of psychosis-risk experiences in the context of normative developmental experiences (Carol & Mittal, 2015; Schiffman et al., 2019). In regard to developmental milestones, psychosis risk experiences have also been positively associated with developmental delays in speech and motor skills (Karcher et al., 2018), as well as various, early neurocognitive delays (Gur et al., 2014), highlighting the importance of a detailed intake and collaboration with collaterals (e.g., caregivers) for historical information.

From a psychological perspective, adolescence and young adulthood are key periods of identity development (Erikson, 1968), and researchers are only just beginning to understand the potentially unique process of identity development for youth at CHR-P (Ben-David & Kealy, 2020; Sarac et al., 2021). In terms of cognitive development, researchers have also shed light on cognitive biases that may be more prevalent among youth at CHR-P, such as jumping to conclusions (Catalan, Tognin, et al., 2020). These factors can directly inform assessment and treatment strategies.

From a biological and neurobiological perspective, several developmental mechanisms have also been proposed, including the role of hormones in modulating brain development for

individuals at CHR-P (Corcoran et al., 2003; Damme et al., 2020; Trotman et al., 2013; Walker, Trotman, Goulding, et al., 2013). One recent study (Damme et al., 2020) reported that early exposure to estrogens (i.e., earlier age at menarche) may have a protective impact on hippocampal connectivity for youth at CHR-P, highlighting the importance of considering developmental factors during evaluation, treatment, and research. As will be noted in a later section, this finding may also have implications for equity in early psychosis services for women, a group that also tends to have a later onset of psychotic experiences (Ferrara & Srihari, 2021). Lastly, biological changes also trigger physical changes for youth, and youth at CHR-P have been found to perceive more discrimination related to their “appearance” than a non-clinical sample (Saleem et al., 2014).

From a social-contextual perspective, most youth at CHR-P live with their families (Addington et al., 2015), and studies have found that stronger youth-reported family functioning (e.g., cohesion, support) may be a protective factor for youth at CHR-P (Izon et al., 2018; Izon et al., 2021; Thompson et al., 2019; Wüsten & Lincoln, 2017). Further, poorer caregiver-reported family functioning has been associated with higher perceived discrimination experiences among youth with psychosis-spectrum symptoms (DeLuca et al., 2021). Research has also found that stigma among families may be an important variable to consider for youth at CHR-P (Colizzi et al., 2020; Wong et al., 2009). In particular, one of these studies (Wong et al., 2009) found that only REM families (of individuals with recent, full psychosis) endorsed feeling shame and a need to conceal their family member’s illness. Thus, outreach and interventions focused on engaging families of youth at CHR-P (particularly REM families) and building coping skills in a culturally sensitive manner appear to be important endeavors (Gerson et al., 2011; Santesteban-Echarri et al., 2018). Further, given the high rate of general social functioning deficits among youth at CHR-P (Cornblatt et al., 2007; Velthorst et al., 2010), as well as high rates of childhood bullying victimization (Mayo et al., 2017), multipronged and transdiagnostic efforts focused on these factors may also be warranted.

Race, Ethnicity, and Culture

From the prenatal period through childhood, it is well-established that the rate of exposure to many of the risk factors for psychosis (e.g., poor prenatal care, trauma, stress, discrimination) is much higher for REM than White youth (Dunkel Schetter et al., 2013). Despite this, REM youth are less likely to receive mental health services compared to White youth (Kataoka et al., 2002). In one of the largest studies conducted to date on adolescent mental health service utilization (Merikangas et al., 2011)—using a US national sample of over 6,000 adolescents and structured interviews to determine mental health diagnoses—it was found that Hispanic and non-Hispanic Black adolescents were significantly less likely to receive services (compared to White adolescents), specifically for mood and anxiety disorders. This is particularly notable in regard to equity in CHR-P services, because anxiety and mood disorders often precede or co-occur with attenuated psychotic symptoms among youth at CHR-P (Addington et al., 2017; Lim et al., 2015) and symptoms related to these disorders may be the first presenting problems for youth considering treatment.

The barriers to care that REM youth experience are often structural and a product of marginalization. REM youth are more likely to be uninsured or underinsured, thus creating financial obstacles to accessing needed mental health services (Kataoka et al., 2002). However, barriers to mental health services are not solely financial; a recent study with youth experiencing FEP found that racial/ethnic disparities in diagnosis and treatment may be present even for insured youth and after controlling for household income (Heun-Johnson et al., 2021).

There are also significant cultural differences in how REM individuals perceive mental health concerns, pathways to mental health services, and help-seeking preferences (Cauce et al., 2002). REM families may have different interpretive frameworks for mental health experiences, such as spiritual or religious explanations (Esterberg & Compton, 2006) that can impact CHR-P help-seeking. REM individuals often seek out non-medical support to deal with emotional concerns such as pastors or religious clergy, family, and friends (Cauce et al., 2002; Novacek et al., 2019). In addition, REM youth often experience aversive pathways to mental health through justice system involvement and involuntary hospital admissions (Compton et al., 2006). REM youth may also have difficulty accessing mental health care services due to the lack of culturally competent providers in their area. In addition, many CHR-P clinics are housed within academic medical centers that often do not serve REM youth. Among Black Americans, there is a skepticism or cultural mistrust of healthcare including mental health services due to medical racism (Whaley, 2001a, 2001b). Given that many CHR-P services are housed within academic medical centers, this distrust, along with “ivory tower” and inaccessibility perceptions, could prevent youth from seeking care (Lynch et al., 2016). Research also demonstrates that REM individuals often hold more stigmatizing beliefs about mental illness and mental health treatment compared to White Americans (Brown et al., 2010; DeFreitas et al., 2018). These stigmatizing attitudes could deter REM youth from seeking and adhering to mental health services.

Faith

Faith, spirituality, and religious beliefs often serve as an integral coping strategy and as protective factors against suicide (Koenig, 2009). However, a double-edge sword is possible in that religious interpretations of legitimate attenuated psychotic experiences could delay treatment seeking. On the other end, clinicians must be cautious of pathologizing religious beliefs without considering how normative the experiences are within an individual’s faith community (O’Connor & Vandenberg; Pierre, 2001). Very little research to date has been conducted on faith and CHR-P, but emerging research in this area suggests that there may be a relation between religiosity and depressive symptomology (Severaid et al., 2019) and religiosity and perceptual abnormalities/hallucinations (Loch et al., 2019) for youth at CHR-P. Loch et al. (2019) also found relations between religiosity and other attenuated experiences (ideational richness, suspiciousness) that require further research. In an earlier study, it was also found that families of patients at CHR-P or in the early stages of a psychotic disorder used religious coping (i.e., seeking God’s help, trying to find comfort in religion, praying) at a moderate level (Gerson et al., 2011), highlighting the importance of considering broader family coping strategies.

Immigration Status

An individual's status as an immigrant may also play an important role in their risk for psychosis and decision or ability to engage in CHR-P services. Research has consistently shown that the incidence of psychosis is higher among refugee and nonrefugee migrant populations compared to native populations (Brandt et al., 2019; Cantor-Graae & Selten, 2005). Research also shows that increased incidence of psychosis for second-generation migrants is comparable to first-generation migrants (Bourque et al., 2011; Weiser et al., 2008), suggesting potential intergenerational transmission of trauma related to migration. It has been suggested that this risk for second-generation migrants may reflect stress within the new country (e.g., discrimination, social exclusion) rather than migration-specific stressors (Henssler et al., 2020), but this effect for second-generation immigrants has not been found in some studies (Corcoran et al., 2009b).

Although there is a wealth of research demonstrating increased risk for psychosis among migrants, there is minimal research on CHR-P migrants. Thus far, researchers have not demonstrated an association between migrant status and transition to psychosis among individuals at CHR-P (Geros et al., 2020; O'Donoghue et al., 2021). However, studies show that duration of untreated psychosis is significantly longer in migrants (Boonstra et al., 2012), indicating that migrants are not being identified early on in the CHR-P state and are thus less likely to receive the benefits of early intervention.

The proposed pathways by which migration may contribute to increased risk for psychosis emphasizes the compounding effect of post-migratory stress, trauma-related factors, social isolation, difficulties re-entering society, stigma/discrimination, family dysfunction, poverty, and poor housing conditions (Henssler et al., 2020; Veling & Susser, 2011). Individuals at CHR-P with a migration history have also shown more severe symptoms, including significantly more formal thought disturbances and sleeping problems compared to those without a migration history (Nogueira et al., 2020). In addition to social and societal challenges, neurochemical changes in the brain of CHR-P migrants may increase risk of psychotic symptoms. In independent CHR-P samples from the United Kingdom and Canada, striatal dopamine function was elevated in immigrants compared with natives (Egerton et al., 2017), which is consistent with findings among individuals who have already developed a psychotic disorder (Henssler et al., 2020). It has been suggested that these neurochemical changes may be caused by psychosocial factors such as victimization, discrimination, social exclusion, and other social and environmental factors (Egerton et al., 2017).

In a study of providers' impressions of the disadvantages experienced by minority groups in early intervention for psychosis services, providers noted higher rates of family disengagement among recent immigrants and those with a history of marginalization (Jones, Kamens, et al., 2021b). Providers indicated that this may be due to limited knowledge about mental health system and lack of resources. The location of CHR-P programs also matters for migrants. In a prospective population-based study, low neighborhood ethnic density (proportion of ethnic minorities in an area) at age 15 was associated with increased incidence of psychosis for migrants, even after controlling for urbanicity, parental SES, and parental psychiatric history (Schofield et al., 2017). This suggests that living in a higher

ethnically dense neighborhood may reduce social stress, and thus highlights the importance of moving care into migrant neighborhoods.

Stigma may present another barrier to mental health treatment for migrants. There is evidence to suggest that the incidence of psychosis is higher when immigrant groups perceive more racial/ethnic discrimination (Veling et al., 2007), and that discrimination may serve as a mediator for psychotic symptom severity in immigrants (Berg et al., 2011). However, unlike studies in psychosis, one study using a CHR-P sample did *not* find associations between immigration status and perceived discrimination (Saleem et al., 2014). More research is needed to understand stigma among youth at CHR-P with a personal or family history of migration.

Geography/Residence

Epidemiological and cohort studies reliably demonstrate that exposure to urban environments is associated with increased risk for psychosis. Risk increases as level of urban exposure increases in a dose-response manner (Pedersen & Mortensen, 2001a), leading many to conclude that urbanicity is an environmental cause of psychosis (Krabbendam & van Os, 2005). This finding is consistent across varied definitions of urbanicity, and holds up strongly when assessing for other important demographic and environmental factors such as age, sex, ethnicity, substance use, social class, genetic risk, distance to hospital, use of mental health services, and countless others (Krabbendam & van Os, 2005; Lewis et al., 1992; McGrath et al., 2004; Pedersen & Mortensen, 2001b; Spauwen et al., 2004; Stefanis et al., 2004). However, this effect may not be present in developing countries (DeVylder et al., 2018).

Urbanicity has been associated with both positive and negative symptoms of psychosis (van Os et al., 2002), and with psychosis-like symptoms, which are more prevalent in the population than psychotic disorders (van Os et al., 2001). The relationship between urbanicity and incidence for schizophrenia suggests some unique environmental impact of urban areas that likely exerts its influence during child and adolescent development rather than close to illness onset (Marcelis et al., 1999; Pedersen & Mortensen, 2001a). What this means for CHR-P youth is that the earlier that psychosis risk symptoms can be identified and treated, the more likely it is that providers can impart protective factors that counter such environmental risk factors.

Research specifically on individuals in the early stages of psychosis or at CHR-P demonstrates the potential adverse impact of urbanicity on important outcomes. In a prospective study of Individuals at CHR-P, urbanicity was among the predictors for transition to psychosis (Dragt et al., 2011). In a study of FEP patients from rural and urban areas, duration of untreated psychosis was significantly longer for individuals from highly urbanized areas (and first-generation immigrants, as noted above) (Boonstra et al., 2012).

Aspects of rural living may also impact risk for psychosis. In a study looking only within rural environments (e.g., consisting of dispersed farms, villages, small towns), higher prevalence of psychotic disorder was associated with material deprivation (i.e., a single measure that combines information regarding unemployment, housing, car ownership, etc.)

(Omer et al., 2014). Similarly, in a population-based study of FEP individuals in rural environments, there was a higher incidence of psychosis in more deprived and socially isolated areas (Richardson et al., 2018). Rural youth at CHR-P also have different help-seeking behaviors compared to urban youth and have greater functional impairment (Stain et al., 2018).

For individuals at CHR-P, social isolation and material deprivation (e.g., lack of car, lack of internet access in the home) may present barriers to seeking and accessing care during the onset of psychotic like symptoms, and indeed one study showed a trend for individuals at CHR-P to reside in relatively more deprived areas (O'Donoghue et al., 2015). Rural youth often must travel longer distances to receive care, and there is less access to specialized CHR-P programs in rural compared to urban areas (Stain et al., 2018). Overall, there is minimal research regarding the specific needs of rural youth and service delivery in rural areas (Welch & Welch, 2007). Lastly, beyond rural and urban environments, there is evidence in the US that there are major quality and racial-ethnic disparities in psychosis-spectrum care depending on which county you live in (Horvitz-Lennon et al., 2015).

Gender Identity

In discussing what is known regarding the role of gender identity in CHR-P, it is important to note that much of our understanding of these risk states and of the illness and treatment of schizophrenia is based on research conducted disproportionately with men and virtually all within binary gender categories or by biological sex, which conflates gender and sex and does not as readily account for environmental factors that may influence people of different genders quite differently. Indeed, meta-analyses of epidemiological and non-epidemiological research conclude that while expected epidemiological incidence rate was 58% male, on average, males comprise over two-thirds of study samples of schizophrenia (Longenecker et al., 2010). While CHR-P recruitment often involves rigorous case finding, meta-analyses on gender differences in symptoms and functioning similarly include mostly males (Tor et al., 2018), with only about one-third of participants categorized as binary female (Rietschel et al., 2017). In light of this, our understanding of etiology, symptom expression and associations, and resulting treatment recommendations may not fully reflect or apply to the experiences of CHR-P youth who do not identify as male-gendered.

Further, the overproportion of identified male-gendered participants may indicate gendered barriers, such as under-identification by referral pathways due to schema, to entrance to treatment settings where such clinical research is conducted. Indeed, compared with men, literature suggests that women tend to present with more mood symptoms which may lead to misdiagnosis of psychosis due to primary affective disorder, often an exclusion criterion for research-based FEP services (Heitz et al., 2019; Ochoa et al., 2012). Additionally, CHR-P clinics may narrow eligible age range to transition-aged-youth, excluding individuals whose biological female sex is associated with later onset (median first contact age 34 for female vs 28 for men) (Ferrara & Srihari, 2021). No gender differences have been found in conversion rates of CHR-P youth across longitudinal follow up (Walder et al., 2013), or within epidemiological studies (Scott et al., 2006). Delays in receiving care due to

misdiagnosis or missed identification may lead to longer duration of untreated psychosis, a predictor for poorer outcomes.

There may also be strengths associated with the differential gender socialization of individuals. For example, it may be more acceptable for individuals socialized as women to disclose experiences, seek help (Mackenzie et al., 2006), or rely on social networks and support (Willhite et al., 2008). Poorer baseline social functioning and positive prodromal symptoms have been shown to predict greater conversion risk in men (Walder et al., 2013).

There remains a scarcity of literature on clinical care for transgender and gender expansive populations with serious mental illness (Barr et al., 2021; Meijer et al., 2017; Mizock & Fleming, 2011). They may avoid participation in such environments due to perceived burden to educate (Lucksted, 2004), transphobic rejection (Rossman et al., 2017; Smith et al., 2019), and historically founded concerns of misjudgment of gender identity as a symptom of psychosis (Hoening & Kenna, 1974) or intervention targets (Mizock, 2008). Preliminary evidence from recent initial explorations demonstrates that many gender-expansive youth do not disclose their gender identity to all of their health care providers, though positive associations are found between disclosure and self-esteem (McKay & Watson, 2020). Previous work demonstrates disclosure as an important protective factor from all psychiatric disorders (Dhejne et al., 2016). Social signaling, such as use of binary categories of gender within care settings and failure to glean specific knowledge of various genders within research explorations or clinic intake, may further this chasm within CHR-P populations.

Given markedly greater relative risk demonstrated for comorbid depression, anxiety and suicidal ideation among gender expansive compared to cisgender youth (Day et al., 2017; Reisner et al., 2015), targeted treatment may be particularly effective, as reduction of mood symptoms alone has demonstrated improvement in CHR-P prognosis (Thompson et al., 2015; van der Gaag et al., 2013). Further, adolescent and transition-age individuals who present to CHR-P clinics may be in the process of discovering gender identity at the time of presentation or may be fluid in presentation over time. As disruption and loss of sense of self are increasingly implicated in schizophrenia spectrum experiences (Ferri et al., 2014), affirmation of presenting identity may be particularly crucial for youth at risk for psychosis in self-integration to reduce a potential predisposition that incongruent treatment may exacerbate.

Sexual Orientation

Not to be conflated with gender identity, consideration of sexual orientation may be an additional important facet in conceptualization and treatment for individuals at CHR-P. Large scale studies indicate an up to three-fold increased risk of psychotic experiences for queer as compared to heterosexual individuals, that may be partially mediated by experience of discrimination, stress, and lack of social support related to sexual orientation (Bolton & Sareen, 2011; Chakraborty et al., 2011; Gevonden et al., 2014; Jacob et al., 2021). Most recently, when accounting for additional sociodemographic factors including sex, age, and ethnicity, a two-fold risk remained (Jacob et al., 2021). Particularly, loneliness has been robustly associated with increased risk of psychosis (Michalska da Rocha et al., 2018) and endorsed commonly in addition to lack of social support amongst queer individuals

(Doyle & Molix, 2016). Some evidence suggests that particular psychotic experiences such as paranoia or mania, rather than hallucinations, may be more prominent for LGBT+ individuals (Jacob et al., 2021; Janssen et al., 2003), supporting a hypothesis that adversity may have specific effects on psychotic experiences which differ in etiology and may not otherwise develop.

Data demonstrates a growing imperative for attention and care to sexual orientation diversity. In 2020, an estimated 1 in 6 transition-aged youth surveyed in the United States identified as LGBT+ (Gallup, 2020), continuing a rising trend in such identification since tracking began in 2012, without accounting for response-bias due to stigma. Estimates of US adolescent populations indicate greater variance in orientation, with about 20-25% of youth in grades 9-12 identifying as LGBT in 2019, according to the CDC (Health, 2019). However, only 1 in 3 LGBT+ adolescents report disclosure of orientation to medical providers due to realistic concerns of stigma, discrimination, or lack of inquiry from providers (Rossman et al., 2017).

Socioeconomic Status/Class

Another social and environmental risk factor for psychosis that may contribute to inequity in CHR-P identification and treatment is socioeconomic status (SES). An individual's SES can be measured in a variety of ways, but is typically measured as household income or education level, as higher levels of education are associated with better economic outcomes and resources. Neighborhood level economic factors, such as neighborhood poverty level, neighborhood crime, or the ranking of a school system, have also been assessed as a proxy for SES. In a systematic review and meta-analysis, lower SES was significantly associated with higher incidence of psychosis (Castillejos et al., 2018) and income inequality has also been associated with increased incidence of schizophrenia (Burns et al., 2014). The impact of SES on incidence of psychosis has been found to be most detrimental for offspring of fathers in the lowest social class (i.e., paternal occupation at time of birth) (Corcoran et al., 2009a). There is also the risk of low SES driving public stigma toward the individual. In an online study of 1,348 adult volunteers, both "uneducated people" and "mentally ill people" elicited negative appraisals of incompetence and abnormality, suggesting a strong bias against individuals with lower formal education and SES (Phelan et al., 2019).

Ability Status

An important consideration regarding equity and access to CHR-P services includes ability status. Ability status refers to differences in intellectual, physical, social, or emotional domains that may or may not present visibly (e.g., use of a wheelchair, speech impediment, major psychological or neurological condition, etc.), and which may be temporary or permanent and dependent on context (Nielsen, 2012). Ableism continues to display and detract from mental health care, through lack of attempts toward inclusion such as physically accessible space, organization and visualization of written documents, infrequent provision of close captioning and use of language depicting individuals as "high or low functioning" based on neurodivergence (Cornwall, 2020). As technology evolves, opportunities for access expand and must be creatively adapted.

Very few studies have been published in the psychosis field on ability status, and the majority of studies published on “disability” and psychosis-risk refer to psychosocial disability (e.g., greater difficulty making or maintaining friendships or joining community activities) (Olvet et al., 2015; Velthorst et al., 2010). Individuals at CHR-P are generally aware of such functional impairment and this awareness is associated with depression (Olvet et al., 2015). Neurocognitive deficits are also well-documented among youth at CHR-P and can directly impact functioning (Carrión et al., 2011). Overall, it is plausible that early intervention psychosis efforts may help to prevent long-term functional (e.g., social and role) disability.

One study has assessed physical disability among youth at CHR-P and found no association between physical difficulties (e.g., moving and getting around) and transition to psychosis (Velthorst et al., 2010). Another study assessed physical disability among a broader group of people with “psychotic experiences” (Oh et al., 2018) and found that individuals who reported a disability (including mobility disabilities) had a 1.5-3x higher chance of reporting lifetime psychotic experiences, even after adjusting for socio-demographics, lifetime mental illness, and lifetime chronic health conditions. In sum, more research is needed in this area, and it may be helpful to partner with other medical clinics/programs. Epidemiological studies, for example, have found that psychosis may be 2-3x higher among young individuals with multiple sclerosis (which can involve physical disability), compared to the general population (Patten et al., 2005). Other potential groups to be considered within this domain may include individuals with intellectual disabilities and individuals with Autism Spectrum Disorder (Cooper et al., 2007; Foss-Feig et al., 2019).

Minority Stress and Trauma

As noted, it is important to emphasize that trauma and stressors are common among individuals at youth at CHR-P (Bardol et al., 2020; Colizzi et al., 2020; Kraan et al., 2015; Loewy et al., 2019; Mayo et al., 2017; Redman et al., 2017; Stowkowy et al., 2016), particularly for those with marginalized identities (Anglin et al., 2021; Anglin et al., 2014; Anglin et al., 2018; Anglin et al., 2015; Saleem et al., 2014). Based on aforementioned minority stress models (Brooks, 1981; Meyer, 2003), individuals who possess minoritized identities face tremendous, often cumulative, stigma-related stressors and traumas, which confer risk for psychopathology. Thus, it is particularly important to be aware of how these factors impact youth at CHR-P.

Our review revealed that discrimination (toward various marginalized identities) is associated with psychotic symptoms (Janssen et al., 2003; Pearce et al., 2019) and is a significant predictor of the conversion to psychosis in CHR-P youth (Colizzi et al., 2020; Stowkowy et al., 2016). In addition to the aforementioned social inequalities, including studies on the positive association between neighborhood crime and CHR-P (e.g., Vargas et al., 2020; Wilson et al., 2016), community studies have also documented associations between direct exposure to community violence and psychotic experiences (Rakhshan Rouhakhtar et al., 2019). CHR-P studies have also shown that trauma exposure is significantly associated with severity of attenuated positive symptoms, specifically among REM patients (Thompson et al., 2009). Such stress and trauma has been found to “get

under the skin” of youth at CHR-P, with research demonstrating that greater exposure to deprivation, such as poverty and neglect, is associated with smaller cortical volume and smaller right hippocampal volume in a CHR-P sample (LoPilato et al., 2019). To this end, researchers have also proposed that the diathesis-stress model can be used to understand the impact of stressors on individuals at CHR-P, with the biological effects of stress being mediated by the hypothalamic-pituitary-adrenal axis (Corcoran et al., 2012; Corcoran et al., 2003; Walker, Trotman, Pearce, et al., 2013).

Call to Action: Recommendations

Our recommendations (see supplemental online material) are based on the narrative review conducted and include evidence-based suggestions based on empirical research and future directions guided by identified gaps in CHR-P work. We separated some of these recommendations by identity status to draw attention to specific concerns within categories, while also providing overarching recommendations to improve equity that apply across identity status. Though not an exhaustive list, we believe that these recommendations can increase awareness of equity issues in CHR-P work and, most importantly, contribute to the development of a more equitable early-stage psychosis system of care. We offer wide-ranging recommendations covering clinical care, research, training, outreach, and policy/systemic factors, some of which have been offered by leading scholars in the field (Anglin et al., 2021; Anglin et al., 2020; Carol & Mittal, 2015; Chapman, 2020; Cornwall, 2020; Deriu et al., 2018; Fusar-Poli, de Pablo, et al., 2020; Gara et al., 2012; Hall et al., 2021; Hankerson et al., 2018; Jones, Atterbury, et al., 2021; Jones, Kamens, et al., 2021a; Kline et al., 2019; Li, Friedman-Yakoobian, et al., 2019; Li, Shapiro, et al., 2019; McGorry et al., 2019; Meijer et al., 2017; Meyer et al., 2020; Mizock & Fleming, 2011; Novacek et al., 2019; Oluwoye et al., 2021; Ozer et al., 2020; Schiffman et al., 2019; Smith et al., 2019)¹ and some of which we have developed after synthesizing the findings of this narrative review and call to action.

Discussion

Worldwide, there continues to be a significant focus on identifying and providing treatment to youth at CHR-P and their families. To fully realize the goal of preventing or delaying the onset of psychosis, it is imperative to have accessible, equitable, and personalized services that address the needs of diverse youth and families. As noted, there is a long history of abuse and racism in psychiatry toward REM groups, culminating in understandable mistrust toward institutions and groups that provide CHR-P services and conduct CHR-P research. The current research, policy, and social climates make it clear that inclusive CHR-P services should be a priority. The work of current CHR-P programs highlights a major success in early intervention in psychiatry; CHR-P programs have proliferated over the past few decades and have provided much-needed services and expertise to youth, families, and professionals. At the same time, there are critical gaps identified by the literature regarding equitable care and there is room for improvement. Throughout this narrative review and call to action, we highlighted critical intersections in this work and provided recommendations

¹Although not exhaustive, several examples are provided here of published work with explicit recommendations, in addition to the many other articles referenced throughout this paper.

for future work in this area – some recommendations specific to an identity, and others more general. Practitioners must be ready to initiate and respond to such conversations throughout the treatment process, and to adapt to family needs and preferences.

Overall, our current call to action took the form of a narrative review of various identities that warrant clinical consideration in the assessment and treatment CHR-P youth. To our knowledge, this was one of the first attempts to comprehensively discuss considerations for historically marginalized and oppressed groups to help promote equity in CHR-P research and services. Although we discussed issues relevant for many aspects of identity, additional research is needed to have a more nuanced understanding how to improve services for youth in which these identities intersect to create unique challenges (e.g., LGBTQ+ REM youth). As research grows in this area, future studies may also wish to use systematic reviews and/or meta-analyses to more accurately define and quantify inequity in CHR-P services, and perhaps prioritize needs for the field moving forward.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments:

We are grateful to the youth, families, staff, and mentors in the CHR-P clinics we have worked in – in addition to the field of dedicated CHR-P and early psychosis researchers – who have all influenced and shaped our thinking on these topics.

Funding

This work was supported by the National Institute of Mental Health under Grant 1T32MH122394-01 (author JSD), and the American Psychological Foundation's Pearson Early Career Grant (author SNH).

References

- Addington J, Cornblatt BA, Cadenhead KS, Cannon TD, McGlashan TH, Perkins DO, Seidman LJ, Tsuang MT, Walker EF, & Woods SW (2011). At clinical high risk for psychosis: outcome for nonconverters. *American Journal of Psychiatry*, 168(8), 800–805. [PubMed: 21498462]
- Addington J, Liu L, Buchy L, Cadenhead KS, Cannon TD, Cornblatt BA, Perkins DO, Seidman LJ, Tsuang MT, & Walker EF (2015). North American prodrome longitudinal study (NAPLS 2): the prodromal symptoms. *The Journal of nervous and mental disease*, 203(5), 328. [PubMed: 25919383]
- Addington J, Piskulic D, Liu L, Lockwood J, Cadenhead KS, Cannon TD, Cornblatt BA, McGlashan TH, Perkins DO, & Seidman LJ (2017). Comorbid diagnoses for youth at clinical high risk of psychosis. *Schizophrenia Research*, 190, 90–95. [PubMed: 28372906]
- Al-Rousan T, Rubenstein L, Sieleni B, Deol H, & Wallace RB (2017). Inside the nation's largest mental health institution: A prevalence study in a state prison system. *BMC public health*, 17(1), 1–9. [PubMed: 28049454]
- Anglin DM, Ereshefsky S, Klaunig MJ, Bridgwater MA, Niendam TA, Ellman LM, DeVylder J, Thayer G, Bolden K, & Musket CW (2021). From Womb to Neighborhood: A Racial Analysis of Social Determinants of Psychosis in the United States. *American Journal of Psychiatry*, appi. ajp. 2020.20071091.
- Anglin DM, Galea S, & Bachman P (2020). Going upstream to advance psychosis prevention and improve public health. *JAMA Psychiatry*, 77(7), 665–666. [PubMed: 32236511]

- Anglin DM, Lighty Q, Greenspoon M, & Ellman LM (2014). Racial discrimination is associated with distressing subthreshold positive psychotic symptoms among US urban ethnic minority young adults. *Social psychiatry and psychiatric epidemiology*, 49(10), 1545–1555. [PubMed: 24695907]
- Anglin DM, Lui F, Espinosa A, Tikhonov A, & Ellman L (2018). Ethnic identity, racial discrimination and attenuated psychotic symptoms in an urban population of emerging adults. *Early Intervention in Psychiatry*, 12(3), 380–390. 10.1111/eip.12314 [PubMed: 26818635]
- Anglin DM, Polanco-Roman L, & Lui F (2015). Ethnic variation in whether dissociation mediates the relation between traumatic life events and attenuated positive psychotic symptoms. *Journal of Trauma & Dissociation*, 16(1), 68–85. [PubMed: 25365538]
- Bardol O, Grot S, Oh H, Poulet E, Zeroug-Vial H, Brunelin J, & Leane E (2020, May). Perceived ethnic discrimination as a risk factor for psychotic symptoms: a systematic review and meta-analysis. *Psychol Med*, 50(7), 1077–1089. 10.1017/s003329172000094x [PubMed: 32317042]
- Barr SM, Roberts D, & Thakkar KN (2021). Psychosis in transgender and gender non-conforming individuals: A review of the literature and a call for more research. *Psychiatry Research*, 306, 114272. [PubMed: 34808496]
- Beck K, Studerus E, Andreou C, Egloff L, Leanza L, Simon AE, Borgwardt S, & Riecher-Rössler A (2019). Clinical and functional ultra-long-term outcome of patients with a clinical high risk (CHR) for psychosis. *European Psychiatry*, 62, 30–37. 10.1016/j.eurpsy.2019.08.005 [PubMed: 31514058]
- Ben-David S, Cole A, Brucato G, Girgis RR, & Munson MR (2019a). Mental health service use decision-making among young adults at clinical high risk for developing psychosis. *Early Intervention in Psychiatry*, 13(5), 1050–1055. 10.1111/eip.12725 [PubMed: 30152169]
- Ben-David S, Cole AR, Brucato G, Girgis R, & Munson MR (2019b, Mar). A conceptual model of mental health service utilization among young adults at clinical high-risk for developing psychosis. *Psychiatr Rehabil J*, 42(1), 17–25. 10.1037/prj0000336 [PubMed: 30489142]
- Ben-David S, & Kealy D (2020, 2020/01/02). Identity in the context of early psychosis: a review of recent research. *Psychosis*, 12(1), 68–78. 10.1080/17522439.2019.1656283
- Berg AO, Melle I, Rossberg JI, Romm KL, Larsson S, Lagerberg TV, Andreassen OA, & Hauff E (2011, May 6). Perceived discrimination is associated with severity of positive and depression/anxiety symptoms in immigrants with psychosis: a cross-sectional study. *BMC Psychiatry*, 11, 77. 10.1186/1471-244X-11-77 [PubMed: 21548949]
- Bolton SL, & Sareen J (2011, Jan). Sexual orientation and its relation to mental disorders and suicide attempts: findings from a nationally representative sample. *Can J Psychiatry*, 56(1), 35–43. 10.1177/070674371105600107 [PubMed: 21324241]
- Boonstra N, Sterk B, Wunderink L, Sytema S, De Haan L, & Wiersma D (2012, Oct). Association of treatment delay, migration and urbanicity in psychosis. *European Psychiatry*, 27(7), 500–505. 10.1016/j.eurpsy.2011.05.001 [PubMed: 21705200]
- Bourque F, van der Ven E, & Malla A (2011, May). A meta-analysis of the risk for psychotic disorders among first- and second-generation immigrants. *Psychological Medicine*, 41(5), 897–910. 10.1017/S0033291710001406 [PubMed: 20663257]
- Boydell J, Van Os J, McKenzie K, & Murray R (2004). The association of inequality with the incidence of schizophrenia. *Social psychiatry and psychiatric epidemiology*, 39(8), 597–599. [PubMed: 15300368]
- Brandt L, Henssler J, Muller M, Wall S, Gabel D, & Heinz A (2019, Nov 1). Risk of Psychosis Among Refugees: A Systematic Review and Meta-analysis. *JAMA Psychiatry*, 76(11), 1133–1140. 10.1001/jamapsychiatry.2019.1937 [PubMed: 31411649]
- Brooke LE, Lin A, Ntoumanis N, & Gucciardi DF (2019). Is sport an untapped resource for recovery from first episode psychosis? A narrative review and call to action. *Early Intervention in Psychiatry*, 13(3), 358–368. [PubMed: 30039635]
- Brooks VR (1981). *Minority stress and lesbian women*. Free Press.
- Brown C, Conner KO, Copeland VC, Grote N, Beach S, Battista D, & Reynolds CF 3rd. (2010, Apr). Depression Stigma, Race, and Treatment Seeking Behavior and Attitudes. *J Community Psychol*, 38(3), 350–368. 10.1002/jcop.20368 [PubMed: 21274407]

- Burns JK, Tomita A, & Kapadia AS (2014). Income inequality and schizophrenia: increased schizophrenia incidence in countries with high levels of income inequality. *International Journal of Social Psychiatry*, 60(2), 185–196. [PubMed: 23594564]
- Calkins ME, Moore TM, Merikangas KR, Burstein M, Satterthwaite TD, Bilker WB, Ruparel K, Chiavacci R, Wolf DH, & Mentch F (2014). The psychosis spectrum in a young US community sample: findings from the Philadelphia Neurodevelopmental Cohort. *World Psychiatry*, 13(3), 296–305. [PubMed: 25273303]
- Cantor-Graae E, & Selten JP (2005, Jan). Schizophrenia and migration: a meta-analysis and review. *American Journal of Psychiatry*, 162(1), 12–24. 10.1176/appi.ajp.162.1.12 [PubMed: 15625195]
- Carol EE, & Mittal VA (2015). Normative adolescent experiences may confound assessment of positive symptoms in youth at ultra-high risk for psychosis. *Schizophrenia Research*, 1(166), 358–359.
- Carión RE, Goldberg TE, McLaughlin D, Auther AM, Correll CU, & Cornblatt BA (2011). Impact of neurocognition on social and role functioning in individuals at clinical high risk for psychosis. *American Journal of Psychiatry*, 168(8), 806–813. [PubMed: 21536691]
- Castillejos MC, Martin-Perez C, & Moreno-Kustner B (2018, Feb 22). A systematic review and meta-analysis of the incidence of psychotic disorders: the distribution of rates and the influence of gender, urbanicity, immigration and socio-economic level. *Psychological Medicine*, 1–15. 10.1017/S0033291718000235
- Catalan A, Salazar de Pablo G, Vaquerizo Serrano J, Mosillo P, Baldwin H, Fernández-Rivas A, Moreno C, Arango C, Correll CU, Bonoldi I, & Fusar-Poli P (2020). Annual Research Review: Prevention of psychosis in adolescents – systematic review and meta-analysis of advances in detection, prognosis and intervention. *Journal of Child Psychology and Psychiatry*, n/a(n/a). 10.1111/jcpp.13322
- Catalan A, Tognin S, Kempton MJ, Stahl D, de Pablo GS, Nelson B, Pantelis C, Riecher-Rössler A, Bressan R, & Barrantes-Vidal N (2020). Relationship between jumping to conclusions and clinical outcomes in people at clinical high-risk for psychosis. *Psychological medicine*, 1–9.
- Cauce AM, Domenech-Rodriguez M, Paradise M, Cochran BN, Shea JM, Srebnik D, & Baydar N (2002, Feb). Cultural and contextual influences in mental health help seeking: a focus on ethnic minority youth. *J Consult Clin Psychol*, 70(1), 44–55. 10.1037//0022-006x.70.1.44 [PubMed: 11860055]
- Chakraborty A, McManus S, Brugha TS, Bebbington P, & King M (2011, Feb). Mental health of the non-heterosexual population of England. *Br J Psychiatry*, 198(2), 143–148. 10.1192/bjp.bp.110.082271 [PubMed: 21282785]
- Chapman R (2020). Defining neurodiversity for research and practice. *Neurodiversity Studies: A New Critical Paradigm*, 14.
- Cicero DC, Krieg A, & Martin EA (2019). Measurement invariance of the Prodromal Questionnaire–Brief among White, Asian, Hispanic, and multiracial populations. *Assessment*, 26(2), 294–304. [PubMed: 28092988]
- Coleman KJ, Yarborough BJ, Beck A, Lynch FL, Stewart C, Penfold RS, Hunkeler EM, Operskalski BH, & Simon GE (2019). Patterns of Health Care Utilization Before First Episode Psychosis in Racial and Ethnic Groups. *Ethnicity & disease*, 29(4), 609. [PubMed: 31641328]
- Colizzi M, Ruggeri M, & Lasalvia A (2020, Apr). Should we be concerned about stigma and discrimination in people at risk for psychosis? A systematic review. *Psychol Med*, 50(5), 705–726. 10.1017/s0033291720000148 [PubMed: 32063250]
- Compton MT, Esterberg ML, Druss BG, Walker EF, & Kaslow NJ (2006, Jul). A descriptive study of pathways to care among hospitalized urban African American first-episode schizophrenia-spectrum patients. *Soc Psychiatry Psychiatr Epidemiol*, 41(7), 566–573. 10.1007/s00127-006-0065-z [PubMed: 16604270]
- Cooper S-A, Smiley E, Morrison J, Allan L, Williamson A, Finlayson J, Jackson A, & Mantry D (2007). Psychosis and adults with intellectual disabilities. *Social psychiatry and psychiatric epidemiology*, 42(7), 530–536. [PubMed: 17502974]
- Corcoran C, Perrin M, Harlap S, Deutsch L, Fennig S, Manor O, Nahon D, Kimhy D, Malaspina D, & Susser E (2009a). Effect of socioeconomic status and parents' education at birth on risk

of schizophrenia in offspring. *Social psychiatry and psychiatric epidemiology*, 44(4), 265–271. [PubMed: 18836884]

- Corcoran C, Perrin M, Harlap S, Deutsch L, Fennig S, Manor O, Nahon D, Kimhy D, Malaspina D, & Susser E (2009b). Incidence of schizophrenia among second-generation immigrants in the Jerusalem perinatal cohort. *Schizophrenia Bulletin*, 35(3), 596–602. [PubMed: 18648022]
- Corcoran C, Smith C, McLaughlin D, Auther A, Malaspina D, & Cornblatt B (2012). HPA axis function and symptoms in adolescents at clinical high risk for schizophrenia. *Schizophrenia Research*, 135(1-3), 170–174. [PubMed: 22226904]
- Corcoran C, Walker E, Huot R, Mittal V, Tessner K, Kestler L, & Malaspina D (2003). The Stress Cascade and Schizophrenia: Etiology and Onset. *Schizophrenia Bulletin*, 29(4), 671–692. 10.1093/oxfordjournals.schbul.a007038 [PubMed: 14989406]
- Cornblatt BA (2002). The New York high risk project to the Hillside recognition and prevention (RAP) program. *American journal of medical genetics*, 114(8), 956–966. [PubMed: 12457393]
- Cornblatt BA, Auther AM, Niendam T, Smith CW, Zinberg J, Bearden CE, & Cannon TD (2007). Preliminary findings for two new measures of social and role functioning in the prodromal phase of schizophrenia. *Schizophrenia Bulletin*, 33(3), 688–702. [PubMed: 17440198]
- Cowan HR (2020). Is schizophrenia research relevant during the COVID-19 pandemic? *Schizophrenia Research*, 220, 271–272. [PubMed: 32345478]
- Crenshaw K (1989). Demarginalizing the intersection of race and sex: A black feminist critique of antidiscrimination doctrine, feminist theory and antiracist politics. *u. Chi. Legal f*, 139.
- Crenshaw K (1990). Mapping the margins: Intersectionality, identity politics, and violence against women of color. *Stan. L. Rev*, 43, 1241.
- Damme KS, Ristanovic I, Vargas T, & Mittal VA (2020). Timing of menarche and abnormal hippocampal connectivity in youth at clinical-high risk for psychosis. *Psychoneuroendocrinology*, 117, 104672. [PubMed: 32388227]
- Day JK, Fish JN, Perez-Brumer A, Hatzenbuehler ML, & Russell ST (2017, Dec). Transgender Youth Substance Use Disparities: Results From a Population-Based Sample. *J Adolesc Health*, 61(6), 729–735. 10.1016/j.jadohealth.2017.06.024 [PubMed: 28942238]
- DeFreitas SC, Crone T, DeLeon M, & Ajayi A (2018). Perceived and Personal Mental Health Stigma in Latino and African American College Students. *Front Public Health*, 6, 49. 10.3389/fpubh.2018.00049 [PubMed: 29536000]
- DeLuca JS, Akouri-Shan L, Jay SY, Redman SL, Petti E, Lucksted A, Rakhshan Rouhakhtar P, Klaunig MJ, Edwards SM, Reeves GM, & Schiffman J (2021). Predictors of internalized mental health stigma in a help-seeking sample of youth: The roles of psychosis-spectrum symptoms and family functioning. *Journal of abnormal psychology*.
- DeLuca JS, Andorko ND, Chibani D, Jay SY, Rakhshan Rouhakhtar PJ, Petti E, Klaunig MJ, Thompson EC, Millman ZB, Connors KM, Akouri-Shan L, Fitzgerald J, Redman SL, Roemer C, Bridgwater MA, DeVyllder JE, King CA, Pitts SC, Reinblatt SP, Wehring HJ, Bussell KL, Solomon N, Edwards SM, Reeves GM, Buchanan RW, & Schiffman J (2020). Telepsychotherapy with youth at clinical high risk for psychosis: Clinical issues and best practices during the COVID-19 pandemic. *Journal of Psychotherapy Integration*, 30(2), 304–331. 10.1037/int0000211 [PubMed: 34305369]
- Deriu V, Moro MR, & Benoit L (2018). Early intervention for everyone? A review of cross-cultural issues and their treatment in ultra-high-risk (UHR) cohorts. *Early Intervention in Psychiatry*, 12(5), 796–810. [PubMed: 29708310]
- DeVyllder JE, Cogburn C, Oh HY, Anglin D, Smith ME, Sharpe T, Jun H-J, Schiffman J, Lukens E, & Link B (2017). Psychotic Experiences in the Context of Police Victimization: Data From the Survey of Police–Public Encounters. *Schizophrenia Bulletin*, 43(5), 993–1001. 10.1093/schbul/sbx038 [PubMed: 28369639]
- DeVyllder JE, Kelleher I, Lalane M, Oh H, Link BG, & Koyanagi A (2018). Association of urbanicity with psychosis in low-and middle-income countries. *JAMA Psychiatry*, 75(7), 678–686. [PubMed: 29799917]

- Dhejne C, Van Vlerken R, Heylens G, & Arcelus J (2016). Mental health and gender dysphoria: A review of the literature. *Int Rev Psychiatry*, 28(1), 44–57. 10.3109/09540261.2015.1115753 [PubMed: 26835611]
- Doyle DM, & Molix L (2016, Aug). Disparities in Social Health by Sexual Orientation and the Etiologic Role of Self-Reported Discrimination. *Arch Sex Behav*, 45(6), 1317–1327. 10.1007/s10508-015-0639-5 [PubMed: 26566900]
- Dragt S, Nieman DH, Veltman D, Becker HE, van de Fliert R, de Haan L, & Linszen DH (2011, Jan). Environmental factors and social adjustment as predictors of a first psychosis in subjects at ultra high risk. *Schizophrenia Research*, 125(1), 69–76. 10.1016/j.schres.2010.09.007 [PubMed: 20884179]
- Dunkel Schetter C, Schafer P, Lanzi RG, Clark-Kauffman E, Raju TN, Hillemeier MM, & Network CCH (2013). Shedding light on the mechanisms underlying health disparities through community participatory methods: The stress pathway. *Perspectives on Psychological Science*, 8(6), 613–633. [PubMed: 26173227]
- Egerton A, Howes OD, Houle S, McKenzie K, Valmaggia LR, Bagby MR, Tseng HH, Bloomfield MA, Kenk M, Bhattacharyya S, Suridjan I, Chaddock CA, Winton-Brown TT, Allen P, Rusjan P, Remington G, Meyer-Lindenberg A, McGuire PK, & Mizrahi R (2017, Mar 1). Elevated Striatal Dopamine Function in Immigrants and Their Children: A Risk Mechanism for Psychosis. *Schizophrenia Bulletin*, 43(2), 293–301. 10.1093/schbul/sbw181 [PubMed: 28057720]
- Erikson E (1968). *Identity: Youth and crisis*. WW Norton & company.
- Esterberg ML, & Compton MT (2006, May-Jun). Causes of schizophrenia reported by family members of urban African American hospitalized patients with schizophrenia. *Compr Psychiatry*, 47(3), 221–226. 10.1016/j.comppsy.2005.07.005 [PubMed: 16635652]
- Ferrara M, & Srihari VH (2021). Early intervention for psychosis in the United States: Tailoring services to improve care for women. *Psychiatric Services*, 72(1), 5–6. [PubMed: 32966169]
- Ferri F, Costantini M, Salone A, Di Iorio G, Martinotti G, Chiarelli A, Merla A, Di Giannantonio M, & Gallese V (2014, Jan). Upcoming tactile events and body ownership in schizophrenia. *Schizophr Res*, 152(1), 51–57. 10.1016/j.schres.2013.06.026 [PubMed: 23835002]
- Foss-Feig JH, Velthorst E, Smith L, Reichenberg A, Addington J, Cadenhead KS, Cornblatt BA, Mathalon DH, McGlashan TH, & Perkins DO (2019). Clinical profiles and conversion rates among young individuals with autism spectrum disorder who present to clinical high risk for psychosis services. *Journal of the American Academy of Child & Adolescent Psychiatry*, 58(6), 582–588.
- Fusar-Poli P, Borgwardt S, Bechdolf A, Addington J, Riecher-Rössler A, Schultze-Lutter F, Keshavan M, Wood S, Ruhrmann S, & Seidman LJ (2013). The psychosis high-risk state: a comprehensive state-of-the-art review. *JAMA Psychiatry*, 70(1), 107–120. [PubMed: 23165428]
- Fusar-Poli P, de Pablo GS, Correll CU, Meyer-Lindenberg A, Millan MJ, Borgwardt S, Galderisi S, Bechdolf A, Pfennig A, & Kessing LV (2020). Prevention of psychosis: advances in detection, prognosis, and intervention. *JAMA Psychiatry*.
- Fusar-Poli P, Salazar de Pablo G, Correll CU, Meyer-Lindenberg A, Millan MJ, Borgwardt S, Galderisi S, Bechdolf A, Pfennig A, Kessing LV, van Amelsvoort T, Nieman DH, Domschke K, Krebs M-O, Koutsouleris N, McGuire P, Do KQ, & Arango C (2020). Prevention of Psychosis: Advances in Detection, Prognosis, and Intervention. *JAMA Psychiatry*, 77(7), 755–765. 10.1001/jamapsychiatry.2019.4779 [PubMed: 32159746]
- Fusar-Poli P, Schultze-Lutter F, Cappucciati M, Rutigliano G, Bonoldi I, Stahl D, Borgwardt S, Riecher-Rössler A, Addington J, & Perkins DO (2016). The dark side of the moon: meta-analytical impact of recruitment strategies on risk enrichment in the clinical high risk state for psychosis. *Schizophrenia Bulletin*, 42(3), 732–743. [PubMed: 26591006]
- Fusar-Poli P, McGorry PD, & Kane JM (2017). Improving outcomes of first-episode psychosis: an overview. *World Psychiatry*, 16(3), 251–265. [PubMed: 28941089]
- Gallup. (2020). *LGBT Identification Rises to 5.6% in Latest U.S. Estimate*.
- Gara MA, Vega WA, Arndt S, Escamilla M, Fleck DE, Lawson WB, Lesser I, Neighbors HW, Wilson DR, Arnold LM, & Strakowski SM (2012, Jun). Influence of patient race and ethnicity on clinical

- assessment in patients with affective disorders. *Arch Gen Psychiatry*, 69(6), 593–600. 10.1001/archgenpsychiatry.2011.2040 [PubMed: 22309972]
- Geros H, Sizer H, Mifsud N, Reynolds S, Kim DJ, Eaton S, McGorry P, Nelson B, & O'Donoghue B (2020, Jan). Migrant status and identification as ultra-high risk for psychosis and transitioning to a psychotic disorder. *Acta Psychiatrica Scandinavica*, 141(1), 52–59. 10.1111/acps.13099 [PubMed: 31520527]
- Gerson R, Wong C, Davidson L, Malaspina D, McGlashan T, & Corcoran C (2011). Self-reported coping strategies in families of patients in early stages of psychotic disorder: an exploratory study. *Early Intervention in Psychiatry*, 5(1), 76–80. [PubMed: 21272279]
- Gevonden MJ, Selten JP, Myin-Germeys I, de Graaf R, ten Have M, van Dorsselaer S, van Os J, & Veling W (2014, Jan). Sexual minority status and psychotic symptoms: findings from the Netherlands Mental Health Survey and Incidence Studies (NEMESIS). *Psychol Med*, 44(2), 421–433. 10.1017/S0033291713000718 [PubMed: 23710972]
- Grant MJ, & Booth A (2009). A typology of reviews: an analysis of 14 review types and associated methodologies. *Health information & libraries journal*, 26(2), 91–108. [PubMed: 19490148]
- Gur RC, Calkins ME, Satterthwaite TD, Ruparel K, Bilker WB, Moore TM, Savitt AP, Hakonarson H, & Gur RE (2014). Neurocognitive growth charting in psychosis spectrum youths. *JAMA Psychiatry*, 71(4), 366–374. [PubMed: 24499990]
- Hall GCN, Berkman ET, Zane NW, Leong FT, Hwang W-C, Nezu AM, Nezu CM, Hong JJ, Chu JP, & Huang ER (2021). Reducing mental health disparities by increasing the personal relevance of interventions. *American psychologist*, 76(1), 91. [PubMed: 32118456]
- Hankerson SH, Svob C, & Jones MK (2018). Partnering with black churches to increase access to care. *Psychiatric Services (Washington, DC)*, 69(2), 125.
- Harrop C, Ellett L, Brand R, & Lobban F (2015). Friends interventions in psychosis: a narrative review and call to action. *Early Intervention in Psychiatry*, 9(4), 269–278. [PubMed: 25130455]
- Hays PA (1996). Addressing the complexities of culture and gender in counseling. *Journal of Counseling & Development*, 74(4), 332–338.
- Health CD o. A. a. S. (2019). Youth Risk Behavior Survey: Number and percentage of students, by sexual identity.
- Heitz U, Studerus E, Menghini-Muller S, Pappmeyer M, Egloff L, Ittig S, Navarra A, Andreou C, & Riecher-Rossler A (2019, Jun). Gender differences in first self-perceived signs and symptoms in patients with an at-risk mental state and first-episode psychosis. *Early Interv Psychiatry*, 13(3), 582–588. 10.1111/eip.12528 [PubMed: 29235240]
- Henssler J, Brandt L, Muller M, Liu S, Montag C, Sterzer P, & Heinz A (2020, Apr). Migration and schizophrenia: meta-analysis and explanatory framework. *European Archives of Psychiatry and Clinical Neuroscience*, 270(3), 325–335. 10.1007/s00406-019-01028-7 [PubMed: 31161262]
- Heun-Johnson H, Menchine M, Axeen S, Lung K, Claudius I, Wright T, & Seabury SA (2021). Association Between Race/Ethnicity and Disparities in Health Care Use Before First-Episode Psychosis Among Privately Insured Young Patients. *JAMA Psychiatry*, 78(3), 311–319. 10.1001/jamapsychiatry.2020.3995 [PubMed: 33355626]
- Hoenig J, & Kenna JC (1974, May). The nosological position of transsexualism. *Arch Sex Behav*, 3(3), 273–287. 10.1007/BF01541490 [PubMed: 4836845]
- Horvitz-Lennon M, Volya R, Garfield R, Donohue JM, Lave JR, & Normand SLT (2015). Where you live matters: quality and racial/ethnic disparities in schizophrenia care in four state Medicaid programs. *Health services research*, 50(5), 1710–1729. [PubMed: 25759240]
- Izon E, Berry K, Law H, & French P (2018, 2018/12/01/). Expressed emotion (EE) in families of individuals at-risk of developing psychosis: A systematic review. *Psychiatry Research*, 270, 661–672. 10.1016/j.psychres.2018.10.065 [PubMed: 30384287]
- Izon E, Berry K, Wearden A, Carter LA, Law H, & French P (2021). Investigating expressed emotion in individuals at-risk of developing psychosis and their families over 12 months. *Clinical Psychology & Psychotherapy*.
- Jacob L, Smith L, McDermott D, Haro JM, Stickley A, & Koyanagi A (2021, Jan). Relationship between sexual orientation and psychotic experiences in the general population in England. *Psychol Med*, 51(1), 138–146. 10.1017/S003329171900309X [PubMed: 31694728]

- Janssen I, Hanssen M, Bak M, Bijl RV, de Graaf R, Vollebergh W, McKenzie K, & van Os J (2003, Jan). Discrimination and delusional ideation. *Br J Psychiatry*, 182, 71–76. 10.1192/bjp.182.1.71 [PubMed: 12509322]
- Jones N, Atterbury K, Byrne L, Carras M, Brown M, & Phalen P (2021). Lived Experience, Research Leadership, and the Transformation of Mental Health Services: Building a Researcher Pipeline. *Psychiatric Services*, appi. ps. 202000468.
- Jones N, Kamens S, Oluwoye O, Mascayano F, Perry C, Manseau M, & Compton MT (2021a). Structural Disadvantage and Culture, Race, and Ethnicity in Early Psychosis Services: International Provider Survey. *Psychiatric Services*, appi. ps. 202000211.
- Jones N, Kamens S, Oluwoye O, Mascayano F, Perry C, Manseau M, & Compton MT (2021b, Mar 1). Structural Disadvantage and Culture, Race, and Ethnicity in Early Psychosis Services: International Provider Survey. *Psychiatric Services*, 72(3), 254–263. 10.1176/appi.ps.202000211 [PubMed: 33430649]
- Karcher NR, Barch DM, Avenevoli S, Savill M, Huber RS, Simon TJ, Leckliter IN, Sher KJ, & Loewy RL (2018). Assessment of the Prodromal Questionnaire–Brief Child Version for measurement of self-reported psychoticlike experiences in childhood. *JAMA Psychiatry*, 75(8), 853–861. [PubMed: 29874361]
- Kataoka SH, Zhang L, & Wells KB (2002). Unmet need for mental health care among US children: Variation by ethnicity and insurance status. *American Journal of Psychiatry*, 159(9), 1548–1555. [PubMed: 12202276]
- Kline ER, Chokran C, Rodenhiser-Hill J, Seidman LJ, & Woodberry KA (2019). Psychosis screening practices in schools: A survey of school-based mental health providers. *Early Intervention in Psychiatry*, 13(4), 818–822. [PubMed: 29726591]
- Kotlicka-Antczak M, Podgórski M, Oliver D, Maric NP, Valmaggia L, & Fusar-Poli P (2020). Worldwide implementation of clinical services for the prevention of psychosis: The IEPA early intervention in mental health survey. *Early Intervention in Psychiatry*, n/a(n/a). 10.1111/eip.12950
- Kraan T, Velthorst E, Smit F, de Haan L, & van der Gaag M (2015, 2015/02/01/). Trauma and recent life events in individuals at ultra high risk for psychosis: Review and meta-analysis. *Schizophrenia Research*, 161(2), 143–149. 10.1016/j.schres.2014.11.026 [PubMed: 25499046]
- Krabbendam L, & van Os J (2005, Oct). Schizophrenia and urbanicity: a major environmental influence--conditional on genetic risk. *Schizophrenia Bulletin*, 31(4), 795–799. 10.1093/schbul/sbi060 [PubMed: 16150958]
- Lång U, Mittal VA, Schiffman J, & Therman S (2020). Measurement Invariance of Psychotic-Like Symptoms as Measured With the Prodromal Questionnaire, Brief Version (PQ-B) in Adolescent and Adult Population Samples. *Frontiers in psychiatry*, 11.
- Laster Pirtle WN (2020). <? covid19?> Racial Capitalism: A Fundamental Cause of Novel Coronavirus (COVID-19) Pandemic Inequities in the United States. *Health Education & Behavior*, 47(4), 504–508. [PubMed: 32338071]
- Leanza L, Studerus E, Mackintosh AJ, Beck K, Seiler L, Andreou C, & Riecher-Rössler A (2020). Predictors of study drop-out and service disengagement in patients at clinical high risk for psychosis. *Social psychiatry and psychiatric epidemiology*, 55(5), 539–548. [PubMed: 31646355]
- Lewis G, David A, Andreasson S, & Allebeck P (1992, Jul 18). Schizophrenia and city life. *Lancet*, 340(8812), 137–140. 10.1016/0140-6736(92)93213-7 [PubMed: 1352565]
- Li H, Friedman-Yakobian M, Hasler VC, Shapiro DI, & Wu E (2019). Attenuated psychosis syndromes among Asian American youth and young adults: A culturally relevant case illustration approach. In *Handbook of Attenuated Psychosis Syndrome Across Cultures* (pp. 219–236). Springer.
- Li H, Shapiro DI, & Seidman LJ (2019). *Handbook of Attenuated Psychosis Syndrome Across Cultures: International Perspectives on Early Identification and Intervention*. Springer Nature.
- Lim J, Rekhi G, Rapisarda A, Lam M, Kraus M, Keefe RS, & Lee J (2015). Impact of psychiatric comorbidity in individuals at Ultra High Risk of psychosis—Findings from the Longitudinal Youth at Risk Study (LYRIKS). *Schizophrenia Research*, 164(1-3), 8–14. [PubMed: 25818728]

- Loch AA, Freitas EL, Hortêncio L, Chianca C, Alves TM, Serpa MH, Andrade JC, van de Bilt MT, Gattaz WF, & Rössler W (2019). Hearing spirits? Religiosity in individuals at risk for psychosis —Results from the Brazilian SSAPP cohort. *Schizophrenia Research*, 204, 353–359. [PubMed: 30266512]
- Loewy RL, Corey S, Amirfathi F, Dabit S, Fulford D, Pearson R, Hua JPY, Schlosser D, Stuart BK, Mathalon DH, & Vinogradov S (2019, 2019/03/01/). Childhood trauma and clinical high risk for psychosis. *Schizophrenia Research*, 205, 10–14. 10.1016/j.schres.2018.05.003 [PubMed: 29779964]
- Longenecker J, Genderson J, Dickinson D, Malley J, Elvevag B, Weinberger DR, & Gold J (2010, Jun). Where have all the women gone?: participant gender in epidemiological and non-epidemiological research of schizophrenia. *Schizophr Res*, 119(1-3), 240–245. 10.1016/j.schres.2010.03.023 [PubMed: 20399612]
- LoPilato AM, Goines K, Addington J, Bearden CE, Cadenhead KS, Cannon TD, Cornblatt BA, Mathalon DH, McGlashan TH, Seidman L, Perkins DO, Tsuang MT, Woods SW, & Walker EF (2019, Nov). Impact of childhood adversity on corticolimbic volumes in youth at clinical high-risk for psychosis. *Schizophrenia Research*, 213, 48–55. 10.1016/j.schres.2019.01.048 [PubMed: 30745068]
- Loring M, & Powell B (1988). Gender, race, and DSM-III: A study of the objectivity of psychiatric diagnostic behavior. *Journal of health and social behavior*, 1–22. [PubMed: 3367027]
- Lucksted A (2004). Lesbian, Gay, Bisexual, and Transgender People Receiving Services in the Public Mental Health System: Raising Issues. *Journal of Gay & Lesbian Psychotherapy*, 8(3-4), 25–42. 10.1300/J236v08n03_03
- Lynch S, McFarlane WR, Joly B, Adelsheim S, Auther A, Cornblatt BA, Migliorati M, Ragland JD, Sale T, & Spring E (2016). Early detection, intervention and prevention of psychosis program: community outreach and early identification at six US sites. *Psychiatric Services*, 67(5), 510–516. [PubMed: 26766751]
- Mackenzie CS, Gekoski WL, & Knox VJ (2006, Nov). Age, gender, and the underutilization of mental health services: the influence of help-seeking attitudes. *Aging Ment Health*, 10(6), 574–582. 10.1080/13607860600641200 [PubMed: 17050086]
- Marcelis M, Takei N, & van Os J (1999, Sep). Urbanization and risk for schizophrenia: does the effect operate before or around the time of illness onset? *Psychological Medicine*, 29(5), 1197–1203. 10.1017/s0033291799008983 [PubMed: 10576311]
- Mayo D, Corey S, Kelly LH, Yohannes S, Youngquist AL, Stuart BK, Niendam TA, & Loewy RL (2017). The role of trauma and stressful life events among individuals at clinical high risk for psychosis: a review. *Frontiers in psychiatry*, 8, 55. [PubMed: 28473776]
- McGorry P, Trethowan J, & Rickwood D (2019). Creating headspace for integrated youth mental health care. *World Psychiatry*, 18(2), 140. [PubMed: 31059618]
- McGorry PD, Mei C, Hartmann J, Yung AR, & Nelson B (2021). Intervention strategies for ultra-high risk for psychosis: Progress in delaying the onset and reducing the impact of first-episode psychosis. *Schizophrenia Research*, 228, 344–356. [PubMed: 33545668]
- McGrath J, Saha S, Welham J, El Saadi O, MacCauley C, & Chant D (2004, Apr 28). A systematic review of the incidence of schizophrenia: the distribution of rates and the influence of sex, urbanicity, migrant status and methodology. *BMC Medicine*, 2, 13. 10.1186/1741-7015-2-13 [PubMed: 15115547]
- McKay TR, & Watson RJ (2020). Gender expansive youth disclosure and mental health: Clinical implications of gender identity disclosure. *Psychology of Sexual Orientation and Gender Diversity*, 7(1), 66–75. 10.1037/sgd0000354 [PubMed: 33855103]
- Meijer JH, Eeckhout GM, van Vlerken RH, & de Vries AL (2017, Apr). Gender Dysphoria and Co-Existing Psychosis: Review and Four Case Examples of Successful Gender Affirmative Treatment. *LGBT Health*, 4(2), 106–114. 10.1089/lgbt.2016.0133 [PubMed: 28170299]
- Merikangas KR, He J. p., Burstein M, Swendsen J, Avenevoli S, Case B, Georgiades K, Heaton L, Swanson S, & Olfson M (2011). Service utilization for lifetime mental disorders in US adolescents: results of the National Comorbidity Survey–Adolescent Supplement (NCS-A). *Journal of the American Academy of Child & Adolescent Psychiatry*, 50(1), 32–45. [PubMed: 21156268]

- Metzl JM (2010). *The protest psychosis: How schizophrenia became a black disease*. Beacon Press.
- Meyer IH (2003). Prejudice, social stress, and mental health in lesbian, gay, and bisexual populations: conceptual issues and research evidence. *Psychological bulletin*, 129(5), 674. [PubMed: 12956539]
- Meyer MS, Rosenthal A, Bolden KA, Loewy RL, Savill M, Shim R, Rodriguez J, Flores V, Pavao E, & Niendam TA (2020). Psychosis screening in schools: Considerations and implementation strategies. *Early Intervention in Psychiatry*, 14(1), 130–136. [PubMed: 31287611]
- Michalska da Rocha B, Rhodes S, Vasilopoulou E, & Hutton P (2018, Jan 13). Loneliness in Psychosis: A Meta-analytical Review. *Schizophr Bull*, 44(1), 114–125. 10.1093/schbul/sbx036 [PubMed: 28369646]
- Millman ZB, Rakhshan Rouhakhtar PJ, DeVlyder JE, Smith ME, Phalen PL, Woods SW, Walsh BC, Parham B, Reeves GM, & Schiffman J (2019). Evidence for differential predictive performance of the prime screen between black and white help-seeking youths. *Psychiatric Services*, 70(10), 907–914. [PubMed: 31310187]
- Mizock L, & Fleming MZ (2011, Apr). Transgender and Gender Variant Populations With Mental Illness: Implications for Clinical Care. *Professional Psychology-Research and Practice*, 42(2), 208–213. 10.1037/a0022522
- Mizock LL,TK (2008). Trauma in transgender populations: Risk, resilience, and clinical care. *Journal of Emotional Abuse* 8(3), 335–354.
- Morgan C, Fisher H, Hutchinson G, Kirkbride J, Craig TK, Morgan K, Dazzan P, Boydell J, Doody GA, & Jones PB (2009). Ethnicity, social disadvantage and psychotic-like experiences in a healthy population based sample. *Acta Psychiatrica Scandinavica*, 119(3), 226–235. [PubMed: 19053965]
- Nagendra A, Schooler NR, Kane JM, Robinson DG, Mueser KT, Estroff SE, Addington J, Marcy P, & Penn DL (2018). Demographic, psychosocial, clinical, and neurocognitive baseline characteristics of Black Americans in the RAISE-ETP study. *Schizophrenia Research*, 193, 64–68. [PubMed: 28709773]
- Nazroo JY, Bhui KS, & Rhodes J (2020). Where next for understanding race/ethnic inequalities in severe mental illness? Structural, interpersonal and institutional racism. *Sociology of health & illness*, 42(2), 262–276. [PubMed: 31562655]
- Nemani K, Li C, Olfson M, Blessing EM, Razavian N, Chen J, Petkova E, & Goff DC (2021). Association of psychiatric disorders with mortality among patients with COVID-19. *JAMA Psychiatry*.
- Newbury JB, Arseneault L, Beevers S, Kitwiroon N, Roberts S, Pariante CM, Kelly FJ, & Fisher HL (2019). Association of air pollution exposure with psychotic experiences during adolescence. *JAMA Psychiatry*, 76(6), 614–623. [PubMed: 30916743]
- Nielsen KE (2012). *A disability history of the United States (Vol. 2)*. Beacon Press.
- NIMH. (2020). Research Highlight: NIMH Part of Collaborative Effort to Advance Early Intervention for Individuals at Risk of Developing Schizophrenia.
- Nogueira AS, Andrade JC, Serpa MH, Alves TM, Freitas EL, Hortencio L, van de Bilt MT, Rossler W, Gattaz WF, & Loch AA (2020, Aug 3). Influence of migration on the thought process of individuals at ultra-high risk for psychosis. *Braz J Psychiatry*.
- Novacek DM, Hampton-Anderson JN, Ebor MT, Loeb TB, & Wyatt GE (2020). Mental health ramifications of the COVID-19 pandemic for Black Americans: Clinical and research recommendations. *Psychological Trauma: Theory, Research, Practice, and Policy*.
- Novacek DM, Lopilato AM, Goines KB, Trotman HD, Compton MT, & Walker EF (2019). Cultural Considerations in the Treatment of African American Youth with Attenuated Psychosis Syndromes: The Importance of Socio-contextual and Clinical Factors. In Shapiro H, Li, D. I., & Seidman LJ (Eds.), *Handbook of Attenuated Psychosis Syndrome Across Cultures* (pp. 199–2018). Springer Nature Switzerland.
- O'Donoghue B, Geros H, Sizer H, Addington J, Amminger GP, Beaden CE, Cadenhead KS, Cannon TD, Cornblatt BA, Berger GE, Chen EYH, de Haan L, Hartmann JA, Hickie IB, Ising HK, Lavoie S, Lin A, Markulev C, Mathalon DH, McGlashan TH, Mifsud NG, Mossaheb N, Nieman DH, Nordentoft M, Perkins DO, Riecher-Rossler A, Schafer MR, Schlogelhofer M, Seidman LJ,

- Smesny S, Thompson A, Tsuang MT, van der Gaag M, Verma S, Walker EF, Wood SJ, Woods SW, Yuen HP, Yung AR, McGorry PD, & Nelson B (2021, Jan 5). The association between migrant status and transition in an ultra-high risk for psychosis population. *Social Psychiatry and Psychiatric Epidemiology*.
- O'Donoghue B, Yung AR, Wood S, Thompson A, Lin A, McGorry P, & Nelson B (2015, Dec). Neighbourhood characteristics and the rate of identification of young people at ultra-high risk for psychosis. *Schizophrenia Research*, 169(1-3), 214–216. [PubMed: 26391282]
- Ochoa S, Usall J, Cobo J, Labad X, & Kulkarni J (2012). Gender differences in schizophrenia and first-episode psychosis: a comprehensive literature review. *Schizophr Res Treatment*, 2012, 916198. [PubMed: 22966451]
- Oh H, Cogburn CD, Anglin D, Lukens E, & DeVlyder J (2016). Major discriminatory events and risk for psychotic experiences among Black Americans. *American journal of Orthopsychiatry*, 86(3), 277. [PubMed: 26963179]
- Oh H, Koyanagi A, Kelleher I, & DeVlyder J (2018). Psychotic experiences and disability: findings from the Collaborative Psychiatric Epidemiology Surveys. *Schizophrenia Research*, 193, 343–347. [PubMed: 28797526]
- Oluwoye O, Davis B, Kuhney FS, & Anglin DM (2021). Systematic review of pathways to care in the US for Black individuals with early psychosis. *npj Schizophrenia*, 7(1), 1–10. [PubMed: 33479257]
- Oluwoye O, Stiles B, Monroe-DeVita M, Chwastiak L, McClellan JM, Dyck D, Cabassa LJ, & McDonnell MG (2018). Racial-ethnic disparities in first-episode psychosis treatment outcomes from the RAISE-ETP study. *Psychiatric Services*, 69(11), 1138–1145. [PubMed: 30152275]
- Olivet DM, Carrión RE, Auther AM, & Cornblatt BA (2015). Self-awareness of functional impairment in individuals at clinical high-risk for psychosis. *Early Intervention in Psychiatry*, 9(2), 100–107. [PubMed: 23968457]
- Omer S, Kirkbride JB, Pringle DG, Russell V, O'Callaghan E, & Waddington JL (2014, Jan). Neighbourhood-level socio-environmental factors and incidence of first episode psychosis by place at onset in rural Ireland: the Cavan-Monaghan First Episode Psychosis Study [CAMFEPS]. *Schizophrenia Research*, 152(1), 152–157. [PubMed: 24342585]
- Ozer EJ, Abraczinskas M, Duarte C, Mathur R, Ballard PJ, Gibbs L, Olivas ET, Bewa MJ, & Afifi R (2020). Youth participatory approaches and health equity: Conceptualization and integrative review. *American journal of community psychology*.
- Patten SB, Svenson LW, & Metz LM (2005). Psychotic disorders in MS: population-based evidence of an association. *Neurology*, 65(7), 1123–1125. [PubMed: 16217073]
- Pearce J, Rafiq S, Simpson J, & Varese F (2019, Sep). Perceived discrimination and psychosis: a systematic review of the literature. *Soc Psychiatry Psychiatr Epidemiol*, 54(9), 1023–1044. [PubMed: 31236631]
- Pedersen CB, & Mortensen PB (2001a, Nov). Evidence of a dose-response relationship between urbanicity during upbringing and schizophrenia risk. *Archives of General Psychiatry*, 58(11), 1039–1046. [PubMed: 11695950]
- Pedersen CB, & Mortensen PB (2001b, Jul). Family history, place and season of birth as risk factors for schizophrenia in Denmark: a replication and reanalysis. *British Journal of Psychiatry*, 179, 46–52.
- Phelan JC, Lucas JW, Teachman B, Braverman BH, Namaky N, & Greenberg M (2019). A Comparison of Status and Stigma Processes: Explicit and Implicit Appraisals of “Mentally Ill People” and “Uneducated People”. *Stigma and health (Washington, D.C.)*, 4(2), 213–224.
- Rakhshan Rouhakhtar PJ, Pitts SC, & Schiffman J (2019). Associations between race, discrimination, community violence, traumatic life events, and psychosis-like experiences in a sample of college students. *Journal of Clinical Medicine*, 8(10), 1573.
- Redman SL, Corcoran CM, Kimhy D, & Malaspina D (2017). Effects of early trauma on psychosis development in clinical high-risk individuals and stability of trauma assessment across studies: a review. *Archives of psychology (Chicago, Ill.)*, 1(3).

- Reisner SL, Pardo ST, Gamarel KE, White Hughto JM, Pardee DJ, & Keo-Meier CL (2015, Dec). Substance Use to Cope with Stigma in Healthcare Among U.S. Female-to-Male Trans Masculine Adults. *LGBT Health*, 2(4), 324–332. [PubMed: 26788773]
- Richardson L, Hameed Y, Perez J, Jones PB, & Kirkbride JB (2018, Jan 1). Association of Environment With the Risk of Developing Psychotic Disorders in Rural Populations: Findings from the Social Epidemiology of Psychoses in East Anglia Study. *JAMA Psychiatry*, 75(1), 75–83. [PubMed: 29188295]
- Rietschel L, Lambert M, Karow A, Zink M, Muller H, Heinz A, de Millas W, Janssen B, Gaebel W, Schneider F, Naber D, Juckel G, Kruger-Ozgurdal S, Wobrock T, Wagner M, Maier W, Klosterkotter J, Bechdolf A, & group P. s. (2017, Aug). Clinical high risk for psychosis: gender differences in symptoms and social functioning. *Early Interv Psychiatry*, 11(4), 306–313. [PubMed: 25808791]
- Rimvall MK, van Os J, Verhulst F, Wolf RT, Larsen JT, Clemmensen L, Skovgaard AM, Rask CU, & Jeppesen P (2020). Mental health service use and psychopharmacological treatment following psychotic experiences in preadolescence. *American Journal of Psychiatry*, 177(4), 318–326. [PubMed: 32098486]
- Rossmann K, Salamanca P, & Macapagal K (2017). A Qualitative Study Examining Young Adults' Experiences of Disclosure and Nondisclosure of LGBTQ Identity to Health Care Providers. *J Homosex*, 64(10), 1390–1410. [PubMed: 28459379]
- Rouhakhtar PR, Pitts SC, Millman ZB, Andorko ND, Redman S, Wilson C, Demro C, Phalen PL, Walsh B, & Woods S (2019). The impact of age on the validity of psychosis-risk screening in a sample of help-seeking youth. *Psychiatry Research*, 274, 30–35. [PubMed: 30780059]
- Ruiz B, Ceccolini CJ, Shah BB, Crump F, Girgis RR, Brucato G, Yang LH, & Corcoran CM (2021). Stigma and coping experiences in Latinx individuals at clinical high-risk for psychosis. *Early Intervention in Psychiatry*.
- Saleem MM, Stowkowy J, Cadenhead KS, Cannon TD, Cornblatt BA, McGlashan TH, Perkins DO, Seidman LJ, Tsuang MT, & Walker EF (2014). Perceived discrimination in those at clinical high risk for psychosis. *Early Intervention in Psychiatry*, 8(1), 77–81. [PubMed: 23773288]
- SAMHSA. (2018). *Community Programs for Outreach and Intervention with Youth and Young Adults at Clinical High Risk for Psychosis*.
- Santesteban-Echarri O, MacQueen G, Goldstein BI, Wang J, Kennedy SH, Bray S, Lebel C, & Addington J (2018). Family functioning in youth at-risk for serious mental illness. *Comprehensive Psychiatry*, 87, 17–24. [PubMed: 30193153]
- Sarac C, DeLuca JS, Bilgrami ZR, Herrera SN, Myers JJ, Dobbs MF, Haas SS, Todd TL, Srivastava A, Jespersen R, Shaik RB, Landa Y, Davidson L, Pavlo AJ, & Corcoran CM (2021). A qualitative study on identity in individuals at clinical high-risk for psychosis: "...Why does it have to be one thing?". under review.
- Schiffman J, Ellman LM, & Mittal VA (2019). Individual differences and psychosis-risk screening: Practical suggestions to improve the scope and quality of early identification. *Frontiers in psychiatry*, 10, 6. [PubMed: 30837898]
- Schofield P, Thygesen M, Das-Munshi J, Becares L, Cantor-Graae E, Pedersen C, & Agerbo E (2017, Dec). Ethnic density, urbanicity and psychosis risk for migrant groups -A population cohort study. *Schizophrenia Research*, 190, 82–87. [PubMed: 28318842]
- Schwartz RC, & Blankenship DM (2014). Racial disparities in psychotic disorder diagnosis: A review of empirical literature. *World journal of psychiatry*, 4(4), 133. [PubMed: 25540728]
- Scott J, Chant D, Andrews G, & McGrath J (2006, Feb). Psychotic-like experiences in the general community: the correlates of CIDI psychosis screen items in an Australian sample. *Psychol Med*, 36(2), 231–238. [PubMed: 16303059]
- Seng JS, Lopez WD, Sperlich M, Hamama L, & Meldrum CDR (2012). Marginalized identities, discrimination burden, and mental health: Empirical exploration of an interpersonal-level approach to modeling intersectionality. *Social science & medicine*, 75(12), 2437–2445. [PubMed: 23089613]
- Severaid KB, Osborne KJ, & Mittal VA (2019). Implications of religious and spiritual practices for youth at clinical high risk for psychosis. *Schizophrenia Research*, 208, 481. [PubMed: 30799216]

- Smith WB, Goldhammer H, & Keuroghlian AS (2019, Jan 1). Affirming Gender Identity of Patients With Serious Mental Illness. *Psychiatr Serv*, 70(1), 65–67. [PubMed: 30332928]
- Spauwen J, Krabbendam L, Lieb R, Wittchen HU, & van Os J (2004, Nov-Dec). Does urbanicity shift the population expression of psychosis? *Journal of Psychiatric Research*, 38(6), 613–618. [PubMed: 15458857]
- Stain HJ, Halpin SA, Baker AL, Startup M, Carr VJ, Schall U, Crittenden K, Clark V, Lewin TJ, & Bucci S (2018, Dec). Impact of rurality and substance use on young people at ultra high risk for psychosis. *Early Interv Psychiatry*, 12(6), 1173–1180. [PubMed: 28744989]
- State, U. D. o. (2019). *Country Reports on Terrorism 2019*.
- Stefanis NC, Delespaul P, Smyrnis N, Lembesi A, Avramopoulos DA, Evdokimidis IK, Stefanis CN, & van Os J (2004, May). Is the excess risk of psychosis-like experiences in urban areas attributable to altered cognitive development? *Social Psychiatry and Psychiatric Epidemiology*, 39(5), 364–368. [PubMed: 15133592]
- Stowkowy J, Liu L, Cadenhead KS, Cannon TD, Cornblatt BA, McGlashan TH, Perkins DO, Seidman LJ, Tsuang MT, Walker EF, Woods SW, Bearden CE, Mathalon DH, & Addington J (2016, Apr). Early traumatic experiences, perceived discrimination and conversion to psychosis in those at clinical high risk for psychosis. *Soc Psychiatry Psychiatr Epidemiol*, 51(4), 497–503. [PubMed: 26851943]
- Thompson E, Millman ZB, Okuzawa N, Mittal V, DeVlyder J, Skadberg T, Buchanan RW, Reeves GM, & Schiffman J (2015). Evidence-based early interventions for individuals at clinical high risk for psychosis: a review of treatment components. *The Journal of nervous and mental disease*, 203(5), 342–351. [PubMed: 25919384]
- Thompson E, Rakhshan P, Pitts SC, Demro C, Millman ZB, Bussell K, DeVlyder J, Kline E, Reeves GM, & Schiffman J (2019). Family functioning moderates the impact of psychosis-risk symptoms on social and role functioning. *Schizophrenia Research*, 204, 337–342. [PubMed: 30205908]
- Thompson JL, Kelly M, Kimhy D, Harkavy-Friedman JM, Khan S, Messinger JW, Schobel S, Goetz R, Malaspina D, & Corcoran C (2009). Childhood trauma and prodromal symptoms among individuals at clinical high risk for psychosis. *Schizophrenia Research*, 108(1-3), 176–181. [PubMed: 19174322]
- Tor J, Dolz M, Sintes A, Munoz D, Pardo M, de la Serna E, Puig O, Sugranyes G, & Baeza I (2018, Jun). Clinical high risk for psychosis in children and adolescents: a systematic review. *Eur Child Adolesc Psychiatry*, 27(6), 683–700. [PubMed: 28914382]
- Trotman HD, Holtzman CW, Ryan AT, Shapiro DI, MacDonald AN, Goulding SM, Brasfield JL, & Walker EF (2013). The development of psychotic disorders in adolescence: a potential role for hormones. *Hormones and behavior*, 64(2), 411–419. [PubMed: 23998682]
- van der Gaag M, Smit F, Bechdolf A, French P, Linszen DH, Yung AR, McGorry P, & Cuijpers P (2013, Sep). Preventing a first episode of psychosis: meta-analysis of randomized controlled prevention trials of 12 month and longer-term follow-ups. *Schizophr Res*, 149(1-3), 56–62. [PubMed: 23870806]
- van Os J, Hanssen M, Bijl RV, & Vollebergh W (2001). Prevalence of Psychotic Disorder and Community Level of Psychotic Symptoms: An Urban-Rural Comparison. *Archives of General Psychiatry*, 58(7), 663–668. [PubMed: 11448373]
- van Os J, Hanssen M, de Graaf R, & Vollebergh W (2002, Oct 2002 2014-08-02). Does the urban environment independently increase the risk for both negative and positive features of psychosis? *Social Psychiatry and Psychiatric Epidemiology*, 37(10), 460–464. [PubMed: 12242624]
- Vargas T, Rakhshan Rouhakhtar PJ, Schiffman J, Zou DS, Rydland KJ, & Mittal VA (2020, Jan). Neighborhood crime, socioeconomic status, and suspiciousness in adolescents and young adults at Clinical High Risk (CHR) for psychosis. *Schizophr Res*, 215, 74–80. [PubMed: 31759810]
- Veling W, Selten JP, Susser E, Laan W, Mackenbach JP, & Hoek HW (2007, Aug). Discrimination and the incidence of psychotic disorders among ethnic minorities in The Netherlands. *International Journal of Epidemiology*, 36(4), 761–768. [PubMed: 17517810]
- Veling W, & Susser E (2011, Jan). Migration and psychotic disorders. *Expert Review of Neurotherapeutics*, 11(1), 65–76. [PubMed: 21158556]

- Velthorst E, Nieman DH, Linszen D, Becker H, de Haan L, Dingemans PM, Birchwood M, Patterson P, Salokangas RK, & Heinimaa M (2010). Disability in people clinically at high risk of psychosis. *The British Journal of Psychiatry*, 197(4), 278–284. [PubMed: 20884950]
- Walder DJ, Holtzman CW, Addington J, Cadenhead K, Tsuang M, Cornblatt B, Cannon TD, McGlashan TH, Woods SW, Perkins DO, Seidman LJ, Heinssen R, & Walker EF (2013, Mar). Sexual dimorphisms and prediction of conversion in the NAPLS psychosis prodrome. *Schizophr Res*, 144(1-3), 43–50. [PubMed: 23340377]
- Walker EF, Trotman HD, Goulding SM, Holtzman CW, Ryan AT, McDonald A, Shapiro DI, & Brasfield JL (2013). Developmental mechanisms in the prodrome to psychosis. *Development and psychopathology*, 25(4 0 2), 1585. [PubMed: 24342857]
- Walker EF, Trotman HD, Pearce BD, Addington J, Cadenhead KS, Cornblatt BA, Heinssen R, Mathalon DH, Perkins DO, & Seidman LJ (2013). Cortisol levels and risk for psychosis: initial findings from the North American prodrome longitudinal study. *Biological psychiatry*, 74(6), 410–417. [PubMed: 23562006]
- Weiser M, Werbeloff N, Vishna T, Yoffe R, Lubin G, Shmushkevitch M, & Davidson M (2008). Elaboration on immigration and risk for schizophrenia. *Psychological medicine*, 38(8), 1113–1119. [PubMed: 17988415]
- Welch M, & Welch T (2007, Jun). Early psychosis in rural areas. *Australian and New Zealand Journal of Psychiatry*, 41(6), 485–494. [PubMed: 17508318]
- Whaley AL (2001a). Cultural Mistrust and Mental Health Services for African Americans: A Review and Meta-Analysis. *The Counseling Psychologist*, 29(4), 513–531.
- Whaley AL (2001b, Apr). Cultural mistrust of white mental health clinicians among African Americans with severe mental illness. *Am J Orthopsychiatry*, 71(2), 252–256. [PubMed: 11347366]
- Whitney DG, & Peterson MD (2019). US national and state-level prevalence of mental health disorders and disparities of mental health care use in children. *JAMA Pediatrics*, 173(4), 389–391. [PubMed: 30742204]
- Wigman J, van Winkel R, Raaijmakers QA, Ormel J, Verhulst FC, Reijneveld S, Van Os J, & Vollebergh WA (2011). Evidence for a persistent, environment-dependent and deteriorating subtype of subclinical psychotic experiences: a 6-year longitudinal general population study. *Psychological medicine*, 41(11), 2317–2329. [PubMed: 21477418]
- Willhite RK, Niendam TA, Bearden CE, Zinberg J, O'Brien MP, & Cannon TD (2008, Sep). Gender differences in symptoms, functioning and social support in patients at ultra-high risk for developing a psychotic disorder. *Schizophr Res*, 104(1-3), 237–245. [PubMed: 18573639]
- Wilson C, Smith ME, Thompson E, Demro C, Kline E, Bussell K, Pitts SC, DeVlylder J, Reeves G, & Schiffman J (2016, Mar). Context matters: The impact of neighborhood crime and paranoid symptoms on psychosis risk assessment. *Schizophr Res*, 171(1-3), 56–61. [PubMed: 26777883]
- Wong C, Davidson L, Anglin D, Link B, Gerson R, Malaspina D, McGlashan T, & Corcoran C (2009, May). Stigma in families of individuals in early stages of psychotic illness: family stigma and early psychosis. *Early Interv Psychiatry*, 3(2), 108–115. [PubMed: 19777087]
- Woods SW, Walsh BC, Powers AR, & McGlashan TH (2019). Reliability, validity, epidemiology, and cultural variation of the Structured Interview for Psychosis-Risk Syndromes (SIPS) and the Scale of Psychosis-Risk Symptoms (SOPS). In *Handbook of attenuated psychosis syndrome across cultures* (pp. 85–113). Springer.
- Wüsten C, & Lincoln TM (2017). The association of family functioning and psychosis proneness in five countries that differ in cultural values and family structures. *Psychiatry Research*, 253, 158–164. [PubMed: 28371682]
- Youman K, Drapalski A, Stuewig J, Bagley K, & Tangney J (2010). Race differences in psychopathology and disparities in treatment seeking: Community and jail-based treatment-seeking patterns. *Psychological Services*, 7(1), 11. [PubMed: 21814487]
- Zhang T, Wang J, & Woodberry KA (2020). Identifying Individuals at Clinical High Risk of Psychosis in Different Cultures and Countries. *Frontiers in psychiatry*, 11, 159. [PubMed: 32180746]