

RESEARCH

Open Access



BMSC-derived exosomes protect against kidney injury through regulating klotho in 5/6 nephrectomy rats

Feng Wan^{1,2}, Ru-chun Yang^{1,2*}, Yue-wen Tang¹, Xuan-li Tang^{1,2}, Tian Ye¹, Jie Zheng¹, Hua-qin Zhang^{1,2} and Yi Lin^{1,2}

Abstract

Aim: The aim of this study was to investigate the renoprotective effects of exosomes derived from rat bone marrow mesenchymal stem cells (rBMSCs) in a rat model of 5/6 nephrectomy (Nx)-induced chronic kidney disease (CKD).

Methods: A rat model of 5/6 Nx-induced CKD was established using conventional method. rBMSC-derived exosomes were isolated using ultracentrifugation and characterized. The exosomes were injected into 5/6 Nx rats through the caudal vein. After 12 weeks, 24 h proteinuria, serum creatinine (SCr), and blood urea nitrogen (BUN) levels were evaluated, and renal pathology was analyzed by H&E and Masson staining, and transmission electron microscopy. The expression of klotho was analyzed and the activity of the klotho promoter was evaluated using a luciferase reporter assay.

Results: The isolated exosomes showed typical morphological features. Exosomes transplantation reduced 24 h urinary protein excretion, and SCr and BUN levels in 5/6 Nx-induced CKD rats. Furthermore, renal pathology was improved in the exosome-treated 5/6 Nx rats. Mechanistically, the exosomes significantly upregulated the activity of klotho promoter and its expression.

Conclusions: Transplantation of rBMSC-derived exosomes may protect against kidney injury, probably by regulating klotho activity and expression. Our results provide a theoretical basis for the application of rBMSC-derived exosomes in CKD therapy.

Keywords: BMSC-derived exosome, Renoprotective, Klotho, 5/6 Nephrectomy rats

Introduction

Chronic kidney disease (CKD) is a global public health concern, and poses a substantial risk of progression to end-stage renal disease [1]. The first-line treatment for CKD is hormonal and immunosuppressant therapy (Additional file 1). However, its clinical use is limited due

to associated adverse effects and high incidence of hormone resistance [2]. Thus, there is an urgent need for a novel treatment strategy for CKD.

Mesenchymal stem cells (MSCs) are pluripotent stem cells that play important roles in treating various diseases, including CKD [3–5]. The mechanisms underlying therapeutic effects of bone marrow MSCs (BMSCs) are rather complex, with cell differentiation and paracrine signaling being part of these mechanisms [6]. As mediators of paracrine effects, exosomes carry active proteins and nucleic acids and have been proposed as alternative

*Correspondence: yangruchunhz@163.com

¹ Department of Nephrology, Hangzhou TCM Hospital Affiliated to Zhejiang Chinese Medical University, Tiyuchang Road 453, Hangzhou 31007, People's Republic of China

Full list of author information is available at the end of the article



therapeutic agents with broad prospects (Additional file 2).

Klotho is identified as an anti-aging gene and is highly expressed in kidney [7]. α -Klotho deficiency has been reported in animal models of CKD as well as in patients with CKD [7–9]. Thus, regulating klotho expression may be a promising strategy for treating CKD.

In this current study, we explored the effects of rat BMSC (rBMSC)-derived exosomes on kidney injury and regulating klotho expression in a rat model of 5/6 nephrectomy (5/6 Nx)-induced CKD. Our study outcomes provide new insights for devising therapeutic strategies for CKD in the future (Additional file 3).

Materials and methods

Exosome isolation and identification

rBMSCs were isolated from the bone marrow, as reported in our previous study [10]. The rBMSCs in passage 3–5 (P3–P5) were used for exosome extraction. Exosome-free medium was used to avoid interference from bovine exosomes [11, 12]. After 48 h culture, the supernatant was collected and the exosomes were isolated by sequential centrifugation at $300\times g$ for 10 min, $2000\times g$ for 20 min, and $10,000\times g$ for 30 min to remove cell debris. Next, the samples were centrifuged at $100,000\times g$ for 3 h. Subsequently, the pellets were resuspended in PBS and passed through a $0.22\ \mu\text{m}$ filter (Additional file 4). CD63 and CD9 were used as markers for exosomes identification. In addition, the size distribution and concentration of exosomes were analyzed using nanoparticle tracking analysis (NTA) by the Umibio Science and Technology Group (Shanghai). The characterized exosomes were either used immediately or stored at $-80\ ^\circ\text{C}$ until further analysis.

Animal study

Sprague–Dawley rats were purchased from the Zhejiang Academy of Medical Sciences and housed under standard conditions, as previously described [10]. After 1 week of acclimation, the rats were subjected to 5/6 nephrectomy as reported earlier [13] and were randomly divided into three groups: control group, 5/6-nephrectomized (5/6 Nx), and exosome group. The rats were then injected with BMSC-derived exosome ($1\times 10^7/\text{mL}$) through the caudal vein on days 30 and 45 (Additional file 5). At week 12, the rats were moved to metabolism cages for 24 h urine collection and urinary protein levels were quantified immediately by a biochemical analyser (HITACHI 7180) after centrifugation (Additional file 6). The rats were then sacrificed, and the blood and remnant kidney tissues were harvested for further analyses. The blood samples were centrifuged at $6000\times g$, $4\ ^\circ\text{C}$ for

10 min to obtain serum and stored at $-80\ ^\circ\text{C}$ for further experiments.

All the procedures were approved by the Institutional Animal Care and Use Committee of the Zhejiang Academy of Medical Sciences (No. ZJCLA-IACUC-20070015).

qRT-PCR

Total RNA was extracted from rat kidney tissues using TRIzol reagent, according to the manufacturer's protocol. cDNA was synthesized using a reverse transcription kit (RR036A, Takara, Japan). The primer sequences were as follows: Klotho-Forward: 5'-AACAACTTTCTTCTGCCCTATTTC-3', Klotho-Reverse: 5'-TGAGCGGTCACTAAGCGAAT-3', α -SMA-Forward: 5'-GCGTGGCTATTCCCTTCGTGACTAC-3', α -SMA-Reverse: 5'-CCATCAGGCAGTTCGTAGCTCTTC-3', desmin-Forward: 5'-AATGACCGCTTCGCCAACTACTTC-3', desmin-Reverse: 5'-GCTCTCGCATCTCCTCCTCGTAG-3', Vimentin-Forward: 5'-GTCCGTGTCCTCCTCCTAC-3', Vimentin-Reverse: 5'-TAGAGGCTGCGGCTAGTGCTG-3', GAPDH-Forward: 5'-ACCACAGTCCATGCCATCAC-3', GAPDH-Reverse: 5'-TCCACCACCCTGTTGCTGTA-3'. The mRNA expression levels were normalized to that of GAPDH.


H&E and Masson staining and TEM analysis

Freshly isolated kidney tissues were fixed in 4% paraformaldehyde, embedded in paraffin, cut into $3\ \mu\text{m}$ thick section, and stained with hematoxylin and eosin (H&E) and Masson's trichrome for kidney histological analysis. A semi-quantitative evaluation of renal tissues pathology was blindly accomplished by an expert pathologist. Masson's trichrome stained sections were used to evaluate interstitial tubular fibrosis and tubular atrophy and the scoring criteria was as follows: (0) denotes no change; grade (1) changes affecting $<25\%$ tubular damage (mild); grade (2) changes affecting $25\text{--}50\%$ of tubules (moderate); grade (3) changes affecting $>50\%$ of tubules (severe). Besides, H&E stained sections were used to evaluate the degree of glomerulosclerosis and the glomerular sclerosis index (GSI) was calculated as described previously [10].

For TEM, kidney tissues of $1\ \text{mm}^3$ were fixed in 2.5% glutaraldehyde for 6 h, rinsed three times in 0.1 M PBS, post-fixed with 1% osmium tetroxide for 1 h, dehydrated in graded solutions of acetone, and embedded in graded Epon 812. Subsequently, $80\text{--}100\ \text{nm}$ sections were cut and stained with uranyl acetate and lead citrate, and observed using a JEM-1400 transmission electron microscope (JEOL, Japan).

Immunohistochemical analysis

$3\ \mu\text{m}$ serial paraffin sections of the kidney were subjected to immunohistochemical staining. The sections



were treated with Tris–EDTA buffer (pH 9.0) for antigen retrieval and incubated by the primary antibodies, including rabbit polyclonal anti- α SMA (14395-1-AP, ProteinTech) (1:500), rabbit monoclonal anti-desmin (ab32362, abcam) (1:500), and rabbit monoclonal anti-vimentin (ab92547, abcam) (1:100). Thereafter, slides were incubated with HRP-conjugated anti-rabbit IgG Ab at 37 $^{\circ}\text{C}$ for 1 h. Finally, the slides were visualized by DAB (brown coloration) or AEC (red coloration) for 10 min

and counterstained with hematoxylin for 10 s. The staining results were obtained via light microscope.

Luciferase reporter gene assay

HEK293T cells were co-transfected with GV238-klotho-p-Luc (klotho promoter, NM_004795, ~2000 bp) and pRL-TK-luc at a 50:1 ratio using lip3000 according to the manufacturer's instructions. The transfected cells were treated with TGF- β 1 (1 ng/ μL), exosomes (0.4 mg),





Fig. 2 Effect of exosomes on renal function. The renal function indexes including 24 h proteinuria, serum creatinine (SCr), blood urea nitrogen (BUN), and ALB were examined by a biochemical analyzer. Data are shown as the mean ± SD. **p* < 0.05, ***p* < 0.01, *****p* < 0.0001

or co-cultured with BMSCs (5×10^4 /well) in a Transwell plate for 24 h. GW4869 (GLPBIO, Montclair, CA, USA) was used as an inhibitor of exosome biogenesis and release. The cells were then harvested and lysed to

determine firefly and Renilla luciferase activities using a dual luciferase reporter gene assay (Promega). Firefly luciferase activity was normalized to that of Renilla luciferase. The experiment was conducted three times.



Western blotting

Rat kidney tissues were lysed using RIPA buffer containing a proteinase inhibitor cocktail (Beyotime). The protein content of cell lysates was quantified using the BCA assay, separated by SDS-PAGE using 10% gels, transferred onto a nitrocellulose membrane, blocked with 5% skimmed milk, and then incubated with the primary antibodies specific for klotho (Abclonal, A12028) and GAPDH (ProteinTech, 60,004-1-Ig). Subsequently, the membranes were washed and incubated with infrared-labeled anti-rabbit/mouse IgG Ab (1:15,000). Finally, the signal was detected using an Odyssey CLx image system (LI-COR).

Statistical analysis

Data are presented as mean \pm standard deviation (SD). Graphs were constructed using the Prime GraphPad software (version 8.0). The data were analyzed using one-way analysis of variance (ANOVA), followed by Tukey's post hoc test for multiple comparisons. Statistical significance was set at $p < 0.05$.


Results

Characterization of exosomes extracted from rBMSCs

Morphologically, rBMSC-derived exosomes appeared round with a central depression and tested positive for its specific markers, CD63 and CD9 (Fig. 1A, B). The mean diameter of exosomes was approximately 130 nm in size, as determined using a ZetaView (Fig. 1C). These results indicated that the isolated exosomes had typical characteristics of exosomes.

rBMSCs-derived exosome improved the kidney function of 5/6 Nx rats

The 5/6 Nx rats exhibited increased 24 h proteinuria, and SCr and BUN levels compared to control rats. After exosome intervention, 24 h proteinuria and SCr and BUN levels decreased markedly compared with those in the untreated 5/6 Nx rats (Fig. 2). These data suggest that rBMSC-derived exosomes partially improved the kidney function.



rBMSCs-derived exosome improved renal pathological injury in 5/6 Nx rats

As shown in Fig. 3, control rats had well-opened glomerular capillary loops, closely arranged renal tubules, and an intact foot process. In contrast, 5/6 Nx rats exhibited glomerulosclerosis, mesangial cell proliferation, interstitial fibrosis, and inflammatory cell infiltration in the renal interstitium, partial fusion, and flattening of the foot process (Fig. 3). However, these lesions were relieved upon exosome treatment. In addition, immunochemical staining results showed that the expression levels of fibrotic markers such as α -SMA, desmin, and vimentin were significantly increased in the renal tubule region in 5/6 Nx rat (Fig. 4A). However, their expression levels were significantly decreased in the exosome treatment group. Similar results were observed at the mRNA level as evaluated by RT-qPCR (Fig. 4B). These results indicated that rBMSC-derived exosomes improved renal pathological injury in 5/6 Nx rats.


rBMSCs-derived exosome could regulate klotho expression

To evaluate the effect of exosomes on klotho activation, 293 T cells were co-transfected with the klotho-luc

reporter and pRL-TK, which encoded Renilla luciferase. The transfected cells were treated with TGF- β 1 for 24 h. Results showed that TGF- β 1 inhibited klotho activity; however, the activity was increased by both BMSC co-culture and exosome treatment group (Fig. 5A). In addition, klotho activity did not increase in the cells treated with both BMSC and exosome biogenesis/release inhibitor group (Fig. 5A). These results suggest that rBMSCs-derived exosomes could regulate the klotho promoter. Similar results were observed at the protein level (Fig. 5B). In addition, the mRNA and protein expression levels of klotho were markedly decreased in the kidneys of 5/6 rats, whereas their expression levels were upregulated in the rBMSC-derived exosome-treated group (Fig. 5C, D). Collectively, these results indicate that rBMSC-derived exosomes may regulate the activity and expression of klotho.

Discussion

In this study, we explored the effect of rBMSC-derived exosomes on kidney injury in 5/6 Nx rats. The 5/6 Nx rats treated with exosomes exhibited significantly decreased proteinuria, improved renal function, and inhibited renal



pathological features. Further experiment showed that exosomes could upregulate *klotho* expression by regulating the promoter activity of *klotho*. Our study outcomes add new information to this research field, thereby providing a theoretical basis for clinical treatment of CKD in the future.

MSCs have been widely explored in experimental models of CKD [14]. In mice with Streptozotocin (STZ)-induced type 1 diabetes, MSC transplantation prevented kidney injury and decreased albuminuria significantly [15]. Another study showed that BMSCs exerted protective effects in rats with diabetic nephropathy by reducing the number and inhibiting maturation of CD103⁺ DCs, thus regulating the CD8 T cell responses [5]. Human umbilical cord mesenchymal stem cells also improved the renal function and prevented the progression of STZ-induced diabetic nephropathy in rats [16]. Amnion-derived mesenchymal stem cells were reported to ameliorate CKD in a 5/6 Nx rat model [17]. Consistently, we previously reported that BMSCs attenuated the progression of renal damage in FSGS rats [10]. Collectively, these findings suggest that MSCs therapeutic approach is effective in animal studies. Interestingly, in our previous study, we observed that a significant number of fluorescence-labeled MSCs, injected via the caudal vein, did not reach the kidneys but were lodged in the lungs instead. These results indicated that the beneficial effects of MSCs may be mainly attributed to the paracrine mechanisms.

However, several challenges need to be overcome before MSCs can be applied for the treatment of kidney diseases. The technical difficulty in directly injecting BMSCs into the renal parenchyma or subcapsular space to exert their therapeutic effects is one of these challenges. Furthermore, biosafety remains a pertinent issue as transplanted cells may increase the risk of generating neoplasms. Thus, in terms of controlling and reducing risk, MSC-derived exosomes are promising alternative agents for treating kidney diseases.

As an “aging-suppressor” gene, *klotho* expression exhibits tissue specificity and is highly expressed in renal tubules [18]. Existing evidence reveals that abnormal *klotho* levels are regarded as the earliest biochemical abnormality of kidney disease [19–22]. Therefore, *klotho* is considered a key target for the treatment of kidney disease. Currently, the regulatory mechanism of *klotho* is complicated and under intensive investigation. Rodrigues et al. reported human umbilical cord-derived mesenchymal stromal cells treatment upregulated *klotho* protein expression in ischaemia/perfusion injury-induced renal senescence in rats [23]. We found BMSC could also upregulate *klotho* expression in our preliminary experiment (data not shown). Studies showed MSC-derived exosomes

contain many proteins, microRNAs, mRNAs, long non-coding RNAs, and phospholipids [24, 25]. Mehi et al. identified that there were many microRNA binding sites within the 3' untranslated region of *klotho* gene, such as microRNA-339, microRNA-556, microRNA-10, and microRNA-199 [26]. Similarly, many studies found other microRNAs could also regulate *klotho* [27–29]. We, therefore, hypothesized that microRNAs in exosomes might play an important role in regulating *klotho*. Here we found rBMSC-derived exosomes could ameliorate kidney function of 5/6 Nx rats, and the effect possibly was related to the modulation of *klotho* expression. However, the limit of this study is the lack of *klotho* knockout rat in our study to further validate the causal relationship of exosomes and *klotho*. We will continue to explore the accurate molecular mechanism in our further research.

In conclusion, we demonstrated that BMSCs-derived exosomes alleviated renal injury in 5/6 Nx rats probably by upregulating *klotho* activity and expression. These results provide new insight into stem cell therapy for nephropathy.

Abbreviations

MSCs: Mesenchymal stem cells; rBMSCs: Rat bone marrow mesenchymal stem cells; Nx: Nephrectomy; CKD: Chronic kidney disease; SCr: Serum creatinine; BUN: Blood urea nitrogen; NTA: Nanoparticle tracking analysis; STZ: Streptozotocin.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s40001-022-00742-8>.

Additional file 1. The western blot membrane for CD63 in Figure 1B.

Additional file 2. The western blot membrane for CD9 in Figure 1B.

Additional file 3. The western blot membrane for β -actin in Figure 1B.

Additional file 4. The western blot membrane for GAPDH in Figure 5B.

Additional file 5. The western blot membrane for *klotho* in Figure 5B.

Additional file 6. The western blot membranes for *klotho* and GAPDH in Figure 5D.

Acknowledgements

Not applicable.

Author contributions

FW and RCY contributed to the conception and design of the study. FW drafted the manuscript. YWT and HQZ conducted the animal experiment. XLT analyzed the renal pathology. TY and JZ performed the immunostaining and TEM. YL participated in the biochemical analyses. All authors read and approved the final manuscript.

Funding

This work was supported by grants from the Natural Science Foundation of Zhejiang Province (Y20H050004, LQ21H290005) and Public Welfare Technology Application Research Program of Zhejiang Province (LGC21H290002).

Availability of data and materials

The data sets generated and analyzed during the current study are available from the corresponding author on reasonable request.

Declarations**Ethics approval and consent to participate**

All animal procedures were approved by the Animal experiment Ethics Committee of Zhejiang Academy of Medical Sciences in accordance with the principles outlined in the NIH Guide for the Care and Use of Laboratory Animals.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Nephrology, Hangzhou TCM Hospital Affiliated to Zhejiang Chinese Medical University, Tiyuchang Road 453, Hangzhou 31007, People's Republic of China. ²Key Laboratory of Kidney Disease Prevention and Control Technology, Hangzhou, Zhejiang, China.

Received: 24 December 2021 Accepted: 27 June 2022

Published online: 11 July 2022

References

- Luyckx VA, Tuttle KR, Garcia-Garcia G, Gharbi MB, Heerspink HJL, Johnson DW, Liu ZH, Massy ZA, Moe O, Nelson RG, Sola L, Wheeler DC, White SL. Reducing major risk factors for chronic kidney disease. *Kidney Int Suppl* (2011). 2017; 7(2):71–87.
- Dogra S, Kaskel F. Steroid-resistant nephrotic syndrome: a persistent challenge for pediatric nephrology. *Pediatr Nephrol*. 2017;32(6):965–74.
- Wise AF, Ricardo SD. Mesenchymal stem cells in kidney inflammation and repair. *Nephrology (Carlton)*. 2012;17(1):1–10.
- Kuppe C, Kramann R. Role of mesenchymal stem cells in kidney injury and fibrosis. *Curr Opin Nephrol Hypertens*. 2016;25(4):372–7.
- Zhang F, Wang C, Wen X, Chen Y, Mao R, Cui D, Li L, Liu J, Chen Y, Cheng J, et al. Mesenchymal stem cells alleviate rat diabetic nephropathy by suppressing CD103(+) DCs-mediated CD8(+) T cell responses. *J Cell Mol Med*. 2020;24(10):5817–31.
- Morigi M, Introna M, Imberti B, Corna D, Abbate M, Rota C, Rottoli D, Benigni A, Perico N, Zoja C, et al. Human bone marrow mesenchymal stem cells accelerate recovery of acute renal injury and prolong survival in mice. *Stem Cells*. 2008;26(8):2075–82.
- Zou D, Wu W, He Y, Ma S, Gao J. The role of klotho in chronic kidney disease. *BMC Nephrol*. 2018;19(1):285.
- Neyra JA, Hu MC, Moe OW. Klotho in clinical nephrology: diagnostic and therapeutic implications. *Clin J Am Soc Nephrol*. 2020;16(1):162–76.
- Zhang XT, Wang G, Ye LF, Pu Y, Li RT, Liang J, Wang L, Lee KKH, Yang X. Baicalin reversal of DNA hypermethylation-associated Klotho suppression ameliorates renal injury in type 1 diabetic mouse model. *Cell Cycle*. 2020;19(23):3329–47.
- Yang RC, Zhu XL, Wang J, Wan F, Zhang HQ, Lin Y, Tang XL, Zhu B. Bone marrow mesenchymal stem cells attenuate the progression of focal segmental glomerulosclerosis in rat models. *BMC Nephrol*. 2018;19(1):335.
- Zhang X, Sai B, Wang F, Wang L, Wang Y, Zheng L, Li G, Tang J, Xiang J. Hypoxic BMSC-derived exosomal miRNAs promote metastasis of lung cancer cells via STAT3-induced EMT. *Mol Cancer*. 2019;18(1):40.
- Kamerkar S, LeBleu VS, Sugimoto H, Yang S, Ruivo CF, Melo SA, Lee JJ, Kalluri R. Exosomes facilitate therapeutic targeting of oncogenic KRAS in pancreatic cancer. *Nature*. 2017;546(7659):498–503.
- Ware K, Brodsky P, Satoskar AA, Nadasdy T, Nadasdy G, Wu H, Rovin BH, Bhatt U, Von Visger J, Hebert LA, et al. Warfarin-related nephropathy modeled by nephron reduction and excessive anticoagulation. *J Am Soc Nephrol*. 2011;22(10):1856–62.
- Lazzeri E, Romagnani P, Lasagni L. Stem cell therapy for kidney disease. *Expert Opin Biol Ther*. 2015;15(10):1455–68.
- Ezquer FE, Ezquer ME, Parrau DB, Carpio D, Yañez AJ, Conget PA. Systemic administration of multipotent mesenchymal stromal cells reverts hyperglycemia and prevents nephropathy in type 1 diabetic mice. *Biol Blood Marrow Transplant*. 2008;14(6):631–40.
- Xiang E, Han B, Zhang Q, Rao W, Wang Z, Chang C, Zhang Y, Tu C, Li C, Wu D. Human umbilical cord-derived mesenchymal stem cells prevent the progression of early diabetic nephropathy through inhibiting inflammation and fibrosis. *Stem Cell Res Ther*. 2020;11(1):336.
- Akan E, Cetinkaya B, Kipmen-Korgun D, Ozmen A, Koksoy S, Mendilcioglu I, Sakinci M, Suleymanlar G, Korgun ET. Effects of amnion derived mesenchymal stem cells on fibrosis in a 5/6 nephrectomy model in rats. *Biotech Histochem*. 2021;96(8):594–607.
- Kuro OM. The Klotho proteins in health and disease. *Nat Rev Nephrol*. 2019;15(1):27–44.
- Hu MC, Shi M, Zhang J, Quiñones H, Kuro-o M, Moe OW. Klotho deficiency is an early biomarker of renal ischemia-reperfusion injury and its replacement is protective. *Kidney Int*. 2010;78(12):1240–51.
- Lin Y, Kuro-o M, Sun Z. Genetic deficiency of anti-aging gene klotho exacerbates early nephropathy in STZ-induced diabetes in male mice. *Endocrinology*. 2013;154(10):3855–63.
- Hu MC, Moe OW. Klotho as a potential biomarker and therapy for acute kidney injury. *Nat Rev Nephrol*. 2012;8(7):423–9.
- Lee EY, Kim SS, Lee JS, Kim IJ, Song SH, Cha SK, Park KS, Kang JS, Chung CH. Soluble α -klotho as a novel biomarker in the early stage of nephropathy in patients with type 2 diabetes. *PLoS ONE*. 2014;9(8):e102984.
- Rodrigues CE, Capcha JM, de Bragança AC, Sanches TR, Gouveia PQ, de Oliveira PA, Malheiros DM, Volpini RA, Santinho MA, Santana BA, et al. Human umbilical cord-derived mesenchymal stromal cells protect against premature renal senescence resulting from oxidative stress in rats with acute kidney injury. *Stem Cell Res Ther*. 2017;8(1):19.
- van Balkom BWM, Gremmels H, Giebel B, Lim SK. Proteomic signature of mesenchymal stromal cell-derived small extracellular vesicles. *Proteomics*. 2019;19(1–2):e1800163.
- Liu H, Chen Y, Yin G, Xie Q. Therapeutic prospects of MicroRNAs carried by mesenchymal stem cells-derived extracellular vesicles in autoimmune diseases. *Life Sci*. 2021;277:119458.
- Mehi SJ, Maltare A, Abraham CR, King GD. MicroRNA-339 and microRNA-556 regulate Klotho expression in vitro. *Age (Dordr)*. 2014;36(1):141–9.
- Liang H, Yang K, Xin M, Liu X, Zhao L, Liu B, Wang J. MiR-130a protects against lipopolysaccharide-induced glomerular cell injury by upregulation of Klotho. *Pharmazie*. 2017;72(8):468–74.
- Ye H, Su B, Ni H, Li L, Chen X, You X, Zhang H. microRNA-199a may be involved in the pathogenesis of lupus nephritis via modulating the activation of NF- κ B by targeting Klotho. *Mol Immunol*. 2018;103:235–42.
- Li Y, Ren D, Xu G. Long noncoding RNA MALAT1 mediates high glucose-induced glomerular endothelial cell injury by epigenetically inhibiting klotho via methyltransferase G9a. *IUBMB Life*. 2019;71(7):873–81.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

