



A family cluster of schistosomiasis acquired in Solenzara River, Corsica (France) — Solenzara River is clearly a transmission site for schistosomiasis in Corsica

Nele Wellinghausen¹ · Hélène Moné² · Gabriel Mouahid² · Axel Nebel³ · Dennis Tappe⁴ · Martin Gabriel⁴

Received: 10 March 2022 / Accepted: 7 June 2022 / Published online: 18 June 2022
© The Author(s) 2022

Abstract

We report a patient with urogenital schistosomiasis and three cases of subclinical infection within one family acquired from Solenzara River, Corsica, in 2019. Our cases confirm that transmission of schistosomiasis in Corsica is ongoing and has been extended from the Cavu River to the Solenzara River. Solenzara River is clearly a transmission site for schistosomiasis in Corsica. Public health efforts are recommended to uncover and prevent further cases.

Keywords Schistosomiasis · Corsica · France · Solenzara River · *Schistosoma*

Introduction

Urogenital schistosomiasis is an emerging disease in Europe. In 2014, a first case of *Schistosoma haematobium* infection in a German boy who acquired infection in 2013 in the Cavu River in Southern Corsica (France) has been reported (Holtfreter et al. 2014), closely followed by another case in a French girl acquired in the same river (Berry et al. 2014). Surveillance studies later revealed 106 documented cases associated with the Cavu River (Noel et al. 2018). From 2015 to 2018 also, sporadic infections acquired from the Cavu River have been recorded (Ramalli et al. 2018). Recently, a first case of urogenital schistosomiasis transmitted in 2019 from a different river in Corsica, likely but not

surely, the Solenzara River, has been published in a German traveller (Rothe et al. 2021). Here, we report a family cluster of schistosomiasis clearly linked to the Solenzara River in the same year.

Case report

A 19-year-old formerly healthy young man presented to his urologist in July 2021 with macrohematuria for 2–3 weeks. Ultrasound and cystoscopy revealed a tumor of 1.5 to 2 cm in diameter on the left upper bladder wall protruding into the bladder cavity. The tumor was completely resected by transurethral resection, and histology showed pronounced granulomatous inflammation and suspicion of egg fragments suggestive of schistosomiasis. Urine microscopy revealed eggs with typical morphology of *S. haematobium*. In addition, elevated *Schistosoma*-specific IgG antibodies were detected in serum by enzyme-linked immunoassay (ELISA) with an index of 60 (cutoff index < 9, DRG Diagnostics, Springfield, NJ, USA) and by indirect immunofluorescence test (IIFT) with a titer > 1:160 (cutoff titer < 1:20, Euroimmun, Lübeck, Germany). The patient was treated with praziquantel for 3 days and fully recovered. Shortly after therapy, polymerase chain reaction targeting a nuclear repeated sequence was performed on the patient serum; for methods see Abbasi et al. (2007), Hamburger et al. (2001), and Moné et al. (2015). The alignment of the obtained, albeit short, sequence of the patient (90 bp) with those of *S.*

Section Editor: Elizabeth Marie Warburton

✉ Nele Wellinghausen
nele.wellinghausen@labor-gaertner.de

¹ MVZ Labor Ravensburg, Elisabethenstr. 11, 88212 Ravensburg, Germany

² UMR 5244 IHPE Interactions Hôtes-Pathogènes-Environnements, Université de Montpellier, CNRS, IFREMER, Université de Perpignan, 66860 Perpignan, France

³ Ravensburg, Germany

⁴ National Reference Centre for Tropical Pathogens, Bernhard Nocht Institute for Tropical Medicine, Hamburg, Germany

Discussion — ongoing endemicity of schistosomiasis in Corsica

After the first published case of schistosomiasis in 2014 and detection of several further patients in Corsica affecting the Cavu River, only sporadic cases were recorded in the following years. Recently, a first case of urogenital schistosomiasis was reported that was acquired in 2019 from the Solenzara River situated north of the Cavu River (Rothe et al. 2021). Interestingly, the causative *Schistosoma* species in both the Cavu and Solenzara Rivers was a hybrid between *Schistosoma haematobium* and *S. bovis* (Moné et al. 2015). Phylogenetic analyses showed that this strain was previously imported from Senegal (Boissier et al. 2016) or Benin (Savassi et al. 2020). The waters of the Solenzara River do not intermingle with those of the Cavu River but the intermediate vector, *Bulinus truncatus* snails, have been detected earlier in the Solenzara River during environmental surveys (Mulero et al. 2019). Apart from contamination of the river by an infected person, animal reservoirs have been discussed as a possible source of transmission of schistosomes in Corsica (Oleaga et al. 2019). However, an animal source has not been proven yet (Oleaga et al. 2019), and transmission by an infected person appears, thus, the most likely source of transmission to the Solenzara River. Broad thermal range of schistosomes indicates that they can overwinter under temperate climates like in Corsica (Mulero et al. 2019) and also in infected humans. In addition, *Bulinus* snails can overwinter in the rivers. Infected persons may thus reinfect the rivers in spring and may lead to new cases in late summer, as observed in our family and in other published cases.

Patient history of the family reported here revealed that all four cases did only have water contact in the Solenzara River during two bathing events, and that they avoided all other rivers in Corsica. The bathing site in the Solenzara River (GPS coordinates 41°50'37.8"N 9°20'36.3"E) was situated close to a restaurant next to a campsite and corresponds to one of the possible sources of infection of the first patient published by Rothe et al. (2021). Infection of our patients occurred 1 to 2 weeks after the infection of the abovementioned patient in August/September 2019.

Conclusion

Our cases confirm that transmission of schistosomiasis in Corsica is ongoing and has been extended from the Cavu River to the Solenzara River. The fact that all family members acquired infection after two bathing events indicates a relevant risk of infection from the Solenzara River in

August/September 2019. Public health awareness should be increased and epidemiological survey investigating the actual infection risk from the Solenzara River should be initiated. In addition, residents and tourists should be informed about the possible risk of transmission of schistosomiasis from the Solenzara River and possible unknown cases should be sought.

Acknowledgements We thank Birgit Muntau for excellent technical work regarding the PCR and NGS analyses.

Declarations

Conflict of interest The authors declare no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Abbasi I, King CH, Sturrock RF, Kariuki C, Muchiri E, Hamburger J (2007) Differentiation of *Schistosoma haematobium* from related schistosomes by PCR amplifying an inter-repeat sequence. *Am J Trop Med Hyg* 76:950–955
- Berry A, Moné H, Iriart X, Mouahid G, Aboo O, Boissier J, Fillaux J, Cassaing S, Debuissou C, Valentin A, Mitta G, Theron A, Magneval JF (2014) Schistosomiasis haematobium, Corsica, France. *Emerg Infect Dis* 20:1595–1597
- Boissier J, Grech-Angelini S, Webster BL, Allienne JF, Huysse T, Mas-Coma S, Toulza E, BarrÃ©-Cardi H, Rollinson D, Kincaid-Smith J, Oleaga A, Galinier R, Foata J, Rognon A, Berry A, Mouahid G, Henneron R, MonÃ© H, Noel H, Mitta G (2016) Outbreak of urogenital schistosomiasis in Corsica (France): an epidemiological case study. *Lancet Infect Dis* 16:971–979
- Hamburger J, He N, Abbasi I, Ramzy RM, Jourdan J, Ruppel A (2001) Polymerase chain reaction assay based on a highly repeated sequence of *Schistosoma haematobium*: a potential tool for monitoring schistosome-infested water. *Am J Trop Med Hyg* 65:907–911
- Holtfreter MC, Moné H, Müller-Stöver I, Mouahid G, Richter J (2014) *Schistosoma haematobium* infections acquired in Corsica, France, August 2013. *Euro Surveill* 19
- Moné H, Holtfreter MC, Allienne JF, Mintsu-Nguema R, Ibikounle M, Boissier J, Berry A, Mitta G, Richter J, Mouahid G (2015) Introgressive hybridizations of *Schistosoma haematobium* by *Schistosoma bovis* at the origin of the first case report of schistosomiasis in Corsica (France, Europe). *Parasitol Res* 114:4127–4133
- Mulero S, Rey O, Arancibia N, Mas-Coma S, Boissier J (2019) Persistent establishment of a tropical disease in Europe: the preadaptation of schistosomes to overwinter. *Parasit Vectors* 12:379

- Noel H, Ruello M, Maccary A, Pelat C, Sommen C, Boissier J, Barré-Cardi H, Fillaux J, Termignon JL, Debruyne M, Chyderiotis G, Durand J, Ramalli L, Chiappini JD, Malfait P, Berry A (2018) Large outbreak of urogenital schistosomiasis acquired in Southern Corsica, France: monitoring early signs of endemicization? *Clin Microbiol Infect* 24:295–300
- Oleaga A, Rey O, Polack B, Grech-Angelini S, Quilichini Y, Perez-Sanchez R, Boireau P, Mulero S, Brunet A, Rognon A, Vallue I, Kincaid-Smith J, Allienne JF, Boissier J (2019) Epidemiological surveillance of schistosomiasis outbreak in Corsica (France): are animal reservoir hosts implicated in local transmission? *PLoS Negl Trop Dis* 13:e0007543
- Ramalli L, Mulero S, Noel H, Chiappini JD, Vincent J, Barré-Cardi H, Malfait P, Normand G, Busato F, Gendrin V, Allienne JF, Fillaux J, Boissier J, Berry A (2018) Persistence of schistosomal transmission linked to the Cavu river in southern Corsica since 2013. *Euro Surveill* 23
- Rothe C, Zimmer T, Schunk M, Wallrauch C, Helfrich K, Gueltekin F, Bretzel G, Allienne JF, Boissier J (2021) Developing endemicity of schistosomiasis, Corsica, France. *Emerg Infect Dis* 27:319–321
- Savassi BAES, Mouahid G, Lasica C, Mahaman SK, Garcia A, Courtin D, Allienne JF, Ibikounlé M, Moné H (2020) Cattle as natural host for *Schistosoma haematobium* (Bilharz, 1852) Weinland, 1858 x *Schistosoma bovis* Sonsino, 1876 interactions, with new cercarial emergence and genetic patterns. *Parasitol Res* 119:2189–2205

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.