



HHS Public Access

Author manuscript

Sci Transl Med. Author manuscript; available in PMC 2022 December 01.

Published in final edited form as:

Sci Transl Med. 2021 December ; 13(622): eabe7430. doi:10.1126/scitranslmed.abe7430.

Repeated *Plasmodium falciparum* infection in humans drives the clonal expansion of an adaptive $\gamma\delta$ T cell repertoire

Anouk von Borstel¹, Priyanka Chevour¹, Daniel Arsovski¹, Jelte M. M. Kroil^{2,3}, Lauren J. Howson^{1,†}, Andrea A. Berry⁴, Cheryl L. Day⁵, Paul Ogongo^{6,7}, Joel D. Ernst⁶, Effie Y. H. Nomicos⁸, Justin A. Boddey^{2,3}, Edward Giles⁹, Jamie Rossjohn^{1,10,11}, Boubacar Traore¹², Kirsten E. Lyke^{4,#}, Kim C. Williamson^{13,#}, Peter D. Crompton^{14,#}, Martin S. Davey^{1,*}

¹Infection and Immunity Program and Department of Biochemistry and Molecular Biology, Biomedicine Discovery Institute, Monash University, Clayton, Victoria 3800, Australia

²The Walter and Eliza Hall Institute of Medical Research, Parkville, VIC 3052, Australia

³University of Melbourne, Melbourne, VIC 3010, Australia

⁴Vaccine Development and Global Health, University of Maryland School of Medicine, Baltimore, MD, USA

⁵Department of Microbiology and Immunology, Emory Vaccine Center and Yerkes National Primate Research Center, Emory University, Atlanta, Georgia, USA

⁶Division of Experimental Medicine, Department of Medicine, UCSF School of Medicine, San Francisco, California, USA

⁷Department of Tropical and Infectious Diseases, Institute of Primate Research, National Museums of Kenya, P.O Box 24481 - 00502, Nairobi, Kenya

⁸Parasitology and International Programs Branch, Division of Microbiology and Infectious Diseases, NIAID, NIH, Bethesda, MD, USA

⁹Department of Paediatrics, Monash University, and Centre for Innate Immunity and Infectious Disease, Hudson Institute of Medicine, Clayton, Victoria 3168, Australia

¹⁰Australian Research Council Centre of Excellence in Advanced Molecular Imaging, Monash University, Clayton, Victoria 3800, Australia

¹¹Institute of Infection and Immunity, Cardiff University School of Medicine, Heath Park, CF14 4XN Cardiff, United Kingdom

^{*}To whom correspondence should be addressed: martin.davey@monash.edu.

[†]Present address: Immunology Division, Walter and Eliza Hall Institute of Medical Research, Parkville, VIC, Australia

Author contributions: Conceptualization and study design, K.E.L., K.C.W., P.D.C., and M.S.D.; Design of Experiments, A.v.B. and M.S.D.; Investigation and analysis, A.v.B., P.C., D.A., L.J.H., and M.S.D.; Resources, C.L.D., P.O., J.D.E., J.M.M.K, J.A.B., E.G., B.T. and P.D.C.; Repeated controlled human malaria infection trial (NCT03014258) funding acquisition, administration and implementation, A.A.B., E.Y.H.M., K.E.L. and K.C.W.; Project administration, J.R., B.T., E.Y.H.M, K.E.L., K.C.W., P.D.C. and M.S.D; A.v.B. and M.S.D. wrote the draft, A.v.B., K.E.L., K.C.W., P.D.C. and M.S.D. wrote the final manuscript and all authors provided critical review; Supervision, J.R. and M.S.D.

[#]Contributed equally

Publisher's Disclaimer: Disclaimers: The opinions and assertions expressed herein are those of the author(s) and do not necessarily reflect the official policy or position of the Uniformed Services University or the Department of Defense.

Competing interests: The authors declare no competing interests.

¹²Malaria Research and Training Center, Department of Epidemiology of Parasitic Diseases, International Center of Excellence in Research, University of Sciences, Techniques and Technologies of Bamako, Bamako, Mali

¹³Department of Microbiology and Immunology, Uniformed Services University of the Health Sciences, Bethesda, MD, USA

¹⁴Malaria Infection Biology and Immunity Section, Laboratory of Immunogenetics, National Institute of Allergy and Infectious Disease, National Institute of Health, Rockville, MD, USA

Abstract

Repeated *Plasmodium falciparum* infections drive the development of clinical immunity to malaria in humans, however, the immunological mechanisms that underpin this response are only partially understood. Here, we investigated the impact of repeated *P. falciparum* infections on human $\gamma\delta$ T cells in the context of natural infection in Malian children and adults, as well as serial controlled human malaria infection (CHMI) of U.S. adults, some of whom became clinically immune to malaria. In contrast to the predominant V δ 2⁺ $\gamma\delta$ T cell population in malaria-naïve Australian individuals, clonally expanded cytotoxic-V δ 1_{effector} T cells were enriched in the $\gamma\delta$ T cell compartment of Malian subjects. Malaria-naïve U.S. adults exposed to four sequential CHMIs defined the precise impact of *P. falciparum* on the $\gamma\delta$ T cell repertoire. Specifically, innate-like V δ 2⁺ $\gamma\delta$ T cells exhibited an initial robust polyclonal response to *P. falciparum* infection that was not sustained with repeated infections, whereas V δ 1⁺ $\gamma\delta$ T cell frequencies increased in frequency with repeated infections. Moreover, repeated *P. falciparum* infection drove waves of clonal selection in the V δ 1⁺ TCR repertoire that coincided with the differentiation of V δ 1_{naïve} cells into cytotoxic-V δ 1_{effector} cells. Finally, V δ 1⁺ T cells of malaria-exposed Malian and U.S. individuals were now licensed for reactivity to *P. falciparum* parasites *in vitro*. Together, our study indicates that repeated *P. falciparum* infection drives the clonal expansion of an adaptive $\gamma\delta$ T cell repertoire and establishes a role for V δ 1⁺ T cells in the human immune response to malaria.

One Sentence Summary:

Malaria drives the adaptive differentiation of the human $\gamma\delta$ T cell repertoire.

Introduction

In malaria-endemic regions, non-sterilizing clinical immunity to blood-stage *Plasmodium falciparum* parasites can be acquired, but this typically only occurs after many years of repeated infections (1). However, the mechanisms underlying this protection are only partially understood (2, 3). Recent observational studies in malaria-endemic areas, as well as clinical trials of attenuated *P. falciparum* sporozoite vaccine *PfSPZ*, have suggested that $\gamma\delta$ T cells may contribute to protection from malaria (4–7).

Human $\gamma\delta$ T cells are an unconventional T cell population that are thought to play an important role in immunity to microbial pathogens and cancer (8). Unlike conventional $\alpha\beta$ T cells, $\gamma\delta$ T cells are not restricted by classical MHC or MHC-I-like molecules to recognize antigens (9–11), but instead respond directly to non-peptidic metabolite antigens and other diverse ligands (12, 13). $\gamma\delta$ T cells were present in the first jawed vertebrates and

co-evolved with pathogenic organisms for millions of years (10, 14). In humans, the major peripheral blood population of $\gamma\delta$ T cells (5–10% of total T cells) express a restricted TCR that consists of paired V δ 2 and V γ 9 chains (15). The V γ 9/V δ 2⁺ T cell population directly responds to a prenyl-pyrophosphate metabolite (PAg) (E)-4-Hydroxy-3-methyl-but-2-enyl pyrophosphate (HMB-PP) produced by the microbial non-mevalonate pathway (16). The V γ 9/V δ 2⁺ TCR repertoire is generated early in gestation and is shaped soon after birth, with a high frequency of public V γ 9 clonotypes (17–19). Innate-like V γ 9/V δ 2⁺ T cells can expand and comprise up to 40% of T cells during blood-stage malaria infection, a response thought to be driven by recognition of *P. falciparum*-derived HMB-PP (20–22) and they may participate in limiting parasite replication by targeting *P. falciparum* blood-stage parasites through granulysin-dependent cytotoxicity (23) and phagocytosis of antibody-coated iRBCs (24).

In contrast to innate-like V γ 9/V δ 2⁺ $\gamma\delta$ T cells, a diverse biology has been established for $\gamma\delta$ T cells that predominantly express the V δ 1⁺ TCR chain and circulate in blood at low frequency but are a major population in peripheral tissues (25). Firstly, V δ 1⁺ T cells that form tissue-associated populations in the intraepithelial lymphocyte (IEL) compartment of gut and breast tissue are thought to provide innate-like immune surveillance through host-encoded Natural Killer-receptors (NKR) (26) and BTN-like (BTNL) 3 proteins (27). Secondly, peripheral blood and liver-resident V δ 1⁺ T cells possess hallmarks of adaptive T cells and comprise naïve-like (V δ 1_{naïve}) and effector (V δ 1_{effector}) populations with diverse or highly focused TCR repertoires, respectively (19, 25, 28). Acute cytomegalovirus (CMV) infection has been associated with the selection of a limited set of V δ 2^{neg} $\gamma\delta$ TCR clonotypes (19, 29). Interestingly, expanded populations of V δ 1⁺ T cells have been observed in both children and adults with symptomatic *P. falciparum* infection (30–32) and in individuals residing in regions of malaria transmission (33, 34). Despite evidence that $\gamma\delta$ T cells contribute to immunity to microbial pathogens, it remains unclear whether *P. falciparum* infection per se or factors associated with malaria transmission in endemic areas are responsible for the expansion of V δ 1⁺ T cells. Moreover, it is also unclear the impact of repeated *P. falciparum* infection on the phenotype, function and clonality of the $\gamma\delta$ T cell compartment.

In this study, we investigated the $\gamma\delta$ T cell immune repertoire response to *P. falciparum* malaria in a cohort of children and adults residing in a malaria-endemic region of Mali, and in malaria-naïve U.S. adults serially infected with *P. falciparum* via mosquito bite in a controlled setting. We found that repeated *P. falciparum* infections drove the clonal selection and expansion of circulating cytotoxic V δ 1_{effector} T cells that reacted to *P. falciparum* blood-stage parasites.

Results

Heterogeneity in the $\gamma\delta$ T cell compartment exists across diverse geographic locations

In general, immune profiles are known to differ between children of high- and low-income countries where the latter typically suffer a disproportionately high burden of infectious disease (35). Here, we compared the circulating $\gamma\delta$ T cell repertoire of Malian children (aged 4 – 17 years) who are exposed to intense seasonal malaria transmission (36), with

that of age and gender matched children from Melbourne, Australia (aged 1 – 17 years) (Table S1). We first analyzed Mali samples collected from uninfected subjects at the end of the dry season when malaria transmission is negligible to assess $\gamma\delta$ T cell repertoires in a relatively unperturbed state. We found that $\gamma\delta$ T cell and $V\delta 1^+$ T cell frequencies were significantly higher in Malian children (Fig. 1A and S1A), whereas the frequency of $V\gamma 9/V\delta 2^+$ T cells were similar between both groups (Fig. 1A and B). We then analyzed $\gamma\delta$ T cells in Malian adults (aged 21 – 26 years) and adults residing in an area of low malaria transmission in Kenya (aged 26 – 49 years) as well as Australian adults with no history of malaria exposure (aged 20 – 71 years). $V\delta 1^+$ T cell frequencies were lower in Kenyan and Australian adults compared to Malian children (Fig. S1B). Next, we assessed $\gamma\delta$ T cell effector subsets in Malian children. From birth, $V\gamma 9/V\delta 2^+$ T cells typically form a stable innate-like T cell population composed of a $CD27^+ CD28^+$ Granzyme (Gzm) $A^+ GzmB^+$ Perforin $^+$ compartment (18, 19, 37). In Malian individuals we found that $V\gamma 9/V\delta 2^+$ T cells had reduced expression of $CD27^+ CD28^+$ (Fig. 1C) and lytic Perforin (Fig. 1D), while GzmA was increased (Fig. 1E and F). In contrast, cord blood $V\delta 1^+$ T cell population is typically composed of naïve-like $CD27^{hi} CX_3CR1^{neg} GzmA/B^{neg}$ Perforin neg cells ($V\delta 1_{naive}$) (28). However, the $V\delta 1^+$ compartment in Malian subjects was predominantly composed of $CD27^{lo} CX_3CR1^+ GzmA/B^+$ Perforin $^+$ effector-like cells ($V\delta 1_{eff}$) (Fig. 1C–F). Interestingly, a $CD16^+ V\gamma 9/V\delta 2^+$ T cell compartment has recently been implicated in antibody-mediated phagocytosis of iRBCs (24). We found that Malian children, when compared to Australian children, tended to have increased frequencies of $CD16^+ V\delta 1^+$ T cells rather than $CD16^+ V\gamma 9/V\delta 2^+$ T cells (Fig. S1C). Together, these data suggest that the composition of the $\gamma\delta$ T cell compartment varies significantly across geographic locations. However, it remained unclear whether high malaria transmission and/or factors associated with malaria transmission drive the proportional expansion of $V\delta 1^+$ T cells and skewing towards a $V\delta 1_{eff}$ phenotype in the Mali cohort.

Episodes of febrile malaria associate with fluctuations in $V\delta 1^+$ $\gamma\delta$ T cell frequencies

To more directly investigate the potential impact of natural malaria infection on the $\gamma\delta$ T cell compartment, we conducted a longitudinal analysis of nine Malian children (aged 8 – 14 years) over three malaria seasons (Fig. 1G). These individuals from Mali were exposed to an annual six-month malaria season in which *P. falciparum* transmission is intense and predictable (36). Consistent with this, *P. falciparum* parasite density increased during each malaria season in a subset of five children whose blood smears were examined longitudinally (Fig. S1CD). Moreover, each subject was selected because they experienced two to three febrile malaria episodes over multiple years, as detected by both passive and active clinical surveillance, allowing for longitudinal analyses of $\gamma\delta$ T cell dynamics in response to acute symptomatic malaria followed by sustained periods without febrile malaria (Fig. 1G). We then investigated $\gamma\delta$ T cell and $CD8^+ \alpha\beta$ T cell frequencies across consecutive episodes of febrile malaria over three seasons, these analyses pooled T cell frequencies from children who had experienced two or three episodes of febrile malaria (Fig. 1H). $\gamma\delta$ T cell and $CD8^+ \alpha\beta$ T cell frequencies were assessed in blood samples collected on the day febrile malaria was diagnosed and again within 3 – 6 months of diagnosis. We found that $V\gamma 9/V\delta 2^+$ T cell frequencies increased after febrile malaria in year 1 but did not consistently change after febrile malaria in years 2 and 3 (Fig. 1H),

although these observations could be due to the different sampling times in each year. CD8⁺ T cell frequencies were unchanged after each febrile episode (Fig. 1H). In contrast, Vδ1⁺ T cell frequencies were consistently decreased upon presentation with febrile malaria and increased after each febrile malaria episode across all three years (Fig. 1H). Moreover, across a sub-set of eight subjects in year 1, we also observed equivalent CD3⁺ lymphocyte and αβ T cell counts, and all γδ T cell populations expanded in number after febrile malaria (Fig. S1E). We then assessed a subset of children at timepoints without infection before a documented period of asymptomatic *P. falciparum* infection but no febrile malaria episodes (Fig. 1I; pooled from data between month 12 – 19 or 24 – 30). Vγ9/Vδ2⁺ and Vδ1⁺ T cell frequencies did not change significantly across this six to seven-month period. We noted previously that Vδ1⁺ T cells in Malian children were predominantly composed of Vδ1_{effector} cells (Fig. 1C and D), however, yearly episodes of febrile malaria had little impact on Vδ1_{effector} frequencies and CD27⁺ CD28⁺ Vγ9/Vδ2⁺ T cell frequencies were reduced in year 3 (Fig. S1F). Together, these data indicate that exposure to seasonal episodes of febrile malaria transiently impacts circulating frequencies of Vδ1⁺ γδ T cells.

Malian γδTCR repertoires are clonally skewed and change after febrile malaria

We next explored the underlying γδTCR repertoires in Malian children and whether febrile malaria could impact individual clonotypes over time. Initially, we conducted a cross-sectional analysis of blood samples collected subjects during periods of no malaria transmission (subjects 066, 521, 766) and from one subject with febrile malaria (subject 269) and compared these repertoires to those of Australian children (Fig. 2A and S3A). We analyzed both Vγ9/Vδ2⁺ (Vδ2⁺) and non-Vγ9/Vδ2 (Vδ2^{neg}) γδ T cell populations, effectively encompassing the total γδ T cell repertoire (Fig. S2). Phenotypically, in both Malian and Australian subjects Vδ2⁺ γδ T cell populations were composed of effector-like populations of CD27⁺ CD28⁺ cells and Vδ2^{neg} γδ T cells were composed CD27^{lo} CX₃CR1⁺ effector cells in Malian subjects and CD27^{hi} CX₃CR1^{neg} naïve cells in Australian subjects (Fig. S3B). The Vγ9/Vδ2⁺ T cell subset displayed γδ TCR repertoires consistent with those seen in children and adults from Europe (18, 19, 37) (Fig. 2A and Fig. S3A), which are almost exclusively composed of Vδ2–Jδ1 (Fig. S3C) paired to Vγ9–JγP (Fig. S3D), with diverse clonotype composition and common CDR3γ9–JγP sequences shared between individuals (Fig. 2A and Fig. S3C). Vδ2^{neg} γδ TCR repertoires were predominantly composed of Vδ1–Jδ1 sequences (Fig. S3D) that were paired to various Vγ–Jγ1/2 regions (Fig. S3D). These Vδ2^{neg} γδTCR repertoires from Malian children exhibited expanded clonotypes, indicated by an increase in the accumulated frequency of the top 10 clonotypes in comparison to Vδ2^{neg} γδ TCR repertoires in Australian children (Fig. 2B). In support of the skew towards expanded Vδ1 clonotypes, Malian Vδ2^{neg} γδ TCR repertoires also showed a reduced diversity of clonotype composition (Fig. 2C) and a reduced frequency of shared sequences compared to Vδ2^{neg} γδTCR repertoires of Australian individuals (Fig. 2D). These data suggest that the Vγ9/Vδ2⁺ T cell repertoires in Malian subjects are highly similar to those of Australian individuals. In contrast, Vδ2^{neg} γδTCR repertoires of Malian individuals showed evidence of reduced clonotype sharing and diversity as a result of expanded private clonotypes.

Next, in a longitudinal analysis we assessed the impact of episodes of acute febrile malaria on $\gamma\delta$ TCR clonotype composition within the $V\gamma 9/V\delta 2^+$ and $V\delta 2^{\text{neg}}$ $\gamma\delta$ T cell populations. $V\gamma 9/V\delta 2^+$ clonotypes remained remarkably stable during and after acute febrile malaria (Fig. 2E, Fig. S3E and F). We and others have previously reported on the stability of $V\delta 2^{\text{neg}}$ and $V\delta 1^+$ $\gamma\delta$ TCR clonotypes over several years (18, 19, 28, 29). Here, $V\delta 2^{\text{neg}}$ $\gamma\delta$ TCR repertoires displayed changes after acute febrile malaria, characterized by contraction and expansion of existing clonotypes or emergence of new prevalent sequences (Fig. 2F, Fig. S3E and G). These changes impacted the frequency of $V\delta 1$ sequence usage (Fig. S3H), $V\gamma 2$ usage (Fig. 2G), the overall repertoire diversity (Fig. 2G) and nucleotide length dynamics (Fig. S3I). To explore the impact of febrile malaria on clonotype composition within $V\delta 1_{\text{effector}}$ cells, we sorted single cells from the $CD27^{\text{lo}} CX_3CR1^+ V\delta 1_{\text{effector}}$ cell compartment from samples collected over 32 months and three separate acute febrile malaria episodes from subject 179 (Fig. 2H). Interestingly, we noted by flow cytometry that $V\delta 1/\gamma\delta$ TCR antibody staining intensity changed over time, with distinct $V\delta 1/\gamma\delta$ TCR antibody populations emerging after each episode of febrile malaria (Fig. 2H and Fig. S3J). Underpinning these observations, single cell $\gamma\delta$ TCR sequencing revealed changes in the frequency and identity of individual $V\delta 1_{\text{effector}}$ clonotypes over time (Fig. 2I and Fig. S3K). Together, these data suggest that Malian individuals have highly stable $V\gamma 9/V\delta 2^+$ T cell repertoires that are retained across episodes of febrile malaria and are shared between individuals. In contrast, $\gamma\delta$ TCR clonotypes in the $V\delta 1_{\text{effector}}$ compartment were composed of clonotypes that varied in frequency and identity over time.

Repeated human controlled malaria infection can establish clinical immunity that correlates with increased $V\delta 1^+$ $\gamma\delta$ T cell frequencies

To understand the precise impact of *P. falciparum* infection on the trajectory of $\gamma\delta$ T cell development and selection, we assessed $\gamma\delta$ T cell subset dynamics in PBMCs collected from five malaria-naïve adults voluntarily exposed to repeated controlled human malaria infection (CHMI). Each volunteer was exposed to the bites of five *Anopheles stephensi* mosquitos infected with *P. falciparum* (strain: NF54) on four separate occasions over 644 days (Fig. 3A). Symptomatic malaria occurs during the blood stage of the *P. falciparum* parasite life cycle, which typically develops after an incubation period of nine to fourteen days (36). Here, we analyzed samples at baseline (malaria naïve), immediately prior to *P. falciparum* infection (day 1; at CHMI1 and 3), and 21 days after infection for all CHMIs (Fig. 3A). We did not observe any noticeable leukopenia measured by white blood cell counts (at day 1 or day 28; Fig. S4A) or by clinical tests prior to apheresis (day 21) at the timepoints sampled in this study. Over the course of the four CHMIs, peak parasitemia measured by blood smear did not significantly change (Fig. S4B). We then assessed the instances of febrile malaria and symptomatic disease (ranging from headaches to vomiting; Table S2). Fever was observed at CHMI1 or 2 in all but one individual and the number of symptoms observed in each individual decreased after repeated CHMIs (Fig. 3B). Three individuals had asymptomatic *P. falciparum* infections following CHMI4, while two volunteers remained symptomatic (Fig. 3B). Next, we analyzed $\gamma\delta$ and $\alpha\beta$ T cell frequencies across all CHMIs. Total $\alpha\beta$ T cell frequencies within $CD3^+$ T cells showed a non-significant decline with repeated CHMI (Fig. 3C). $CD8^+$ $\alpha\beta$ T cells increased in frequency and peaked prior to CHMI3 (Fig. 3C), coinciding with an increase in $CD8^+$

T_{naive} cells and $CD8^+ T_{CM}$ on day 21 after CHMI2–4 (Fig. S4C). In contrast, $\gamma\delta$ T cells frequencies increased across all CHMIs, and this was largely driven by an increase in $V\gamma9/V\delta2^+$ T cells (Fig. 3D). We also found an increase in $V\delta1^+$ T cell frequencies across repeated CHMIs (Fig. 3D). We then analysed the relationship between $\gamma\delta$ T cell frequencies and the risk of developing symptomatic malaria. Overall, $\alpha\beta$ T cell and $CD8^+$ T cell frequencies were similar in asymptomatic and symptomatic individuals (Fig. S4D). However, volunteers that progressed to asymptomatic malaria with serial CHMIs displayed robust profiles of increasing $V\delta1^+$ and $V\gamma9/V\delta2^+$ T cells frequencies across CHMIs, while symptomatic volunteers retained frequencies of $V\delta1^+$ and $V\gamma9/V\delta2^+$ T cells that were similar to their baseline samples (Fig. 3E). $V\gamma9/V\delta2^+$ T cells frequencies decreased between CHMIs and were not durably maintained at CHMI4 (Fig. 3E). Repeated measures correlations found a significant inverse association between the number of malaria symptoms and $V\delta1^+$ T cell frequencies ($P=0.007$) (Fig. 3F), but not with $\alpha\beta^+$ ($P=0.431$), $\gamma\delta^+$ ($P=0.109$), $CD8^+$ ($P=0.391$) or $V\gamma9/V\delta2^+$ T cell frequencies ($P=0.572$) (Fig. S4E). Together, these data from a highly controlled human malaria challenge model confirm that repeated *in vivo* *P. falciparum* infections drive changes in both $V\delta2^+$ and $V\delta1^+$ T cell frequencies. Increased $V\delta1^+$ T cell frequencies correlated with the development of asymptomatic malaria after CHMI4, while $V\gamma9/V\delta2^+$ T cell frequencies decreased between infections and were not durably maintained after CHMI4 in asymptomatic subjects, suggesting that regulation of $V\gamma9/V\delta2^+$ T cells may contribute to symptom reduction, a hypothesis that is consistent with previous reports in the context of natural infection (38–40).

Repeated *P. falciparum* infection initiates $V\delta1_{naive}$ to $V\delta1_{effector}$ T cell differentiation

Next, we investigated the impact of repeated *P. falciparum* infections on the differentiation of $\gamma\delta$ T cell subsets. Although $CD27^{hi} CD28^+ V\delta1_{naive}$ T cells were the main population of $V\delta1^+$ $\gamma\delta$ T cells in subjects prior to CHMI (malaria naïve), this cell population decreased after repeated *P. falciparum* infections (Fig. 4A). Conversely, $CD27^{lo} CX_3CR1^+ V\delta1_{effector}$ cells became the dominant population within total $V\delta1^+$ T cells (Fig. 4B). The increase in the $CD27^{lo} CX_3CR1^+ V\delta1_{effector}$ T cell population also correlated with a reduction in malaria symptoms (Fig. S4F). In response to a combination of inflammatory cytokines and HMB-PP stimulation, it has been proposed that $V\gamma9/V\delta2^+$ T cells switch phenotype from $CD27^+ CD28^+$ to $CD27^- CD28^-$ (41); however, we found no significant changes in these populations across repeated *P. falciparum* infections (Fig. 4C). As noted earlier, $V\gamma9/V\delta2^+$ T cells can control parasite replication through CD16-mediated antibody-dependent cytotoxicity (24), we found that CD16 expression was upregulated on $V\delta1^+$ T cells, but not $V\gamma9/V\delta2^+$ T cells after repeated *P. falciparum* infections (Fig. 4D). Interestingly, subject 17 displayed a major $CD27^- CD28^- CD16^+ V\gamma9/V\delta2^+$ T cell population that persisted over time (Fig. 4C and D). $V\delta1^+$ T cells consistently expressed the T cell activation marker CD38 after each *P. falciparum* infection, while $V\gamma9/V\delta2^+$ T cells only significantly upregulated CD38 after CHMI1 and 2 (Fig. 4E). We previously showed that $V\delta1_{effector}$ cells possess significant cytotoxic potential (19, 28). Here, we found that repeated *P. falciparum* infection drove $V\delta1^+$ T cells to express Gzm A, B, perforin, but not Gzm K (Fig. 4F), whereas $CD8^+$ T cells had no significant increase in Gzm A, B, perforin or Gzm K (Fig. S4G). In keeping with their pre-formed cytotoxic potential, $V\gamma9/V\delta2^+$ T cells retained robust levels of Gzm A, B, K and perforin after repeated *P. falciparum*

infections (Fig. 4F). Our data indicate that *in vivo* *P. falciparum* infection in humans drives the differentiation of human V δ ₁^{effector} $\gamma\delta$ T cells.

Repeated *P. falciparum* infections drive diverse waves of $\gamma\delta$ TCR selection

Next, we sought to understand whether repeated CHMIs impacted $\gamma\delta$ TCR repertoires. We used the approach described above (Fig. 2) and sorted V δ 2⁺ and V δ 2^{neg} $\gamma\delta$ T cell populations from longitudinal timepoints from all five CHMI subjects. We then analyzed the relationship between CD27^{lo} CX₃CR1⁺ V δ 2^{neg}_{effector} cells and V δ 2^{neg} TCR repertoires prior to CHMI1 and at CHMI4 + 21d in subject 2 (Fig. 5A). At baseline, V δ 2^{neg} $\gamma\delta$ T cells were predominantly CD27^{hi} CX₃CR1^{neg} and displayed a reasonably diverse $\gamma\delta$ TCR repertoire (Fig. 5A), but then after repeated CHMIs we observed a shift toward a CD27^{lo} CX₃CR1⁺ effector phenotype (Fig. 5A). Alongside these phenotypic changes, clonotypes found prior to CHMI1 remained stable or contracted over time, and new V δ 1⁺ clonotypes expanded, suggesting the potential recruitment of specific TCR sequences into the $\gamma\delta$ T cell immune repertoire after repeated CHMIs (Fig. 5B). Analysis of the V δ 2^{neg} $\gamma\delta$ T cell repertoires indicated V γ 9 and V δ 1 chain usage to be the most prevalent (Fig. 5C). Overall, diversity within V δ 2^{neg} or V δ 2⁺ $\gamma\delta$ T cell repertoires did not show any significant change (Fig. S5A). Next, we assessed if CDR3 clonotype changes were occurring in V δ 2⁺ TCR repertoires. We found that V δ 2⁺ clonotypes remained stable over time, despite significant changes in the frequencies of the total population (Fig. 5D). Interestingly, in subject 17, the V δ 2⁺ TCR repertoire was already dominated by hyperexpanded CDR3 γ and δ sequences at baseline (Fig. S5B), in contrast to all other V δ 2⁺ TCR repertoires in this study. Given the stability of V δ 2⁺ TCR clonotype repertoires at each CHMI, we then assessed the potential for dynamic changes in V δ 2^{neg} $\gamma\delta$ T cell repertoires at each CHMI and over time. Analysis of the $\gamma\delta$ TCR repertoire of subject 4 from baseline and over subsequent CHMI's 1, 3 and 4 indicated dynamic changes in the TCR repertoire, with an increase in low frequency clonotypes at CHMI1 and establishment of a broader immune repertoire over time (Fig. 5E). We then analyzed the 20 most prevalent clonotypes at baseline (subject 4, 10, 17) or at CHMI1 (subject 9; Fig. S5C). We found that prevalent baseline clonotypes declined with each CHMI and we observed waves of new clonotypes that expanded into the most abundant 20 clonotypes after each CHMI (Fig. 5F). In many cases these clonotypes were found at low frequency in the preceding timepoint, suggesting that each CHMI drove rounds of $\gamma\delta$ TCR selection (Fig. 5F). V δ 2^{neg} $\gamma\delta$ T cell repertoire clonotypes possessed few overlapping clonotypes between individuals, while there were many inter-individual overlapping TCR γ sequences in V δ 2⁺ TCR repertoires (Fig. S5D). Although subject 10 and 17 were symptomatic at CHMI4 and did not have a significant increase in V δ 1⁺ T cell frequencies, the repertoire of their V δ 2^{neg} $\gamma\delta$ TCR repertoire also displayed waves of clonotype selection (Fig. 5F). Together, the V γ 9/V δ 2⁺ T cell response to *P. falciparum* infection displays a highly stable polyclonal immune repertoire over time and infection. In contrast, V δ 2^{neg} $\gamma\delta$ T cell repertoires underwent dramatic remodeling of the $\gamma\delta$ TCR repertoire and displayed waves of clonal selection after each *P. falciparum* infection.

Previous *P. falciparum* exposure licenses V δ 1⁺ T cells reactivity to blood-stage parasites

Finally, we explored the reactivity of $\gamma\delta$ T cell subsets towards *P. falciparum* blood-stage parasites. PBMCs from Australian adults with no history of malaria exposure were co-

cultured with *P. falciparum* infected red blood cells (*PRBC*) or trophozoite/schizont extracts (*PTSE*) or intact uninfected RBCs (uRBC) or extracts (uRBCE) as controls. $V\delta 1^+$ T cells from Australian adults were unresponsive to *PRBC*s or *PTSE*, whereas $V\gamma 9/V\delta 2^+$ T cell populations were responsive (Fig. 6A), corroborating prior studies (24, 33, 42). Our *in vivo* results (Fig. 3, 4 and 5) prompted us to re-challenge PBMCs of Australian subjects twice over the 5-day culture period. Upon re-challenge we found that $V\delta 1^+$ T cells showed varying levels of proliferation after the second re-stimulation (Fig. 6B) but only in response to *PTSE* and not *PRBC*s. Previous studies have reported that $V\delta 1^+$ T cells from individuals living in malaria endemic regions of Gambia or Tanzania were unresponsive to *PRBC* *in vitro* (33, 42, 43). Using PBMCs from two Malian subjects and a malaria-naïve subject after 2 CHMIs (subject 10 at CHMI3+1d), we found that $V\delta 1^+$ T cells proliferated in response to *PTSE* but not *PRBC* after a single stimulation (Fig. 6C and D). This differential responsiveness to *PRBC* or *PTSE* was not consistently seen in paired $V\gamma 9/V\delta 2^+$ T cell populations or in malaria unexposed Australian subjects (Fig. 6D). These data indicate that prior *P. falciparum* infection primes $V\delta 1^+$ T cells for proliferate upon re-challenge with *P. falciparum* parasites.

Discussion

$\gamma\delta$ T cells have been implicated in the immune response to pathogenic microbes, including bacteria, viruses and parasites (9). These responses in mice and humans appear to be mediated by innate-like $\gamma\delta$ T cell populations, often utilizing semi- or invariant $\gamma\delta$ TCR repertoires that allow rapid effector responses to be mounted during the acute phases of microbial infection (9). Emerging evidence is currently re-shaping our understanding of the immunobiology of human $\gamma\delta$ T cell populations and $\gamma\delta$ T cells have the potential for both innate and adaptive properties (44). However, the adaptive-like features of $V\delta 2^{\text{neg}}$ $\gamma\delta$ T cell subsets are only partially understood (19, 28, 29, 45), and the establishment of this arm of the immune response to infectious disease has remained unclear.

Here, we show that repeated *in vivo* *P. falciparum* infection impacts populations of circulating innate-like $V\delta 2^+$ and adaptive-like $V\delta 1^+$ $\gamma\delta$ T cells. We found that repeated *P. falciparum* infection triggers the differentiation of $V\delta 1^+$ T cells from a $V\delta 1^{\text{naive}}$ phenotype into a distinct $V\delta 1^{\text{effector}}$ subset, concomitant with dynamic clonotype selection in the $\gamma\delta$ TCR repertoire with each *P. falciparum* infection. Together, our data indicate that *P. falciparum* infection drives the selection and differentiation of the $\gamma\delta$ T cell repertoire.

The association of human $\gamma\delta$ T cells and malaria has been largely attributed to the remarkable responsiveness of innate-like $V\gamma 9/V\delta 2^+$ T cells to *P. falciparum* infection (43, 46, 47). In line with this, we found that $V\gamma 9/V\delta 2^+$ T cells were retained after natural infection in Malian subjects and increased in frequency upon exposure to repeated CHMI, an observations that is likely due to encounter with blood stage *P. falciparum* merozoite-derived HMB-PP (15, 24, 48), and possibly also be in response to liver stage infection (4, 7, 49). However, notwithstanding hypotheses that $V\gamma 9/V\delta 2^+$ T cells mount oligoclonal responses to microbial encounters (50, 51), we found that public $V\gamma 9/V\delta 2^+$ TCR repertoires remained highly stable over time despite significant changes in cellular frequency. The composition of these repertoires were very similar to those seen in gestation (52), cord blood, and after birth

(17–19). Moreover, the cellular phenotype of V γ 9/V δ 2⁺ T cells after repeated *P. falciparum* infection was highly stable. Thus, the $\gamma\delta$ TCR repertoire of innate-like V γ 9/V δ 2⁺ T cells appears to allow sustained responsiveness upon *P. falciparum* infection.

In contrast to V γ 9/V δ 2⁺ T cells, the exact nature of human V δ 2⁻ $\gamma\delta$ T cells, and in particular V δ 1⁺ T cells, in the immune response to microbial pathogens is poorly defined, with recent studies identifying both innate- (26) and adaptive-like potential for these cells (28). Moreover, how V δ 1⁺ T cells participate in the complex immune response to *P. falciparum* is largely unknown (30, 33, 34, 53, 54). Current paradigms for conventional memory $\alpha\beta$ T cells indicate that T_{effector} arise from T_{naïve} cells driven by antigen-specific challenge to provide a rapid memory-response upon re-exposure to the same pathogen (55). Whether a similar paradigm applies to human $\gamma\delta$ T cells is unclear (56). Here, we demonstrate that V δ 1_{effector} T cells are a major population in Malian children, and that V δ 1_{naïve} cells differentiate into V δ 1_{effectors} after repeated *P. falciparum* infections in malaria-naïve adults. Given that V δ 1_{effector} $\gamma\delta$ T cells may infiltrate peripheral tissues (25), we speculate that *P. falciparum*-reactive V δ 1⁺ $\gamma\delta$ T cells will subsequently infiltrate the liver (25), and spleen (57). Therefore, *P. falciparum*-reactive V δ 1⁺ $\gamma\delta$ T cells may exert cytotoxic and/or immunoregulatory functions in peripheral tissues during malaria infection and may be possible to explore under certain clinical circumstances (58). Moreover, we also found that $\gamma\delta$ TCR repertoires undergo dynamic clonotype selection after each *P. falciparum* infection. While only a handful of the antigenic targets are known for V δ 2⁻ $\gamma\delta$ TCRs, nearly all identified ligands to date are endogenous host proteins (13, 59). In the case of malaria, it has been proposed that the V δ 1⁺ T cell response during *P. falciparum* infection is also driven by unknown endogenous host factors, based on the observation that V δ 1⁺ T cells from malaria-exposed individuals do not respond to *P. falciparum* antigens *in vitro* (33, 42). Our findings, that V δ 1⁺ T cells from malaria-exposed individuals react to *P. falciparum* lysate *in vitro*, suggests that V δ 1⁺ T cells may also have the potential to recognize parasite-derived antigens.

In malaria endemic regions, non-sterilizing immunity to symptomatic malaria is gradually acquired with repeated *P. falciparum* infections (60). It is hypothesized that the acquisition of immunity to malaria in humans involves resistance to severe disease followed by resistance to uncomplicated disease (3). Our study provides a window into the dynamic evolution of innate-and adaptive-like $\gamma\delta$ T cells in the context of natural *P. falciparum* infection and indicates that these cells may represent an important component of the cellular immune response that contributes to immunity to malaria (4–6). However, we cannot conclude from the current study that there is an association between V δ 1_{effector} T cell expansion and protection from febrile malaria in the context of natural infection, as Malian children who still experience febrile malaria show evidence of V δ 1_{effector} T cell expansion. Our previous analysis of the same cohort in Mali shows that the risk of febrile malaria gradually decreases with age over years of repeated malaria exposures (1). Subjects in the age range (7–14 years) included in the longitudinal portion of the current study are at lower risk of febrile malaria than younger children in the same cohort, but generally, even 7–14-year-olds have yet to acquire immunity that fully protects against febrile malaria from year to year, leaving open the possibility that V δ 1_{effector} T cell expansion with repeated infections may contribute to the gradual acquisition of immunity to malaria in endemic areas. Studies

with larger sample sizes that encompass a broader age range and include more frequent assessments of $\gamma\delta$ T cells relative to incident *P. falciparum* infections (both symptomatic and asymptomatic), will be required to assess the relationship between $V\delta 1_{\text{effector}}$ T cells and the risk of febrile malaria in the context of natural infection.

The findings from our CHMI study suggest that initial $V\gamma 9/V\delta 2^+$ T cell activation may contribute to the early priming and activation of naïve $V\delta 1^+$ $\gamma\delta$ T cells, potentially involving the capacity of $V\gamma 9/V\delta 2^+$ T cells to phagocytose and present parasite antigens (22, 24). Moreover, we noted a reduction in $V\gamma 9/V\delta 2^+$ T cells at the fourth CHMI, consistent with prior studies showing that loss and dysfunction of $V\delta 2^+$ T cells is associated with clinical immunity to malaria (40). How the emerging population of $V\delta 1^+$ $\gamma\delta$ T cells may contribute to protection from symptomatic malaria is unclear. While the regulatory functions of $\gamma\delta$ T cells in response to infectious diseases remains poorly understood, there is mounting evidence that these cells may play a role in regulating inflammation in the context of cancer (61, 62). Therefore, it seems plausible that repeated febrile malaria episodes could drive the expansion of a regulatory population of $V\delta 1^+$ $\gamma\delta$ T cells that dampen inflammation through IL-10 (31), TGF- β 1 (63) or other mechanisms (64).

There are several limitations of this study. First, the Mali cohort was conducted in a small rural village where the population is predominantly of a single ethnic group, limiting the generalizability of our findings. Nonetheless, we observed lower frequencies of $CD16^+ V\gamma 9/V\delta 2^+$ T cells in the Mali cohort relative to studies in Uganda and Brazil (24, 65). Thus, it will be of interest to further investigate the impact of genetics and/or environmental factors underlying regional differences. Secondly, the number of subjects included in the CHMI study was relatively small, precluding a rigorous analysis of the factors that underlie the inter-individual variability we observed in $\gamma\delta$ T cell responses.

In summary, our study shows that both innate and adaptive-like properties of the human $\gamma\delta$ T cell repertoire are driven by *P. falciparum* infection *in vivo*. $V\delta 2^+$ $\gamma\delta$ T cells mount a rapid innate-like polyclonal immune response to acute *P. falciparum* infection. Alongside these innate-like $V\delta 2^+$ $\gamma\delta$ T cell responses, repeated *P. falciparum* infection established clonally selected populations of adaptive-like $V\delta 1_{\text{effector}}$ $\gamma\delta$ T cells. Together, our study suggests the importance of future studies exploring the role of the $\gamma\delta$ T cell repertoire in contributing to the establishment of clinical immunity to malaria.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments:

We thank the study volunteers in Kenya and Mali, as well as the U.S. CHMI study volunteers for their contribution and commitment to malaria research. We thank Dr. Gregory Deye, of the National Institutes of Allergy and Infectious Diseases at the National Institutes of Health for service as program medical officer of the repetitive challenge study at the University of Maryland, Baltimore (UMB), Faith Pa'ahana-Brown, RN, Lisa Chrisley, RN, Alyson Kwon, Brenda Dorsey, Ana Raquel Da Costa, Jeffrey Crum, Kathleen Strauss, and Biraj Shrestha for their roles in the repetitive challenge study at UMB. We also thank Sanaria, Inc. for providing mosquitoes for human malaria infections. We thank FlowCore (Monash University) for assistance with cell sorting, and the Medical Genomics Facility (Hudson Institute) for their services. We also thank Dr. Eldho Paul from Monash Biostatistics

Platform for statistical analyses. We also thank Prof. Benjamin E. Willcox, Dr Carrie R. Willcox, Dr Robert Seder, Prof. Matthias Eberl and Prof. Adrian Hayday for helpful discussions.

Funding:

M.S.D. is supported by an Australian Research Council (ARC) Discovery Early Career Researcher Award (DE200100292) and this work was funded by a Rebecca L. Cooper Medical Research Foundation Project Grant (PG2020668) and ARC Discovery Project (DP210103327). C.L.D., P.O., and J.D.E. are supported by grants from the U.S. National Institutes of Health (U19 AI11211 and R01 AI11948). K.C.W. and K.E.L. are supported by a National Institutes of Health (NIH), Division of Allergy and Infectious Diseases (NIAID) U01 (AI-110852), distributed by the Henry M. Jackson Foundation (#1701447C). K.E.L. is further supported by additional funding from the NIAID (U01-HD092308, R01-AE141900, AI110820-06), The Geneva Foundation (V-12VAXHRFS-03), Medical Technology Enterprise Consortium (MTEC-17-01) and Pfizer Inc (C4591001, site 1002). The Mali study was funded by the Division of Intramural Research, National Institute of Allergy and Infectious Diseases, National Institutes of Health. J.R. is supported by an ARC Laureate Fellowship.

Data and materials availability:

All data are available in Data File S1. The T cell receptor (TCR) sequence data that support the findings of this study have been deposited in the Open Science Framework (OSF) and is accessible from <https://osf.io/7rdm9/> and <https://osf.io/3qvmh/>.

References:

1. Tran TM, Li S, Doumbo S, Doumtabe D, Huang CY, Dia S, Bathily A, Sangala J, Kone Y, Traore A, Niangaly M, Dara C, Kayentao K, Ongoiba A, Doumbo OK, Traore B, Crompton PD, An intensive longitudinal cohort study of Malian children and adults reveals no evidence of acquired immunity to Plasmodium falciparum infection. *Clin Infect Dis* 57, 40–47 (2013); published online EpubJul (10.1093/cid/cit174). [PubMed: 23487390]
2. Good MF, Doolan DL, Immune effector mechanisms in malaria. *Curr Opin Immunol* 11, 412–419 (1999); published online EpubAug (10.1016/S0952-7915(99)80069-7). [PubMed: 10448141]
3. Crompton PD, Moebius J, Portugal S, Waisberg M, Hart G, Garver LS, Miller LH, Barillas-Mury C, Pierce SK, Malaria immunity in man and mosquito: insights into unsolved mysteries of a deadly infectious disease. *Annu Rev Immunol* 32, 157–187 (2014)10.1146/annurev-immunol-032713-120220). [PubMed: 24655294]
4. Ishizuka AS, Lyke KE, DeZure A, Berry AA, Richie TL, Mendoza FH, Enama ME, Gordon IJ, Chang LJ, Sarwar UN, Zephir KL, Holman LA, James ER, Billingsley PF, Gunasekera A, Chakravarty S, Manoj A, Li M, Ruben AJ, Li T, Eappen AG, Stafford RE, N. K C, Murshedkar T, DeCederfelt H, Plummer SH, Hendel CS, Novik L, Costner PJ, Saunders JG, Laurens MB, Plowe CV, Flynn B, Whalen WR, Todd JP, Noor J, Rao S, Sierra-Davidson K, Lynn GM, Epstein JE, Kemp MA, Fadle GA, Mikolajczak SA, Fishbaugher M, Sack BK, Kappe SH, Davidson SA, Garver LS, Bjorkstrom NK, Nason MC, Graham BS, Roederer M, Sim BK, Hoffman SL, Ledgerwood JE, Seder RA, Protection against malaria at 1 year and immune correlates following PfSPZ vaccination. *Nat Med* 22, 614–623 (2016); published online EpubJun (10.1038/nm.4110). [PubMed: 27158907]
5. Zaidi I, Diallo H, Conteh S, Robbins Y, Kolasny J, Orr-Gonzalez S, Carter D, Butler B, Lambert L, Brickley E, Morrison R, Sissoko M, Healy SA, Sim BKL, Doumbo OK, Hoffman SL, Duffy PE, gammadelta T Cells Are Required for the Induction of Sterile Immunity during Irradiated Sporozoite Vaccinations. *J Immunol* 199, 3781–3788 (2017); published online EpubDec 1 (10.4049/jimmunol.1700314). [PubMed: 29079696]
6. de Jong SE, van Unen V, Manurung MD, Stam KA, Goeman JJ, Jochems SP, Holtt T, Pezzotti N, Mouwenda YD, Betouke Ongwe ME, Lorenz FR, Kruize YCM, Azimi S, Konig MH, Vilanova A, Eisemann E, Lelieveldt BPF, Roestenberg M, Sim BKL, Reinders MJT, Fendel R, Hoffman SL, Kremsner PG, Koning F, Mordmuller B, Lell B, Yazdanbakhsh M, Systems analysis and controlled malaria infection in Europeans and Africans elucidate naturally acquired immunity. *Nat Immunol* 22, 654–665 (2021); published online EpubMay (10.1038/s41590-021-00911-7). [PubMed: 33888898]

7. Mwakingwe-Omari A, Healy SA, Lane J, Cook DM, Kalhori S, Wyatt C, Kolluri A, Marte-Salcedo O, Imeru A, Nason M, Ding LK, Decederfelt H, Duan J, Neal J, Raiten J, Lee G, Hume JCC, Jeon JE, Ikpeama I, Kc N, Chakravarty S, Murshedkar T, Church LWP, Manoj A, Gunasekera A, Anderson C, Murphy SC, March S, Bhatia SN, James ER, Billingsley PF, Sim BKL, Richie TL, Zaidi I, Hoffman SL, Duffy PE, Two chemoattenuated PfSPZ malaria vaccines induce sterile hepatic immunity. *Nature* 595, 289–294 (2021); published online EpubJul (10.1038/s41586-021-03684-z). [PubMed: 34194041]
8. Bonneville M, O'Brien RL, Born WK, Gammadelta T cell effector functions: a blend of innate programming and acquired plasticity. *Nat Rev Immunol* 10, 467–478 (2010); published online EpubJul (10.1038/nri2781). [PubMed: 20539306]
9. Chien YH, Meyer C, Bonneville M, gammadelta T cells: first line of defense and beyond. *Annu Rev Immunol* 32, 121–155 (2014)10.1146/annurev-immunol-032713-120216. [PubMed: 24387714]
10. Hayday AC, [gamma][delta] cells: a right time and a right place for a conserved third way of protection. *Annu Rev Immunol* 18, 975–1026 (2000)10.1146/annurev.immunol.18.1.975. [PubMed: 10837080]
11. Rossjohn J, Gras S, Miles JJ, Turner SJ, Godfrey DI, McCluskey J, T cell antigen receptor recognition of antigen-presenting molecules. *Annu Rev Immunol* 33, 169–200 (2015)10.1146/annurevimmunol-032414-112334. [PubMed: 25493333]
12. Tanaka Y, Sano S, Nieves E, De Libero G, Rosa D, Modlin RL, Brenner MB, Bloom BR, Morita CT, Nonpeptide ligands for human gamma delta T cells. *Proc Natl Acad Sci U S A* 91, 8175–8179 (1994); published online EpubAug 16 (10.1073/pnas.91.17.8175). [PubMed: 8058775]
13. Willcox BE, Willcox CR, gammadelta TCR ligands: the quest to solve a 500-million-year-old mystery. *Nat Immunol* 20, 121–128 (2019); published online EpubFeb (10.1038/s41590-018-0304-y). [PubMed: 30664765]
14. Hirano M, Guo P, McCurley N, Schorpp M, Das S, Boehm T, Cooper MD, Evolutionary implications of a third lymphocyte lineage in lampreys. *Nature* 501, 435–438 (2013); published online EpubSep 19 (10.1038/nature12467). [PubMed: 23934109]
15. Morita CT, Jin C, Sarikonda G, Wang H, Nonpeptide antigens, presentation mechanisms, and immunological memory of human Vgamma2Vdelta2 T cells: discriminating friend from foe through the recognition of prenyl pyrophosphate antigens. *Immunol Rev* 215, 59–76 (2007); published online EpubFeb (10.1111/j.1600-065X.2006.00479.x). [PubMed: 17291279]
16. Hintz M, Reichenberg A, Altincicek B, Bahr U, Gschwind RM, Kollas AK, Beck E, Wiesner J, Eberl M, Jomaa H, Identification of (E)-4-hydroxy-3-methyl-but-2-enyl pyrophosphate as a major activator for human gammadelta T cells in *Escherichia coli*. *FEBS Lett* 509, 317–322 (2001); published online EpubDec 7 (10.1016/s0014-5793(01)03191-x). [PubMed: 11741609]
17. Papadopoulou M, Dimova T, Shey M, Briel L, Veldtsman H, Khomba N, Africa H, Steyn M, Hanekom WA, Scriba TJ, Nemes E, Vermijlen D, Fetal public Vgamma9Vdelta2 T cells expand and gain potent cytotoxic functions early after birth. *Proceedings of the National Academy of Sciences of the United States of America*, (2020); published online EpubJul 14 (10.1073/pnas.1922595117).
18. Ravens S, Fichtner AS, Willers M, Torkornoo D, Pirr S, Schoning J, Deseke M, Sandrock I, Bubke A, Wilharm A, Doodoo D, Egyir B, Flanagan KL, Steinbruck L, Dickinson P, Ghazal P, Adu B, Viemann D, Prinz I, Microbial exposure drives polyclonal expansion of innate gammadelta T cells immediately after birth. *Proc Natl Acad Sci U S A* 117, 18649–18660 (2020); published online EpubAug 4 (10.1073/pnas.1922588117). [PubMed: 32690687]
19. Davey MS, Willcox CR, Hunter S, Kasatskaya SA, Remmerswaal EBM, Salim M, Mohammed F, Bemelman FJ, Chudakov DM, Oo YH, Willcox BE, The human Vdelta2(+) T-cell compartment comprises distinct innate-like Vgamma9(+) and adaptive Vgamma9(–) subsets. *Nat Commun* 9, 1760 (2018); published online EpubMay 2 (10.1038/s41467-018-04076-0). [PubMed: 29720665]
20. Roussillon C, Agrapart M, Guglielmi P, Bensussan A, Bresseur P, Ballet JJ, Human TcR gamma delta+ lymphocyte response on primary exposure to *Plasmodium falciparum*. *Clin Exp Immunol* 95, 91–97 (1994); published online EpubJan (10.1111/j.1365-2249.1994.tb06020.x). [PubMed: 8287613]

21. Riganti C, Massaia M, Davey MS, Eberl M, Human gammadelta T-cell responses in infection and immunotherapy: common mechanisms, common mediators? *Eur J Immunol* 42, 1668–1676 (2012); published online EpubJul (10.1002/eji.201242492). [PubMed: 22806069]
22. Howard J, Loizon S, Tyler CJ, Duluc D, Moser B, Mechain M, Duvignaud A, Malvy D, Troye-Blomberg M, Moreau JF, Eberl M, Mercereau-Puijalon O, Dechanet-Merville J, Behr C, Mamani-Matsuda M, The Antigen-Presenting Potential of Vgamma9Vdelta2 T Cells During Plasmodium falciparum Blood-Stage Infection. *J Infect Dis* 215, 1569–1579 (2017); published online EpubMay 15 (10.1093/infdis/jix149). [PubMed: 28368498]
23. Costa G, Loizon S, Guenot M, Mocan I, Halary F, de Saint-Basile G, Pitard V, Dechanet-Merville J, Moreau JF, Troye-Blomberg M, Mercereau-Puijalon O, Behr C, Control of Plasmodium falciparum erythrocytic cycle: gammadelta T cells target the red blood cell-invasive merozoites. *Blood* 118, 6952–6962 (2011); published online EpubDec 22 (10.1182/blood-2011-08-376111). [PubMed: 22045985]
24. Junqueira C, Polidoro RB, Castro G, Absalon S, Liang Z, Sen Santara S, Crespo A, Pereira DB, Gazzinelli RT, Dvorin JD, Lieberman J, gammadelta T cells suppress Plasmodium falciparum blood-stage infection by direct killing and phagocytosis. *Nat Immunol* 22, 347–357 (2021); published online EpubMar (10.1038/s41590-020-00847-4). [PubMed: 33432229]
25. Hunter S, Willcox CR, Davey MS, Kasatskaya SA, Jeffery HC, Chudakov DM, Oo YH, Willcox BE, Human liver infiltrating gammadelta T cells are composed of clonally expanded circulating and tissue-resident populations. *J Hepatol*, (2018); published online EpubMay 18 (10.1016/j.jhep.2018.05.007).
26. Wu Y, Kyle-Cezar F, Woolf RT, Naceur-Lombardelli C, Owen J, Biswas D, Lorenc A, Vantourout P, Gazinska P, Grigoriadis A, Tutt A, Hayday A, An innate-like Vdelta1(+) gammadelta T cell compartment in the human breast is associated with remission in triple-negative breast cancer. *Sci Transl Med* 11, (2019); published online EpubOct 9 (10.1126/scitranslmed.aax9364).
27. Di Marco Barros R, Roberts NA, Dart RJ, Vantourout P, Jandke A, Nussbaumer O, Deban L, Cipolat S, Hart R, Iannitto ML, Laing A, Spencer-Dene B, East P, Gibbons D, Irving PM, Pereira P, Steinhoff U, Hayday A, Epithelia Use Butyrophilin-like Molecules to Shape Organ-Specific gammadelta T Cell Compartments. *Cell* 167, 203–218 e217 (2016); published online EpubSep 22 (10.1016/j.cell.2016.08.030). [PubMed: 27641500]
28. Davey MS, Willcox CR, Joyce SP, Ladell K, Kasatskaya SA, McLaren JE, Hunter S, Salim M, Mohammed F, Price DA, Chudakov DM, Willcox BE, Clonal selection in the human Vdelta1 T cell repertoire indicates gammadelta TCR-dependent adaptive immune surveillance. *Nat Commun* 8, 14760 (2017); published online EpubMar 1 (10.1038/ncomms14760). [PubMed: 28248310]
29. Ravens S, Schultze-Florey C, Raha S, Sandrock I, Drenker M, Oberdorfer L, Reinhardt A, Ravens I, Beck M, Geffers R, von Kaisenberg C, Heuser M, Thol F, Ganser A, Forster R, Koenecke C, Prinz I, Human gammadelta T cells are quickly reconstituted after stem-cell transplantation and show adaptive clonal expansion in response to viral infection. *Nat Immunol* 18, 393–401 (2017); published online EpubApr (10.1038/ni.3686). [PubMed: 28218745]
30. Hviid L, Kurtzhals JA, Adabayeri V, Loizon S, Kemp K, Goka BQ, Lim A, Mercereau-Puijalon O, Akanmori BD, Behr C, Perturbation and proinflammatory type activation of V delta 1(+) gamma delta T cells in African children with Plasmodium falciparum malaria. *Infect Immun* 69, 3190–3196 (2001); published online EpubMay (10.1128/IAI.69.5.3190-3196.2001). [PubMed: 11292740]
31. Taniguchi T, Md Mannoor K, Nonaka D, Toma H, Li C, Narita M, Vanisaveth V, Kano S, Takahashi M, Watanabe H, A Unique Subset of gammadelta T Cells Expands and Produces IL-10 in Patients with Naturally Acquired Immunity against Falciparum Malaria. *Front Microbiol* 8, 1288 (2017)10.3389/fmicb.2017.01288. [PubMed: 28769886]
32. Worku S, Bjorkman A, Troye-Blomberg M, Jemaneh L, Farnert A, Christensson B, Lymphocyte activation and subset redistribution in the peripheral blood in acute malaria illness: distinct gammadelta+ T cell patterns in Plasmodium falciparum and P. vivax infections. *Clin Exp Immunol* 108, 34–41 (1997); published online EpubApr (10.1046/j.1365-2249.1997.d01-981.x). [PubMed: 9097908]
33. Goodier M, Krause-Jauer M, Sanni A, Massougbdji A, Sadeler BC, Mitchell GH, Modolell M, Eichmann K, Langhorne J, Gamma delta T cells in the peripheral blood of individuals from

- an area of holoendemic *Plasmodium falciparum* transmission. *Trans R Soc Trop Med Hyg* 87, 692–696 (1993); published online EpubNov-Dec (10.1016/0035-9203(93)90299-6). [PubMed: 8296383]
34. Hviid L, Smith-Togobo C, Willcox BE, Human Vdelta1(+) T Cells in the Immune Response to *Plasmodium falciparum* Infection. *Front Immunol* 10, 259 (2019)10.3389/fimmu.2019.00259. [PubMed: 30837999]
 35. Hill DL, Carr EJ, Rutishauser T, Moncunill G, Campo JJ, Innocentin S, Mpina M, Nhabomba A, Tumbo A, Jairoce C, Moll HA, van Zelm MC, Dobano C, Daubenberger C, Linterman MA, Immune system development varies according to age, location, and anemia in African children. *Sci Transl Med* 12, (2020); published online EpubFeb 5 (10.1126/scitranslmed.aaw9522).
 36. Portugal S, Tran TM, Ongoiba A, Bathily A, Li S, Doumbo S, Skinner J, Doumtabe D, Kone Y, Sangala J, Jain A, Davies DH, Hung C, Liang L, Ricklefs S, Homann MV, Felgner PL, Porcella SF, Farnert A, Doumbo OK, Kayentao K, Greenwood BM, Traore B, Crompton PD, Treatment of Chronic Asymptomatic *Plasmodium falciparum* Infection Does Not Increase the Risk of Clinical Malaria Upon Reinfection. *Clin Infect Dis* 64, 645–653 (2017); published online EpubMar 1 (10.1093/cid/ciw849). [PubMed: 28362910]
 37. Papadopoulou M, Dimova T, Shey M, Briel L, Veldtsman H, Khomba N, Africa H, Steyn M, Hanekom WA, Scriba TJ, Nemes E, Vermijlen D, Fetal public Vgamma9Vdelta2 T cells expand and gain potent cytotoxic functions early after birth. *Proc Natl Acad Sci U S A* 117, 18638–18648 (2020); published online EpubAug 4 (10.1073/pnas.1922595117). [PubMed: 32665435]
 38. Jagannathan P, Lutwama F, Boyle MJ, Nankya F, Farrington LA, McIntyre TI, Bowen K, Naluwu K, Nalubega M, Musinguzi K, Sikyomu E, Budker R, Katureebe A, Rek J, Greenhouse B, Dorsey G, Kanya MR, Feeney ME, Vdelta2+ T cell response to malaria correlates with protection from infection but is attenuated with repeated exposure. *Sci Rep* 7, 11487 (2017); published online EpubSep 13 (10.1038/s41598-017-10624-3). [PubMed: 28904345]
 39. Farrington LA, Jagannathan P, McIntyre TI, Vance HM, Bowen K, Boyle MJ, Nankya F, Wamala S, Auma A, Nalubega M, Sikyomu E, Naluwu K, Bigira V, Kapisi J, Dorsey G, Kanya MR, Feeney ME, Frequent Malaria Drives Progressive Vdelta2 T-Cell Loss, Dysfunction, and CD16 Up-regulation During Early Childhood. *J Infect Dis* 213, 1483–1490 (2016); published online EpubMay 1 (10.1093/infdis/jiv600). [PubMed: 26667315]
 40. Jagannathan P, Kim CC, Greenhouse B, Nankya F, Bowen K, Eccles-James I, Muhindo MK, Arinaitwe E, Tappero JW, Kanya MR, Dorsey G, Feeney ME, Loss and dysfunction of Vdelta2(+) gammadelta T cells are associated with clinical tolerance to malaria. *Sci Transl Med* 6, 251ra117 (2014); published online EpubAug 27 (10.1126/scitranslmed.3009793).
 41. Wragg KM, Tan HX, Kristensen AB, Nguyen-Robertson CV, Kelleher AD, Parsons MS, Wheatley AK, Berzins SP, Pellicci DG, Kent SJ, Juno JA, High CD26 and Low CD94 Expression Identifies an IL-23 Responsive Vdelta2(+) T Cell Subset with a MAIT Cell-like Transcriptional Profile. *Cell Rep* 31, 107773 (2020); published online EpubJun 16 (10.1016/j.celrep.2020.107773). [PubMed: 32553157]
 42. Rutishauser T, Lepore M, Di Blasi D, Dangy JP, Abdulla S, Jongo S, Ramadhani K, Sim BKL, Hoffman SL, Tanner M, Daubenberger C, De Libero G, Activation of TCR Vdelta1(+) and Vdelta1(-)Vdelta2(-) gammadelta T Cells upon Controlled Infection with *Plasmodium falciparum* in Tanzanian Volunteers. *J Immunol* 204, 180–191 (2020); published online EpubJan 1 (10.4049/jimmunol.1900669). [PubMed: 31801816]
 43. Goodier M, Fey P, Eichmann K, Langhorne J, Human peripheral blood gamma delta T cells respond to antigens of *Plasmodium falciparum*. *Int Immunol* 4, 33–41 (1992); published online EpubJan (10.1093/intimm/4.1.33). [PubMed: 1531764]
 44. Hayday AC, gammadelta T Cell Update: Adaptate Orchestrators of Immune Surveillance. *J Immunol* 203, 311–320 (2019); published online EpubJul 15 (10.4049/jimmunol.1800934). [PubMed: 31285310]
 45. Kaminski H, Menard C, El Hayani B, Adjibabi AN, Marseres G, Courant M, Zouine A, Pitard V, Garrigue I, Burrell S, Moreau JF, Couzi L, Visentin J, Merville P, Dechanet-Merville J, Characterization of a unique gammadelta T cell subset as a specific marker of CMV infection severity. *J Infect Dis*, (2020); published online EpubJul 5 (10.1093/infdis/jiaa400).

46. Schmalzer M, Orlova-Fink N, Rutishauser T, Abdulla S, Daubenberger C, Human unconventional T cells in *Plasmodium falciparum* infection. *Semin Immunopathol* 42, 265–277 (2020); published online EpubJun (10.1007/s00281-020-00791-3). [PubMed: 32076813]
47. Behr C, Dubois P, Preferential expansion of V gamma 9 V delta 2 T cells following stimulation of peripheral blood lymphocytes with extracts of *Plasmodium falciparum*. *Int Immunol* 4, 361–366 (1992); published online EpubMar (10.1093/intimm/4.3.361). [PubMed: 1533150]
48. Hernandez-Castaneda MA, Happ K, Cattalani F, Wallimann A, Blanchard M, Fellay I, Scolari B, Lannes N, Mbagwu S, Fellay B, Filgueira L, Mantel PY, Walch M, gammadelta T Cells Kill *Plasmodium falciparum* in a Granzyme- and Granulysin-Dependent Mechanism during the Late Blood Stage. *J Immunol* 204, 1798–1809 (2020); published online EpubApr 1 (10.4049/jimmunol.1900725). [PubMed: 32066596]
49. Seder RA, Chang LJ, Enama ME, Zephir KL, Sarwar UN, Gordon IJ, Holman LA, James ER, Billingsley PF, Gunasekera A, Richman A, Chakravarty S, Manoj A, Velmurugan S, Li M, Ruben AJ, Li T, Eappen AG, Stafford RE, Plummer SH, Hendel CS, Novik L, Costner PJ, Mendoza FH, Saunders JG, Nason MC, Richardson JH, Murphy J, Davidson SA, Richie TL, Sedegah M, Sutamihardja A, Fahle GA, Lyke KE, Laurens MB, Roederer M, Tewari K, Epstein JE, Sim BK, Ledgerwood JE, Graham BS, Hoffman SL, Team VRCS, Protection against malaria by intravenous immunization with a nonreplicating sporozoite vaccine. *Science* 341, 1359–1365 (2013); published online EpubSep 20 (10.1126/science.1241800). [PubMed: 23929949]
50. Willcox CR, Davey MS, Willcox BE, Development and Selection of the Human Vgamma9Vdelta2(+) T-Cell Repertoire. *Front Immunol* 9, 1501 (2018)10.3389/fimmu.2018.01501. [PubMed: 30013562]
51. Pauza CD, Cairo C, Evolution and function of the TCR Vgamma9 chain repertoire: It's good to be public. *Cell Immunol* 296, 22–30 (2015); published online EpubJul (10.1016/j.cellimm.2015.02.010). [PubMed: 25769734]
52. Dimova T, Brouwer M, Gosselin F, Tassignon J, Leo O, Donner C, Marchant A, Vermijlen D, Effector Vgamma9Vdelta2 T cells dominate the human fetal gammadelta T-cell repertoire. *Proc Natl Acad Sci U S A* 112, E556–565 (2015); published online EpubFeb 10 (10.1073/pnas.1412058112). [PubMed: 25617367]
53. Hviid L, Akanmori BD, Loizon S, Kurtzhals JA, Ricke CH, Lim A, Koram KA, Nkrumah FK, Mercereau-Puijalon O, Behr C, High frequency of circulating gamma delta T cells with dominance of the v(delta)1 subset in a healthy population. *Int Immunol* 12, 797–805 (2000); published online EpubJun (10.1093/intimm/12.6.797). [PubMed: 10837407]
54. Deroost K, Langhorne J, Gamma/Delta T Cells and Their Role in Protection Against Malaria. *Front Immunol* 9, 2973 (2018)10.3389/fimmu.2018.02973. [PubMed: 30619330]
55. Omilusik KD, Goldrath AW, The origins of memory T cells. *Nature* 552, 337–339 (2017); published online EpubDec 21 (10.1038/d41586-017-08280-8).
56. Davey MS, Willcox CR, Baker AT, Hunter S, Willcox BE, Recasting Human Vdelta1 Lymphocytes in an Adaptive Role. *Trends Immunol* 39, 446–459 (2018); published online EpubJun (10.1016/j.it.2018.03.003). [PubMed: 29680462]
57. Falini B, Flenghi L, Pileri S, Pelicci P, Fagioli M, Martelli MF, Moretta L, Ciccone E, Distribution of T cells bearing different forms of the T cell receptor gamma/delta in normal and pathological human tissues. *J Immunol* 143, 2480–2488 (1989); published online EpubOct 15 ([PubMed: 2477444]
58. Kho S, Qotrunnada L, Leonardo L, Andries B, Wardani PAI, Fricot A, Henry B, Hardy D, Margyaningsih NI, Apriyanti D, Puspitasari AM, Prayoga P, Trianty L, Kenangalem E, Chretien F, Safeukui I, Del Portillo HA, Fernandez-Becerra C, Meibalan E, Marti M, Price RN, Woodberry T, Ndour PA, Russell BM, Yeo TW, Minigo G, Noviyanti R, Poespoprodjo JR, Siregar NC, Buffet PA, Anstey NM, Hidden Biomass of Intact Malaria Parasites in the Human Spleen. *N Engl J Med* 384, 2067–2069 (2021); published online EpubMay 27 (10.1056/NEJMc2023884). [PubMed: 34042394]
59. Willcox BE, Mohammed F, Willcox CR, gammadelta TCR Recognition of MR1: Adapting to Life on the Flip Side. *Trends Biochem Sci* 45, 551–553 (2020); published online EpubJul (10.1016/j.tibs.2020.03.012). [PubMed: 32299647]

60. Marsh K, Kinyanjui S, Immune effector mechanisms in malaria. *Parasite Immunol* 28, 51–60 (2006); published online EpubJan-Feb (10.1111/j.1365-3024.2006.00808.x). [PubMed: 16438676]
61. Silva-Santos B, Serre K, Norell H, gammadelta T cells in cancer. *Nat Rev Immunol* 15, 683–691 (2015); published online EpubNov (10.1038/nri3904). [PubMed: 26449179]
62. Peng G, Wang HY, Peng W, Kiniwa Y, Seo KH, Wang RF, Tumor-infiltrating gammadelta T cells suppress T and dendritic cell function via mechanisms controlled by a unique toll-like receptor signaling pathway. *Immunity* 27, 334–348 (2007); published online EpubAug (10.1016/j.immuni.2007.05.020). [PubMed: 17656116]
63. Bhagat G, Naiyer AJ, Shah JG, Harper J, Jabri B, Wang TC, Green PH, Manavalan JS, Small intestinal CD8+TCRgammadelta+NKG2A+ intraepithelial lymphocytes have attributes of regulatory cells in patients with celiac disease. *J Clin Invest* 118, 281–293 (2008); published online EpubJan (10.1172/JCI30989). [PubMed: 18064301]
64. Siegers GM, Lamb LS Jr., Cytotoxic and regulatory properties of circulating Vdelta1+ gammadelta T cells: a new player on the cell therapy field? *Mol Ther* 22, 1416–1422 (2014); published online EpubAug (10.1038/mt.2014.104). [PubMed: 24895997]
65. Farrington LA, Callaway PC, Vance HM, Baskevitch K, Lutz E, Warriar L, McIntyre TI, Budker R, Jagannathan P, Nankya F, Musinguzi K, Nalubega M, Sikyomu E, Naluwu K, Arinaitwe E, Dorsey G, Kamya MR, Feeney ME, Opsonized antigen activates Vdelta2+ T cells via CD16/FCgammaRIIIa in individuals with chronic malaria exposure. *PLoS Pathog* 16, e1008997 (2020); published online EpubOct (10.1371/journal.ppat.1008997). [PubMed: 33085728]




Figure 1. Increased V δ 1⁺ $\gamma\delta$ T cells frequencies in Malian subjects exposed to *P. falciparum* infection.

In age and gender matched Malian (n=23) or Australian subjects (n=20): **A**. Frequencies of total $\gamma\delta$, V γ 9/V δ 2⁺ and V δ 1⁺ T cells within CD3⁺ T cells, **B**. Frequencies of V γ 9/V δ 2⁺ and V δ 1⁺ T cells in total CD3⁺ T cells, **C**. Frequencies of CD27^{lo} CX₃CR1⁺ cells within V δ 1⁺ or CD27⁺ CD28⁺ cells within V γ 9/V δ 2⁺ T cells, **D**. Frequencies of perforin⁺, **E**. Gzm A⁺, **F**. Gzm B⁺ cells within V δ 1⁺ or V γ 9/V δ 2⁺ T cells (**D**, **E** and **F**: Malian n=19 and Australian n=15). **G**. Schematic of samples and malaria exposure for Malian subjects included in the longitudinal arm of our study. Subjects are stratified based on presentation with a confirmed febrile malaria episode in all three years (n=4) or two febrile episodes (n=5), with either one episode in year 2 (n=3) or 3 (n=2). **H**. Frequencies of V δ 1⁺,

V γ 9/V δ 2⁺ and CD8⁺ T cells in total CD3⁺ T cells during febrile malaria and 3–6 months following treatment over the 3-year seasonal transmission periods. Year 1 (n=9), Year 2 (n=7) and Year 3 (n=6). **I.** Frequencies of V δ 1⁺ and V γ 9/V δ 2⁺ T cells within CD3⁺ T cells over a 6-month period without a febrile malaria episode (n=5; 12 – 18 months, n=2, or 24 – 30 months, n=3). Bars show the mean and error bars indicate means \pm SEM. Normality was tested using the Shapiro-Wilk test.; *P < 0.05; **P < 0.01; ***P < 0.001; ****P < 0.0001; p-values were determined by Mann-Whitney test (**a - f**) and Wilcoxon matched-pairs signed rank test (**h, i**).




Figure 2. $\gamma\delta$ TCR repertoires in Malian subjects evolve over time.

A. TCR γ clonotype tree plot analysis of V δ 2^{neg} and V γ 9/V δ 2⁺ T cell populations from Australian children or Malian children during stable periods without malaria transmission. Tree plots show unique clonotypes (coloured segments) and their proportion within the total repertoire (size), in general coloured clonotypes do not match between plots unless indicated. **B.** Pooled accumulated frequency curves of the top 10 most prevalent clonotypes in V δ 2^{neg} or V γ 9/V δ 2⁺ TCR repertoires (Australian, n=3; Mali, n=4). **C.** Diversity index of V δ 2^{neg} and V δ 2⁺ $\gamma\delta$ T cell repertoires in Malian (n=4) or Australian (n=3) subjects. **D.** Frequency of shared CDR3 γ (a.a.) sequences in V δ 2^{neg} and V γ 9/V δ 2⁺ T cell repertoires (Australian, n=3; Mali, n=4). **E.** Longitudinal tracking of the 20 most abundant TCR γ clonotypes in V γ 9/V δ 2⁺ and **F.** V δ 2^{neg} T cell repertoires over time in subject 066. (M)

indicates acute febrile malaria. **G.** Longitudinal analysis of V γ chain usage and diversity index for V δ 2^{neg} (red) and V γ 9/V δ 2⁺ (blue) T cell repertoires from subject 066. **H.** $\gamma\delta$ TCR expression patterns within (CD27^{lo} CX₃CR1⁺) V δ 1⁺_{effector} populations in donor 179. Each flow cytometry plot has two time points overlaid, indicated by an arrow, together covering three febrile *P. falciparum* infections (months 0, 17 and 30). **I.** TCR δ clonotypes sequencing relative to total CD3⁺ T cells from subject 179. Error bars indicate means \pm SEM. Normality was tested using the Shapiro-Wilk test.; *P < 0.05; **P < 0.01; ***P < 0.001; p-values were determined by two-way ANOVA with Sidaks post hoc testing (**b**) and one-way ANOVA with Holm-Sidaks post hoc testing (**c**, **d**).

p-values were determined by Kruskal-Wallis test (**b**), linear mixed effects modelling with Bonferroni's correction (**c, d, e**) and repeated measures correlation (**f**).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 4. Repeated *P. falciparum* infection drives the differentiation of cytotoxic Vδ1 effector T cells.

A. Representative flow cytometry plot and graph showing the frequencies of CD27⁺ CD28⁺ cells in Vδ1⁺ T cells after repeated CHMIs (n=5). **B.** Representative flow cytometry plot and graph showing the differentiation of CD27^{lo} CX₃CR1⁺ Vδ1⁺ effector cells after repeated CHMIs (n=5). **C.** Frequencies of CD27⁺ CD28⁺ and CD27^{neg} CD28^{neg} cells within Vγ9/Vδ2⁺ T cells. **D-F.** Within Vδ1⁺ (red) and Vγ9/Vδ2⁺ (blue) T cells, the frequencies of: **D.** CD16⁺, **E.** CD38⁺, **F.** Gzm A⁺, B⁺, K⁺ and perforin⁺ cells. Bars show the mean. *P < 0.05; **P < 0.01; ***P < 0.001; ****P < 0.0001; p-values were determined by linear mixed effects modelling with Bonferroni's correction.




Figure 5. Repeated *P. falciparum* infection drives waves of Vδ1 TCR clonotype selection.
A. Flow cytometry plots showing frequencies of CD27^{lo} CX₃CR1⁺ Vδ2^{neg} γδ T cells after repeated CHMI challenge in subject 2. TCRδ tree plots of the corresponding total Vδ2^{neg} γδ T cells and DI are given for each tree plot. **B.** Increase in new Vδ1 sequences between baseline and CHMI4 within the top 20 clonotypes in TCRγ from subject 2. **C.** Vγ and Vδ usage in Vδ2^{neg} T cell repertoires from baseline to CHMI 4 (n=4–5). **D.** Longitudinal tracking of the top 20 CDR3γ clonotypes in Vδ2⁺ T cell repertoires as a frequency of total CD3⁺ T cell populations. **E.** TCRγ tree plots showing Vδ2^{neg} TCR repertoires at baseline and after repeated CHMIs in subject 4. The D75 and DI metrics are indicated. The graphs show the accumulated frequency of the top 20 clonotypes for each repertoire. **F.** Longitudinal tracking of the top 20 CDR3γ clonotypes in Vδ2^{neg} TCR repertoires from

subject 4, 9, 10, 17; displayed as a proportion of the total TCR γ repertoire (left) or within the total CD3⁺ T cell population (right).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 6. Previous *P. falciparum* exposure licenses Vδ1⁺ T cells for parasite reactivity. Vδ1⁺ and Vγ9/Vδ2⁺ T cells were assessed for proliferation in Australian adult donors with no history of malaria, PBMCs were labelled with cell trace and incubated for 6 days with **A.** One or **B.** two stimulations (at day 0 and 3 of culture) with *P. falciparum* trophozoite/early schizont extract (*PfTSE*) or infected red blood cells (RBCs) and uninfected RBCs (uRBC) or extract (uRBCE) (uRBCE/*PfTSE*: n=6; uRBC/*PfRBC*: n=10). **C.** Representative flow cytometry plots show Vδ1⁺ (blue) and Vδ2⁺ (black) T cells assessed for proliferation in the PBMCs from a Malian subject after co-culture with *PfTSE*. *PfRBC*s, uRBC or uRBCE controls. **D.** Graphs show the proliferation of Vδ1⁺ and Vγ9/Vδ2⁺ T cells from two Malian subjects with a history of repeated prior exposure to *P. falciparum* malaria, subject 10 at CHMI3 + 1d and three independent Australian donors with no history of

malaria exposure. Each data point represents the proportion of proliferating cells in cultures exposed to *PRBCs* or *PfTSE* minus the response to uRBC or uRBCE controls. Bars show the mean \pm SEM. Normality was tested using the Shapiro-Wilk test.; **P < 0.01; p-values were determined by one-way ANOVA with Holm-Sidak's post hoc testing (**d**).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript