



Correction for Kouamou et al., “Diagnostic Accuracy of Pan-Degenerate Amplification and Adaptation Assay for HIV-1 Drug Resistance Mutation Analysis in Low- and Middle-Income Countries”

 Vinie Kouamou,^a Justen Manasa,^b David Katzenstein,^c Alan M. McGregor,^a Chiratidzo E. Ndhlovu,^a Tariro Makadzange^{a,d}

^aDepartment of Medicine, College of Health Sciences, University of Zimbabwe, Harare, Zimbabwe

^bDepartment of Medical Microbiology, College of Health Sciences, University of Zimbabwe, Harare, Zimbabwe

^cDepartment of Molecular Biology, Biomedical Research and Training Institute, Harare, Zimbabwe

^dHarvard Medical School, Boston, Massachusetts, USA

Volume 58, no. 9, e01045-20, 2020, <https://doi.org/10.1128/JCM.01045-20>. In Table 2, the revised kappa (95% confidence interval) values for NRTI DRMs and NNRTI DRMs are as follows: for NRTI DRMs, 0.97 (0.94–1.00); for NNRTI DRMs, 0.81 (0.71–0.92).

Copyright © 2022 American Society for Microbiology. All Rights Reserved.

Address correspondence to Vinie Kouamou, kvinie2005@yahoo.fr.

Published 13 June 2022