



Meal frequency and incidence of type 2 diabetes: a prospective study

Xiaowen Wang^{1,2}, Yonghua Hu^{1,3}, Li-Qiang Qin⁴ and Jia-Yi Dong^{2*}

¹Department of Epidemiology and Biostatistics, School of Public Health, Peking University Health Science Center, Beijing 100191, People's Republic of China

²Public Health, Department of Social Medicine, Osaka University Graduate School of Medicine, Osaka 5650871, Japan

³Medical Informatics Center, Peking University Health Science Center, Beijing 100191, People's Republic of China

⁴Department of Nutrition and Food Hygiene, School of Public Health, Soochow University, Suzhou 215000, People's Republic of China

(Submitted 25 March 2021 – Final revision received 8 August 2021 – Accepted 18 August 2021 – First published online 23 August 2021)

Abstract

Dietary habits play an important role in the development of obesity and type 2 diabetes. However, evidence on association between diet frequency and type 2 diabetes was limited and inconclusive. We aimed to examine the association between meal frequency and risk of type 2 diabetes. The cohort study used data from China Health and Retirement Longitudinal Study of 8874 community-dwelling people aged over 45 years. Participants were classified as eating two meals per day, three meals per day and four meals per day. Multiple Poisson regression models were used to examine risk of 4-year incident type 2 diabetes among people who ate more or less than three meals per day compared with people who ate three meals per day. We documented 706 type 2 diabetes cases during follow-up. After adjustment for known risk factors for type 2 diabetes, except for BMI, participants who ate four meals per day were at a lower risk of type 2 diabetes than those who ate three meals per day (relative risk (RR) = 0.73 (0.58, 0.92)). After further adjustment for baseline BMI, the association was slightly attenuated but remained statistically significant (RR = 0.76 (0.60, 0.97)). Subgroup analysis showed that the fully adjusted RR of type 2 diabetes for people eating four meals per day were 0.66 (0.48, 0.91) and 0.93 (0.65, 1.34) among those had a BMI < 25 and ≥ 25 kg/m², respectively. Eating four meals per day, compared with eating three meals per day was associated with lower risk of developing type 2 diabetes in a Chinese population, particularly in those with a BMI < 25 kg/m².

Key words: Meal frequency; Type 2 diabetes; Prospective study; Chinese

Dietary habits, including eating speed, breakfast skipping and meal frequency, have received increasing interest because of their potential impacts on human health, especially metabolic-related diseases^(1–4). For instance, a number of observational studies have shown an association between a fast eating speed and increased risk of obesity⁽⁴⁾ and type 2 diabetes^(5,6). Breakfast skipping was also associated with an increased risk of type 2 diabetes⁽³⁾.

Observational studies suggested that increased meal frequency was associated with a lower risk of obesity⁽⁷⁾ and improved lipid profiles^(8–10). One meta-analysis of randomised controlled trials (RCT) investigating the effects of meal frequency on weight loss and body composition found that increased meal frequency had beneficial effects on body composition, including reductions in fat mass and increases in fat-free mass⁽¹¹⁾. Such evidence indicated that increased meal frequency may be helpful in diabetes prevention. However, another recent meta-analysis of

RCT indicated that body weight and waist circumference might be affected by meal frequency, but not fat mass and energy intake⁽¹²⁾. Also, cohort studies examining the association between meal frequency and type 2 diabetes posed controversial results. Two prospective cohort studies reported the non-significant association between eating frequency and risk of type 2 diabetes^(13,14), while one study found an increased risk of type 2 diabetes in people eating four meals per day compared with those eating 1–3 meals per day⁽¹⁵⁾.

Of note, all the three cohort studies^(13–15) were performed in the USA, where the prevalence of overweight and obesity is high, and no evidence existed from Asian population on this topic. Therefore, the objective of this study was to examine the association between meal frequency and risk of type 2 diabetes in a Chinese population. We also performed stratified analyses by BMI to test whether BMI could modify the association.

Abbreviations: CHARLS, China Health and Retirement Longitudinal Study; RR, relative risks; RCT, randomised controlled trials.

* **Corresponding author:** Jia-Yi Dong, email dongjy@mail3.sysu.edu.cn

Methods

Participants

The study was based on a nationally representative cohort study derived from the China Health and Retirement Longitudinal Study (CHARLS). The design, sampling procedure, and data collection methods of the CHARLS have been previously described⁽¹⁶⁾. Briefly, CHARLS recruited residents aged over 45 years from 150 county-level units across 28 provinces in China. The baseline survey was conducted in 2011–2012 (wave 1) among 17 708 participants with a response rate of 80.5%, and the study was followed up every 2 years. In wave 3 (2015–2016), 15 331 participants successfully responded. People with diabetes or high blood glucose at baseline were excluded: reporting a diagnosed history of diabetes, fasting blood glucose ≥ 126 mg/dl or HbA1c ≥ 6.5 %. We further excluded people who reported a history of heart disease, stroke or cancer. In addition, people with a BMI < 14 or > 40 kg/m² or no data on meal frequency were further excluded. Finally, 8874 Chinese men and women were eligible for this analysis (Fig. 1).

All the participants completed a written informed consent before the survey. Ethics approval for CHARLS was obtained from the Biomedical Ethics Review Committee of Peking University (IRB00001052-11015).

Assessment of meal frequency

Participants were asked about meal frequency by the question 'how many meals do you normally eat every day'. Six options were provided: 'more than 4 meals per day', '4 meals per day', '3 meals per day', '2 meals per day', '1 meal per day' and '< 1 meal per day'. Because very limited people eating more than four meals per day or less than two meals per day, the main analysis was restricted in people eating 2–4 meals per day. In a sensitivity analysis, we divided the participants into three groups: ≤ 2 meals per day, 3 meals per day and ≥ 4 meals per day. In this questionnaire, meal frequency was defined based on main meals with staple food at customary or regular occasions. Snacks without staple food, such as beverages, desserts and chips, were not considered as a meal.

Covariates measurement

Data on demographic background, physical and mental health status, and lifestyles were based on interviewer-administered questionnaires. Body weight, height and waist circumference were obtained from physical examination by well-trained clinicians. Participants erect on the floor board of the stadiometer and stand on scale, with shoes off. Waist circumference was measured by a soft measuring tape around waist, over clothing, holding securely at the level of navel. The BMI was calculated as weight (kg)/height (m)². Venous blood samples were collected and measured for biomarkers, including fasting blood glucose, HbA1c and lipid profiles at the KingMed laboratory. The details of the assay used have been described previously⁽¹⁷⁾.




Fig. 1. Flow chart of participant selection.

Outcome assessment

Participants were considered as cases of type 2 diabetes if they were diagnosed to have diabetes by doctors, had fasting blood glucose ≥ 126 mg/dl or had HbA1c ≥ 6.5 % at wave 3 survey^(18–20). Validity of outcome assessment has been established elsewhere^(18–20).

Statistical analyses

One-way ANOVA and Pearson χ^2 tests were used to compare the mean values and percentages of characteristics at wave 1. To examine the association between meal frequency and type 2 diabetes risk, we estimated relative risks (RR) and 95% CI by using the multiple Poisson regression models, which could be used to estimate the incidence of a disease among exposed and unexposed people^(21,22). People eating three meals per day were used as the reference group. Model 1 adjusted for age only. Model 2 further adjusted for sex, study area (urban *v.* rural areas), education level (primary school or lower, middle school, high school, college or higher education), marital status (married, divorced, widowed or never married), level of physical activity (0, 1–3, 4–6 and 7 d/week), history of hypertension (yes or no), history of dyslipidemia (yes or no), smoking status (never, past or current smoking) and drinking status (never drinking, past drinking, less than once/month or more than once/month). Baseline BMI was further adjusted in additional analysis. We further conducted stratified analyses according to baseline BMI (< 25 *v.* ≥ 25 kg/m²) to test whether the association could be modified by BMI. All analyses performed using SAS version 9.4 (SAS Institute Inc.). *P* values < 0.05 were considered statistically significant.

Results

Table 1 shows the baseline characteristics of 8874 Chinese men and women according to their meal frequency. The numbers of

Table 1. Baseline characteristics of Chinese people according to meal frequency

	Two times per day (n 110)			Three times per day (n 7574)			Four times per day (n 1190)		
	%	Mean	SD	%	Mean	SD	%	Mean	SD
Age (year)		60.0	8.9		58.5	9.2		59.6	10.2
Men	53.4			49.0			48.5		
BMI (kg/m ²)		22.1	3.4		23.2	3.4		22.6	3.4
Waist circumference (cm)		79.5	12.2		83.5	12.2		82.8	12.2
BMI ≥ 25 (kg/m ²)	16.3			28.5			22.4		
Urban	16.4			34.2			23.4		
High school or higher education	6.5			10.3			6.3		
Married	83.8			89.1			82.6		
Hypertension	7.4			18.3			13.8		
Dyslipidemia	2.9			5.7			4.4		
Current smoker	30.1			32.2			37.6		
Current drinker	24.5			35.6			30.6		
Vagarious activity every day	47.7			25.2			24.9		
Moderate activity every day	60.1			47.2			40.3		
HbA1c (%)		5.12	0.5		5.10	0.5		5.06	0.5
Fasting blood glucose (mg/dl)		96	19		100	17		99	17
TAG (mg/dl)		105	51		118	53		115	50
TC (mg/dl)		193	37		193	38		189	38
LDL-cholesterol (mg/dl)		117	33		118	34		114	34
HDL-cholesterol (mg/dl)		55	14		52	15		51	15

TC, total cholesterol.

All values were means (standard deviation) or percentage.

participants who ate meals two times per day, three times per day and four times per day were 110 (1.2%), 7574 (85.4%) and 1190 (13.4%), respectively. Compared with people who ate three meals per day, people eating four meals per day appeared to be slightly older and have a lower BMI and a lower waist circumference. Also, they were more likely to smoke and live in rural area but less likely to drink, be married, and have a higher education level and a history of hypertension. The baseline HbA1c, fasting blood glucose and lipid profile did not significantly differ by meal frequency (Table 1).

We identified 706 cases of type 2 diabetes among 8874 Chinese men and women over 4 years of follow-up. As is shown in Table 2, compared with people who ate three meals per day, those eating four meals per day had a lower risk of developing type 2 diabetes in the age-adjusted model (RR = 0.73 (0.57, 0.92)). After adjustment for sex, study area, education level, marital status, level of physical activity, history of hypertension, history of dyslipidemia, smoking status and drinking status, the association changed little (RR = 0.73 (0.58, 0.92)). After further adjustment for baseline BMI, the association was slightly attenuated but remained statistically significant (RR = 0.76 (0.60, 0.97)). Eating two meals per day was not associated with risk of type 2 diabetes in all models. In the sensitivity analysis with three groups (≤ 2 , 3 and ≥ 4 meals per day) yielded similar results (online Supplementary Table 1).

We next conducted a stratified analysis by baseline BMI. People eating two meals per day were excluded from this analysis because of a small number in this group. The fully-adjusted RR of type 2 diabetes for people eating four meals per day were 0.66 (0.48, 0.91) and 0.93 (0.65, 1.34) among those had a BMI < 25 and ≥ 25 kg/m², respectively (Table 3). Interaction tests suggested that baseline BMI did not modify the association between meal frequency and incidence of type 2 diabetes ($P_{\text{interaction}} = 0.11$).

Discussion

In this prospective study of Chinese people, we observed that eating four meals per day, compared with eating three meals per day, was associated with lower risk of developing type 2 diabetes. This association was evident among people with a BMI < 25 kg/m² but was non-significant among people with a BMI ≥ 25 kg/m². Eating two meals per day was not associated with type 2 diabetes.

Potential mechanisms underlying the association between meal frequency and the risk of type 2 diabetes warrant discussion. Several lines of evidence from cross-sectional studies have suggested an inverse association between increased eating frequency and prevalence of obesity⁽⁷⁾. Specifically, there was evidence that four meals per day compared with three meals per day was associated with a lower prevalence of general and central obesity^(23,24). Furthermore, a RCT found an increase in fat mass among young lean male participants who switches from four meals per day to three meals per day (the usual afternoon meal was skipped)⁽²⁵⁾. In addition, higher eating frequency may also improve blood lipid concentrations. A recent meta-analysis of twenty-one RCT suggested that higher meal frequency may help improve total cholesterol and low-density lipoprotein cholesterol⁽²⁶⁾, though blood glucose was not affected. In a subgroup analysis of trials > 12 weeks, higher meal frequency may also reduce serum TAG and increase HDL-cholesterol. Overall, eating frequency may be one of the contributors affecting body weight, fat deposition and serum lipid concentrations, which involves in the pathogenesis of type 2 diabetes⁽²⁷⁾.

To our knowledge, only three prospective cohort studies^(13–15) have reported the association between meal frequency and risk of type 2 diabetes. The Nurses' Health Study (NHS) of US women showed that in comparison with women who ate three times per day, the RR for type 2 diabetes were 1.09 (0.84, 1.41) for women

Table 2. Association between meal frequency and type 2 diabetes

	Two times per day		Three times per day		Four times per day	
	RR	95 % CI	RR	95 % CI	RR	95 % CI
No. of cases		9		626		71
No. of participants	110		7574		1190	
RR, model 1	1.11	0.61, 2.02	1.00		0.73	0.57, 0.92
RR, model 2	1.17	0.64, 2.14	1.00		0.73	0.58, 0.92
RR, model 2 + BMI	1.25	0.68, 2.27	1.00		0.76	0.60, 0.97

RR, relative risk.

Model 1: Age.

Model 2: Further adjusted for sex, study area, highest education level, marital status, level of vigorous activity, level of moderate activity, history of hypertension, history of dyslipidemia, smoking status and drinking status.

Table 3. Association between meal frequency and type 2 diabetes by baseline BMI

	Three times per day	Four times per day		
		RR	95 % CI	<i>P</i> _{interaction}
Baseline BMI < 25 (kg/m ²)				
No. of cases	342	38		
No. of participants	5407	930		
RR, model 1	1.00	0.65	0.47, 0.90	0.13
RR, model 2	1.00	0.64	0.47, 0.89	0.07
RR, model 2 + BMI	1.00	0.66	0.48, 0.91	0.11
Baseline BMI ≥ 25 (kg/m ²)				
No. of cases	284	33		
No. of participants	2167	260		
RR, model 1	1.00	0.95	0.67, 1.36	
RR, model 2	1.00	0.94	0.66, 1.35	
RR, model 2 + BMI	1.00	0.93	0.65, 1.34	

RR, relative risk.

Model 1: Age.

Model 2: Further adjusted for sex, study area, highest education level, marital status, level of vigorous activity, level of moderate activity, history of hypertension, history of dyslipidemia, smoking status and drinking status.

who ate 1–2 times per day and 1.13 (1.00, 1.27) for women who ate 4–5 times per day. However, after further adjusting for baseline BMI, no significant association remained (RR: 1.13 (0.87, 1.46) and 1.04 (0.92, 1.17), respectively)⁽¹³⁾. The Health Professionals Follow-Up Study (HPFS) of US men demonstrated that participants who ate 1–2 times per day or four times per day were at higher risk of type 2 diabetes than those who ate three times per day (RR: 1.26 (1.09, 1.46) and 1.19 (1.07, 1.32), respectively). After further adjustment for BMI, the risk for men who ate 4–5 times per day became non-significant (RR: 1.09 (0.99, 1.21))⁽¹⁴⁾. Another cohort study from the Women's Health Initiative Dietary Modification Trial (WHI-DM) showed similar results that four meals per day compared with 1–3 meals per day was associated with an increased risk of type 2 diabetes in postmenopausal women (HR: 1.38 (1.03, 1.84))⁽¹⁵⁾. Overall, these three cohort studies of US populations showed an increase in risk of type 2 diabetes among people eating four or five times per day, which may be partly explained by baseline BMI.

Of note, the above three studies were all conducted in the USA, where the prevalence of overweight and obesity is high. The baseline BMI in US adults was substantially greater than that observed in our study. Our study included lean participants with

a mean BMI ranging from 22 to 23 kg/m² across groups, whereas the mean BMI in the US studies was greater than 25 kg/m² (and even >28 kg/m² in WHI-DM study). Our subgroup analysis observed no significant association among those with a BMI ≥ 25 kg/m², which was in line with the results from NHS and HPFS study. Participants with a higher BMI may have chronic inflammation, dyslipidemia, impaired postprandial metabolism or insulin resistance^(28,29). These metabolic disorders may attenuate the potentially beneficial effects of a higher meal frequency. Moreover, the questionnaires about eating frequency in the US studies included meals and snacks, while in our study, only meals with staple food were considered as diet frequency. There was evidence that additional snacks beyond the main meals were associated with type 2 diabetes risk^(14,30). Taken together, differences in participant baseline BMI and diet frequency measurements may be responsible for the discrepant results between this study and previous ones.

Our study was not powered to evaluate the association of eating two meals per day with type 2 diabetes due to a small number of participants in this group. A 23-year cohort study demonstrated that skipping meals to lose weight was associated with higher risk of type 2 diabetes among men, elevated cholesterol, high blood pressure and restless sleep⁽³¹⁾. In addition, emerging evidence suggested that skipping breakfast was associated with increased risks of type 2 diabetes and CVD^(3,32), probably by maintaining a healthy weight and lipid profile⁽³³⁾. Unfortunately, we had no data on which meal was skipped and could, therefore, not examine the association for breakfast skipping.

This is the first prospective study that showed an inverse association of meal frequency with the risk of type 2 diabetes incidence. We used nationally representative data and a prospective cohort design. However, limitations of the present study should be acknowledged. First, the CHARLS study did not conduct dietary assessment and we were therefore unable to control for food, nutrient and total energy intakes. Although a high-quality diet was associated with lower risk of type 2 diabetes⁽³⁴⁾, evidence on association between meal frequency and diet quality was inconsistent^(35–37). Furthermore, a recent meta-analysis of RCT also found that no significant effects were observed on energy intake comparing the different meal frequencies⁽¹²⁾. In line with these reports, the NHS study showed that further adjustment for total energy intake, cereal fibre intake and the Alternative Health Eating Index 2010 changed the results little⁽¹³⁾. Nevertheless, residual confounding due to dietary

intakes (e.g. snacks) and other unmeasured risk factors (e.g. family history of diabetes) remained a possible explanation for the observed association. For example, snacks are usually fatty, sugary, salt and fried foods with low nutrient profiles and high in energies, which are associated with weight gain and higher risk of type 2 diabetes^(14,30,38). Thus, it might bias the association towards a higher risk without adjusting snacks. Second, the validity of meal frequency has not been carried out. The self-report dietary information was subject to misreporting and measurement error. However, such measurement error was likely to be non-differential in the prospective study and led to associations being biased towards the null. Moreover, lifestyle changes (e.g. smoking, drinking status changes) could also have an impact on the outcomes. Third, our observations were conducted in a Chinese population, caution should be paid to generalise the results to other populations with different genetic backgrounds, diet habits and lifestyles.

In conclusion, eating four meals per day, compared with eating three meals per day, was associated with lower risk of developing type 2 diabetes in a Chinese population, particularly in those with a BMI < 25kg/m². The results from our study suggested that meal frequency might be a potential modifiable factor for type 2 diabetes and having an appropriate meal frequency may have a role in the prevention of type 2 diabetes.

Acknowledgements

The authors would like to thank all the participants in the CHARLS study. Also, the authors appreciate the National School of Development of Peking University for providing the data.

The CHARLS study was supported by the National Institute on Aging of the National Institute of Health (No. R01-AG037031-01, R01AG037031-03S1, R03-TW008358-01 and R21-AG031372-01), the Natural Science Foundation of China (No. 70910107022 and 71130002), and the Knowledge for Change Program at the World Bank (No. 7159234). The sponsors had no roles in study design, data collection, analysis, and interpretation, or manuscript writing.

X. W. and J.-Y. D. designed the study and analysed the data. X. W. drafted the manuscript. All authors conducted the technique review and edited the manuscript.

The authors declared no competing interests.

Supplementary material

For supplementary material/s referred to in this article, please visit <https://doi.org/10.1017/S0007114521003226>

References

- Zarrinpar A, Chaix A & Panda S (2016) Daily eating patterns and their impact on health and disease. *Trends Endocrinol Metab* **27**, 69–83.
- Paoli A, Tinsley G, Bianco A, *et al.* (2019) The influence of meal frequency and timing on health in humans: the role of fasting. *Nutrients* **11**, 719.
- Ballon A, Neuenschwander M & Schlesinger S (2019) Breakfast skipping is associated with increased risk of type 2 diabetes among adults: a systematic review and meta-analysis of prospective cohort studies. *J Nutr* **149**, 106–113.
- Ohkuma T, Hirakawa Y, Nakamura U, *et al.* (2015) Association between eating rate and obesity: a systematic review and meta-analysis. *Int J Obes* **39**, 1589–1596.
- Kudo A, Asahi K, Satoh H, *et al.* (2019) Fast eating is a strong risk factor for new-onset diabetes among the Japanese general population. *Sci Rep* **9**, 8210.
- Radzeviciene L & Ostrauskas R (2013) Fast eating and the risk of type 2 diabetes mellitus: a case-control study. *Clin Nutr* **32**, 232–235.
- Wang YQ, Zhang YQ, Zhang F, *et al.* (2016) Increased eating frequency is associated with lower obesity risk, but higher energy intake in adults: a meta-analysis. *Int J Environ Res Public Health* **13**, 603.
- Maugeri A, Kunzova S, Medina-Inojosa JR, *et al.* (2018) Association between eating time interval and frequency with ideal cardiovascular health: results from a random sample Czech urban population. *Nutr Metab Cardiovasc Dis* **28**, 847–855.
- Titan SM, Bingham S, Welch A, *et al.* (2001) Frequency of eating and concentrations of serum cholesterol in the Norfolk population of the European prospective investigation into cancer (EPIC-Norfolk): cross sectional study. *BMJ* **323**, 1286–1288.
- Ha K & Song Y (2019) Associations of meal timing and frequency with obesity and metabolic syndrome among Korean Adults. *Nutrients* **11**, 2437.
- Schoenfeld BJ, Aragon AA & Krieger JW (2015) Effects of meal frequency on weight loss and body composition: a meta-analysis. *Nutr Rev* **73**, 69–82.
- Schwingshackl L, Nitschke K, Zahringer J, *et al.* (2020) Impact of meal frequency on anthropometric outcomes: a systematic review and network meta-analysis of randomized controlled trials. *Adv Nutr* **11**, 1108–1122.
- Mekary RA, Giovannucci E, Cahill L, *et al.* (2013) Eating patterns and type 2 diabetes risk in older women: breakfast consumption and eating frequency. *Am J Clin Nutr* **98**, 436–443.
- Mekary RA, Giovannucci E, Willett WC, *et al.* (2012) Eating patterns and type 2 diabetes risk in men: breakfast omission, eating frequency, and snacking. *Am J Clin Nutr* **95**, 1182–1189.
- Neuhouser ML, Wertheim BC, Perrigue MM, *et al.* (2020) Associations of number of daily eating occasions with type 2 diabetes risk in the women's health initiative dietary modification trial. *Curr Dev Nutr* **4**, nzaa126.
- Zhao Y, Hu Y, Smith JP, *et al.* (2014) Cohort profile: the China Health and Retirement Longitudinal Study (CHARLS). *Int J Epidemiol* **43**, 61–68.
- Chen X, Crimmins E, Hu PP, *et al.* (2019) Venous blood-based biomarkers in the China health and retirement longitudinal study: rationale, design, and results from the 2015 wave. *Am J Epidemiol* **188**, 1871–1877.
- American Diabetes A (2017) 2. Classification and diagnosis of diabetes. *Diabetes Care* **40**, S11–S24.
- Chen G, Yi Q, Hou L, *et al.* (2021) Transition of hypertriglyceridemic-waist phenotypes and the risk of type 2 diabetes mellitus among middle-aged and older Chinese: a National Cohort Study. *Int J Environ Res Public Health* **18**, 3664.
- Peng J, Zhao F, Yang X, *et al.* (2021) Association between dyslipidemia and risk of type 2 diabetes mellitus in middle-aged and older Chinese adults: a secondary analysis of a nationwide cohort. *BMJ Open* **11**, e042821.
- Spiegelman D & Hertzmark E (2005) Easy SAS calculations for risk or prevalence ratios and differences. *Am J Epidemiol* **162**, 199–200.

22. McNutt L-A, Wu C, Xue X, *et al.* (2003) Estimating the relative risk in cohort studies and clinical trials of common outcomes. *Am J Epidemiol* **157**, 940–943.
23. Ma Y, Bertone ER, Stanek EJ, *et al.* (2003) Association between eating patterns and obesity in a free-living US adult population. *Am J Epidemiol* **158**, 85–92.
24. Holmback I, Ericson U, Gullberg B, *et al.* (2010) A high eating frequency is associated with an overall healthy lifestyle in middle-aged men and women and reduced likelihood of general and central obesity in men. *Br J Nutr* **104**, 1065–1073.
25. Chapelot D, Marmonier C, Aubert R, *et al.* (2006) Consequence of omitting or adding a meal in man on body composition, food intake, and metabolism. *Obesity* **14**, 215–227.
26. Abdollahi S, Kazemi A, de Souza RJ, *et al.* (2021) The effect of meal frequency on biochemical cardiometabolic factors: a systematic review and meta-analysis of randomized controlled trials. *Clin Nutr* **40**, 3170–3181.
27. Stumvoll M, Goldstein BJ & van Haeften TW (2005) Type 2 diabetes: principles of pathogenesis and therapy. *Lancet* **365**, 1333–1346.
28. Monteiro R & Azevedo I (2010) Chronic inflammation in obesity and the metabolic syndrome. *Mediators Inflamm* **2010**, 289645.
29. Kumar AA, Sathesh G, Vijayakumar G, *et al.* (2020) Postprandial metabolism is impaired in overweight normoglycemic young adults without family history of diabetes. *Sci Rep* **10**, 353.
30. McNaughton SA, Mishra GD & Brunner EJ (2008) Dietary patterns, insulin resistance, and incidence of type 2 diabetes in the Whitehall II Study. *Diabetes Care* **31**, 1343–1348.
31. Carew AS, Mekary RA, Kirkland S, *et al.* (2021) Prospective study of skipping meals to lose weight as a predictor of incident type 2 diabetes with potential modification by cardiometabolic risk factors: the Canadian 1995 Nova Scotia Health Survey. *Can J Diabetes* **45**, 306–312.
32. Rong S, Snetelaar LG, Xu G, *et al.* (2019) Association of skipping breakfast with cardiovascular and all-cause mortality. *J Am Coll Cardiol* **73**, 2025–2032.
33. Uzhova I, Fuster V, Fernandez-Ortiz A, *et al.* (2017) The importance of breakfast in atherosclerosis disease: insights from the PESA Study. *J Am Coll Cardiol* **70**, 1833–1842.
34. Yu D, Zheng W, Cai H, *et al.* (2018) Long-term diet quality and risk of type 2 diabetes among urban Chinese adults. *Diabetes Care* **41**, 723–730.
35. Murakami K & Livingstone MB (2016) Associations between meal and snack frequency and diet quality in US Adults: National Health and Nutrition Examination Survey 2003–2012. *J Acad Nutr Diet* **116**, 1101–1113.
36. Murakami K & Livingstone MB (2016) Associations between meal and snack frequency and diet quality and adiposity measures in British adults: findings from the National Diet and Nutrition Survey. *Public Health Nutr* **19**, 1624–1634.
37. Leech RM, Livingstone KM, Worsley A, *et al.* (2016) Meal frequency but not snack frequency is associated with micronutrient intakes and overall diet quality in Australian men and women. *J Nutr* **146**, 2027–2034.
38. McCrory MA & Campbell WW (2011) Effects of eating frequency, snacking, and breakfast skipping on energy regulation: symposium overview. *J Nutr* **141**, 144–147.