





MONOGENIC DIABETES AND OTHER GENETIC DISORDERS OF GLUCOSE METABOLISM

Increased referrals for congenital hyperinsulinism genetic testing in children with trisomy 21 reflects the high burden of non-genetic risk factors in this group

Thomas I. Hewat¹  | Thomas W. Laver¹ | Jayne A. L. Houghton² |
Jonna M. E. Männistö³ | Sabah Alvi⁴ | Stephen P. Brearey⁵ | Declan Cody⁶ |
Antonia Dastamani⁷ | Miguel De los Santos La Torre⁸  | Nuala Murphy⁹ |
Birgit Rami-Merhar¹⁰  | Birgit Wefers¹¹ | Hanna Huopio¹² |
Indraneel Banerjee¹³ | Matthew B. Johnson¹ | Sarah E. Flanagan¹ 

¹Institute of Biomedical and Clinical Science, University of Exeter Medical School, Exeter, UK

²Exeter Genomics Laboratory, Royal Devon and Exeter NHS Foundation Trust, Exeter, UK

³Department of Pediatrics, University of Eastern Finland and Kuopio University Hospital, Kuopio, Finland

⁴Leeds Children's Hospital, Leeds, UK

⁵Countess of Chester Hospital, Chester, UK

⁶Children's Health Ireland at Crumlin, Dublin, Ireland

⁷Endocrinology Department, Great Ormond Street Hospital for Children NHS Foundation Trust, London, UK

⁸Department of Endocrinology and Metabolism of the Instituto Nacional de Salud del Niño, Lima, Peru

⁹Children's University Hospital, Dublin, Ireland

¹⁰Department of Pediatric and Adolescent Medicine, Medical University of Vienna, Vienna, Austria

¹¹Ninewells Hospital, Dundee, UK

¹²Department of Pediatrics, Kuopio University Hospital, Kuopio, Finland

¹³Department of Paediatric Endocrinology, Royal Manchester Children's Hospital, Manchester, UK

Correspondence

Sarah E. Flanagan, University of Exeter Medical School, Exeter EX2 5DW, UK.
Email: s.flanagan@exeter.ac.uk

Funding information

Research England; Royal Society; Wellcome Trust

Abstract

Background: Hyperinsulinism results from inappropriate insulin secretion during hypoglycaemia. Down syndrome is causally linked to a number of endocrine disorders including Type 1 diabetes and neonatal diabetes. We noted a high number of individuals with Down syndrome referred for hyperinsulinism genetic testing, and therefore aimed to investigate whether the prevalence of Down syndrome was increased in our hyperinsulinism cohort compared to the population.

Methods: We identified individuals with Down syndrome referred for hyperinsulinism genetic testing to the Exeter Genomics Laboratory between 2008 and 2020. We sequenced the known hyperinsulinism genes in all individuals and investigated their clinical features.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *Pediatric Diabetes* published by John Wiley & Sons Ltd.

Results: We identified 11 individuals with Down syndrome in a cohort of 2011 patients referred for genetic testing for hyperinsulinism. This represents an increased prevalence compared to the population (2.5/2011 expected vs. 11/2011 observed, $p = 6.8 \times 10^{-5}$). A pathogenic *ABCC8* mutation was identified in one of the 11 individuals. Of the remaining 10 individuals, five had non-genetic risk factors for hyperinsulinism resulting from the Down syndrome phenotype: intrauterine growth restriction, prematurity, gastric/oesophageal surgery, and asparaginase treatment for leukaemia. For five individuals no risk factors for hypoglycaemia were reported although two of these individuals had transient hyperinsulinism and one was lost to follow-up.

Conclusions: Down syndrome is more common in patients with hyperinsulinism than in the population. This is likely due to an increased burden of non-genetic risk factors resulting from the Down syndrome phenotype. Down syndrome should not preclude genetic testing as coincidental monogenic hyperinsulinism and Down syndrome is possible.

1 | INTRODUCTION

Hyperinsulinism (HI) is a disorder of the pancreatic beta-cell where inappropriately high levels of insulin are secreted leading to hypoglycaemia. Prolonged neonatal HI can be transient, often remitting within 6 months, with risk factors including male sex, low birth weight, and perinatal stress.¹ In contrast, persistent HI is likely to be genetic with disease-causing mutations in single genes identified in 50%–70% of cases.^{2,3} HI has also been reported as a rare feature in patients with aneuploidies. For example, HI can present in females with Turner syndrome resulting from a complete or partial monosomy of the X chromosome and in children with Patau syndrome resulting from mosaic trisomy 13.^{4,5}

The most common aneuploidy is trisomy 21, causing Down syndrome, which affects 1 in 794 live births in the USA.⁶ Down syndrome is characterized by intellectual disability, microcephaly, congenital heart defects, gastrointestinal disorders, and endocrine disorders which include Type 1 diabetes or neonatal diabetes.^{7–9} Whilst HI has not been reported as a feature of Down syndrome, we noted a high number of individuals with the co-existence of these two conditions being referred to our laboratory for genetic testing. Our aim was to assess whether the prevalence of children with HI and Down syndrome was higher than expected in our cohort and if so to determine the reason(s) for this.

2 | METHODS

We studied 2011 individuals referred for HI genetic testing to the Exeter Genomics Laboratory between 2008 and 2020. Clinical information was provided at referral using a standardized request form. Follow-up data by case note review were requested for all individuals with HI and Down syndrome.

We performed targeted next-generation sequencing of 13 known HI genes including *ABCC8*, *CACNA1D*, *CDKN1C*, *GCK*, *GLUD1*, *HADH*, *HNF1A*, *HNF4A*, *INSR*, *KCNJ11*, *PMM2*, *SLC16A1*, and *TRMT10A* in all individuals with HI and Down syndrome using previously described methods.¹⁰ We used Stata/SE v16.0 to perform a one-sample binomial test to assess if the prevalence of Down syndrome in our cohort was significantly higher than the population prevalence (Stata Corp, College Station, TX, USA).

Informed consent was obtained from the parents or guardians of all probands. This study was approved by the North Wales Research Ethics Committee (517/WA/0327).

3 | RESULTS

Within our international cohort of 2011 individuals, we identified 11 cases with Down syndrome ($n = 11/2011$ [0.55%]). This represents a minimal prevalence as we do not routinely screen for aneuploidies, and some clinicians may not have provided this information on the genetic request form. The number of children with Down syndrome was significantly higher than expected by chance given the population prevalence of Down syndrome of 12.6/10,000⁶ (2.5/2011 expected vs. 11/2011 observed, $p = 6.8 \times 10^{-5}$).

We identified a mutation in a known HI gene in 1/11 (9%) patients. This individual had a pathogenic paternally inherited *ABCC8* mutation.¹¹ Of the 10 individuals without a mutation in a known gene, two were born with intrauterine growth retardation (IUGR) (birth weight Z-score < -2). The median age at diagnosis at HI of the 10 individuals was 101 days (IQR 1–581 days) with insulin detected at the time of hypoglycaemia (plasma glucose <2.8 mmol/L) in all cases. Persistent HI (defined here as requiring treatment for >6 months) was confirmed in four of the 10 genetically unsolved individuals. In the remaining six individuals the HI was transient ($n = 5$) or follow-up

TABLE 1 Clinical features of patients diagnosed with Down syndrome and Hyperinsulinism

	Patient 1	Patient 2	Patient 3	Patient 4	Patient 5	Patient 6	Patient 7	Patient 8	Patient 9	Patient 10	Patient 11
Genetic results	No mutation detected	No mutation detected	No mutation detected	No mutation detected	No mutation detected	No mutation detected	No mutation detected	No mutation detected	No mutation detected	No mutation detected	ABCC8 p.G25fs/N**
Sex	Female	Female	Male	Male	Female	Female	Male	Female	Male	Female	Male
Birth weight SDS	-2.55	Not available	1.35	0.25	-0.77	0.22	-1.07	0.89	-3.74	1.25	0.65
Age at HI diagnosis (weeks)	36	20	120	0.14	83	0.14	0.14	0.14	9	208	0.43
Glucose (mmol/L) (Insulin [pmol/L]) at diagnosis	2 (108)	1.5 (60.5)	1 (18.3)	0.4 (26)	1.6 (88)	1.9 (12.8)	2.0 (347)	1.8 (12.5)	2.8 (4.5)	2.4 (141)	1.1 (56.1)
Post-prandial hypoglycaemia	Not noted	Not noted	No	No	Yes	Not noted	Not noted	Not noted	Not noted	Not noted	Not noted
Transient/persistent HI	Persistent Diazoxide treatment ongoing at 13 years (3 mg/kg/day)	Transient No treatment required	Persistent Pancreatectomy due to side effects of diazoxide and no response to octreotide	Transient Treated transiently with I.V. glucose and increased feeds	Persistent Diazoxide unresponsive, managed with continuous feeds until remission at 3 years following VSD correction	Transient Treated with diazoxide until 4 months	Transient No treatment required	Unknown Lost to follow-up (10 mg/kg/day diazoxide at referral)	Persistent Treated with diazoxide until 12 months	Transient Treated transiently with I.V. glucose	Persistent Octreotide (20ug/kg/day) ongoing at 5 years
Gastric or oesophageal surgery (age)	No	Yes, prior to HI diagnosis (16 weeks)	Yes, following HI diagnosis (3 years)	No	Yes, prior to HI diagnosis (1st, 4 weeks, 2nd 1 year)	Yes, following HI diagnosis (>0.14 weeks)	No	Yes (age unknown)	Yes, following HI diagnosis (26 weeks)	No	Yes, following HI diagnosis (0.86 weeks)
Additional features	ASD	GORD, VSD	West syndrome, GORD, asthma	Mild hypoventilation, ASD	Tracheo-oesophageal fistula, GORD, VSD, jaundice, portosystemic shunt	Duodenal atresia, ASD, PDA	Prematurity (31/40), perinatal compromise (poor CTG, reduced movements, at birth: raised lactate, biochemical evidence of liver & renal compromise) Cerebral palsy with right hemiplegia 2nd to left Periventricular Leukomalacia, Cataracts, Hearing loss 2nd auditory neuropathy	Duodenal stenosis, haematuria	Tracheomalacia	Acute lymphoblastic leukaemia treated with L-asparaginase at 3.8 years	Duodenal atresia

Note: Gray-filled boxes represent risk factors for Hyperinsulinism (HI). ASD = atrial septal defect. VSD = ventricular septal defect. PDA = patent ductus arteriosus. GORD = gastro-oesophageal reflux disease. I.V. intravenous. For the purposes of this study persistent disease is defined as HI requiring treatment for >6 months and transient disease is defined as HI requiring treatment for <6 months. Patient 3 previously reported in.¹² Patient 5 previously reported in.¹³ ** indicates that mutation was previously reported in.¹¹

information was not available ($n = 1$). One individual with persistent HI demonstrated side-effects to diazoxide and did not respond to octreotide, necessitating a near-total pancreatectomy.¹² Consanguinity was reported in this individual.

Seven individuals, including the child with an *ABCC8* mutation, had undergone gastric or oesophageal surgery for duodenal atresia, duodenal stenosis, tracheomalacia, or gastro-oesophageal reflux disease (GORD). In two cases surgery had been performed prior to the onset of HI. One of these cases had also undergone surgery to repair a portosystemic shunt.¹³ A further individual had been diagnosed with acute lymphoblastic leukaemia and had received L-asparaginase treatment prior to the onset of HI. An overview of the clinical features of the cohort are provided in Table 1.

4 | DISCUSSION

We identified 11 individuals with HI and Down syndrome. Given that Down syndrome has an approximate incidence of one in 794 live births, we would have expected two or three individuals with Down syndrome in our cohort of 2011 individuals.⁶ The statistically significant enrichment and higher prevalence therefore suggest that the two conditions are related.

The prevalence of mutations in the known genes was low in the Down syndrome and HI cohort ($n = 1/11$, 9%) although this increased to 20% in those with confirmed persistent HI ($n = 1/5$). This pick-up rate is lower than anticipated given previous studies have reported mutations in the known genes in 50%–70% of HI cases.^{2,3} While this may reflect the small sample size, it is also possible that the Down syndrome is increasing the risk of the child developing HI.

We identified risk factors for developing HI in five of the 10 individuals without a mutation in a known gene. Two children had surgery to correct a gastrointestinal (GI) disorder prior to the onset of HI (Table 1). GI disorders are common in individuals with Down syndrome and surgical management of this can lead to iatrogenic hypoglycaemia as a result of dumping syndrome.^{14,15} Furthermore, one of these individuals had confirmed post-prandial hypoglycaemia following surgery lending further support to this diagnosis.¹⁶ This patient also had a portosystemic shunt, with surgical closure resulting in a resolution of the hypoglycaemia.¹³ In four further cases, gastric surgery was performed but this occurred after the onset of HI in three cases suggesting that the HI was unlikely to be due to gastric surgery induced post-prandial hypoglycaemia. The age at gastric surgery in the remaining patient was unknown.

IUGR or biochemical evidence of perinatal and postnatal stress associated with prematurity, was reported in two individuals. These are well-recognized risk factors for prolonged neonatal hypoglycaemia.¹ IUGR was reported in a second individual however the HI was ongoing at the age of 13 years suggesting it was not causative of the hypoglycaemia.¹⁷

One individual had been diagnosed with acute lymphoblastic leukaemia that had been treated with an L-asparaginase based chemotherapy prior to the onset of HI at 4 years. Children with Down

syndrome are at increased risk of developing acute lymphoblastic leukaemia and previous studies have shown that treatment with L-asparaginase can cause hypoglycaemia in younger patients.^{18,19} This could explain the transitory hypoglycaemia observed in this child.

Of the five individuals without an identifiable risk factor for HI, two had persistent HI, two had transient HI and one case was lost to follow-up which might suggest that the HI was transient and not severe. It is also possible that in this patient risk factors for HI were present but not reported at referral for genetic testing. The finding of two individuals with Down syndrome and persistent HI within our cohort is expected based on the population prevalence of Down syndrome. Interestingly, consanguinity was reported in one of these individuals, supporting the possibility of a recessively inherited monogenic etiology.

Recently, a study of HI in Finland identified five cases with Down syndrome in a cohort of 238 individuals. The authors noted that this was a statistically significant increase compared to the population prevalence of Down syndrome.²⁰ In keeping with our findings, screening of the known genes identified an *ABCC8* mutation in a single individual whilst the four mutation negative individuals had non-genetic risk factors for HI which could be attributed to the Down syndrome phenotype: extreme prematurity and cardiac insufficiency, IUGR, gastric surgery/fundoplication, and stress due to congenital heart defects (personal communication Huopio and Männistö). In two individuals the HI remitted before the age of 4 months.

Genetic testing identified an *ABCC8* mutation in one individual with Down syndrome in our cohort and this, together with the finding of an *ABCC8* mutation in an individual within the Finnish cohort, highlights the need to perform genetic testing in all individuals with persistent HI.²⁰ Whilst a diagnosis of Down syndrome does not preclude co-incidental monogenic HI, our study suggests HI in Down syndrome is most likely to be due to non-genetic risk factors.

In conclusion, we have identified an increased referral rate for HI genetic testing for individuals with Down syndrome. Our findings suggest that HI is not a feature of Trisomy 21 but a consequence of the high burden of non-genetic risk factors resulting from the Down syndrome phenotype.

FUNDING STATEMENT

TWL is the recipient of a Lectureship and MBJ an Independent Fellowship from the Exeter Diabetes Centre of Excellence funded by Research England's Expanding Excellence in England (E3) fund. SEF has a Sir Henry Dale Fellowship jointly funded by the Wellcome Trust and the Royal Society (Grant Number: 105636/Z/14/Z). This research was funded in whole, or in part, by Wellcome [105636/Z/14/Z]. For the purpose of open access, the author has applied a CC BY public copyright license to any Author accepted Manuscript version arising from this submission.

CONFLICT OF INTEREST

The authors have nothing to disclose.

AUTHOR CONTRIBUTION

T.I. Hewat, M.B. Johnson, and S.E. Flanagan designed the study. S. Alvi, S.P. Brearey, D. Cody, A. Dastamani, M. de los Santos la Torre, N. Murphy, B. Rami-Merhar, B. Wefers, I. Banerjee and S.E. Flanagan recruited patients to the study and with J.M.E Männistö and H. Huopio analysed the clinical data. J.A.L. Houghton and S.E. Flanagan performed the molecular genetic studies. T.I. Hewat, T.W. Laver, M.B. Johnson and S.E. Flanagan performed the data analysis. T.I. Hewat, M. B. Johnson and S.E. Flanagan prepared the draft manuscript. All authors contributed to the discussion of the results and to the manuscript preparation.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

ETHICAL APPROVAL

This study was approved by the North Wales Research Ethics Committee (517/WA/0327).

PATIENT CONSENT STATEMENT

Informed consent was obtained from the parents or guardians of all probands.

ORCID

Thomas I. Hewat  <https://orcid.org/0000-0002-5330-760X>

Miguel De los Santos La Torre  <https://orcid.org/0000-0002-9176-2988>

Birgit Rami-Merhar  <https://orcid.org/0000-0001-5575-5222>

Sarah E. Flanagan  <https://orcid.org/0000-0002-8670-6340>

REFERENCES

- Hoe FM, Thornton PS, Wanner LA, Steinkrauss L, Simmons RA, Stanley CA. Clinical features and insulin regulation in infants with a syndrome of prolonged neonatal hyperinsulinism. *J Pediatr*. 2006; 148(2):207-212.
- Güemes M, Rahman SA, Kapoor RR, et al. Hyperinsulinemic hypoglycemia in children and adolescents: recent advances in understanding of pathophysiology and management. *Rev Endocr Metab Disord*. 2020; 21(4):577-597.
- Snider KE, Becker S, Boyajian L, et al. Genotype and phenotype correlations in 417 children with congenital hyperinsulinism. *J Clin Endocrinol Metab*. 2013;98(2):E355-E363.
- Gibson CE, Boodhansingh KE, Li C, et al. Congenital Hyperinsulinism in infants with turner syndrome: possible association with monosomy X and KDM6A Haploinsufficiency. *Horm Res Paediatr*. 2018;89(6): 413-422.
- Tamame T, Hori N, Homma H, et al. Hyperinsulinemic hypoglycemia in a newborn infant with trisomy 13. *Am J Med Genet A*. 2004;129A (3):321-322.
- de Graaf G, Buckley F, Skotko BG. Estimates of the live births, natural losses, and elective terminations with down syndrome in the United States. *Am J Med Genet A*. 2015;167A(4):756-767.
- Startin CM, D'Souza H, Ball G, et al. Health comorbidities and cognitive abilities across the lifespan in down syndrome. *J Neurodev Disord*. 2020;12(1):4.
- Aitken RJ, Mehers KL, Williams AJ, et al. Early-onset, coexisting autoimmunity and decreased HLA-mediated susceptibility are the characteristics of diabetes in down syndrome. *Diabetes Care*. 2013; 36(5):1181-1185.
- Johnson MB, De Franco E, Greeley SAW, et al. Trisomy 21 is a cause of permanent neonatal diabetes that is autoimmune but not HLA associated. *Diabetes*. 2019;68(7):1528-1535.
- Ellard S, Lango Allen H, De Franco E, et al. Improved genetic testing for monogenic diabetes using targeted next-generation sequencing. *Diabetologia*. 2013;56(9):1958-1963.
- De Franco E, Saint-Martin C, Brusgaard K, et al. Update of variants identified in the pancreatic beta-cell KATP channel genes KCNJ11 and ABCC8 in individuals with congenital hyperinsulinism and diabetes. *Hum Mutat*. 2020;41(5):884-905.
- Shi Y, Avatapalle HB, Skae MS, et al. Increased plasma incretin concentrations identifies a subset of patients with persistent congenital hyperinsulinism without KATP channel gene defects. *J Pediatr*. 2015; 166(1):191-194.
- Senniappan S, Pitt K, Shah P, et al. Postprandial hyperinsulinaemic hypoglycaemia secondary to a congenital portosystemic shunt. *Horm Res Paediatr*. 2015;83(3):217-220.
- Chicoine B, Rivelli A, Fitzpatrick V, Chicoine L, Jia G, Rzhetsky A. Prevalence of common disease conditions in a large cohort of individuals with down syndrome in the United States. *J Patient Cent Res Rev*. 2021;8(2):86-97.
- Chesser H, Abdulhussein F, Huang A, Lee JY, Gitelman SE. Continuous glucose monitoring to diagnose hypoglycemia due to late dumping syndrome in children after gastric surgeries. *J Endocr Soc*. 2021; 5(3):bvaa197.
- Calabria AC, Charles L, Givler S, De Leon DD. Postprandial hypoglycemia in children after gastric surgery: clinical characterization and pathophysiology. *Horm Res Paediatr*. 2016;85(2):140-146.
- Fafoula O, Alkhayyat H, Hussain K. Prolonged hyperinsulinaemic hypoglycaemia in newborns with intrauterine growth retardation. *Arch Dis Child Fetal Neonatal Ed*. 2006;91(6):F467.
- Panigrahi M, Swain TR, Jena RK, Panigrahi A. L-asparaginase-induced abnormality in plasma glucose level in patients of acute lymphoblastic leukemia admitted to a tertiary care hospital of Odisha. *Indian J Pharmacol*. 2016;48(5):595-598.
- Brown AL, de Smith AJ, Gant VU, et al. Inherited genetic susceptibility to acute lymphoblastic leukemia in down syndrome. *Blood*. 2019; 134(15):1227-1237.
- Männistö JME, Jaaskelainen J, Otonkoski T, Huopio H. Long-term outcome and treatment in persistent and transient congenital Hyperinsulinism: a Finnish population-based study. *J Clin Endocrinol Metab*. 2021;106(4):e1542-e1551.

How to cite this article: Hewat TI, Laver TW, Houghton JAL, et al. Increased referrals for congenital hyperinsulinism genetic testing in children with trisomy 21 reflects the high burden of non-genetic risk factors in this group. *Pediatr Diabetes*. 2022; 23(4):457-461. doi:10.1111/pedi.13333