

Cloning and Characterization of a Tetracycline Resistance Determinant Present in *Agrobacterium tumefaciens* C58

ZHAO-QING LUO¹ AND STEPHEN K. FARRAND^{1,2*}

Departments of Crop Sciences¹ and Microbiology,² University of Illinois at Urbana-Champaign, Urbana, Illinois 61801

Received 14 September 1998/Accepted 7 November 1998

Agrobacterium tumefaciens C58 and its derivatives give rise to spontaneous mutants resistant to tetracycline at a high frequency. We observed that a mutation affecting a tRNA processing function significantly affected the emergence of such mutants, suggesting that C58 contained a positively acting gene conferring resistance to tetracycline. A cosmid clone conferring resistance to tetracycline in *Escherichia coli* and *Agrobacterium* was isolated from a genomic bank of one such mutant. Subcloning, transposon mutagenesis, and DNA sequence analysis revealed that this DNA fragment contained two divergently transcribed genes, *tetA* and *tetR*, encoding products that were very similar to proteins of the Tet(A) class of tetracycline resistance systems. In the clone from this mutant, *tetR* was disrupted by an IS426. The homologous region from wild-type NT1 contained an intact *tetR* gene and did not confer resistance to tetracycline. Hybridization analysis showed that of 22 members of the genus *Agrobacterium* surveyed, only strains C58 and T37 contained the *tet* determinant. Moreover, only these two strains mutated to resistance to this antibiotic. Unlike other Tet(A) systems, neither tetracycline nor a series of its derivatives induced the expression of this *tet* gene unit. Other polycyclic compounds, including many of plant origin, also did not induce this *tet* gene system. The divergent promoter region of this *tet* system contained a single inverted repeat element identical to one such operator repeat in the promoter region of the *tet* determinant from the IncP1 α R plasmid RP4. TetR repressor proteins from the *Agrobacterium tet* system and from RP4 interacted with the heterologous operators. While the repressive effect of the TetR protein from strain C58 (TetR_{C58}) on the *tetA* gene from strain RP4 (*tetA*_{RP4}) was not relieved by tetracycline, repression of *tetA*_{C58} by TetR_{RP4} was lifted by this antibiotic.

Genes encoding resistance to antibiotics have been identified in a wide spectrum of prokaryotic organisms, including gram-negative and gram-positive bacteria. Although most of the genes characterized to date have been isolated from clinical isolates, these determinants are believed to have originated in soil bacteria (38). More recently, genes encoding proteins that confer resistance to several different drugs, the multidrug systems, have been identified and characterized (37). Horizontal transfer of these determinants among diverse bacteria are challenging the effectiveness of antibiotics in infectious disease control (39, 48).

Genes conferring resistance to tetracycline are among the most abundant of the identified drug resistance elements (29). Although mechanisms such as target site protection and drug inactivation are known (7, 44), most tetracycline resistance determinants function through the efflux mechanism, pumping a drug-metal ion complex out of the bacterial cell by a process energized by the membrane proton gradient (37, 45).

It is a well-recognized but unpublished observation that *Agrobacterium tumefaciens* C58 and its derivatives give rise at a high frequency to spontaneous mutants resistant to tetracycline. Most genetic and molecular analyses concerning *A. tumefaciens* and its plasmids are conducted in the C58 chromosomal background. Since many broad-host-range cloning vectors encode resistance to tetracycline as the selectable marker, the propensity of this strain to mutate to tetracycline resistance poses problems to the genetic analysis of this bacterium and to its use as a plant transformation agent.

During a study of mutants affecting the induction of *tra* genes of pTiC58 in NT1 (51), a Ti plasmid-cured derivative of C58 (53), we observed that one mutant, NTM7, while remaining able to mutate to tetracycline resistance, expressed the phenotype at a much lower level than did its wild-type parent. NTM7 later was found to carry a Tn5 insertion in a locus encoding a protein homologous to the product of the *Escherichia coli rnd* gene (30). In *E. coli*, a *miaA* mutant is unable to express Tet(M)-mediated tetracycline resistance (8). Since both the *rnd* and *miaA* genes are involved in tRNA modification (12, 52), we reasoned that there may be a positively acting tetracycline resistance element in the genome of strain C58 which can function normally in the wild-type bacterium but not in the *rnd* mutant. A study was initiated to investigate the nature of such tetracycline-resistant mutants. We confirmed our hypothesis and report here the molecular cloning, nucleotide sequence, and genetic characterization of a tetracycline resistance gene unit from this *A. tumefaciens* strain.

MATERIALS AND METHODS

Bacterial strains and growth conditions. The strains of *E. coli* and *A. tumefaciens* and the plasmids used in this study are listed in Table 1. *E. coli* strains were grown at 37°C in L broth (LB), on L agar plates (Difco Laboratories, Detroit, Mich.), or in A medium (33). *Agrobacterium* strains were grown at 28°C in LB or on nutrient agar (NA) (Difco Laboratories). AB medium (11) supplemented with 0.2% mannitol as the sole carbon source was used as the defined minimal medium for *Agrobacterium* strains. Biotin was added to a final concentration of 2 µg/ml for biovar 2 strains. Agar (Difco Laboratories) was added at 1.5% to make solid medium. Except when specified in the text, antibiotics were used at the following concentrations: for *E. coli*, kanamycin at 50 µg/ml, tetracycline at 10 µg/ml, and ampicillin at 100 µg/ml; for *Agrobacterium* spp., kanamycin at 50 µg/ml, carbenicillin at 50 or 100 µg/ml, and tetracycline at 2 µg/ml. X-Gal (5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside; Sigma Chemical Co., St. Louis, Mo.) was included in the medium at 40 µg/ml to monitor the production of β-galactosidase.

DNA manipulations. Plasmids were isolated by an alkaline lysis method (40). Cloning followed standard recombinant DNA techniques (40). Restriction di-

* Corresponding author. Mailing address: Department of Crop Sciences, University of Illinois at Urbana-Champaign, 240 Edward R. Madigan Laboratory, 1201 West Gregory Dr., Urbana, IL 61801. Phone: (217) 333-1526. Fax: (217) 244-7830. E-mail: stephenf@uiuc.edu.

TABLE 1. Bacterial strains and plasmids used in this study

Strain or plasmid	Relevant characteristic(s) ^a	Source
<i>E. coli</i>		
DH5 α	F ⁻ ϕ 80dlacZ Δ M15 <i>endA1 recA1 hsdR17</i> (r _K ⁻ m _K ⁺) <i>supE44 thi-1 gyrA96</i> Δ (<i>lacZYA-argF</i>)U169	40
S17-1	Pro ⁻ Res ⁻ Mod ⁺ <i>recA</i> ; integrated RP4-Tet::Mu-Kan::Tn7, Mob ⁺	43
S17-1(pHoHo1, pSShe)	Amp ^r Cm ^r ; source for Tn3HoHo1	46
LE392	F ⁻ <i>hsdR514</i> (r _K ⁻ m _K ⁺) <i>supE44 supF58 lacY1 galK2 galT22 metB1 trpR55</i>	Promega
<i>A. tumefaciens</i>		
C58	Wild-type <i>A. tumefaciens</i> strain containing the nopaline-type Ti plasmid pTiC58	53
NT1	pTiC58-cured derivative of strain C58	51
NT1TcR1	Randomly picked tetracycline-resistant mutant of NT1	This study
Plasmids		
pBluescriptSK(+)	Amp ^r ; cloning vector	Stratagene
pCP13/B	Tc ^r ; IncP1 α ; broad-host-range cosmid vector	15
pRK415K	Tc ^r Km ^r ; IncP1 α ; broad-host-range cloning vector	14
pDSK519	Km ^r ; IncQ; broad-host-range cloning vector	25
pBAD22	Amp ^r ; expression vector controlled by P _{BAD}	19
pRG970b	Sp ^r /Str ^r Amp ^r ; promoter selection vector	12
pJB3	Amp ^r ; broad-host-range cloning vector	3
pZL1	1.7-kb <i>Bgl</i> II <i>cos</i> fragment from pCP13/B cloned into pJB3	This study
pBBRMCS-2	Km ^r ; broad-host-range cloning vector of unknown Inc group	27
pDLB4	Km ^r ; P _{BAD} from pBAD22 cloned into pBBRMCS-2	This study
pZLT1	Cosmid clone containing Tc ^r trait from NT1TcR1 in pZL1	This study
pZLE4.5	4.5-kb <i>Eco</i> RI fragment containing the Tc ^r trait in pZL1	This study
pDLE4.5	4.5-kb <i>Eco</i> RI fragment containing the Tc ^r trait in pDSK519	This study
pZLT2	Cosmid clone containing the wild-type <i>tet</i> _{C58} locus from NT1	This study
pZLHE1.6	1.6-kb <i>Hind</i> III/ <i>Eco</i> RI fragment containing IS426-disrupted <i>tet</i> R _{C58} from pZLE4.5 cloned into pBluescript SK(+)	This study
pZLHS2.9	2.9-kb <i>Hind</i> III/ <i>Sma</i> I fragment containing <i>tet</i> A _{C58} from pZLE4.5 cloned into pBluescript SK(+)	This study
pZLE4.8	4.8-kb <i>Eco</i> RI fragment containing portions of IS426 and <i>tet</i> R _{C58} from pZLT1 cloned into pBluescript SK(+)	This study
pZLE8.5	8.5-kb <i>Eco</i> RI fragment containing the wild-type <i>tet</i> _{C58} locus from pZLT2 cloned into pBluescript SK(+)	This study
pZLOP1	66-bp <i>tet</i> _{C58} intergenic region cloned into promoter selection vector pRG970b	This study
pBetR	<i>tet</i> R _{C58} cloned as a 2.7-kb <i>Nco</i> I fragment into pBAD22	This study
pDLB4-tetR	<i>tet</i> R _{C58} cloned as a 2.7-kb <i>Eco</i> RI/ <i>Bam</i> HI fragment from pBetR into pDLB4	This study

^a Amp^r, ampicillin resistant; Cm^r, chloramphenicol resistant; Km^r, kanamycin resistant; Sp^r, spectinomycin resistant; Tet^r or Tc^r, tetracycline resistant; Str^r, streptomycin resistant.

gestions were carried out in accordance with the instructions of the manufacturers, and digestion products were separated by electrophoresis in agarose gels with Tris-borate-EDTA buffer (40). DNA fragments were recovered from agarose gels by using GenElute spin columns (Supelco Inc., Bellefonte, Pa.). Plasmids were introduced into *E. coli* by CaCl₂-mediated transformation (40) and into *A. tumefaciens* by S17-1-mediated mating (43) or by electroporation (9).

Genomic bank construction. Cosmid pZL1 was constructed by cloning the *cos* site from pCP13 (15) as a 1.7-kb *Bgl*II fragment into pJB3 (6). Genomic DNA was digested with *Sau*3AI to give an average fragment size of 40 kb and ligated to pZL1 that had been digested with *Bam*HI and treated with alkaline phosphatase. The ligation mixture was packaged into phage λ by using the Packagene system (Promega Corporation, Madison, Wis.) and transduced into *E. coli* LE392.

Southern hybridization. Genomic DNA was prepared from overnight cultures of *Agrobacterium* spp. as described previously (18). After digestion with restriction endonuclease, DNA fragments were separated on 0.7% agarose gels and transferred to a nylon membrane. DNA probes were randomly labeled by using the Genius digoxigenin kit (Boehringer GmbH, Mannheim, Germany). Hybridization, washing, and detection followed protocols provided by the manufacturer. Hybridization and washing were performed under conditions of medium stringency.

Tn3HoHo1 mutagenesis. The insert in clone pDLE4.5 (Table 1) was mutagenized with Tn3HoHo1 as described previously (46). Sites and orientations of the insertions were determined by restriction mapping. Each mutant was tested for tetracycline resistance on LB medium containing this antibiotic.

DNA sequence analysis. DNA fragments containing the *tet* region were sequenced by use of the chain termination method (41) by the Genetic Engineering Facility at the University of Illinois. DNA sequences were analyzed by using DNA Strider (32). The BLAST (4) protocols were used to search the GenBank database for related sequences. The GAP subroutine of the GCG program (Genetics Computer Group, Madison, Wis.) was used to compare sequences for similarity.

Isolation of tetracycline-resistant mutants. *Agrobacterium* spp. were grown in LB liquid medium to saturation. A 200- μ l volume of culture from each strain was spread onto NA medium containing tetracycline at 5 or 10 μ g/ml. The capacity to give tetracycline-resistant mutants was assessed by the appearance of colonies on the plates after 72 h of incubation at 28°C.

Gene expression assays. *E. coli* and *A. tumefaciens* strains were grown in A and AB-mannitol media, respectively. Overnight cultures were washed and diluted

40-fold in the same respective media, and incubation was continued until the optical density at 600 nm (OD₆₀₀) reached about 0.7 for *E. coli* and 0.3 for *A. tumefaciens*. When needed, cultures were divided, and inducers (arabinose or tetracycline) were added to one subculture at the concentrations specified in Results. β -Galactosidase activity was determined quantitatively as described previously (33). β -Glucuronidase activity was measured as described previously (13, 24, 49). The activity of this enzyme was calculated by using the Miller formula (33), with OD₄₂₀ replaced by OD₄₁₅. All enzymatic activities were expressed as units per 10⁹ CFU.

Induction assays. The ability of tetracycline, its derivatives, and related compounds to induce the *tet* system of strain C58 (*tet*_{C58} system) was assessed two ways. In the first, reporter strains were inoculated onto medium containing X-Gal and the compound being tested. The appearance of blue colonies signalled the induction of the reporter gene. In the second, reporter strains were grown in 30 ml of liquid medium with appropriate antibiotics to saturation and mixed with 50 ml of warm 0.8% agar with X-Gal, and 6-ml volumes were layered over a base of LB or AB medium in 100-mm-diameter petri dishes. Discs of Whatman 3MM paper (0.5-cm diameter), onto which a few crystals of the compound to be tested were sprinkled, were laid on the surface of the soft agar. Dissolution and diffusion of the chemical give a concentration gradient around the disc. The plates were incubated at 28°C for *A. tumefaciens* reporter strains and at 37°C for *E. coli* reporter strains and examined at intervals over a 3-day period. Susceptibility to a given compound was indicated by a zone of growth inhibition around the disc. Induction of the reporter by the compound was indicated by a blue zone around the disc or, in cases where the compound also was inhibitory, by a blue zone in the area of growth at the edge of the zone of inhibition.

Nucleotide sequence accession number. DNA sequences of wild-type *tet*_{C58} were deposited in the GenBank database under accession no. AF090987.

RESULTS

Cloning a tetracycline resistance determinant from *A. tumefaciens* C58. Plating strain C58 onto medium containing tetracycline at concentrations ranging from 0.5 to 20 μ g/ml results




FIG. 1. A mutation in a tRNA processing gene in NTM7 (29) affects expression of resistance in spontaneous tetracycline-resistant mutants of NT1. Approximately 10^8 cells of strains NT1 (A) and NTM7 (B) were spread on NA plates supplemented with tetracycline at $5 \mu\text{g/ml}$. The plates were incubated for 3 (A) or 6 (B) days at 28°C . Note that the tetracycline-resistant mutants of NTM7 arising on the plate grow considerably more slowly than do those of strain NT1 on medium containing the antibiotic.

in the appearance at a high frequency of mutants resistant to this antibiotic (Fig. 1A and data not shown). However, we observed that a C58 mutant with a Tn5 insertion in a gene homologous to *md* (30) yielded tetracycline-resistant mutants that grew much more slowly on medium containing the antibiotic (Fig. 1B). Since a *miaA* mutant of *E. coli* does not express Tet(M)-mediated tetracycline resistance (8) and both *md* and *miaA* genes encode products involved in tRNA modification, we reasoned that strain C58 contains an active tetracycline resistance element that functions in the *md*⁺ but not in the *md* mutant background. To test this, a genomic library of NT1TcR1, a tetracycline-resistant mutant of the otherwise-wild-type strain NT1, was constructed in the broad-host-range cosmid vector pZL1 (Table 1). This bank, in *E. coli* LE392,

consists of 856 clones containing cosmids with an average insert size of 42 kb. The bank was screened on LB medium containing tetracycline, and one cosmid that confers resistance to this antibiotic on its *E. coli* host was isolated. This clone, named pZLT1, also conferred resistance to high levels of this antibiotic on several *Agrobacterium* strains, including 15955, R10, and K84 (data not shown).

The tetracycline resistance trait was subcloned from pZLT1 as a 4.5-kb *EcoRI* fragment into pZL1 to generate pZLE4.5 (Fig. 2). To localize the *tet* determinant, this DNA fragment was recloned into pDSK519 (25) to give pDLE4.5, and this plasmid was mutagenized with Tn3HoHo1 (46). The location and orientation of the insertion in each mutant plasmid were determined, and each was tested for its ability to confer resis-




FIG. 2. Genetic organization of the *tet* locus from the tetracycline-resistant and wild-type genomes of *A. tumefaciens* C58. A restriction map and the locations of the independent Tn3HoHo1 insertions, identified in clone pDLE4.5, are shown. The response of each mutant to tetracycline was assessed as described in Materials and Methods and is expressed as resistant to the antibiotic (+) or susceptible to the antibiotic (-). The horizontal crossbar capping each insertion site indicates the direction of transcription of the *lacZ* reporter of Tn3HoHo1 as determined by restriction mapping. β -Galactosidase activity was assessed on medium containing X-Gal: the presence of blue colonies (+) or white colonies (-) is indicated. The IS426 element is indicated by the hatched portion of the bar. S, *Sma*I; E, *Eco*RI; K, *Kpn*I; H, *Hind*III; Tc, tetracycline.

tance to tetracycline. Of six mutations mapped, the three contiguous insertions that are located in the middle of the fragment abolished resistance to this antibiotic (Fig. 2).

DNA sequence analysis. DNA fragments corresponding to the region defined by the Tn3HoHo1 insertion were subcloned and subjected to sequence analysis. BLAST searches revealed that one such *EcoRI-HindIII* fragment from plasmid pZLHE1.6 (Table 1) contained a portion of a *tetR(A)* homologue. As sequencing continued, however, this gene was abruptly disrupted by a totally different DNA sequence identical to the sequence of IS426, an IS element identified from *A. tumefaciens* T37 (50). Apparently, the 4.5-kb *EcoRI* fragment from the mutant contains an incomplete *tet* locus; insertion of IS426 introduced an *EcoRI* site which divided this region into two fragments when this enzyme was used for cloning. To obtain the complete sequence of the mutant version, the adjacent 4.9-kb *EcoRI* fragment was cloned into pBluescript SK(+) from pZLT1 to generate pZLE4.8 (Table 1). Analysis of the insert identified sequences corresponding to the remainder of IS426 followed by the 3' end of the *tetR* gene of strain C58 (*tetR*_{C58}).

Analysis of sequence data from pZLHS2.9 (Table 1), a plasmid harboring an insert from the opposite end of the region encoding the *tetR(A)* homologue, identified a single open reading frame of 1183 bp (Fig. 2 and data not shown). BLAST analysis indicated that this open reading frame encodes a protein very similar to the *tetA(A)* gene product from other *tet* systems.

To obtain the wild-type version of the *tet* locus from C58, a portion of the *tetA(A)* homologue was used as a probe to screen a genomic bank of the wild-type, tetracycline-sensitive strain NT1 (16). A cosmid hybridizing strongly to the probe was identified and designated pZLT2 (Fig. 2). The hybridizing region was subcloned as an 8.5-kb *EcoRI* fragment into pBluescript SK(+) to generate pZLE8.5 (Fig. 2), and the insert in this plasmid was used as the template to sequence a 2.6-kb region containing the wild-type *tet* locus.

Examination of the sequence indicated that this region contains two divergently transcribed open reading frames (Fig. 2 and data not shown). Database searches revealed that the two open reading frames can code for products that are very similar to the Tet(A) class of tetracycline resistance genes from *E. coli* transposons, with the highest similarity to that of the Tn1721 *tet* system (2). One we call *tetA*_{C58} could code for a protein with an M_r of 41,458 and having 46% identity (55% similarity) with TetA from the IncP1 α R plasmid RP4 (35) (Fig. 3A). The second, which could code for a protein with an M_r of 24,389 and having 46% identity (55% similarity) with TetR from RP4 (35), was named *tetR*_{C58} (Fig. 3B). The nucleotide sequences of *tetA*_{C58} and *tetR*_{C58} from the mutant and wild-type strains are identical. However, in the mutant, the sequence is interrupted by the IS426 insertion at bp 520 of *tetR*_{C58} (data not shown). This insertion generated a 5-bp duplication of the sequence 5' GGGGG 3' at the target site.

A hydropathy plot generated by the Kyte-Doolittle algorithm (28) revealed that TetA_{C58} contains 12 putative hydrophobic segments (data not shown). Such domains, which could represent transmembrane segments, are a typical feature of the major facilitator superfamily of proteins (31, 36). Sequence analysis indicated that, like other class A and class B TetR proteins, TetR_{C58} contains a typical helix-turn-helix domain in its N-terminal portion (data not shown). The *tetA*_{C58} and *tetR*_{C58} genes are separated by an 88-bp intergenic region with characteristics of regulated divergent promoters (Fig. 4A). A 15-bp inverted repeat (IR) element overlaps the putative -10 region of the *tetA*_{C58} promoter. This IR also overlaps the po-

tential -35 site of the *tetR*_{C58} promoter. As shown in Fig. 4B, the sequence of this IR is identical to those of the operators found between the *tetA* and *tetR* genes of other class A and class B *tet* systems, including those of RP4 and Tn1721 (2, 35). However, the promoter region of *tet*_{C58} lacks the second IR element present in the corresponding region of *tet*_{RP4} and of *tet*_{Tn1721}. Furthermore, the intergenic region of *tet*_{C58} is considerably shorter than those of the *tet* elements of RP4 and Tn1721 (Fig. 4B).

Expression of *tet*_{C58} is not induced by tetracycline or related compounds. In class A and class B *tet* systems, TetR represses expression of *tetA* and *tetR*, and this repression is relieved by subinhibitory levels of the antibiotic (3, 21, 23). Tetracycline does not induce the *Agrobacterium tet* system; as noted above, when a large number cells are spread onto medium containing this antibiotic, only a very small fraction of the cells grow to produce colonies. Moreover, of those such colonies tested, all now are constitutively resistant to tetracycline (data not shown). To study the regulation of the *tetA* and *tetR* genes of *Agrobacterium* spp., a 66-bp fragment of the intergenic region, which comprises sequence between the putative ribosome binding sites for the *tetA*_{C58} and *tetR*_{C58} genes, was cloned into the divergent promoter detection vector pRG970b (13) to generate pZLOP1. In this construct, *tetA*_{C58} and *tetR*_{C58} are transcriptionally fused to the *lacZ* and *uidA* genes, respectively, of the vector. In addition, the *tetR*_{C58} gene was cloned as a *NcoI* fragment from pZLT2 into the expression vector pBAD22 (19) to generate pBetR. This construct allows us to regulate the expression of *tetR*_{C58} with arabinose. To express *tetR*_{C58} in *Agrobacterium* spp., this gene was excised from pBetR as an *EcoRI-BamHI* fragment and cloned into pDBL4 (Table 1), a broad-host-range expression vector containing the arabinose promoter and downstream transcriptional terminator from pBAD22 cloned into pBBRMCS-2 (27), to construct pDBL4-*tetR*. Expression of *tetR*_{C58} in these two constructs is inducible in *A. tumefaciens* and *E. coli* by addition of arabinose to the culture medium at a final concentration of 0.4%.

Under conditions in which *tetR*_{C58} is repressed, expression levels of the *tetA*_{C58}::*lacZ* and *tetR*_{C58}::*uidA* fusions were lower in strain NT1(pZLOP1, pDLB4-*tetR*) than in strain NT1TcR1 (pZLOP1, pDLB4-*tetR*). We attribute the lower expression levels of these fusions in NT1 to the chromosomal copy of *tetR*_{C58}. However, when the cloned copy of *tetR*_{C58} was induced by adding arabinose, expression of the reporter fusions in both strains was inhibited to similar levels (Table 2). Similarly, expression of the *tetA*_{C58}::*lacZ* reporter on pZLOP1 was repressed by TetR_{C58} in an *E. coli* host but only upon addition of arabinose (Table 2).

Tetracycline, when added at final concentrations ranging in 10-fold increments from 0.0001 to 1 μ g/ml, had no effect on the expression of *tetA*_{C58} or *tetR*_{C58} in either host (data not shown). Although tetracycline does not induce the expression of *tetA*_{C58}, we consider the possibility that compounds similar to this antibiotic could be the ligands for TetR_{C58}. To test this, a total of 25 polycyclic compounds with structures similar to tetracycline were tested for their ability to induce the expression of the *tetA*_{C58}::*lacZ* fusion. These included active derivatives of tetracycline as well as various nontoxic compounds, most of plant origin. None of these agents induced the expression of the *tetA*_{C58}::*lacZ* fusion at a detectable level (data not shown). When the tetracycline-sensitive strains NT1(pZLOP1) and NT1(pZLOP1, pDLB4-TetR) were used as reporters, resistant mutants formed in the zones of growth inhibition caused by chlortetracycline and oxytetracycline. However, no such mutants appeared in the zone of growth inhibition caused by minocycline (data not shown).




FIG. 3. Relatedness of TetA and TetR from strain C58 to the gene products of *tetA* and *tetR* from RP4. Alignments between the predicted amino acid sequence of TetA_{C58} and TetA_{RP4} (A) and of TetR_{C58} and TetR_{RP4} (B) were performed by using the GAP algorithm from the GCG package as described in Materials and Methods. Identical amino acids are shown as white letters on a black background, and conserved amino acids are shown as black letters on a shaded background.

Cross-operator recognition by TetR_{C58} and TetR_{RP4}. By sequence analysis, the putative 15-bp operator of *tet*_{C58} is identical to those of other *tet* systems (Fig. 4B). Furthermore, the sequence from residues 26 to 47 of TetR_{C58} is virtually identical to that of the N-terminal helix-turn-helix domain of several related TetR proteins, including TetR_{RP4}, TetR_{Tn10}, and TetR_{Tn1721} (Fig. 3B and data not shown). Given these similarities, we determined whether TetR from *A. tumefaciens* and from RP4 will recognize the operators of the noncognate homologues. Strains DH5 α (pRK415K, pBetR) and DH5 α (pRK415K, pZLOP1) were constructed for this study. Plasmid pRK415K carries the RP4 tetracycline resistance gene unit, and expression of this determinant is inducible by tetracycline (14, 25). If TetR_{C58} can repress the expression of *tetA*_{RP4}, upon induction of *tet*_{C58}, strain DH5 α (pRK415K, pBetR) should fail to grow on medium containing tetracycline. As shown in Table 3, this strain grows well on LB medium with tetracycline at a concentration of 10 μ g/ml. However, when the expression of *tetR*_{C58} on pBetR was induced by arabinose, this strain was unable to grow on the same medium. These results indicate that TetR_{C58} can repress at the *tetA*_{RP4} promoter and that

this repression is not responsive to tetracycline. To determine if TetR_{RP4} can repress *tetA*_{C58}, the expression of the *tetA*_{C58}::*lacZ* reporter was examined in strain DH5 α (pRK415K, pZLOP1). Results from this test indicated that TetR_{RP4} from pRK415K indeed repressed *tetA*_{C58} expression, and like the repressive effect on its cognate *tetA*, the repression was responsive to tetracycline (Table 3). Thus, TetR_{RP4} can recognize and repress expression from the promoter of the *A. tumefaciens tet* system. Furthermore, unlike the *tet*_{C58} system, this repression of *tetA*_{C58} by TetR_{RP4} is relieved by tetracycline.

Nature of the tetracycline-resistant mutants. Our data indicate that a *tet* unit in the genome of strain C58 accounts for the appearance of tetracycline-resistant mutants when this strain is grown in the presence of this antibiotic. In the particular mutant used in this study, the tetracycline resistance phenotype arose from the transposition of IS426 into *tetR*_{C58}. Since tetracycline does not induce the expression of *tetA*_{C58}, we were interested in determining whether all tetracycline-resistant mutants were caused by the same type of mutation. To examine this, genomic DNA from eight independent tetracycline-




FIG. 4. Organization of the intergenic regulatory regions from several related *tet* systems. (A) Locations of putative promoter and operator sequences. (B) Comparison of the DNA sequences of operators from the *tet* systems of C58, RP4, and Tn1721. Inverted repeat sequences are boxed. The transcriptional directions of the *tetA* and *tetR* genes are indicated by arrows. A-10 and A-35, the -10 and -35 sites for the *tetA* promoters; R-10 and R-35, the -10 and -35 sites for the *tetR* promoters; O_{C58}, putative operator of *tet*_{C58}; O_L and O_R, the left and right operators, respectively, of the *tet* systems of Tn1721 and RP4.

resistant mutants was probed with a DNA fragment containing the *tet*_{C58} genes. These eight mutants grouped into four types based on the size of the hybridizing fragment (data not shown). One class, represented by three mutants, contained a fragment indistinguishable in size from that of the wild type. In the second, two of the eight mutants gave a fragment indistinguishable in size from that of NT1TcR1, the original mutant from which the *tet* locus was cloned. The remaining three mutants contained two restriction fragment length polymorphism (RFLP) patterns different from that of the wild type and of NT1TcR1 (data not shown).

Surveying for the presence of the *tet* locus in the genomes of *Agrobacterium* strains. We examined a collection of wild-type isolates of *Agrobacterium* spp. for the *tet* locus by Southern analysis using a probe containing part of *tetA* and most of *tetR*. We also examined some commonly used derivatives of strain C58. Of those not derived from C58, only strain T37 contains DNA with detectable relatedness to the *tet* locus from strain C58 (Table 4). Moreover, the probe hybridized to an *Eco*RI fragment of the same size in the two strains (data not shown). Consistent with these results, of the 26 strains tested, only T37 and those derived from C58 gave rise to tetracycline-resistant mutants (Table 4). The probe hybridized with all derivatives of strain C58 tested, including strain UIA5, which lacks both pTiC58 and the 450-kb catabolic plasmid pAtC58 (Table 4). Consistent with this, UIA5 gives rise to mutants resistant to tetracycline.

DISCUSSION

We report here the characterization of a *tet* system from *A. tumefaciens* C58. This active resistance system accounts for the long-known observation that C58 and its derivatives give rise to tetracycline-resistant mutants at a high frequency.

This resistance determinant is structurally similar to that of *tet* systems found in plasmids and transposons from *E. coli* and other gram-negative bacteria, with the highest relatedness being to the class A tetracycline resistance gene system from Tn1721. The *tet*_{C58} system confers resistance to tetracycline and derivatives such as chlortetracycline and oxytetracycline but not to minocycline. This pattern of resistance suggests that *tet*_{C58} does not belong to the Tet(S) class of tetracycline resistance genes, which confers resistance to minocycline as well as to tetracycline, chlortetracycline, and oxytetracycline (10).

Like other class A systems, *tet*_{C58} is organized as two genes, a regulatory gene and a gene coding for an efflux pump, transcribed divergently from an intergenic region. Potential promoter elements for both genes are overlapped by a single putative operator composed of a 15-bp perfect IR element. Our genetic analysis indicates that this intergenic region contains all of the *cis*-acting information required for the regulated expression of these two genes. The DNA sequence of the putative operator is identical to that of operators from other *tet* systems. However, the promoter region of *tet*_{C58} contains only one repeat while the promoter regions of the *tet* determinants

TABLE 2. TetR_{C58} controls expression of *tetA*_{C58} and *tetR*_{C58}

Strain	<i>tetR</i> _{C58} source	β-Galactosidase activity ^a (<i>tetA</i> _{C58} ::lacZ)		β-Glucuronidase activity ^a (<i>tetR</i> _{C58} ::uidA)	
		No ara	With ara	No ara	With ara
NT1TcR1(pZLOP1, pDLB4)	None	77	85	53	45
NT1TcR1(pZLOP1, pDLB4-tetR)	pDLB4-tetR	65	2	18	6
NT1(pZLOP1, pDLB4)	Chromosome	19	14	14	20
NT1(pZLOP1, pDLB4-tetR)	Chromosome/pDLB4-tetR	25	2	14	7
DH5α(pZLOP1, pDLB4)	None	2,213	2,243	NT ^b	NT
DH5α(pZLOP1, pDLB4-tetR)	pDLB4-tetR	1,678	29	NT	NT

^a β-Galactosidase and β-glucuronidase activities are expressed as units per 10⁹ CFU as described in Materials and Methods. Activities were determined in cultures with or without 0.4% arabinose (ara) added as an inducer.

^b NT, not tested (*E. coli* DH5α has an endogenous β-glucuronidase activity).

TABLE 3. TetR_{C58} and TetR_{RP4} recognize noncognate *tet* operators

Strain	β-Galactosidase activity ^a		Growth on TC medium ^b	
	No TC	With TC	No ara	With ara
DH5α(pRK415K, pBetR)	NT ^c	NT	++	–
DH5α(pRK415K, pBAD22)	NT	NT	++	++
DH5α(pZLOP1)	2,311	2,211	NT	NT
DH5α(pZLOP1, pRK415K)	387	2,154	NT	NT

^a β-Galactosidase activity is expressed as units per 10⁹ CFU as described in Materials and Methods. When needed, tetracycline (TC) was added to the culture at 10 μg/ml as an inducer.

^b Growth was assessed on LB medium supplemented with tetracycline at 5 and 10 μg/ml (TC medium) and with or without 0.4% arabinose (ara) added as an inducer. ++, good growth; –, no growth.

^c NT, not tested.

of RP4 and Tn1721 each contain two copies of the operators (Fig. 4B).

Heterologous repressor-operator recognition occurs between different classes of *tet* systems, even those with dissimilar operators (26). The high degree of sequence similarity between operators from the *tet*_{C58} and *tet*_{RP4} systems suggested that TetR proteins from these two systems could recognize the noncognate operator. This proved to be the case: TetR of RP4 repressed the *tet* system of C58 and TetR of C58 repressed tetracycline resistance conferred by the RP4 *tet* system. The capacity of TetR_{RP4} to repress expression of the *tetA*_{C58}::*lacZ* fusion suggests that the 15-bp IR is the operator and indicates that a single copy of this element is sufficient for regulation of both genes.

The expression of almost all tetracycline resistance gene systems examined to date is under strict control (29, 45). Among these, repression of the Tet(B) class by TetR is the best understood (21, 45). In the absence of tetracycline, the repressors bind to operators overlapping the promoters of both the *tetA* and *tetR* genes (21). Binding of tetracycline to TetR(B) induces a conformational change that leads to release from the operators and concomitant expression of *tetA* (22, 34). Tetracycline-inducible TetR-mediated regulation of other *tet* systems is believed to function by a similar mechanism (21). Two lines of evidence indicate that tetracycline does not induce the *Agrobacterium tet* system. First, when cells were spread onto medium containing tetracycline at concentrations ranging from 10 to 1 μg/ml, only a small portion of the cells grew into colonies (data not shown). Moreover, those that did grow now expressed constitutive resistance to tetracycline. If tetracycline functioned as the inducer for *tet*_{C58}, virtually all of the cells should be resistant to this antibiotic. Furthermore, *E. coli* and *Agrobacterium* strains harboring pZLE8.5 (Table 1), which carries the wild-type *tet*_{C58} locus, are sensitive to tetracycline, even at concentrations to which *tet*_{RP4} confers resistance (data not shown). Second, while the *tetA*_{C58}::*lacZ* and the *tetR*_{C58}::*uidA* reporter fusions were expressed constitutively in cells lacking TetR_{C58}, the expression of both fusions was depressed considerably in cells harboring *tetR*_{C58}. Furthermore, neither reporter was inducible by tetracycline. Repression of *tetA*_{RP4} by TetR_{C58} is not relieved by tetracycline. On the other hand, repression of *tet*_{C58} by TetR_{RP4} does respond to tetracycline. These two sets of results indicate that the failure of the *tet* system of C58 to respond to tetracycline is a function of TetR and not of the *tet* operator. Mutants of the related repressor from Tn10 that do not respond to tetracycline have been isolated and characterized. These noninducible mutants repress the expression of

*tetA*_{Tn10}::*lacZ* fusions by binding to the operators even in the presence of tetracycline (5, 20, 34). We propose that TetR_{C58} represents such a noninducible form of TetR. Clearly, this repressor can bind DNA. Why it no longer reacts to tetracycline remains to be determined. We considered the possibility that tetracycline is not the true ligand for TetR_{C58}. However, none of the 25 compounds we tested induced the system. Certainly, this was not an exhaustive survey. But on balance, that the locus of strain C58 can be regulated by TetR_{RP4} in response to tetracycline strongly suggests that this gene unit once was responsive to this antibiotic.

The interchangeability with respect to operator recognition exhibited by the two TetR proteins could explain another observation concerning tetracycline resistance in strain C58. Derivatives of this *A. tumefaciens* strain harboring RP4 and vectors carrying *tet*_{RP4} express only low level resistance to tetracycline. We propose that TetR_{C58} is partially dominant over TetR_{RP4}. Thus, since TetR_{C58} does not respond to tetracycline, the *tet*_{RP4} determinant is not fully expressed in its *A. tumefaciens* host.

Expression of tetracycline resistance in C58 apparently occurs only following a mutation affecting regulation. These mutations can be of several types; in the particular mutant used in this study, the tetracycline resistance phenotype resulted from the insertion of IS426 in *tetR*_{C58}. In genetic analysis carried out with C58 and its derivatives, this element is commonly associated with phenotypes in which the selection gives derepressed mutants (17). However, diverse RFLP patterns in Southern

TABLE 4. Tetracycline resistance characteristics of *Agrobacterium* spp.

<i>Agrobacterium</i> species	Strain ^c	Appearance of TC-resistant mutants on medium with TC at ^a :		Homology with <i>tet</i> _{C58} ^b
		5 μg/ml	10 μg/ml	
<i>A. tumefaciens</i>	C58 (NT1, UIA5, A281)	++	++	++
	T37	++	++	++
	K108	–	–	–
	Ach5 (LBA4404)	–	–	–
	A6NC	–	–	–
	R10	–	–	–
	15955	–	–	–
	B6	–	–	–
	Bo542	–	–	–
	Chry5	–	–	–
	ANT4	–	–	–
<i>A. rhizogenes</i>	2655	–	–	–
	2657	–	–	–
	K599	–	–	–
	8196	–	–	–
	A4	–	–	–
<i>A. vitis</i>	15834	–	–	–
	K308	–	–	–
	Hm1	–	–	–
	Ag57	–	–	–
<i>A. radiobacter</i>	Tm4	–	–	–
	K84	–	–	–

^a Tetracycline (TC)-resistant mutants either appeared (++) or did not appear (–).

^b Determined by Southern analysis as described in Materials and Methods. Homology was (++) or was not (–) found.

^c Strains shown in parentheses are derivatives of the type strain that also were tested.

analysis of eight independent mutants indicated that other types of mutations can result in the tetracycline resistance phenotype. While mutants with a pattern indistinguishable from that of NT1TcR1 may have arisen from IS426 transposition, those with a pattern similar to that of the wild type could represent point mutations or very small deletions or insertions in the *tetR*_{C58} gene. They also could represent operator-constitutive mutations that no longer allow TetR_{C58} to bind at the promoter.

Among the agrobacteria we examined, only strain T37 contains DNA with detectable homology to the *tet* locus from C58. Furthermore, only C58 and T37 give rise to tetracycline-resistant mutants (Table 4). These two strains were isolated from different parts of the United States (42). Moreover, although strain C58 induces unorganized tumors, strain T37 causes teratomas. Both strains contain a nopaline/agrocinopine-type Ti plasmid, but RFLP analysis indicates that these two elements have diverged considerably from each other (42). Our survey also shows that generation of tetracycline-resistant mutants is associated with the presence of a *tet*_{C58} homologue in the genome of the particular isolate; those lacking this element do not yield mutants at a detectable level. The presence of the *tet* locus in strain UIA5, a derivative of C58 lacking both pTiC58 and pAtC58, indicates that this resistance determinant is associated with one of the two chromosomes of strain C58 (1). Interestingly, this is not the only antibiotic resistance determinant in strain C58. This strain, which also mutates to resistance to chloramphenicol at a high frequency, contains a gene coding for an atypical chloramphenicol acetyltransferase (47). Whether this *cat* gene is present in strain T37 or any other isolate of *Agrobacterium* spp. is not known.

ACKNOWLEDGMENTS

We thank Abigail Salyers for supplying the tetracyclines and related chemicals and Stanton Gelvin and Ramon Peñalver for helpful discussions. We also thank Malcolm Winkler, whose seminar on tRNA processing got us started on the problem.

Portions of this research were supported by grants R01 GM52465 from the NIH and NC CNTRL SOY SKF ANTC from the North Central Soybean Association to S.K.F.

REFERENCES

- Allardet-Servent, A., S. Michaux-Charachon, E. Jumas-Bilak, L. Karayan, and M. Ramuz. 1993. Presence of one linear and one circular chromosome in the *Agrobacterium tumefaciens* C58 genome. *J. Bacteriol.* **175**:7869–7874.
- Allmeier, H., B. Cresnar, M. Greck, and R. Schmitt. 1992. Complete nucleotide sequence of Tn721: gene organization and a novel gene product with features of a chemotaxis protein. *Gene* **111**:11–20.
- Altenbuchner, J., K. Schmid, and R. Schmitt. 1983. Tn721-encoded tetracycline resistance: mapping of structural and regulatory genes mediating resistance. *J. Bacteriol.* **153**:116–123.
- Altschul, S. F., W. Gish, W. Miller, E. W. Myers, and D. J. Lipman. 1990. Basic local alignment search tool. *J. Mol. Biol.* **215**:403–410.
- Biburger, M., C. Berens, T. Lederer, T. Crec, and W. Hillen. 1998. Intragenic suppressors of induction-deficient TetR mutants: localization and potential mechanism of action. *J. Bacteriol.* **180**:737–741.
- Blanty, J. M., T. Brautaset, H. C. Winther-Larsen, K. Haugan, and S. Valla. 1997. Construction and use of a versatile set of broad-host-range cloning and expression vectors based on the RK2 replicon. *Appl. Environ. Microbiol.* **63**:370–379.
- Burdett, V. 1991. Purification and characterization of Tet(M), a protein that renders ribosomes resistant to tetracycline. *J. Biol. Chem.* **266**:2872–2877.
- Burdett, V. 1993. tRNA modification activity is necessary for Tet(M)-mediated tetracycline resistance. *J. Bacteriol.* **175**:7209–7215.
- Cangelosi, G. A., E. A. Best, G. Marinetti, and E. W. Nester. 1991. Genetic analysis of *Agrobacterium*. *Methods Enzymol.* **204**:384–397.
- Charpentier, E., G. Gerbaud, and P. Courvalin. 1993. Characterization of a new class of tetracycline-resistance gene tet(S) in *Listeria monocytogenes* BM4210. *Gene* **131**:27–34.
- Chilton, M. D., T. C. Currier, S. K. Farrand, A. J. Bendich, M. P. Gordon, and E. W. Nester. 1974. *Agrobacterium tumefaciens* DNA and PS8 bacteriophage DNA not detected in crown gall tumors. *Proc. Natl. Acad. Sci. USA* **71**:3672–3676.
- Connolly, D. M., and M. E. Winkler. 1989. Structure of *Escherichia coli* K-12 *miaA* and characterization of the mutator phenotype caused by *miaA* insertion mutations. *J. Bacteriol.* **171**:3233–3246.
- Cook, D. M. 1995. Ph.D. dissertation. University of Illinois at Urbana-Champaign, Urbana.
- Cook, D. M., and S. K. Farrand. 1992. The *oriT* region of the *Agrobacterium tumefaciens* Ti plasmid pTiC58 shares DNA sequence identity with the transfer origins of RSF1010 and RK2/RP4 and with T-region borders. *J. Bacteriol.* **174**:6238–6246.
- Dazines, A., and A. M. Chakraborty. 1984. Cloning of genes controlling alginate biosynthesis from a mucoid cystic fibrosis isolate of *Pseudomonas aeruginosa*. *J. Bacteriol.* **159**:9–18.
- Farrand, S. K., S. P. O'Morchoe, and J. McCutchan. 1989. Construction of an *Agrobacterium tumefaciens* C58 *recA* mutant. *J. Bacteriol.* **171**:5314–5321.
- Farrand, S. K., J. Tempé, and Y. Dessaux. 1990. Localization and characterization of the region encoding catabolism of mannopinic acid from the octopine-type Ti plasmid pTi15955. *Mol. Plant Microbe Interact.* **3**:259–267.
- Glickmann, E., L. Gardan, S. Jacquet, S. Hussain, M. Elasri, A. Petit, and Y. Dessaux. 1998. Auxin production is a common feature of most pathogens of *Pseudomonas syringae*. *Mol. Plant Microbe Interact.* **11**:156–162.
- Guzman, L. M., D. Belin, M. J. Carson, and J. Beckwith. 1995. Tight regulation, modulation, and high-level expression by vectors containing the arabinose PBAD promoter. *J. Bacteriol.* **177**:4121–4130.
- Hecht, B., G. Muller, and W. Hillen. 1993. Noninducible Tet repressor mutations map from the operator binding motif to the C terminus. *J. Bacteriol.* **175**:1206–1210.
- Hillen, W., and C. Berens. 1994. Mechanisms underlying expression of the Tn70 encoded tetracycline resistance. *Annu. Rev. Microbiol.* **48**:345–369.
- Hinrichs, W., C. Kisker, M. Duvel, A. Muller, K. Tovar, W. Hillen, and W. Saenger. 1994. Structure of the Tet repressor-tetracycline complex and regulation of antibiotic resistance. *Science* **264**:418–420.
- Izaki, K., K. Kiuchi, and K. Arima. 1966. Specificity and mechanism of tetracycline resistance in a multiple-drug-resistant strain of *Escherichia coli*. *J. Bacteriol.* **91**:628–633.
- Jefferson, R. A. 1987. Assaying chimeric genes in plants: the GUS gene fusion system. *Plant Mol. Biol. Rep.* **5**:387–405.
- Keen, N. T., S. Tamaki, D. Kobayashi, and D. Trollinger. 1988. Improved broad-host-range plasmids for DNA cloning in gram-negative bacteria. *Gene* **70**:191–197.
- Klock, G., B. Unger, C. Gatz, W. Hillen, J. Altenbuchner, K. Schmid, and R. Schmitt. 1985. Heterologous repressor-operator recognition among four classes of tetracycline resistance determinants. *J. Bacteriol.* **161**:326–332.
- Kovach, M. E., P. H. Elzer, D. S. Hill, G. T. Robertson, M. A. Farris, R. M. I. Roop, and K. M. Peterson. 1995. Four new derivatives of the broad-host-range cloning vector pBBR1MCS, carrying different antibiotic-resistance cassettes. *Gene* **166**:175–176.
- Kyte, J., and R. F. Doolittle. 1992. A simple method for displaying the hydrophobic character of a protein. *J. Mol. Biol.* **157**:105–132.
- Levy, S. B. 1988. Tetracycline resistance determinants are widespread. *ASM News* **54**:418–421.
- Luo, Z.-Q., and S. K. Farrand. Unpublished data.
- Marger, M. D., and M. H. Saier, Jr. 1993. A major superfamily of transmembrane facilitators that catalyze uniport, symport and antiport. *Trends Biochem. Sci.* **18**:13–20.
- Mark, C. 1988. "DNA Strider": a "C" program for fast analysis of DNA and protein sequences on the Apple Macintosh family of computers. *Nucleic Acids Res.* **16**:1829–1836.
- Miller, J. 1972. Experiments in molecular genetics. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
- Muller, G., B. Hecht, V. Helbl, W. Saenger, and W. Hillen. 1995. Characterization of non-inducible Tet repressor mutants suggests conformational changes necessary for induction. *Nat. Struct. Biol.* **2**:693–703.
- Pansegrau, W., E. Lanka, P. T. Barth, D. H. Figurski, D. G. Guiney, D. Haas, D. R. Helinski, H. Schwab, V. A. Stanisich, and C. M. Thomas. 1994. Complete nucleotide sequence of Birmingham IncP alpha plasmids, compilation and comparative analysis. *J. Mol. Biol.* **239**:623–663.
- Pao, S. S., I. Paulsen, and J. M. H. Saier. 1998. Major facilitator superfamily. *Microbiol. Mol. Biol. Rev.* **6**:1–34.
- Paulsen, I. T., M. H. Brown, and R. A. Skurray. 1996. Proton-dependent multidrug efflux systems. *Microbiol. Rev.* **60**:575–608.
- Robert, M. C. 1994. Epidemiology of tetracycline resistance determinants. *Trends Microbiol.* **2**:353–357.
- Salyers, A. A., B. S. Speer, and N. B. Shoemaker. 1990. New perspectives on tetracycline resistance. *Mol. Microbiol.* **4**:151–156.
- Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
- Sanger, F., S. Nicklen, and A. R. Coulson. 1977. DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* **74**:5463–5467.
- Sciaky, D., A. L. Montoya, and M.-D. Chilton. 1977. Fingerprints of *Agrobacterium* Ti plasmids. *Plasmid* **1**:238–253.
- Simon, R., U. Priefer, and A. Pühler. 1983. A broad host range mobilization

- system for *in vivo* genetic engineering: transposon mutagenesis in gram negative bacteria. *Bio/Technology* **1**:37-45.
44. **Speer, B. S., and A. A. Salyers.** 1989. Novel aerobic tetracycline resistance gene that chemically modifies tetracycline. *J. Bacteriol.* **171**:148-153.
 45. **Speer, B. S., N. B. Shoemaker, and A. A. Salyers.** 1992. Bacterial resistance to tetracycline: mechanisms, transfer, and clinical significance. *Clin. Microbiol. Rev.* **5**:387-399.
 46. **Stachel, S. E., G. An, C. Flores, and E. W. Nester.** 1985. A *Tn3lacZ* transposon for the random generation of β -galactosidase gene fusions: application to the analysis of gene expression in *Agrobacterium*. *EMBO J.* **4**:891-898.
 47. **Tennigkeit, J., and H. Matzura.** 1991. Nucleotide sequence analysis of a chloramphenicol-resistance determinant from *Agrobacterium tumefaciens* and identification of its gene product. *Gene* **98**:113-116.
 48. **Travis, J.** 1994. Reviving the antibiotic miracle. *Science* **264**:360-362.
 49. **Van den Eede, G., R. Deblaere, K. Goethals, M. Van Montagu, and M. Holsters.** 1992. Broad host range and promoter selection vectors for bacteria that interact with plants. *Mol. Plant Microbe Interact.* **5**:228-234.
 50. **Vanderleyden, J., J. Desair, C. De Meirman, K. Michiels, A. P. Van Gool, M.-D. Chilton, and G. C. Jen.** 1986. Nucleotide sequence of an insertion sequence (IS) element identified in the T-DNA region of a spontaneous variant of the Ti-plasmid pTiT37. *Nucleic Acids Res.* **14**:6699-6709.
 51. **Waston, B., T. C. Currier, M. P. Gordon, M.-D. Chilton, and E. W. Nester.** 1975. Plasmid required for virulence of *Agrobacterium tumefaciens*. *J. Bacteriol.* **123**:255-264.
 52. **Zhang, J., and M. P. Deutscher.** 1988. Cloning, characterization, and over-expression of the *Escherichia coli rna* gene encoding RNase D. *J. Bacteriol.* **170**:522-527.
 53. **Zimmerer, R. P., R. H. Hamilton, and C. Pootjes.** 1966. Isolation and morphology of temperate *Agrobacterium tumefaciens* bacteriophage. *J. Bacteriol.* **92**:746-750.