


INVITED REVIEW

INVITED REVIEW THEMED ISSUE

Immunological memory to SARS-CoV-2 infection and COVID-19 vaccines

Alessandro Sette^{1,2} | Shane Crotty^{1,2} 

¹Center for Infectious Disease and Vaccine Research, La Jolla Institute for Immunology (LJI), La Jolla, California, USA

²Division of Infectious Diseases and Global Public Health, Department of Medicine, University of California, San Diego (UCSD), La Jolla, California, USA

Correspondence

Shane Crotty, Center for Infectious Disease and Vaccine Research, La Jolla Institute for Immunology (LJI), La Jolla, CA 92037, USA.
Email: shane@lji.org

Funding information

National Institute of Allergy and Infectious Diseases, Grant/Award Number: AI142742 and AI144462

Abstract

Immunological memory is the basis of protective immunity provided by vaccines and previous infections. Immunological memory can develop from multiple branches of the adaptive immune system, including CD4 T cells, CD8 T cells, B cells, and long-lasting antibody responses. Extraordinary progress has been made in understanding memory to SARS-CoV-2 infection and COVID-19 vaccines, addressing development; quantitative and qualitative features of different cellular and anatomical compartments; and durability of each cellular component and antibodies. Given the sophistication of the measurements; the size of the human studies; the use of longitudinal samples and cross-sectional studies; and head-to-head comparisons between infection and vaccines or between multiple vaccines, the understanding of immune memory for 1 year to SARS-CoV-2 infection and vaccines already supersedes that of any other acute infectious disease. This knowledge may help inform public policies regarding COVID-19 and COVID-19 vaccines, as well as the scientific development of future vaccines against SARS-CoV-2 and other diseases.

KEYWORDS

adenoviral vectors, coronavirus, hybrid immunity, memory B cells, memory T cells, mRNA vaccines, natural immunity, protein vaccine

1 | INTRODUCTION

Protective immunity provided by vaccines is predicated on the existence of immunological memory: the capacity of the adaptive immune system to not only recognize a novel pathogen but to also remember it. Only in the past few decades have the cellular and molecular sources of immunological memory been defined, and much remains to be determined. The three main branches of the adaptive immune system are B cells (the source of antibodies, “Abs”), CD4 T cells, and CD8 T cells. Immune memory is encoded in four main compartments of adaptive immunity: memory CD8 T cells, memory CD4 T cells, memory B cells (B_{Mem}), and circulating Abs¹ (Figure 1). There

is evidence of roles for B cells (including Abs), CD4 T cells, and CD8 T cells in protective immunity to SARS-CoV-2, and thus, it is important to study immune memory to SARS-CoV-2 and COVID-19 vaccines to understand protective immunity against COVID-19. Because of the size and scope of immunological studies of SARS-CoV-2 in humans, the large number of first-time infections, the large number of first-time vaccinations, and the diversity of COVID-19 vaccines developed in a short period of time, there are now more data on human antigen-specific immune responses to SARS-CoV-2 than any other acute pathogen. As a result, immune memory to SARS-CoV-2 is now a benchmark in human immunology for understanding antigen-specific T cell and B cell memory.

*This article introduces a series of reviews covering SARS-CoV-2 Vaccines appearing in Volume 310 of *Immunological Reviews*.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial](https://creativecommons.org/licenses/by-nc/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2022 The Authors. *Immunological Reviews* published by John Wiley & Sons Ltd.

Immune memory to SARS-CoV-2 can be generated by infection (classically referred to as “natural immunity”), vaccination, or hybrid immunity. Hybrid immunity is the combination of infection-induced immunity and vaccine-induced immunity.² Each of these causes of immune memory is discussed in each section of this review. Overall, immune memory from prior infection, vaccination, or hybrid immunity each have distinctive characteristics. Previous infection can generate robust immune memory,^{3,4} including memory CD8 T cell, CD4 T cell, B_{Mem}, durable Abs, and local immune memory (Figure 2). Epidemiological data on protective immunity in previously infected individuals are consistent with the immune memory measurements. Multiple large studies observe that prior infection provides approximately 80%–95% protection against symptomatic COVID-19 re-infections for 8+ months, for SARS-CoV-2 ancestral strain and the Alpha through Delta VOCs,^{5–11} and significant protection against disease with Omicron.^{12–14}

COVID-19 vaccines, focusing for the moment on 2-dose mRNA vaccines (Moderna mRNA-1273, Pfizer/BioNTech BNT162b2), clearly provide high levels of protective immunity against SARS-CoV-2 ancestral strain and the Alpha through Delta VOCs.^{9,15,16} However, the high levels of vaccine immunity against detectable SARS-CoV-2 infections wanes over a period of months.^{17,18} Immunity provided by 2-dose ChAdOx1 (AstraZeneca/Oxford ChAdOx1-nCoV-19 AZD1222 “ChAdOx1”) is somewhat lower and wanes faster than the mRNA vaccines.¹⁹ With hybrid immunity, neutralizing Ab (nAb) titers and breadth of recognition of SARS-CoV-2 variants are dramatically higher in previously infected individuals receiving at least one dose of a COVID-19 RNA vaccine² (Figure 2). Hybrid immunity from vaccination plus subsequent infection (breakthrough infection) also results in similarly robust immune responses.^{20,21} Multiple epidemiological studies have now validated those immunological findings, by observing that hybrid immunity results in more robust protection against COVID-19 than either previous infection immunity or vaccine-induced immunity.^{9,11} These types of antigen exposure(s)—infection-induced memory, vaccine-induced memory, and hybrid immunity—each have distinct characteristics of immune memory (Figure 2), which are key to the observed protective immunity in each case. High circulating nAb titers can clearly provide protective immunity. However, such high nAb titers are not present in many cases, particularly after SARS-CoV-2 infection, and there is substantial evidence for roles of T cells, B_{Mem}, and local tissue immunity in protection from COVID-19^{22,23} (Goldblatt et al., this volume).²⁴ For example, previous infection provides substantial immunity with significantly lower circulating nAb titers than after

BNT162b2 or ChAdOx1 vaccination.¹⁹ Local tissue immunity contributes to protective immunity after infection (Figure 3). Different adaptive immunity mechanisms may be involved in protective immunity to differing degrees with immune memory generated by previous infection, vaccination, or hybrid immunity. Thus, each is discussed in this review for CD8 T cells, CD4 T cells, B_{Mem}, and Abs.

2 | CD8 T CELL MEMORY TO SARS-CoV-2

In general, CD8 T cells are important in the control and clearance of viral infections.²⁵ In particular, several lines of evidence suggest that CD8 T cells are a relevant and valuable component of the overall adaptive immune response to SARS-CoV-2. One line of evidence is derived from studies in acute SARS-CoV-2 infection, which observed that early CD8 T cell responses were significantly associated with milder disease.²⁶ An inverse association between CD8 T cell response magnitude and disease severity was also reported in 4 of 5 additional independent studies.^{4,27,28,29,30} Additional evidence for a role of CD8 T cells comes from studies in non-human primates. McMahan et al.³¹ directly showed that depletion of CD8 T cells in COVID convalescent animals affected immunity against SARS-CoV-2 re-challenge. An important role for CD8 T cell responses was also reported in an antibody-independent COVID-19 vaccine study.³² See companion article by Goldblatt et al. for a review of T cells in protection²⁴ and ref. 33.

Multiple techniques are commonly utilized to measure antigen-specific T cell responses, with antigen specificity ensured by the use of SARS-CoV-2 derived peptides or defined epitopes, utilized as pools or isolated epitopes. The techniques utilized included activation induced marker (AIM), ICS (intracellular cytokine staining), ELISPOT, and tetramer staining assays and their strengths and weaknesses are reviewed elsewhere.^{34,35}

2.1 | CD8 T cell memory to SARS-CoV-2 infection

SARS-CoV-2-specific CD8 T cells are detectable in approximately 70% of COVID-19 cases 1 month after infection.^{3,4} The frequency of responders then declines to approximately 50% by 8 months post-infection.^{3,4} The Dan et al. study included SARS-CoV-2-specific CD8 T cell measurements from 169 COVID-19 case subjects and the Cohen et al. study included 114 subjects, making them the two largest studies of CD8 T cell memory to an acute viral infection




FIGURE 1 Components of immune memory. Virus-specific CD4 T cells, CD8 T cells, Abs, and B_{Mem} cells constitute the four major components of immune memory to a viral infection

examining a 6+ month period. The estimated SARS-CoV-2-specific memory CD8 T cell kinetics from the Dan et al. study was $t_{1/2} = 125\text{--}190$ days, while the Cohen et al. $t_{1/2}$ was 196 days. That is strong concordance between the two studies, given that the studies utilized different CD8 T cell assays (AIM and ICS) and the calculations were based predominantly on cross-sectional sampling. By comparison, a rigorous study using in vivo deuterium labeling found yellow fever virus (YFV) vaccine CD8 T cell responses to have an initial $t_{1/2}$ of 123 days.³⁶ Given that the YFV vaccine is highly effective, elicits robust CD8 T cells as a live attenuated viral vaccine, and deuterium labeling determined a subsequent $t_{1/2}$ of 460 days,³⁶ the observation of similar initial $t_{1/2}$ for memory CD8 T cells after a SARS-CoV-2

infection indicates the generation of long-lasting memory SARS-CoV-2-specific CD8 T cells^{3,4} (Figure 2). Memory CD8 T cells to SARS-CoV were detected 17 years post-infection.³⁷

SARS-CoV-2-specific CD8 T cells generated in response to SARS-CoV-2 infection predominantly express IFN γ and granzyme B (GzB),^{4,38,39} with some expression of TNF and IL-2.⁴ Eight months post-infection, effector memory (T_{EM}) and CD45RA+ effector memory (T_{EMRA}) phenotype memory CD8 T cells predominate, with a smaller fraction of central memory (T_{CM}) phenotype cells.^{3,4} Other studies have corroborated these central findings, with fewer COVID-19 cases, shorter study periods, or inferred CD8 T cells from PBMC ELISPOT assays.^{40,41} Bystander CD8 T




FIGURE 2 Kinetics of immune memory to SARS-CoV-2 infection and COVID-19 vaccines. Schematics of immune memory components against SARS-CoV-2. (A) Memory CD8 T cells, (B) memory CD4 T cells, (C) memory T_{FH} cells, (D) neutralizing antibodies, and (E) B_{Mem} cells. For T cell memory, with vaccines memory is to spike, and with infection, memory is to the entire virus. For B cell memory, spike-specific is shown in all cases. "Inf" = SARS-CoV-2 infected. "Hybrid" = Hybrid immunity, infected and then vaccinated. "mRNA" = Moderna mRNA-1273 or Pfizer/BioNTech BNT162b2, a 3 dose regimen. "NVX" = Novavax NVX-CoV2373, given as the 2-dose regimen in the main clinical trials. "J&J" = Janssen Ad26.COVS.2, given as the 1-dose approved by EUAs. Lines are color coded by vaccine. CD8 T cell % indicates the % of individuals with detectable CD8 T cell memory at 3–6 months. Scales are non-quantitative, but the antibody scale approximates log₁₀ and the cellular scales approximate log₂. For hybrid immunity, in this schematic, the vaccination occurs at approximately 6 months, indicated by the blue triangle. For mRNA vaccines, in this schematic, the 1st two dose are given at d1 and d21–28, with the 3rd dose ("booster") given at approximately 8 months, indicated by the red arrows




FIGURE 3 Components of local tissue immunity. Human immune responses are most often measured in blood, but immune responses at local sites of infection and/or portals of entry are important and may not be directly reflected by blood measurements

cell activation can occur during COVID-19,⁴² and some AIM markers can represent bystander activation of CD8 T cells and must be used carefully, depending on the experiment context, to avoid miscalculation of antigen-specific CD8 T cells. SARS-CoV-2-specific memory CD8 T cells have also been identified with MHC I tetramers/multimers.^{27,43,44,45,46,47} There has been some confusion about the expression of PD-1 on CD8 T cells in COVID-19. PD-1 is expressed on virtually all activated CD8 T cells. PD-1 is also a marker of exhausted CD8 cells in some contexts. Expression of PD-1 by itself does not indicate an exhausted T cell. While CD8 T cells have been observed to express PD-1 after COVID-19, and increased activation/exhaustion markers were noted in mild as compared to hospitalization-level or severe disease^{48,49}; subsequent studies using tetramers reported that at later time points the memory PD-1-expressing CD8 T cells are not exhausted, and appear to be highly functional.⁴⁶

Notably, SARS-CoV-2-specific memory CD8 T cell responses are undetectable in approximately 30% of COVID-19 cases,^{3,4,38} even when testing the full SARS-CoV-2 ORFeome of epitopes.³ It is unknown whether those subjects have CD8 T cell responses to untested epitopes, CD8 T cell responses just below the technical limit of detection of the assays used, or whether those subjects truly did not develop CD8 T cell response to SARS-CoV-2 infection.

Lower SARS-CoV-2-specific CD8 T cell responses have been observed in hospitalized COVID-19 cases in multiple studies,^{4,26,29} including memory, consistent with a weak CD8 T cell response predisposing for more severe COVID-19, particularly in older adults possessing fewer naive CD8 T cells.²⁶ This contrasts with CD8 T cell responses to YFV vaccine, which positively correlate with viral load. This also contrasts with positive correlations in SARS-CoV-2 infections between spike Ab titers, B_{Mem} cell frequencies, and disease severity, indicating that the humoral immune response and memory development correlates with acute SARS-CoV-2 viral loads (see B_{Mem} cell and Antibody sections). One interpretation of these data is that acute T cell responses are important for the control and clearance of SARS-CoV-2 infection.²²

Many SARS-CoV-2 antigens are recognized by human CD8 T cell responses in SARS-CoV-2-infected individuals,³⁸ with an estimated median of 17 epitopes per individual.⁵⁰ Recognized class I epitopes are distributed throughout the SARS-CoV-2 ORFeome, but structural proteins (spike, nucleocapsid, and M) are relatively immunodominant.^{38,50} CD8 T cell epitopes are in general well conserved between variants.⁵¹⁻⁵³ SARS-CoV-2 CD8 T cell specificities have been recently reviewed elsewhere.^{54,55}

The frequencies of circulating memory CD8 T cells after SARS-CoV-2 infection are lower than the frequencies reported for influenza in humans,⁵⁶ though influenza memory reflects the accumulation of many exposures throughout life. The frequencies of circulating SARS-CoV-2 memory CD8 T cells can be considered relatively low in comparison with frequencies generated to viruses in animal models associated with high levels of CD8 T cell-mediated protective immunity. However, the frequency of memory CD8 T cell required for control of an infection is generally highly dependent on the speed of the clinical disease. Most animal models of CD8 T cell-mediated protection require CD8 T cell immediate effector functions (e.g., cell killing at the site of infection within hours), or viral clearance within a few days. However, once activated CD8 T cells proliferate rapidly and migrate to infected tissues (sites of inflammation), with the cells potentially expanding in number 10-fold every 24 hours. Thus, within 72 hours of activation memory CD8 T cell can increase in number close to 1000-fold. Importantly, progression of hospitalization with COVID-19 is relatively slow, often occurring 10 days post-infection or later.^{57,58} Thus, a role for CD8 T cell in prevention of severe COVID-19 may require quite low levels of circulating memory CD8 T cells at the time of infection. In contrast, since CD8 T cells must exert their functions directly on infected cells, a role for CD8 T cells in prevention of SARS-CoV-2 transmission (half of which occurs within the first 5 days), before reported symptoms⁵⁹⁻⁶¹ is a high bar and likely requires a large number of tissue-resident memory (T_{RM}) CD8 T cells in the URT (Figure 3), which nevertheless can be achieved by immunization.³²

T_{RM} are an important category of T cells for protective immunity. They are underreported in the human immunology literature because it is much more difficult to obtain human tissue samples and isolate T cells from diverse tissues in comparison with blood. It is also much more difficult to identify antigen-specific T cells from such samples, due to limited cell numbers recovered and different cellular phenotypic characteristics.⁶¹ In the context of SARS-CoV-2, immune memory in tissues is particularly relevant in the nasal passages, oral cavity, throat, and lungs (Figure 3). SARS-CoV-2-specific CD8 T cell T_{RM} have been identified in humans.^{62,63} Lung tissue was examined from four subjects up to 6+ months after unremarkable (non-hospitalized) cases of COVID-19. Lymphoid tissues such as blood and lymph nodes (LNs) were examined for comparison. While SARS-CoV-2-specific CD8 T cell was identified in blood from only 2 out of 4 subjects (consistent with the detection rate in much larger studies^{3,4}), SARS-CoV-2-specific CD8 T cell was identified in lung LNs from 4 out of 4 subjects, and

SARS-CoV-2-specific CD8 T cell was identified in lung tissue from 3 or 4 out of 4 subjects.⁶² Additionally, substantial fractions of the CD8 T cells in both lungs and lung LNs were T_{RM} cells. Notably, in two studies SARS-CoV-2-specific CD8 T cell frequencies in lung tissue or bronchoalveolar lavage (BAL) were substantially higher than blood.^{62,63} SARS-CoV-2 is also present in gut tissues. While intestinal tissue was not available, intestinal LNs were examined, and SARS-CoV-2-specific CD8 T cell and CD8 T cell T_{RM} were found in all three samples tested, suggesting that substantial CD8 T cell T_{RM} may be present in intestinal tissue of previously infected individuals.⁶² Thus, blood samples may significantly underreport SARS-CoV-2-specific CD8 T cell memory in previously infected individuals. SARS-CoV-2-specific T_{RM} from throat, nasal passages, or the oral cavity after COVID-19 are less well studied,⁶⁴ representing a major knowledge gap for immune memory.

Development of pediatric T cell memory is of particular interest in the context of the well-known lower disease susceptibility of children to severe COVID-19, and in the context of the many differences in adult versus pediatric immune reactivity.⁶⁵ Conflicting data have been reported regarding pediatric T cell responses to COVID-19. In one study, significantly stronger acute and memory T cell responses were reported.⁶⁶ In another study, significantly lower acute and memory CD8 T cell responses were reported.⁶⁷ Of potential interest is also the observation that T cell memory in children may have differences in immunodominance, perhaps as a result of different exposure to common cold coronaviruses.^{66,67} A small fraction of pediatric SARS-CoV-2 infections later develop a serious hyperinflammatory condition, multisystem inflammatory syndrome in children (MIS-C). Conflicting data have been reported regarding the relative strength of T cell memory in MIS-C versus non-MIS-C cases.⁶⁸⁻⁷³

Long COVID (aka post-acute sequelae of SARS-CoV-2 infection [PASC]) is an important collection of conditions, with unclear etiology.⁷⁴ Given that SARS-CoV-2 viral RNA and protein have been detected 90+ days post-infection in gut biopsies from unremarkable COVID-19 cases (not hospitalized, and not long COVID), it is plausible that significant SARS-CoV-2 viral persistence occurs in some individuals in some tissues. This could be a source of inflammation and symptoms for at least some cases of long COVID. It is unclear why CD8 T cell would not clear SARS-CoV-2 infection from intestines within a few weeks [Goldblatt et al].²⁴ Efficacy of CD8 T cell clearance of virus in humans remains an important knowledge gap. Notably, SARS-CoV-2-specific CD8 T cell responses are undetectable in approximately 30% of COVID-19 cases,^{3,4,38} even when testing the full SARS-CoV-2 ORFome of epitopes.³ It is unknown whether those subjects have CD8 T cell responses to untested epitopes, CD8 T cell responses just below the technical limit of detection of the assays used, or whether those subjects truly have no CD8 T cell response to SARS-CoV-2 infection. It is also possible that some subjects in which SARS-CoV-2-specific CD8 T cell responses are undetectable in the periphery, might nevertheless have T_{RM} antigen-specific cells residing in the lung or upper respiratory tract tissues.^{62,75} Separately, CXCR6⁺ CD8 T cells in lungs have been

associated with extended periods of COVID-19 inflammation and may have a role in long COVID.⁶³

2.2 | CD8 T cell memory to vaccination

Spike-specific CD8 T cell responses are detected in approximately 70%–90% of individuals weeks after receiving 2-dose mRNA COVID-19 vaccines,⁷⁶⁻⁷⁹ and memory CD8 T cells are detectable in approximately 41%–65% of individuals at 6 months after the 2nd dose (7 months from 1st dose).^{76,77,78,80} A low dose (25 µg) of mRNA-1273 was found to generate memory CD8 T cells at similar frequencies as previous infection, comparing 6 months after the 2nd dose to 6 months after infection, indicating similar spike-specific CD8 T cell responses between mRNA vaccination and infection. A separate study also found similar spike-specific CD8 T cell responses at earlier times.⁸¹ Among individuals with detectable CD8 T cell memory to mRNA vaccines months after immunization, the magnitude of the memory is generally observed to be low,^{76,77,79,80,82} both in comparison with spike-specific CD4 T cell memory, and memory to influenza.⁵⁶

There was initial confusion about whether both BNT162b2 and mRNA-1273 mRNA COVID-19 vaccines generated CD8 T cell responses.^{43,83,84} More recently, multiple groups have observed similar CD8 T cell responses to both vaccines,⁷⁷ including in head-to-head comparisons.^{77,79,82} Methodological differences measuring antigen-specific T cells can result in different findings, as studies not detecting memory CD8 T cells usually utilized a less than optimal short stimulation ICS protocol, or less sensitive ELISPOT formats. The mRNA vaccine-elicited memory CD8 T cells detected predominantly have a T_{EM} surface phenotype, consistently express IFN γ , and have proliferative capacity.^{44,79}

The first 6 months are likely to be the period with the fastest decline in T cell memory.³⁶ The observation of approximately twofold declines at 6 months in spike-specific CD8 T cell memory from peak cytokine-positive CD8 T cell frequencies is encouraging evidence that CD8 T cell memory to mRNA vaccines is long-lived and may last many years^{76,78,79} (Figure 2). Memory CD4 T cells exhibit similar kinetics, as discussed in the CD4 T cell section below.

The adenoviral vector vaccines ChAdOx1 and Ad26.COVS.2.S elicit spike-specific CD8 T cell responses in 51%–64% of individuals in immunogenicity clinical trials,⁸⁵ though the response rate drops to 24%–36% in individuals >65 years old.⁸⁵ Stable CD8 T cell memory to Ad26.COVS.2.S is observed to 8 months.⁸⁶ Some comparisons between mRNA vaccines and adenoviral vector vaccines are available for CD8 T cell memory. Similar (within approximately twofold) spike-specific CD8 T cell responses to mRNA vaccines and adenoviral vector vaccines are observed approximately 1 month after immunization, including the 1-dose Ad26.COVS.2.S,^{77,79,82,87} 2-dose Ad26.COVS.2.S,⁸⁷ or 2-dose ChAdOx1.⁸⁸⁻⁹⁰ Two studies that assessed CD8 T cell memory at 5+ months determined that mRNA-1273 elicited larger spike-specific CD8 T cell memory than Ad26.COVS.2.S.^{77,79} Two other studies reported the opposite,^{91,92}

with a clinical trial reporting 45/57 Ad26.COVID.2.S subjects and only 20/116 BNT162b2 or mRNA-1273 subjects positive for CD8 T cell memory.⁹² In PBMC IFN γ ELISPOT assays, T cell memory 2–4 months after ChAdOx1 and BNT162b2 appears to be equivalent, though CD8 and CD4 T cells were not distinguished^{93,94} (bulk PBMC IFN γ ELISPOT assay signal comes from a mixture of CD8 T cells and CD4 T cells.⁴³). A similar observation was made for Ad.COVID.2.S.⁹⁵ In sum, CD8 T cell memory to mRNA and adenoviral vector COVID-19 vaccines appears to be similar in magnitude and % responders (Figure 2), but conclusions vary depending on the study. Additional head-to-head studies or memory are warranted, including examination of CD8 T cell functionality.

Memory CD8 T cells elicited by mRNA vaccines recognize diverse spike epitopes.^{50,51,77} Memory CD8 T cells largely have conserved recognition of variants, including Omicron.^{77,81,87,96,97,98} Mix & match adenoviral vector + mRNA vaccine approaches may increase CD8 T cell responses^{88,89,95} and thereby may alter CD8 T cell memory. Perplexingly, lower CD8 T cell responses were reported to vaccine extended dose intervals, with the caveat that minimal CD8 T cells were measurable in any group.⁹⁹

2.3 | CD8 T cell memory in hybrid immunity

Modest differences have been observed between vaccination only and hybrid immunity for circulating spike-specific CD8 T cells in most studies,⁹⁹ as well as no difference based on symptomatic or asymptomatic infection¹⁰⁰ (Figure 2). In one study, no difference in memory IFN γ ⁺ spike-specific CD8 T cells was observed between vaccination only and hybrid immunity after 6 months.⁸⁰ Multiple studies observed increased T cell responses in hybrid immunity compared to infection or vaccination alone without distinguishing between CD4 and CD8 T cells.^{81,99} CD8 T cell repertoire diversity is maintained after multiple exposures.¹⁰¹

3 | CD4 T CELL MEMORY TO SARS-CoV-2

Memory CD4 T cells are important in the control and clearance of viral infections, both directly and by the effects exerted in the support and amplification of antibody responses. Several different subsets of CD4 T cells can differentiate in antigen-specific responses to infections. This heterogeneity is manifested at the level of different memory subsets, each associated with distinctive patterns of cytokine secretion, transcription factors, and differentiation profiles (T_{H1}, T_{H2}, T_{H17}, T_{FH}, and others¹⁰²). This heterogeneity is further amplified by a diverse array of functional roles. T_{FH} cells play a key role in orchestrating the development and maturation of antibody responses,^{103,104} while Th1 and cytotoxic CD4 T cells (CD4-CTL) can exert direct antiviral functions.^{105–107} Multiple lines of evidence suggest that CD4 T cell is a relevant and valuable component of the overall adaptive immune response to SARS-CoV-2.^{22,33,108} Protective effects of CD4 T cells against

COVID-19 are fully reviewed in the companion article [Goldblatt et al., this volume].²⁴

3.1 | CD4 T cell memory to SARS-CoV-2 infection

Dan et al. found that antibody, CD4 T cell, CD8 T cell, and B cell memory responses were durable over 8 months after infection, with 95% of the subjects still retaining multiple measurable memory responses, including memory CD4 T cells.³ Notably, memory CD4 T cells are detectable in 93% of COVID-19 cases 1 month after infection³ and still 92% at >6 months post-infection.³ Multiple studies have reported similar memory CD4 T cell findings,^{4,39,40,109} with a notable large longitudinal study.⁴ The estimated $t_{1/2}$ of memory SARS-CoV-2-specific CD4 T cells is 94–207 days during the first 8 months,^{3,4} with the $t_{1/2}$ likely increasing substantially over time, based on a study determining a $t_{1/2}$ of 377 days for memory SARS-CoV-2-specific CD4 T cells at 6–15 months post-infection¹¹⁰ (Figure 2), as well as similar data on CD8 T cell memory against a different virus.³⁶ These SARS-CoV-2 infection data are consistent with T cell memory to SARS-CoV being detected 17 years post-infection.^{37,111,112,113}

SARS-CoV-2-specific memory CD4 T cells after SARS-CoV-2 infection predominantly are T_{H1}, T_{FH}, and CD4-CTL cells.^{3,4,110} Eight months post-infection, the memory CD4 T cells predominantly have central memory (T_{CM}) and effector memory (T_{EM}) surface phenotypes.^{3,4} Virus-specific T_{H2} cells and T_{H17} cells are generally not detectable.^{4,26,38} Regarding the T_{H1} memory cells, they are a stably maintained population,⁴ predominantly expressing CD40L and IFN γ ,⁴ with significant expression of TNF and some expression of IL-2.^{4,29,56,100,110,114} The CD4-CTL cells express CD40L and granzyme B (GzB).^{4,79,115,116} CD4-CTL cells are of interest because SARS-CoV-2-infected epithelial cells upregulate class II expression,¹¹⁷ and CD8 T cell responses appear to be low in many individuals, leaving open the possibility that memory CD4-CTL may compensate.^{117,118} Memory T_{FH} cells are generated after SARS-CoV-2 infection and are stably maintained after a brief decline^{3,110,119} (Figure 2). The memory T_{FH} cells are highly functional, as they greatly enhance nAb responses to COVID-19 vaccines.^{2,120,121} Some memory T_{FH} cells express CCR6, which is associated with lung homing.^{3,119} Other memory T_{FH} cells express CXCR3, which is associated with rapid anamnestic antibody responses,^{104,122} while the CXCR3^{neg} memory T_{FH} population is associated with higher quality germinal center and antibody responses.^{123,124} Germinal centers are discussed in the B cell memory section.

In terms of anatomical location of memory T cells, memory CD4 T cells are detectable in the bone marrow, spleen, lung, and multiple lymph nodes (LNs) for 6+ months after infection.⁶² Memory T_{FH} cells are observed in LNs.⁶² CD4 T_{RM} cells were present at substantial frequencies in lungs⁶² and BAL⁶³ (Figure 3). Less is known regarding SARS-CoV-2-specific CD4 T_{RM} in the URT and oral cavity.

Many SARS-CoV-2 antigens are recognized by human memory CD4 T cells in previously infected individuals,³⁸ with an estimated median of 19 epitopes per individual.⁵⁰ Recognized class II epitopes

are distributed throughout the SARS-CoV-2 ORFome, but structural proteins (spike, nucleocapsid, and M) are relatively immunodominant.^{4,38,50,125} CD4 T cell epitopes are in general well conserved between variants.⁵¹⁻⁵³ SARS-CoV-2 CD4 T cell specificities have been recently reviewed elsewhere.^{54,55}

Human antiviral immune memory may be influenced by variables such as viral factors related to the infection event such as viral dose and tissue distribution, and host factors such as age, sex, and general health of the host.¹²⁶ A distinctive feature of SARS-CoV-2 infection and COVID-19 disease is the wide range of clinical outcomes, ranging from fully asymptomatic infection to severe disease and death. The heterogeneity in clinical outcomes associated with SARS-CoV-2 infection and COVID-19 disease is paralleled by large variations in heterogeneity at the level of CD4 T cell responses.^{3,26} Specifically, SARS-CoV-2 T cell responses are influenced by older age and the size of the pool of naive T cells.²⁶ No significant difference in CD4 T cell memory is observed between males and females.^{3,4} Pre-existing comorbidities in COVID patients have been reported to affect magnitude and helper T cell subset composition.¹²⁷ T cell responses of a homogenous group of healthy young males were still widely heterogeneous,¹²⁸ suggesting that heterogeneity of responses is driven by variables other than predisposing conditions, age, and sex. As noted above, conflicting data have been reported regarding pediatric T cell responses to COVID-19. In one study, stronger acute and memory CD4 T cell responses were reported in one study,⁶⁶ but significantly lower acute and memory CD4 T cell responses were reported in another study.⁶⁷ Conflicting data have been reported regarding CD4 T cell memory in MISC versus non-MISC cases.⁶⁸⁻⁷³

Asymptomatic SARS-CoV-2 infections may reflect shorter infections with lower overall viral loads in tissues, which could be expected to have lower T cell memory as a result of lower antigen exposure. Comparison between asymptomatic and mild symptomatic COVID-19 cases revealed slightly lower T cell memory among asymptomatic COVID-19 cases.^{39,129} The relation between COVID-19 severity and quality of memory CD4 T cells is still an open topic of debate and investigation.^{115,125,130,131} In the context of long COVID,⁷⁴ alterations of T cell responses lasting several months post-infection have been reported,⁶³ as have increased total T cell accumulation in BAL.¹³²

3.2 | Crossreactive memory T cells

Memory CD4 T cells able to recognize SARS-CoV-2 have been demonstrated in unexposed subjects, with the clearest evidence coming from blood samples obtained during pre-pandemic times.^{37,38,115,133,134} It was hypothesized that these cells may predominantly be memory T cell to previous common cold coronavirus (CCC) infections.^{54,135} Indeed, at least in some cases, these memory T cells cross-recognize SARS-CoV-2 and CCCs.^{28,46,125,136,137,138,139} Detailed studies suggest that cross-recognition across distant viral species can occur, but rather infrequently,^{140,141} and is observed for SARS-CoV-2 sequences.^{129,142} In general, SARS-CoV-2 crossreactive

memory T cells have been most often described in the case of CD4 T cells, and less often for CD8 T cells.³⁸ In terms of antigen specificity, the sequences associated with crossreactive memory are often derived from non-structural antigens encoded in the Orf1ab, which correlates with the higher degree of conservation of across the genome of CCCs and other coronaviruses.⁵⁴

It has been debated to what extent this pre-existing crossreactive T cell memory is functional and biologically relevant.^{135,143} The T cells associated with this pre-existing immunity display classical memory markers,¹³⁶ and were detected by a variety of assays. However, this crossreactive recognition can be of low affinity, particularly in the case of more distant unrelated viruses.¹⁴² Additionally, SARS-CoV-2 infection is associated with development of T cell response that is largely focus on novel epitopes.⁵⁰ Nevertheless, it has now been demonstrated that the crossreactive memory T cells are biologically functional. Pre-existing crossreactive memory T cells exert a positive influence on COVID-19 vaccination outcomes.^{76,144,145} This is consistent with two reports that persons with CCC infections within recent years preferentially had less severe COVID-19 outcomes,^{146,147} while a different study found no association.¹⁴⁸ Healthcare workers were observed to have high levels of SARS-CoV-2 crossreactive memory T cells, and CCC-specific T cells.¹⁴⁹ It was further shown that the presence of these crossreactive T cells was linked to favorable outcomes in a large healthcare worker cohort during the first wave of the pandemic, with crossreactive memory CD4 T cells possibly providing protection resulting in abortive SARS-CoV-2 infection.¹⁵⁰ Evidence of protection was also observed in a household contacts study.¹⁵¹ Overall, crossreactive memory CD4 T cells recognizing SARS-CoV-2 existed in approximately 50% of individuals pre-pandemic, those crossreactive CD4 T cells have functional properties in vivo, and they have been associated with some degree of protection from COVID-19.

3.3 | CD4 T cell memory to vaccination

COVID-19 vaccines can elicit robust CD4 T cell memory. Spike-specific CD4 T cell responses are detected in close to 100% of individuals weeks after receiving 2-dose mRNA COVID-19 vaccines,⁷⁶⁻⁸⁰ and memory CD4 T cells are detectable in approximately 100% of individuals at 6 months after the 2nd dose (7 months from 1st dose).^{76,77,78,80} mRNA-1273 generated spike-specific memory CD4 T cell frequencies higher than seen in previously infected individuals,⁷⁹ while BNT162b2 generate spike-specific memory CD4 T cell frequencies similar to infection.^{79,80} A dose of mRNA-1273 similar to that of BNT162b2 generated spike-specific memory CD4 T cells at frequencies comparable to previous infection,⁷⁶ indicating that differences between memory CD4 T cells after the two mRNA vaccines most likely predominantly relate to the different doses of the two vaccines. Reductions in memory CD4 T cell frequencies over 6 months were modest, and half-lives of memory CD4 T cells after mRNA COVID-19 vaccines appear to be at least as long as after infection^{76,79,80} (Figure 2). Memory CD4 T cells are generated after a

single dose of mRNA vaccine or Ad26.COv2.S that maintained at least several months.^{79,86,95,99} In the context of vaccine interval extension protocols, IL-2⁺ memory CD4 T cells were increased in vaccinees who waited longer before the 2nd mRNA vaccine dose.⁹⁹

After vaccination, memory T_{H1} cells are stably maintained,^{77,78,79,80,97} and express CD40L, IFN γ , TNF, and IL-2.^{78,79} Memory T_{FH} cells represent approximately 25% of CD4 T cell memory after mRNA immunization,⁷⁹ and the abundance of the cT_{FH} cells is associated with the magnitude of the nAb response.^{76,79,120,121,152} Vaccine-elicited memory T_{FH} cells in blood are stably maintained with minimal decline over 6+ months,⁷⁹ though the cT_{FH} may change phenotype during the first months.^{80,110,153} Active germinal center T_{FH} (GC-T_{FH}) cells are found in LNs for at least 6 months and appear to be critical for maintaining germinal centers and development of nAbs after vaccination.^{152,153} Durable T_{FH} cell memory in blood was observed for mRNA vaccines, Ad26.COv2.S, and NVX-CoV2373.⁷⁹ Memory CD4-CTL cells are also generated in response to several COVID-19 vaccines and are stably maintained for at least 6 months.⁷⁹ Overall, each major subset of memory CD4 T cells is maintained for at least 6 months after vaccination with BNT162b2, mRNA-1273, Ad26.COv2.S, and NVX-CoV2373, with kinetics that indicate the memory CD4 T cells will be substantially maintained for years (Figure 2).

After ChAdOx1-nCoV-19 immunization, polyfunctional T_{H1} memory CD4 T cells are induced.^{154,155} Similar (within approximately twofold) spike-specific CD4 T cell responses to mRNA vaccines and ChAdOx1-nCoV-19 are observed approximately 1 month after 2 doses.⁸⁸ Immunization with 2 doses of the inactivated SARS-CoV-2 alum and imidazoquinolin adjuvanted vaccine BBV152 (Covaxin) generates memory CD4 T cell responses comparable to that seen in infected individuals, stably maintained for over 6 months.¹⁵⁶ Limited T cell memory data are available for several other vaccines, including Coronavac, Sinopharm, and Sputnik.

Memory CD4 T cells elicited by mRNA vaccines recognize diverse spike epitopes.^{50,51,77} Recognition of variants by memory CD4 T cells is maintained in mRNA, Ad26.COv2.S, and NVX-CoV2373 vaccinees.^{77,81,96,97}

3.4 | CD4 T cell memory in hybrid immunity

Modest differences have been observed between vaccination only and hybrid immunity for circulating spike-specific memory CD4 T cells in most studies,^{99,114,120,157} as well as no difference based on symptomatic or asymptomatic infection.¹⁰⁰ In one study, no difference in memory spike-specific CD4 T cells was observed between vaccination only and hybrid immunity after 6 months⁸⁰ (Figure 2). Multiple studies observed increased T cell responses in hybrid immunity compared to infection or vaccination alone using IFN γ ELISPOTs that do not distinguish between CD4 and CD8 T cells,^{81,99} suggesting functional changes may occur. Indeed, a distinct population of spike-specific IFN γ ⁺ IL-10⁺ T_{H1} memory cells is observed in hybrid immunity but not after vaccination alone, demonstrating a function

of imprinting on the memory T_{H1} cells by infection.¹¹⁴ There is dramatic enhancement of antibody and B cell responses in persons with hybrid immunity, demonstrating a strong functional role for memory T_{FH} cells in hybrid immunity, discussed elsewhere.

An additional important aspect of hybrid immunity is the location of the T cell memory. Intramuscular vaccination is expected to generate almost exclusively circulating T cell memory. In contrast, SARS-CoV-2 infection generates both circulating T cell memory and T_{RM} (Figure 3). Thus, hybrid immunity is expected to result in both circulating and T_{RM}, but it is unclear if the vaccines enhance T_{RM} already present from infection. Lastly, if the order is vax+infection, it is unknown whether the T_{RM} are qualitatively or quantitatively different than what occurs after infection alone.

4 | B CELL MEMORY TO SARS-CoV-2

Human B_{Mem} cells can be exceptionally long-lived, with smallpox vaccine B_{Mem} lasting >50 years,¹⁵⁸ and B_{Mem} cells generated from infections during the 1918 pandemic lasting at least 90 years.¹⁵⁹ B_{Mem} cells are re-activated upon an infection and are the source of classic anamnestic antibody responses. B_{Mem} cells serve two purposes. The first is a cellular source for the anamnestic antibody response. B_{Mem} cells can plausibly reactivate and generate an anamnestic antibody response within 3–5 days.¹⁶⁰ The second important value of B_{Mem} cells is to serve as a library of predictions by the immune system of possible future viral variants.^{2,161} The COVID-19 pandemic has dramatically demonstrated the importance of B_{Mem} cell diversity in the recognition of a pathogen and variants, also highlighting the brilliance of the immune system at predicting viral mutations, embedding those predictions in the B_{Mem} cell repertoire. B_{Mem} cells likely play a role in protective immunity against SARS-CoV-2 infection by both of the mechanisms above, and protection by B_{Mem} cells is reviewed in the accompanying article [Goldblatt et al.].²⁴

4.1 | B cell memory to SARS-CoV-2 infection

Detectable B_{Mem} cells develop within two weeks of symptom onset after SARS-CoV-2 infection.^{3,4} Strikingly, B_{Mem} cell frequencies continuously increase over the course of 3–6 months post-infection.^{3,4,162} Spike-, RBD-, and nucleocapsid-specific B_{Mem} cells all exhibit this increase, in a cohort of 160 individuals.³ SARS-CoV-2-specific B_{Mem} cell frequencies stabilize approximately 4 months post-infection³ and are maintained for at least 15 months^{162,163} (Figure 2). These spike- and RBD-binding B_{Mem} cell frequency increases are associated with substantial somatic hypermutation (SHM) for 6 months,^{162,164,165} continuing for at least 12 months.¹⁶² The B_{Mem} cell antibody mutations accumulated over 6–12 months demonstrated increased affinity maturation and increased neutralization potency, particularly against variants.^{162,165} These patterns are all indicative of long-lasting germinal centers after SARS-CoV-2 infection; an exception, however, is fatal COVID-19, in which profound

disruption of germinal centers can be observed in autopsies.^{166,167} The high quality of the B_{Mem} cells after SARS-CoV-2 infection is also evidenced by the anamnestic nAb responses to variants after a subsequent vaccination or infection, as discussed in the "Antibody durability" sections below.

While $IgM^+ B_{Mem}$ cells initially comprise approximately 1/3rd of SARS-CoV-2-specific B_{Mem} cells, IgM^+ cells decline rapidly and are mostly undetectable after 5 months.^{3,4} $IgA^+ B_{Mem}$ cells are uncommon, comprising only approximately 5% of spike or RBD-specific B_{Mem} cells on average,^{3,4,165} but the $IgA^+ B_{Mem}$ cells are stably maintained over 8 months post-infection, in contrast to the $IgM^+ B_{Mem}$ cells.³ B_{Mem} cells can have diverse phenotypes. After SARS-CoV-2 infection, activated SARS-CoV-2-specific B_{Mem} cell frequencies are initially high, but decline over the course of 7 months, with a reciprocal increase in resting B_{Mem} cells.¹⁶⁴

COVID-19 severity does impact the magnitude of the B_{Mem} cell response. Patients with hospitalization-level COVID-19 develop higher RBD-specific B_{Mem} cell frequencies compared to individuals with mild COVID-19,^{3,164} similar to what is observed for antibody titers.¹⁶⁸ Asymptomatic cases develop similar Spike-specific B_{Mem} cell frequencies compared to symptomatic but non-hospitalized COVID-19 cases [Crotty manuscript in prep].

The detailed study of SARS-CoV-2-specific B_{Mem} cells in response to infection, over periods of 6–12 months, in multiple large independent cohorts, with a range of disease severities, and intensive BCR sequencing, amounts to the most detailed understanding of the development of B cell memory to any acute infection. In a small data set from two YFV vaccine (a live viral vaccine) recipients, increases in B_{Mem} cell frequencies were observed for 6 months, increases in affinity maturation were observed for over 6 months, and declining frequencies of IgM^+ or activated B_{Mem} cells were observed over 6+ months.¹⁶⁹ All of those features are commonalities shared with B_{Mem} cell responses to SARS-CoV-2 infection. Ebola infection B_{Mem} cell responses also have some commonalities, though the severity of Ebola disease and longevity of high viral loads may alter that response.¹⁷⁰ Overall, the B_{Mem} cell response to SARS-CoV-2 infection is quite impressive, with substantial RBD- and spike-specific B_{Mem} cell generation; and with only exposure to a single viral strain, the B_{Mem} cell compartment develops over several months to contain B_{Mem} cells with high neutralization potency and B_{Mem} cells capable of recognizing and neutralizing a range of variants.

Tissue-resident B_{Mem} cells (B_{RM}) can exist in some cases. Pathogen-specific tissue B_{RM} have been observed in lungs of mouse models,¹⁷¹ and B_{RM} phenotype B_{Mem} cells are found in multiple human tissues.¹⁷² B_{RM} have now been demonstrated in humans after SARS-CoV-2 infection⁶² (Figure 3). By studying organ donors with a previous history of unremarkable COVID-19 (non-hospitalized), it was shown that spike/RBD-specific B_{RM} were observed in the lungs of all subjects.⁶² Notably, the frequencies of spike/RBD-specific B_{Mem} cells in lungs were significantly higher than in spleen, indicating enrichment of local memory in lungs after SARS-CoV-2-infection. Spike/RBD-specific B_{Mem} cells were also found in bone marrow, lung LNs, and gut LNs.⁶² RBD-specific B_{RM} have also been observed in

BAL.⁶³ Local reactivation of B_{RM} in lungs could result in faster anamnestic antibody responses than from circulating B_{Mem} cells, based on data from a mouse influenza model.¹⁷¹ URT tissues were not available for B_{Mem} cell analysis, and this remains a knowledge gap, given that the URT is the primary site of SARS-CoV-2 replication and transmission.

Regarding long COVID, there is no SARS-CoV-2-specific B_{Mem} cell study reported for long COVID. As noted in the memory T cell sections, it is plausible that at least some long COVID cases are due to persistent infection in some tissue sites. As such, some of the extended SHM of B_{Mem} cells may be due to new SARS-CoV-2 antigen generation after the acute phase of the disease was resolved. These are significant remaining knowledge gaps.

4.2 | B cell memory to vaccination

B_{Mem} cells are generated in response to COVID-19 vaccines. Similar frequencies of RBD-binding $IgG^+ B_{Mem}$ cells are generated after 2-dose RNA vaccines or SARS-CoV-2 infection⁸⁰ (Figure 2). SHM levels are also substantial, and comparable between 2-dose RNA vaccines and SARS-CoV-2 infection at 5 months.^{80,173} A substantial fraction of RBD-binding $IgG^+ B_{Mem}$ cells from 2-dose RNA-vaccinated individuals also bind to VOC RBDs.^{77,80} Thus, 2-dose RNA vaccines generate substantial affinity matured B_{Mem} cells. Nevertheless, the affinity maturation after a standard 2-dose RNA vaccine regimen is qualitatively poorer than that after SARS-CoV-2 infection. Substantial improvements in nAb breadth were observed months after infection but not after RNA vaccination (e.g., 69% of nAbs from previously infected subjects had improved potency, but on 19% of nAbs did from 2-dose RNA vaccinees).¹⁷³ These qualitative differences may be related to the narrow time between dose 1 and dose 2 of the RNA vaccines. The priming period can be important for the quality of a B cell response. Extending the priming period can result in better nAb responses to HIV.^{174–178} Extending the dose interval between RNA vaccine immunizations from 3 to 10 weeks significantly improves nAb titers and nAb breadth,⁹⁹ most likely by impacting affinity maturation.

Spike and RBD $IgG^+ B_{Mem}$ cell frequencies increase between 3 and 6 months after immunization with mRNA vaccines,^{79,80,173} an adenoviral vector vaccine,⁷⁹ or a recombinant protein vaccine.⁷⁹ 1-dose Ad26.COVS vaccine elicits significantly lower spike and RBD $IgG^+ B_{Mem}$ cell frequencies than the mRNA vaccines⁷⁹ (Figure 2). NVX-CoV2373 also elicits lower spike and RBD $IgG^+ B_{Mem}$ cell frequencies than the mRNA COVID-19 vaccines⁷⁹ (Figure 2). Additionally, B_{Mem} cell frequencies are somewhat higher after mRNA-1273 compared to BNT162b2 vaccination.⁷⁹ Less is known regarding B_{Mem} cells after ChAdOx1 immunizations.

Germinal centers appear to be central to the immune responses to COVID-19 vaccines. Most nAb responses, class-switched B_{Mem} cells, and durable antibody responses to viral infections and vaccines are dependent on germinal centers.¹⁰⁴ For COVID-19 vaccines, nAb responses are substantially reduced in many immunocompromised

individuals, such as kidney transplant recipients. Direct examination of germinal centers in healthy subjects compared to kidney transplant recipients revealed dramatically weaker germinal center responses in kidney transplant recipients after mRNA vaccination.¹⁵² The smaller germinal center responses may be due to weaker GC-T_{FH} cell responses, as GC-T_{FH} cell frequencies were severely reduced and associated with poor nAb titers.¹⁵² Germinal centers are observed to continue in draining LNs of healthy vaccinated individuals for at least 6 months after BNT162b2 immunization.^{153,179,180} This is associated with presence of vaccine mRNA in germinal centers for at least a month, as well as detectable spike protein in the germinal centers.¹⁶⁷ Since the vast majority of the B_{Mem} cell response to COVID-19 vaccines is class-switched and contains SHMs, the data indicate that the vast majority of the B_{Mem} cell response, and nAb response, to COVID-19 vaccines is T_{FH}-dependent and germinal center-dependent.

4.3 | B cell memory in hybrid immunity

Circulating spike and RBD IgG⁺ B_{Mem} cell frequencies increase substantially in hybrid immunity,^{80,162,181} but become similar to 2-dose mRNA vaccination after 6 months⁸⁰ (Figure 2). In hybrid immunity, the RBD-binding B_{Mem} cells have substantially more SHM and affinity maturation than after vaccination alone.^{80,162,181} Functionally, this is observed most clearly with the significantly higher potency and variant breadth of nAbs from B_{Mem} cells in people with hybrid immunity compared to vaccination alone or infection alone.^{162,181} The robustness and quality of these responses are likely driven by memory T_{FH} cells and B_{Mem} cells, and can occur after infection + vax or vax + infection ("breakthrough"), discussed in the "Antibody durability" section below.

5 | ANTIBODY DURABILITY TO SARS-CoV-2 INFECTION OR COVID-19 VACCINES

Abs are key components of protective immunity against SARS-CoV-2. Thus, durability of Abs is a major topic of interest for protective immunity against SARS-CoV-2 for previously infected, vaccinated, or person with hybrid immunity. Acute Ab responses are primarily generated by B cells differentiating into short-lived plasma cells (short-lived B_{PC}). These short-lived B_{PC} only live for a few days. IgG protein has a long half-life of 21–28 days in the blood, and thus, a large short-lived B_{PC} response can result in detectable antibody titers in blood for months. Long-lived B_{PC} can survive for many years producing large quantities of Abs daily. Long-lived B_{PC} are typically the product of germinal center B (B_{GC}) cells.

5.1 | Antibody durability to infection

The vast majority of SARS-CoV-2-infected individuals seroconvert and develop nAbs (91%–99%).^{182–184} While nAb titers decline during

the first few months post-infection, nAb titers stabilize between 4 and 6 months post-infection, with little evidence of decline thereafter. After the initial decay phase (dominated by short-lived B_{PC}), the estimate SARS-CoV-2 nAb t_{1/2} is 254 days.⁴ This may further stabilize over time (Figure 2). NAb titers are detectable in approximately 80%–90% of SARS-CoV-2-infected individuals at 6 months and 12 months post-infection.^{3,4} Nevertheless, SARS-CoV-2 nAb titers in previously infected individuals are relatively low, resulting in enhanced interest in understanding all of the other compartments of immune memory nAb titers in previously infected individuals are relatively low. This led to concern that low circulating nAb titers would be insufficient for protection, and increased interest in defining other branches of potential immunity to SARS-CoV-2, such as T cells. Nevertheless, immune memory overall in previously infected individuals was robust,³ leading to a conclusion that natural immunity was likely sufficient to prevent reinfections of significant clinical concern in the majority of people for years.^{3,185}

SARS-CoV-2 spike- and RBD-binding IgG titers exhibit similar kinetics to that of nAbs,^{3,4,186,187,188,189} though not identical, depending on the study, likely due to affinity differences between the assays. A multi-phasic decay kinetic is observed, with a t_{1/2} of >700 days by 6–9 months post-infection.^{187,188} Long-lived B_{PC} are found in bone marrow 7–8 months after infection.¹⁸⁹ SARS-CoV-2-specific IgM responses are not durable, consistent with IgM responses being short-lived for most antigen exposures. SARS-CoV-2-specific serum IgA responses are relatively low but are durable at low levels in most individuals,^{3,4,190} with SARS-CoV-2-specific IgA long-lived B_{PC} detected in approximately 50% of individuals.¹⁸⁹

Long-term antibody titers are lower in asymptomatic cases at 6–16 months post-infection,^{191,192} with some individuals being seronegative, though some amount of this difference is due to false-positive PCR tests with high C_t values.¹⁹²

Antibody titers against other HCoVs are also relatively stable over time.⁴ This is consistent with human immunology findings for multiple acute viral infections and the live attenuated YFV, measles, and smallpox vaccines.^{158,193,194}

Nucleocapsid antibody assays have been found to not be trustworthy indicators of previous infection at timepoints >6 months post-infection. This may be due to a faster decay kinetics of nucleocapsid Abs,¹⁹⁵ or a high background signal from crossreactive nucleocapsid Abs against other HCoVs^{187,196} which may be more problematic in certain assay formats, making it more challenging to definitively distinguish SARS-CoV-2 nucleocapsid IgG. Thus, RBD IgG is more widely used as a serodiagnostic marker, though it cannot distinguish infection from vaccination.

Local immunity is important, and Abs are a key factor of local immunity, as they are the only component of adaptive immunity capable of providing sterilizing immunity (Figure 3). Circulating IgG is transudated into most mucosal tissues, and circulating IgG can provide protective immunity at mucosal surfaces. The most dramatic example of this is the human papillomavirus (HPV) vaccine, which provides 99% protective immunity in the vaginal tract, even though the vaccine is a conventional intramuscular immunization and elicits

circulating IgG. For SARS-CoV-2 previously infected individuals, the titers of circulating IgG correlate with saliva IgG,^{56,190,197} and the correlation was sustained over a period of 9 months.⁵⁶ Correlation between circulating IgA and saliva IgA in previously infected individuals was also substantial over a 9-month period.⁵⁶ A more rapid decay of IgG and IgA titers was observed in saliva compared to blood, possibly indicating local production of Abs in salivary tissue for a limited number of months post-infection.⁵⁶ Few studies have examined nasal passage Abs, but spike IgG is detectable 6 months after infection in the majority of individuals.¹⁰⁰ Overall, the evidence suggests that nAbs at nasal, oral, and lung tissues are important for protective immunity against SARS-CoV-2 and are likely present proportionally to serum IgG titers after infection.

5.2 | Antibody durability to vaccination

Two doses of an COVID-19 mRNA vaccine are incredibly successful at eliciting high titers of nAbs. However, the biggest shortcoming of the mRNA COVID-19 vaccines has been that the nAb titers decline continuously over a period of months. Vaccination with the mRNA-1273 mRNA vaccine generates peak RBD IgG and SARS-CoV-2 nAb titers twofold higher than the BNT162b2 mRNA vaccine, on average^{92,198}; as a result, Ab durability analyses are confounded in studies that mix vaccinees receiving the two vaccines. In one large study of 2600 recipients of the 2-dose BNT162b2 vaccine in Israel, RBD IgG continuously declined over 7 months from peak after the 2nd dose, with a 16-fold reduction in RBD IgG from peak¹⁹⁹ (Figure 2). In that study, a vaccinated cohort and a previous-infected cohort were directly compared; the RBD IgG titers declined extensively in the vaccinated individuals but were largely stable in the previously infected individuals.¹⁹⁹ NAb titers and RBD IgG similarly declined fivefold and 10-fold to the mRNA-1273 vaccine from peak over 6 months after the 2nd dose, ending with low but detectable levels of nAbs in 100% of subjects.²⁰⁰⁻²⁰² Ninefold to 10-fold nAb declines were also observed for a low dose (25 µg) of the mRNA-1273 mRNA COVID vaccine (instead of 100 µg),⁷⁶ comparable to the BNT162b2 dose (30 µg), indicating that the durability of the Ab responses to 2-dose mRNA vaccines is consistent, and the kinetics are not determined by the vaccine dose, though the absolute magnitude of the Ab response is higher with a higher vaccine dose.

Long-lived plasma cells specific for SARS-CoV-2 spike are detectable in a majority of individuals 6 months after 2-dose mRNA vaccination¹⁷⁹; however, given that nAb and RBD IgG titers continue to decline for at least 8 months after 2-dose RNA vaccination, these long-lived BPCs apparently represent low frequencies, or do not have the durability observed for B_{PC}s generated to other antigen exposures.

Due to the precipitous drop in nAb titers over 6–8 months after two doses, and the emergence of VOCs Delta and Omicron, 3-dose mRNA vaccine regimens have been implemented as the norm in many countries (i.e., 2-dose regimen plus a “booster” at approximately 6 months) (Figure 3). A critical question about 3-dose mRNA

regimens is whether they induce more durable Ab responses than 2-dose regimens. Given that the 2-dose mRNA vaccines were immunogenic and elicited substantial memory CD8 T cells, memory CD4 T cells, B_{Mem} cells, and at least a few long-lived PCs, it was reasonable to predict that 3-dose mRNA vaccine regimens would induce substantially more durable Abs than the 2-dose regimen. Results from previously infected individuals cleared demonstrated that the human immune system is capable of making durable Ab responses to SARS-CoV-2, and hybrid immunity also demonstrated that the human immune system is clearly capable of making high nAb titers to SARS-CoV-2. Additionally, many vaccines are three dose regimens, with durable Abs only being developed after the third dose. Teleologically, one can consider that this is because the immune system performs a cost-benefit analysis of durable memory to each antigen exposure. Durable Ab responses for 10 years or more have a high caloric resource cost commitment, whereas durable B_{Mem} cells or T cells have significantly lower caloric costs. As such, it frequently takes multiple antigen exposures to trigger significant durable Ab responses. Nevertheless, it was unclear if mRNA vaccines were capable of triggering durable Ab responses.

Acute Ab responses to a 3rd dose of mRNA vaccine were strong, with peak nAb titers above that of 2-dose immunization.^{203,204} Two studies found much more durable nAb titers at 4 or 6 months after a 3rd dose of mRNA vaccine compared to 2-doses.^{202,205} NAb titers against Ac SARS-CoV-2 only declined 1.6-fold for the BNT162b2 vaccine at 4 months and 2.3-fold for the mRNA-1273 vaccine at 6 months after a 3rd dose.^{202,205} Those findings indicate robust long-lived Ab production after 3 doses (Figure 2). However, not all results agree. In the study of mRNA-1273 vaccinees, there was a substantial discordance between the durability of nAbs against Ac SARS-CoV-2 or Omicron, with nAbs against Ac SARS-CoV-2 only declining 2.3-fold after 6 months, but nAbs against Omicron declining 6.3-fold.²⁰² In a third study, of an Israeli population receiving the BNT162b2 vaccine, Ac SARS-CoV-2 nAbs declined 5.5-fold over approximately 4 months after 3 doses.²⁰⁶ Thus, conclusions about durability of Abs after 3-dose mRNA vaccination remain uncertain.

Adenoviral vector COVID-19 vaccines ChAdOx1 (2-dose) and Ad26.COVS.2 (1-dose) initially elicit substantially lower Ab responses than mRNA vaccines. Spike or RBD IgG titers after 1-dose Ad26.COVS.2 are approximately 70-fold to 355-fold lower than 2-dose mRNA vaccines.^{79,82,207} NAb titers are approximately 10- to 70-fold lower 30 to 60 days after Ad26.COVS.2 compared to mRNA vaccines.^{79,82,92} NAb titers increase some over time after Ad26.COVS.2 in some individuals, but this is variable^{79,82,86,208} (Figure 2). NAb titers are at least stably maintained in most individuals who receive Ad26.COVS.2 vaccination, such that 6–8 months after Ad26.COVS.2 vaccination nAb titers are detectable in almost all individuals, albeit at low levels,^{79,86} though not all studies agree, with some cohorts reporting many negative individuals (table 40 ref. ²⁰⁸) and one study reporting higher levels.⁹¹ ChAdOx1 vaccine (2-dose) generates early nAb titers that are also substantially lower than mRNA vaccine generated nAb titers. nAb titers are 8.3-fold lower after ChAdOx1 compared to BNT162b2 one month after immunization.⁹³

Spike IgG titers after 2-dose ChAdOx1 decline with a $t_{1/2}$ comparable to that of 2-dose BNT162b2,¹⁹ with no evidence of greater durability. However, in one study, the difference between ChAdOx1 and BNT162b2 nAb titers was only approximately twofold at 3 months after the 2nd dose.⁹⁴ There are limited data for 2-dose ChAdOx1 at longer time points with head-to-head comparisons, but when comparing ChAdOx1/ChAdOx1 to ChAdOx1/ BNT162b2, spike Ab titers at 6 months were sixfold lower in the ChAdOx1/ChAdOx1 group.

Protein-based COVID-19 vaccines are in two categories, recombinant spike vaccines, such as NVX-CoV2373, and inactivated virus vaccines such as BBV152 and Sinovac. Minimal data on Ab durability are available for inactivated vaccines. NVX-CoV2373 generates significant nAb titers after two doses.^{209,210} NAb titers 6 months after the 2nd dose are approximately equivalent between individuals receiving NVX-CoV2373, mRNA-1273, or BNT162b2 vaccines (Figure 2), suggesting that high nAb titers may wane after 2-dose NVX-CoV2373.⁷⁹

Mix & match vaccine strategies could potentially elicit more durable Ab responses. Limited data are available on immune memory after mix & match vaccination.⁹²⁻⁹⁴ Given that peak nAb titers are higher with 2-dose mRNA or recombinant protein vaccination compared to adenoviral vaccines, but that adenoviral vaccines might elicit more stable Ab responses at 6 months, it is plausible that mix & match approaches may combine the best of both and result in higher level durable nAb titers. This may also be relevant for boosters.⁹⁴

Regarding local Abs in the respiratory tract and oral mucosa, the mRNA vaccines do elicit some circulating IgA.^{100,173} IgA and IgG are detectable in saliva and nasal swabs in a fraction of vaccinated individuals, but both the IgA and IgG decline substantially over the course of 6 months after 2-dose mRNA vaccination, mirroring the declines in circulating IgG and IgA.^{100,211,212} In immunized non-human primates, RBD IgG and IgA in bronchoalveolar lavage and nasal swabs directly correlated with circulating IgG and IgA levels.²¹³ Exploring development of intranasal vaccines is of substantial interest for potentially improving mucosal immune memory.²¹⁴

5.3 | Antibody durability in hybrid immunity

The most prominent characteristic of hybrid immunity is the impressive improvement in nAb titers and the breadth of neutralization of SARS-CoV-2 variants. In some individuals, SARS-CoV-2 nAb titers increase 100-fold after a single mRNA vaccination. Equally impressive, the nAbs are not only able to neutralize every known SARS-CoV-2 variant, including Omicron, they are also able to neutralize a different viral species, SARS-CoV. B_{Mem} cells and memory CD4 T cells are at the root of these impressive outcomes. While circulating nAb titers are frequently low in previously infected individuals, without much evidence of breadth, some B_{Mem} cells from those same individuals encode Abs with impressive potency and breadth.^{162,165,215,216} Those B_{Mem} cells are then recalled after vaccination to generate an anamnestic Ab response, now composed of

Abs capable of neutralizing breadth against VOCs such as Omicron, and even neutralization of SARS-CoV,^{20,80,157,162,217,218} irrespective of original COVID-19 severity.²¹⁹

Hybrid immunity can also occur in the reverse order—vaccination and then infection—with similarly high titer and broad nAb responses, irrespective of whether the infection was Alpha, Delta, or Omicron, and irrespective of disease severity.^{20,21,212,220,221,222} These responses are again derived from B_{Mem} cells, in this case B_{Mem} cells generated in response to vaccination.^{173,223}

Ab durability at 6 months is robust in a majority of individuals with hybrid immunity, as measured by nAb titers (Figure 2). NAb titers were stable in a majority of hybrid immunity individuals, declined less than twofold over 6 months.^{20,80} Of note, RBD-binding titers exhibited larger declines, for unclear reasons.^{80,100} After 6 months, people with hybrid immunity maintained fivefold to 17-fold higher nAb titers against ancestral SARS-CoV-2, Beta, or Delta compared to individuals who were 2-dose mRNA vaccinated^{20,80}; compared to individuals who were previously infected alone, people with hybrid immunity maintained 10- to 51-fold higher nAb titers against ancestral SARS-CoV-2, Beta, Delta, or Omicron.²⁰

Higher nasal RBD IgG and IgA are found in individuals with hybrid immunity (either inf+vax or vax+inf) when sampled up to 10 months after vaccination.^{100,212}

6 | INTERRELATIONSHIPS BETWEEN IMMUNE MEMORY COMPARTMENTS

Studies of SARS-CoV-2 memory are the first time that large datasets have been collected of multiple antigen-specific memory cell compartments over a period of 6+ months after an acute infection. This provides key opportunities to understand relationships between different aspects of immune memory. It was observed that each compartment of immune memory after infection exhibit distinct kinetics over time, and different quantitative relationships to the other compartments of immune memory.³ Some of the relationships changed dramatically over time.³ Perhaps most importantly from a practical perspective, serum RBD IgG titers were not quantitatively predictive of the other components of immune memory, notably memory T cells.³ Nevertheless, other relationships were observed.³ T_{FH} cells, B_{Mem} cells, and circulating Ab titers are functionally associated.¹⁰⁴ However, cT_{FH} cell frequencies after infection were not quantitatively predictive of germinal centers,¹⁵³ or nAb titers,¹¹⁰ suggesting more complex relationships between circulating T cell memory, germinal centers, and nAbs.

For mRNA vaccines, relationships between nAbs and memory CD4 T cells are clear, and early T_{FH} cell responses do correlate with subsequent nAb titers.⁷⁹ However, at any given memory timepoint, no clear association is observed between serum Ab titers and memory CD4 T cell and memory CD8 T cell frequencies.⁷⁹ CD4 T cells provide help for CD8 T cell differentiation and memory CD8 T cells in multiple contexts.²²⁴ Nevertheless, memory CD4 T cell frequencies and memory CD8 T cell frequencies do not show a strong

relationship in mRNA-vaccinated individuals.⁷⁹ Overall, interrelationships between immune memory compartments exist, but much remains to be learned.

7 | IMMUNE MEMORY IN SPECIAL POPULATIONS

7.1 | Immune memory in the immunocompromised or suppressed

Immune responses to COVID-19 vaccines in immunocompromised or immunosuppressed individuals vary depending on the specific immunocompromised or immunosuppressed condition. B cell depleted individuals (i.e., anti-CD20 mAb treatment) have defective Ab, B_{Mem} cell, and T_{FH} cell responses to mRNA COVID-19 vaccines, but their T_{H1} and CD8 T cell responses are normal or elevated.²²⁵ Thus, it is expected that immune memory will be substantially defect in those individuals for durable Abs, B_{Mem} cells, and memory T_{FH} cells; however, memory T_{H1} cells and CD8 T cells may or may not be compromised. Solid organ transplant patients frequently have reduced responses to COVID-19 vaccines because of their immunosuppressive drug therapies. Immune memory in such individuals is not well understood, but based on the severity of the germinal center, T_{FH} cell, T_{H1} cell, and CD8 T cell response defects in kidney transplant individuals responding to COVID-19 mRNA vaccines,¹⁵² it is likely that there are severe immune memory defects in those patients. Certain categories of cancer patients on similar immunosuppressive drug regimens are likely to also have immune memory defects. Fingolimod, the S1P receptor antagonist, appears to cause an almost complete block of Ab and T cell responses to COVID-19 vaccines,²²⁶ and would be expected to result in severe immune memory defects to COVID-19 vaccines. More information is needed about immune memory to COVID-19 vaccines in a diverse range of immunocompromised or immunosuppressed individuals, given that the efficacy of the vaccines is predicated on immune memory.

7.2 | Boosters in persons with hybrid immunity

Many papers show substantial immunological and epidemiological evidence that hybrid immunity is the most robust immunity against COVID-19.^{2,9} This includes vax+vax+inf,^{20,212,227} inf+vax, and inf+vax+vax. These individuals have the best neutralizing Ab breadth—able to recognize every single known variant, include Omicron, and even able to recognize another species of virus (SARS)^{20,157,228}—and they also have substantially better local immunity in the nose and mouth,^{62,212} which is not generated by intramuscular vaccination. They also have more durable Ab responses, based on the available data.^{3,4,100,186,200,229,230} However, many governments have booster vaccination requirements within 90 days of an infection. For people who had breakthrough infections with Delta or Omicron after being double vaccinated, this is most likely to be far

sooner than needed, and may be counterproductive. It is plausible that such individuals may have such good immune memory that they do not need a booster for years. The quality of the Ab response needs time to develop. The immune system has done an amazing job making Ab responses and memory B cells against SARS-CoV-2 that are educated guesses about potential future variants.^{2,20,162,165,215,229,231,232} That is important for immunity against this virus, but takes time in germinal centers,²³³ and it is likely disrupted by a new immunization. Hence, immunizations too close together are shortsighted and result in poorer quality immunity. We also know that memory B cell frequencies increase for almost 6 months after infection,^{3,215,229} or after vaccination.^{162,229} We know that germinal centers can persist for more than 6 months after SARS-CoV-2 infection.⁶² We know that germinal centers can persist and be productive for more than 6 months after two doses of COVID-19 mRNA vaccines.^{180,234,235} In addition, we know that the quality of neutralizing Abs can improve over 3–6 months,^{86,162,173,215,236} reflective of outcomes of these long processes of developing higher quality immune memory. Boosters too close together may disrupt those processes of generating broader protection against future variants.

8 | CONCLUDING REMARKS

A remarkable amount has been learned about immune memory after SARS-CoV-2 infection. A remarkable amount has also been learned about a multitude of COVID-19 vaccines, and hybrid immunity. Increasing our understanding of the deterministic relationships between early immune responses and immune memory outcomes remain a major knowledge gap for further research. There is much to be learned about local immune memory in mucosal tissues such as nasal passages, oral cavity, the URT broadly, the intestinal tract, and lungs. Investigation of the relationships between local immune memory and systemic immune memory is of particular significance. Longer term durability of each compartment of immune memory after SARS-CoV-2 infection or COVID-19 vaccination of course remains to be empirically determined. However, the wealth of scientific literature already accumulated regarding immune memory provides strong predictions regarding the durability of T cell memory, B cell memory, and long-lasting antibody responses that can be extrapolated for several years, if not decades, and may provide determining factors of sustained protection against disease. Lastly, clearly this knowledge and experience can also be leveraged toward vaccines against other diseases that affect humanity now, and prevent future plagues.

ACKNOWLEDGEMENTS

Thanks to Zeli Zhang and Parham Ramezani-Rad for helpful input. Thanks to graphical assistance by Christina Corbaci. This work was funded by the NIH NIAID under award AI142742 (Cooperative Centers for Human Immunology, CCHI) and CHAVD UM1 AI144462, and was additionally supported in part by La Jolla Institute for Immunology Institutional Funds and an anonymous donor.

CONFLICT OF INTEREST

SC has consulted for GSK, JP Morgan, Citi, Morgan Stanley, Avalia NZ, Nutcracker Therapeutics, University of California, California State Universities, United Airlines, and Roche. A.S. is a consultant for Gritstone Bio, Flow Pharma, ImmunoScape, Moderna, AstraZeneca, Avalia, Fortress, Repertoire, Gilead, Gerson Lehrman Group, RiverVest, MedaCorp, and Guggenheim.

DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

ORCID

Shane Crotty  <https://orcid.org/0000-0002-6484-6262>

REFERENCES

- Schoenberger SP, Crotty S. Immunological memory. In: Paul WE, ed. *Immunological Memory*. 6th ed. Lippincott Williams & Wilkins; 2010.
- Crotty S. Hybrid immunity. *Science*. 2021;372(6549):1392-1393. doi:10.1126/science.abj2258
- Dan JM, Mateus J, Kato Y, et al. Immunological memory to SARS-CoV-2 assessed for up to 8 months after infection. *Science*. 2021;371(6529):eabf4063. doi:10.1126/science.abf4063
- Cohen KW, Linderman SL, Moodie Z, et al. Longitudinal analysis shows durable and broad immune memory after SARS-CoV-2 infection with persisting antibody responses and memory B and T cells. *Cell Rep Med*. 2021;2(7):100354. doi:10.1016/j.xcrm.2021.100354
- Lumley SF, O'Donnell D, Stoesser NE, et al. Antibody status and incidence of SARS-CoV-2 infection in health care workers. *N Engl J Med*. 2020;384(6):533-540. doi:10.1056/nejmoa2034545
- Abu-Raddad LJ, Chemaitelly H, Coyle P, et al. SARS-CoV-2 antibody-positivity protects against reinfection for at least seven months with 95% efficacy. *EClinicalMedicine*. 2021;35:100861. doi:10.1016/j.eclinm.2021.100861
- Hall VJ, Foulkes S, Charlett A, et al. SARS-CoV-2 infection rates of antibody-positive compared with antibody-negative health-care workers in England: a large, multicentre, prospective cohort study (SIREN). *Lancet*. 2021;397(10283):1459-1469. doi:10.1016/s0140-6736(21)00675-9
- Leidi A, Koegler F, Dumont R, et al. Risk of reinfection after seroconversion to SARS-CoV-2: a population-based propensity-score matched cohort study. *Clin Infect Dis*. 2021;74:622-629. doi:10.1093/cid/ciab495
- Pouwels KB, Pritchard E, Matthews PC, et al. Effect of Delta variant on viral burden and vaccine effectiveness against new SARS-CoV-2 infections in the UK. *Nat Med*. 2021;27:2127-2135. doi:10.1038/s41591-021-01548-7
- Altarawneh HN, Chemaitelly H, Hasan MR, et al. Protection against the omicron variant from previous SARS-CoV-2 infection. *N Engl J Med*. 2022;386:1288-1290. doi:10.1056/nejmc2200133
- León TM, Dorabawila V, Nelson L, et al. COVID-19 cases and hospitalizations by COVID-19 vaccination status and previous COVID-19 diagnosis – California and New York, May–November 2021. *MMWR Morb Mortal Wkly Rep*. 2022;71(4):125-131. doi:10.15585/mmwr.mm7104e1
- Nyberg T, Ferguson NM, Nash SG, et al. Comparative analysis of the risks of hospitalisation and death associated with SARS-CoV-2 omicron (B.1.1.529) and delta (B.1.617.2) variants in England: a cohort study. *Lancet Lond Engl*. 2022;399:1303-1312.
- Lewnard JA, Hong VX, Patel MM, Kahn R, Lipsitch M, Tartof SY. Clinical outcomes among patients infected with Omicron (B.1.1.529) SARS-CoV-2 variant in southern California. *medRxiv*. 2022;2022.01.11.22269045. doi:10.1101/2022.01.11.22269045
- Madhi SA, Kwatra G, Myers JE, et al. Population immunity and Covid-19 severity with omicron variant in South Africa. *N Engl J Med*. 2022;386:1314-1326. doi:10.1056/nejmoa2119658
- Polack FP, Thomas SJ, Kitchin N, et al. Safety and efficacy of the BNT162b2 mRNA Covid-19 vaccine. *New Engl J Med*. 2020;383(27):2603-2615. doi:10.1056/nejmoa2034577
- Sahly HME, Baden LR, Essink B, et al. Efficacy of the mRNA-1273 SARS-CoV-2 vaccine at completion of blinded phase. *N Engl J Med*. 2021;385(19):1774-1785. doi:10.1056/nejmoa2113017
- Tartof SY, Slezak JM, Fischer H, et al. Effectiveness of mRNA BNT162b2 COVID-19 vaccine up to 6 months in a large integrated health system in the USA: a retrospective cohort study. *Lancet*. 2021;398(10309):1407-1416. doi:10.1016/s0140-6736(21)02183-8
- Goldberg Y, Mandel M, Bar-On YM, et al. Waning immunity after the BNT162b2 vaccine in Israel. *N Engl J Med*. 2021;385(24):NEJMoa2114228. doi:10.1056/nejmoa2114228
- Wei J, Pouwels KB, Stoesser N, et al. Antibody responses and correlates of protection in the general population after two doses of the ChAdOx1 or BNT162b2 vaccines. *Nat Med*. 2022;28:1072-1082. doi:10.1038/s41591-022-01721-6
- Walls AC, Sprouse KR, Bowen JE, et al. SARS-CoV-2 breakthrough infections elicit potent, broad and durable neutralizing antibody responses. *Cell*. 2022;185(5):872-880.e3. doi:10.1016/j.cell.2022.01.011
- Wratis PR, Stern M, Priller A, et al. Three exposures to the spike protein of SARS-CoV-2 by either infection or vaccination elicit superior neutralizing immunity to all variants of concern. *Nat Med*. 2022;28(3):496-503. doi:10.1038/s41591-022-01715-4
- Sette A, Crotty S. Adaptive immunity to SARS-CoV-2 and COVID-19. *Cell*. 2021;184(4):861-880. doi:10.1016/j.cell.2021.01.007
- Sadarangani M, Marchant A, Kollmann TR. Immunological mechanisms of vaccine-induced protection against COVID-19 in humans. *Nat Rev Immunol*. 2021;21(8):475-484. doi:10.1038/s41577-021-00578-z
- Goldblatt D, Alter G, Crotty S, Plotkin S. Correlates of protection against SARS-CoV-2 infection and COVID-19 disease. *Immunological Reviews*. 2022. (This volume).
- Chang JT, Wherry EJ, Goldrath AW. Molecular regulation of effector and memory T cell differentiation. *Nat Immunol*. 2014;15(12):1104-1115. doi:10.1038/ni.3031
- Moderbacher CR, Ramirez SI, Dan JM, et al. Antigen-specific adaptive immunity to SARS-CoV-2 in acute COVID-19 and associations with age and disease severity. *Cell*. 2020;183(4):996-1012.e19. doi:10.1016/j.cell.2020.09.038
- Peng Y, Felce SL, Dong D, et al. An immunodominant NP105-113-B*07:02 cytotoxic T cell response controls viral replication and is associated with less severe COVID-19 disease. *Nat Immunol*. 2022;23(1):50-61. doi:10.1038/s41590-021-01084-z
- Sekine T, Perez-Potti A, Rivera-Ballesteros O, et al. Robust T cell immunity in convalescent individuals with asymptomatic or mild COVID-19. *Cell*. 2020;183(1):158-168.e14. doi:10.1016/j.cell.2020.08.017
- Peng Y, Mentzer AJ, Liu G, et al. Broad and strong memory CD4+ and CD8+ T cells induced by SARS-CoV-2 in UK convalescent individuals following COVID-19. *Nat Immunol*. 2020;21(11):1336-1345. doi:10.1038/s41590-020-0782-6
- Thieme CJ, Anft M, Paniskaki K, et al. Robust T cell response toward spike, membrane, and nucleocapsid SARS-CoV-2 proteins is not associated with recovery in critical COVID-19 patients. *Cell Rep Med*. 2020;1(6):100092. doi:10.1016/j.xcrm.2020.100092

31. McMahan K, Yu J, Mercado NB, et al. Correlates of protection against SARS-CoV-2 in rhesus macaques. *Nature*. 2020;590:630-634. doi:10.1038/s41586-020-03041-6
32. Ishii H, Nomura T, Yamamoto H, et al. Neutralizing antibody-independent SARS-CoV-2 control correlated with intranasal vaccine-induced CD8+ T-cell responses. *Cell Rep Med*. 2022;3(2):100520. doi:10.1016/j.xcrm.2022.100520
33. Kedzierska K, Thomas PG. Count on us: T cells in SARS-CoV-2 infection and vaccination. *Cell Rep Med*. 2022;3(3):100562-100562. doi:10.1016/j.xcrm.2022.100562
34. Sidney J, Peters B, Sette A. Epitope prediction and identification- adaptive T cell responses in humans. *Semin Immunol*. 2020;50:101418. doi:10.1016/j.smim.2020.101418
35. Reiss S, Baxter AE, Cirelli KM, et al. Comparative analysis of activation induced marker (AIM) assays for sensitive identification of antigen-specific CD4 T cells. *PLoS One*. 2017;12(10):e0186998. doi:10.1371/journal.pone.0186998
36. Akondy RS, Fitch M, Edupuganti S, et al. Origin and differentiation of human memory CD8 T cells after vaccination. *Nature*. 2017;552(7685):362-367. doi:10.1038/nature24633
37. Bert NL, Tan AT, Kunasegaran K, et al. SARS-CoV-2-specific T cell immunity in cases of COVID-19 and SARS, and uninfected controls. *Nature*. 2020;584(7821):457-462. doi:10.1038/s41586-020-2550-z
38. Grifoni A, Weiskopf D, Ramirez SI, et al. Targets of T cell responses to SARS-CoV-2 coronavirus in humans with COVID-19 disease and unexposed individuals. *Cell*. 2020;181(7):1489-1501.e15. doi:10.1016/j.cell.2020.05.015
39. Zuo J, Dowell AC, Pearce H, et al. Robust SARS-CoV-2-specific T-cell immunity is maintained at 6 months following primary infection. *Nat Immunol*. 2021;22(5):620-626. doi:10.1038/s41590-021-00902-8
40. Rodda LB, Netland J, Shehata L, et al. Functional SARS-CoV-2-specific immune memory persists after mild COVID-19. *Cell*. 2020;184:169-183.e17. doi:10.1016/j.cell.2020.11.029
41. Bilich T, Nelde A, Heitmann JS, et al. T cell and antibody kinetics delineate SARS-CoV-2 peptides mediating long-term immune responses in COVID-19 convalescent individuals. *Sci Transl Med*. 2021;13(590):eabf7517. doi:10.1126/scitranslmed.abf7517
42. Bergamaschi L, Mescia F, Turner L, et al. Longitudinal analysis reveals that delayed bystander CD8+ T cell activation and early immune pathology distinguish severe COVID-19 from mild disease. *Immunity*. 2021;54(6):1257-1275.e8. doi:10.1016/j.immuni.2021.05.010
43. Sahin U, Muik A, Vogler I, et al. BNT162b2 vaccine induces neutralizing antibodies and poly-specific T cells in humans. *Nature*. 2021;595:572-577. doi:10.1038/s41586-021-03653-6
44. Oberhardt V, Luxemburger H, Kemming J, et al. Rapid and stable mobilization of CD8+ T cells by SARS-CoV-2 mRNA vaccine. *Nature*. 2021;597(7875):268-273. doi:10.1038/s41586-021-03841-4
45. Gangaev A, Ketelaars SLC, Isaeva OI, et al. Identification and characterization of a SARS-CoV-2 specific CD8+ T cell response with immunodominant features. *Nat Commun*. 2021;12(1):2593. doi:10.1038/s41467-021-22811-y
46. Rha MS, Jeong HW, Ko JH, et al. PD-1-expressing SARS-CoV-2-specific CD8+ T cells are not exhausted, but functional in patients with COVID-19. *Immunity*. 2020;54(1):44-52.e3. doi:10.1016/j.immuni.2020.12.002
47. Nguyen THO, Rowntree LC, Petersen J, et al. CD8+ T cells specific for an immunodominant SARS-CoV-2 nucleocapsid epitope display high naive precursor frequency and TCR promiscuity. *Immunity*. 2021;54(5):1066-1082.e5. doi:10.1016/j.immuni.2021.04.009
48. Files JK, Boppana S, Perez MD, et al. Sustained cellular immune dysregulation in individuals recovering from SARS-CoV-2 infection. *J Clin Invest*. 2020;131(1):e140491. doi:10.1172/jci140491
49. Kusnadi A, Ramirez-Suástegui C, Fajardo V, et al. Severely ill COVID-19 patients display impaired exhaustion features in SARS-CoV-2-reactive CD8+ T cells. *Sci Immunol*. 2021;6(55):eabe4782. doi:10.1126/sciimmunol.abe4782
50. Tarke A, Sidney J, Kidd CK, et al. Comprehensive analysis of T cell immunodominance and immunoprevalence of SARS-CoV-2 epitopes in COVID-19 cases. *Cell Rep Med*. 2021;2(2):100204. doi:10.1016/j.xcrm.2021.100204
51. Tarke A, Sidney J, Methot N, et al. Impact of SARS-CoV-2 variants on the total CD4+ and CD8+ T cell reactivity in infected or vaccinated individuals. *Cell Rep Med*. 2021;2(7):100355. doi:10.1016/j.xcrm.2021.100355
52. Woldemeskel BA, Garliss CC, Blankson JN. SARS-CoV-2 mRNA vaccines induce broad CD4+ T cell responses that recognize SARS-CoV-2 variants and HCoV-NL63. *J Clin Invest*. 2021;131(10):e149335. doi:10.1172/jci149335
53. Redd AD, Nardin A, Kared H, et al. CD8+ T-cell responses in COVID-19 convalescent individuals target conserved epitopes from multiple prominent SARS-CoV-2 circulating variants. *Open Forum Infect Dis*. 2021;8(7):ofab143. doi:10.1093/ofid/ofab143
54. Grifoni A, Sidney J, Vita R, et al. SARS-CoV-2 human T cell epitopes: adaptive immune response against COVID-19. *Cell Host Microbe*. 2021;29(7):1076-1092. doi:10.1016/j.chom.2021.05.010
55. Quadeer AA, Ahmed SF, McKay MR. Landscape of epitopes targeted by T cells in 852 individuals recovered from COVID-19: Meta-analysis, immunoprevalence, and web platform. *Cell Rep Med*. 2021;2(6):100312. doi:10.1016/j.xcrm.2021.100312
56. Law JC, Girard M, Chao GYC, et al. Persistence of T cell and antibody responses to SARS-CoV-2 up to 9 months after symptom onset. *J Immunol*. 2022;208(2):429-443. doi:10.4049/jimmunol.2100727
57. Garg S, Patel K, Pham H, et al. Clinical trends among U.S. adults hospitalized with COVID-19, March to December 2020. *Ann Intern Med*. 2021;174(10):1409-1419. doi:10.7326/m21-1991
58. Lauer SA, Grantz KH, Bi Q, et al. The incubation period of coronavirus disease 2019 (COVID-19) from publicly reported confirmed cases: estimation and application. *Ann Intern Med*. 2020;172(9):577-582. doi:10.7326/m20-0504
59. Killingley B, Mann A, Kalinova M, et al. Safety, tolerability and viral kinetics during SARS-CoV-2 human challenge in young adults. *Nat Med*. 2022;28:1031-1041.
60. He X, Lau EHY, Wu P, et al. Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nat Med*. 2020;26(5):672-675. doi:10.1038/s41591-020-0869-5
61. Steinert EM, Schenkel JM, Fraser KA, et al. Quantifying memory CD8 T cells reveals regionalization of immunosurveillance. *Cell*. 2015;161(4):737-749. doi:10.1016/j.cell.2015.03.031
62. Poon MML, Rybkina K, Kato Y, et al. SARS-CoV-2 infection generates tissue-localized immunological memory in humans. *Sci Immunol*. 2021;6(65):eab19105. doi:10.1126/sciimmunol.ab19105
63. Cheon I, Li C, Son YM, et al. Immune signatures underlying post-acute COVID-19 lung sequelae. *Sci Immunol*. 2021;6:eabk1741. doi:10.1126/sciimmunol.abk1741
64. Roukens AHE, Pothast CR, König M, et al. Prolonged activation of nasal immune cell populations and development of tissue-resident SARS-CoV-2-specific CD8+ T cell responses following COVID-19. *Nat Immunol*. 2022;23(1):23-32. doi:10.1038/s41590-021-01095-w
65. Chou J, Thomas PG, Randolph AG. Immunology of SARS-CoV-2 infection in children. *Nat Immunol*. 2022;23(2):177-185. doi:10.1038/s41590-021-01123-9
66. Dowell AC, Butler MS, Jinks E, et al. Children develop robust and sustained cross-reactive spike-specific immune responses to SARS-CoV-2 infection. *Nat Immunol*. 2022;23(1):40-49. doi:10.1038/s41590-021-01089-8

67. Cohen CA, Li APY, Hachim A, et al. SARS-CoV-2 specific T cell responses are lower in children and increase with age and time after infection. *Nat Commun.* 2021;12(1):4678. doi:10.1038/s41467-021-24938-4
68. Consiglio CR, Cotugno N, Sardh F, et al. The immunology of multisystem inflammatory syndrome in children with COVID-19. *Cell.* 2020;183(4):968-981.e7. doi:10.1016/j.cell.2020.09.016
69. Ramaswamy A, Brodsky NN, Sumida TS, et al. Immune dysregulation and autoreactivity correlate with disease severity in SARS-CoV-2-associated multisystem inflammatory syndrome in children. *Immunity.* 2021;54(5):1083-1095.e7. doi:10.1016/j.immuni.2021.04.003
70. Hsieh L, Grifoni A, Sidney J, et al. Characterization of SARS-CoV-2 and common cold coronavirus-specific T cell responses in MIS-C and Kawasaki disease children. *Eur J Immunol.* 2021;52:123-137. doi:10.1002/eji.202149556
71. Singh V, Obregon-Perko V, Lapp SA, et al. Limited induction of SARS-CoV-2-specific T cell responses in children with multisystem inflammatory syndrome compared with COVID-19. *JCI Insight.* 2022;7(4):e155145. doi:10.1172/jci.insight.155145
72. Conway SR, Lazarski CA, Field NE, et al. SARS-CoV-2-specific T cell responses are stronger in children with multisystem inflammatory syndrome compared to children with uncomplicated SARS-CoV-2 infection. *Front Immunol.* 2022;12:793197. doi:10.3389/fimmu.2021.793197
73. Sacco K, Castagnoli R, Vakkilainen S, et al. Immunopathological signatures in multisystem inflammatory syndrome in children and pediatric COVID-19. *Nat Med.* 2022;28:1050-1062. doi:10.1038/s41591-022-01724-3
74. Merad M, Blish CA, Sallusto F, Iwasaki A. The immunology and immunopathology of COVID-19. *Science.* 2022;375(6585):1122-1127. doi:10.1126/science.abm8108
75. Niessl J, Sekine T, Lange J, et al. Identification of resident memory CD8+ T cells with functional specificity for SARS-CoV-2 in unexposed oropharyngeal lymphoid tissue. *Sci Immunol.* 2021;6(64):eabk0894. doi:10.1126/sciimmunol.abk0894
76. Mateus J, Dan JM, Zhang Z, et al. Low-dose mRNA-1273 COVID-19 vaccine generates durable memory enhanced by cross-reactive T cells. *Science.* 2021;374(6566):eabj9853. doi:10.1126/science.abj9853
77. Tarke A, Coelho CH, Zhang Z, et al. SARS-CoV-2 vaccination induces immunological T cell memory able to cross-recognize variants from Alpha to Omicron. *Cell.* 2022;185(5):847-859.e11. doi:10.1016/j.cell.2022.01.015
78. Guerrero G, Picozza M, D'Orso S, et al. BNT162b2 vaccination induces durable SARS-CoV-2 specific T cells with a stem cell memory phenotype. *Sci Immunol.* 2021;6:eabl5344. doi:10.1126/sciimmunol.abl5344
79. Zhang Z, Mateus J, Coelho CH, et al. Humoral and cellular immune memory to four COVID-19 vaccines. *bioRxiv.* 2022;2022.03.18.484953. doi:10.1101/2022.03.18.484953
80. Goel RR, Painter MM, Apostolidis SA, et al. mRNA vaccines induce durable immune memory to SARS-CoV-2 and variants of concern. *Science.* 2021;374(6572):abm0829. doi:10.1126/science.abm0829
81. Naranbhai V, Nathan A, Kaseke C, et al. T-cell reactivity to the SARS-CoV-2 Omicron variant is preserved in most but not all individuals. *Cell.* 2022;185:1041-1051.e6. doi:10.1016/j.cell.2022.01.029
82. Naranbhai V, Garcia-Beltran WF, Chang CC, et al. Comparative immunogenicity and effectiveness of mRNA-1273, BNT162b2 and Ad26.COV2.S COVID-19 vaccines. *J Infect Dis.* 2021;225(7):1141-1150. doi:10.1093/infdis/jiab593
83. Jackson LA, Anderson EJ, Roupael NG, et al. An mRNA vaccine against SARS-CoV-2 — preliminary report. *N Engl J Med.* 2020;383(20):1920-1931. doi:10.1056/nejmoa2022483
84. Corbett KS, Flynn B, Foulds KE, et al. Evaluation of the mRNA-1273 Vaccine against SARS-CoV-2 in nonhuman primates. *N Engl J Med.* 2020;383(16):1544-1555. doi:10.1056/nejmoa2024671
85. Sadoff J, Gars ML, Shukarev G, et al. Interim results of a phase 1-2a trial of Ad26.COV2.S Covid-19 vaccine. *N Engl J Med.* 2021;384:1824-1835. doi:10.1056/nejmoa2034201
86. Barouch DH, Stephenson KE, Sadoff J, et al. Durable humoral and cellular immune responses 8 months after Ad26.COV2.S vaccination. *N Engl J Med.* 2021;385:951-953. doi:10.1056/nejmc2108829
87. Keeton R, Tincho MB, Ngomti A, et al. T cell responses to SARS-CoV-2 spike cross-recognize Omicron. *Nature.* 2022;603:488-492. doi:10.1038/s41586-022-04460-3
88. Schmidt T, Klemis V, Schub D, et al. Immunogenicity and reactogenicity of heterologous ChAdOx1 nCoV-19/mRNA vaccination. *Nat Med.* 2021;27(9):1530-1535. doi:10.1038/s41591-021-01464-w
89. Pozzetto B, Legros V, Djebali S, et al. Immunogenicity and efficacy of heterologous ChAdOx1-BNT162b2 vaccination. *Nature.* 2021;600(7890):701-706. doi:10.1038/s41586-021-04120-y
90. Schmidt T, Klemis V, Schub D, et al. Cellular immunity predominates over humoral immunity after homologous and heterologous mRNA and vector-based COVID-19 vaccine regimens in solid organ transplant recipients. *Am J Transplant.* 2021;21(12):3990-4002. doi:10.1111/ajt.16818
91. Collier ARY, Yu J, McMahan K, et al. Differential kinetics of immune responses elicited by Covid-19 vaccines. *N Engl J Med.* 2021;385(21):2010-2012. doi:10.1056/nejmc2115596
92. Atmar RL, Lyke KE, Deming ME, et al. Homologous and heterologous Covid-19 booster vaccinations. *New Engl J Med.* 2022;386(11):1046-1057. doi:10.1056/nejmoa2116414
93. Liu X, Shaw RH, Stuart ASV, et al. Safety and immunogenicity of heterologous versus homologous prime-boost schedules with an adenoviral vectored and mRNA COVID-19 vaccine (Com-COV): a single-blind, randomised, non-inferiority trial. *Lancet.* 2021;398(10303):856-869. doi:10.1016/s0140-6736(21)01694-9
94. Munro APS, Janani L, Cornelius V, et al. Safety and immunogenicity of seven COVID-19 vaccines as a third dose (booster) following two doses of ChAdOx1 nCoV-19 or BNT162b2 in the UK (COV-BOOST): a blinded, multicentre, randomised, controlled, phase 2 trial. *Lancet.* 2021;398(10318):2258-2276. doi:10.1016/s0140-6736(21)02717-3
95. Khoo NKH, Lim JME, Gill US, et al. Differential immunogenicity of homologous versus heterologous boost in Ad26.COV2.S vaccine recipients. *Med (N Y).* 2022;3(2):104-118.e4. doi:10.1016/j.medj.2021.12.004
96. Gao Y, Cai C, Grifoni A, et al. Ancestral SARS-CoV-2-specific T cells cross-recognize the Omicron variant. *Nat Med.* 2022;28:472-476. doi:10.1038/s41591-022-01700-x
97. Liu J, Chandrashekar A, Sellers D, et al. Vaccines elicit highly conserved cellular immunity to SARS-CoV-2 Omicron. *Nature.* 2022;603(7901):493-496. doi:10.1038/s41586-022-04465-y
98. Geers D, Shamier MC, Bogers S, et al. SARS-CoV-2 variants of concern partially escape humoral but not T cell responses in COVID-19 convalescent donors and vaccine recipients. *Sci Immunol.* 2021;6(59):eabj1750. doi:10.1126/sciimmunol.abj1750
99. Payne RP, Longet S, Austin JA, et al. Immunogenicity of standard and extended dosing intervals of BNT162b2 mRNA vaccine. *Cell.* 2021;184(23):5699-5714.e11. doi:10.1016/j.cell.2021.10.011
100. Sajadi MM, Myers A, Logue J, et al. Mucosal and systemic responses to SARS-CoV-2 vaccination in infection naïve and experienced individuals. *bioRxiv.* 2021;2021.12.13.472159. doi:10.1101/2021.12.13.472159
101. Minervina AA, Pogorelyy MV, Kirk AM, et al. SARS-CoV-2 antigen exposure history shapes phenotypes and specificity of memory CD8+ T cells. *Nat Immunol.* 2022;23:781-790. doi:10.1038/s41590-022-01184-4

102. Weaver C, Murphy K. *Janeway's Immunobiology*. W. W. Norton; 2016. <https://wwnorton.com/books/9780815345053/about-the-book>. Accessed October 31, 2020.
103. Crotty S. A brief history of T cell help to B cells. *Nat Rev Immunol*. 2015;15(3):185-189. doi:10.1038/nri3803
104. Crotty S. T follicular helper cell biology: a decade of discovery and diseases. *Immunity*. 2019;50(5):1132-1148. doi:10.1016/j.immuni.2019.04.011
105. Wilkinson TM, Li CKF, Chui CSC, et al. Preexisting influenza-specific CD4+ T cells correlate with disease protection against influenza challenge in humans. *Nat Med*. 2012;18(2):274-280. doi:10.1038/nm.2612
106. Weiskopf D, Bangs DJ, Sidney J, et al. Dengue virus infection elicits highly polarized CX3CR1+ cytotoxic CD4+ T cells associated with protective immunity. *Proc Natl Acad Sci USA*. 2015;112(31):E4256-E4263. doi:10.1073/pnas.1505956112
107. Juno JA, van Bockel D, Kent SJ, Kelleher AD, Zaunders JJ, Munier CM. Cytotoxic CD4 T cells-friend or foe during viral infection? *Front Immunol*. 2017;8(2-463):19. doi:10.3389/fimmu.2017.00019
108. Vardhana S, Baldo L, Moricell WG, Wherry EJ. Understanding T-cell responses to COVID-19 is essential for informing public health strategies. *Sci Immunol*. 2022:eabo1303. doi:10.1126/sciimmunol.abo1303
109. Breton G, Mendoza P, Hägglöf T, et al. Persistent cellular immunity to SARS-CoV-2 infection. *J Exp Med*. 2021;218(4):e20202515. doi:10.1084/jem.20202515
110. Wrapp KM, Lee WS, Koutsakos M, et al. Establishment and recall of SARS-CoV-2 spike epitope-specific CD4+ T cell memory. *Nat Immunol*. 2022;23:768-780. doi:10.1038/s41590-022-01175-5
111. Ng OW, Chia A, Tan AT, et al. Memory T cell responses targeting the SARS coronavirus persist up to 11 years post-infection. *Vaccine*. 2016;34(17):2008-2014. doi:10.1016/j.vaccine.2016.02.063
112. Li CK, Wu H, Yan H, et al. T cell responses to whole SARS coronavirus in humans. *J Immunol*. 2008;181(8):5490-5500. doi:10.4049/jimmunol.181.8.5490
113. Yang LT, Peng H, Zhu ZL, et al. Long-lived effector/central memory T-cell responses to severe acute respiratory syndrome coronavirus (SARS-CoV) S antigen in recovered SARS patients. *Clin Immunol*. 2006;120(2):171-178. doi:10.1016/j.clim.2006.05.002
114. Rodda LB, Morawski PA, Pruner KB, et al. Imprinted SARS-CoV-2-specific memory lymphocytes define hybrid immunity. *Cell*. 2022;185:1588-1601.e14. doi:10.1016/j.cell.2022.03.018
115. Meckiff BJ, Ramirez-Suástegui C, Fajardo V, et al. Imbalance of regulatory and cytotoxic SARS-CoV-2-reactive CD4+ T cells in COVID-19. *Cell*. 2020;183(5):1340-1353.e16. doi:10.1016/j.cell.2020.10.001
116. Kaneko N, Boucau J, Kuo HH, et al. Expansion of cytotoxic CD4+ T cells in the lungs in severe COVID-19. *SSRN Electron J*. 2021. doi:10.2139/ssrn.3813278
117. Kaneko N, Boucau J, Kuo HH, et al. Expansion of Cytotoxic CD4+ T cells in the lungs in severe COVID-19. *medRxiv*. 2021;2021.03.23.21253885. doi:10.1101/2021.03.23.21253885
118. Dong T, Liu G, Felce S, et al. Memory cytotoxic SARS-CoV-2 spike protein-specific CD4+ T cells associate with viral control. *Int Arch Allergy Immunol*. 2022;183:350-359. doi:10.21203/rs.3.rs-1317569/v1
119. Juno JA, Tan HX, Lee WS, et al. Humoral and circulating follicular helper T cell responses in recovered patients with COVID-19. *Nat Med*. 2020;26:1428-1434. doi:10.1038/s41591-020-0995-0
120. Painter MM, Mathew D, Goel RR, et al. Rapid induction of antigen-specific CD4+ T cells is associated with coordinated humoral and cellular immunity to SARS-CoV-2 mRNA vaccination. *Immunity*. 2021;54(9):2133-2142.e3. doi:10.1016/j.immuni.2021.08.001
121. Tazuin A, Nayrac M, Benlarbi M, et al. A single dose of the SARS-CoV-2 vaccine BNT162b2 elicits Fc-mediated antibody effector functions and T-cell responses. *Cell Host Microbe*. 2021;29(7):1137-1150.e6. doi:10.1016/j.chom.2021.06.001
122. Benteibibel SE, Khurana S, Schmitt N, et al. ICOS(+)/PD-1(+)/CXCR3(+) T follicular helper cells contribute to the generation of high-avidity antibodies following influenza vaccination. *Sci Rep*. 2016;6:26494. doi:10.1038/srep26494
123. Locci M, Havenar-Daughton C, Landais E, et al. Human circulating PD-1+CXCR3-CXCR5+ memory Tfh cells are highly functional and correlate with broadly neutralizing HIV antibody responses. *Immunity*. 2013;39(4):758-769. doi:10.1016/j.immuni.2013.08.031
124. Crotty S. Do memory CD4 T cells keep their cell-type programming: plasticity versus fate commitment? Complexities of interpretation due to the heterogeneity of memory CD4 T cells, including T follicular helper cells. *Cold Spring Harb Perspect Biol*. 2018;10(3):a032102. doi:10.1101/cshperspect.a032102
125. Nelde A, Bilich T, Heitmann JS, et al. SARS-CoV-2-derived peptides define heterologous and COVID-19-induced T cell recognition. *Nat Immunol*. 2021;22(1):74-85. doi:10.1038/s41590-020-00808-x
126. Poon MML, Byington E, Meng W, et al. Heterogeneity of human anti-viral immunity shaped by virus, tissue, age, and sex. *Cell Rep*. 2021;37(9):110071-110071. doi:10.1016/j.celrep.2021.110071
127. Sattler A, Angermair S, Stockmann H, et al. SARS-CoV-2 specific T-cell responses and correlations with COVID-19 patient predisposition. *J Clin Invest*. 2020;130(12):6477-6489. doi:10.1172/jci140965
128. Bert NL, Chia WN, Wan WY, et al. Widely heterogeneous humoral and cellular immunity after mild SARS-CoV-2 infection in a homogeneous population of healthy young men: Heterogenous immunity to SARS-CoV-2. *Emerg Microbes Infect*. 2021;10(1):2141-2150. doi:10.1080/22221751.2021.1999777
129. Bert NL, Clapham HE, Tan AT, et al. Highly functional virus-specific cellular immune response in asymptomatic SARS-CoV-2 infection. *J Exp Med*. 2021;218(5):e20202617. doi:10.1084/jem.20202617
130. Boyton RJ, Altmann DM. The immunology of asymptomatic SARS-CoV-2 infection: what are the key questions? *Nat Rev Immunol*. 2021;21(12):1-7. doi:10.1038/s41577-021-00631-x
131. Reynolds CJ, Swadling L, Gibbons JM, et al. Discordant neutralizing antibody and T cell responses in asymptomatic and mild SARS-CoV-2 infection. *Sci Immunol*. 2020;5(4):eabf3698. doi:10.1126/sciimmunol.abf3698
132. Vijayakumar B, Boustani K, Ogger PP, et al. Immuno-proteomic profiling reveals aberrant immune cell regulation in the airways of individuals with ongoing post-COVID-19 respiratory disease. *Immunity*. 2022;55(3):542-556.e5. doi:10.1016/j.immuni.2022.01.017
133. Braun J, Loyal L, Frentsch M, et al. SARS-CoV-2-reactive T cells in healthy donors and patients with COVID-19. *Nature*. 2020;587(7833):270-274. doi:10.1038/s41586-020-2598-9
134. Weiskopf D, Schmitz KS, Raadsen MP, et al. Phenotype and kinetics of SARS-CoV-2-specific T cells in COVID-19 patients with acute respiratory distress syndrome. *Sci Immunol*. 2020;5(48):eabd2071. doi:10.1126/sciimmunol.abd2071
135. Sette A, Crotty S. Pre-existing immunity to SARS-CoV-2: the knowns and unknowns. *Nat Rev Immunol*. 2020;20(8):457-458. doi:10.1038/s41577-020-0389-z
136. Mateus J, Grifoni A, Tarke A, et al. Selective and cross-reactive SARS-CoV-2 T cell epitopes in unexposed humans. *Science*. 2020;370(6512):89-94. doi:10.1126/science.abd3871
137. Schulien I, Kemming J, Oberhardt V, et al. Characterization of pre-existing and induced SARS-CoV-2-specific CD8+ T cells. *Nat Med*. 2020;27:78-85. doi:10.1038/s41591-020-01143-2
138. Shomuradova AS, Vagida MS, Sheetikov SA, et al. SARS-CoV-2 epitopes are recognized by a public and diverse repertoire of human T cell receptors. *Immunity*. 2020;53(6):1245-1257.e5. doi:10.1016/j.immuni.2020.11.004

139. Prakash S, Srivastava R, Coulon PG, et al. Genome-wide B cell, CD4+, and CD8+ T cell epitopes that are highly conserved between human and animal coronaviruses, identified from SARS-CoV-2 as targets for preemptive pan-coronavirus vaccines. *J Immunol.* 2021;206(11):ji2001438. doi:10.4049/jimmunol.2001438
140. Ishizuka J, Grebe K, Shenderov E, et al. Quantitating T cell cross-reactivity for unrelated peptide antigens. *J Immunol.* 2009;183(7):4337-4345. doi:10.4049/jimmunol.0901607
141. Grifoni A, Voic H, Dhanda SK, et al. T Cell responses induced by attenuated flavivirus vaccination are specific and show limited cross-reactivity with other flavivirus species. *J Virol.* 2020;94(10):e00089-20. doi:10.1128/jvi.00089-20
142. Bacher P, Rosati E, Esser D, et al. Low-avidity CD4+ T cell responses to SARS-CoV-2 in unexposed individuals and humans with severe COVID-19. *Immunity.* 2020;53(6):1258-1271.e5. doi:10.1016/j.immuni.2020.11.016
143. Lipsitch M, Grad YH, Sette A, Crotty S. Cross-reactive memory T cells and herd immunity to SARS-CoV-2. *Nat Rev Immunol.* 2020;20(11):709-713. doi:10.1038/s41577-020-00460-4
144. Loyal L, Braun J, Henze L, et al. Cross-reactive CD4 + T cells enhance SARS-CoV-2 immune responses upon infection and vaccination. *Science.* 2021;374:eabh1823. doi:10.1126/science.abh1823
145. Meyer-Arndt L, Schwarz T, Loyal L, et al. Cutting edge: serum but not mucosal antibody responses are associated with pre-existing SARS-CoV-2 spike cross-reactive CD4 + T cells following BNT162b2 vaccination in the elderly. *J Immunol.* 2022;208(5):1001-1005. doi:10.4049/jimmunol.2100990
146. Sagar M, Reifler K, Rossi M, et al. Recent endemic coronavirus infection is associated with less severe COVID-19. *J Clin Invest.* 2020;131(1):e143380. doi:10.1172/jci143380
147. Aran D, Beachler DC, Lanes S, Overhage JM. Prior presumed coronavirus infection reduces COVID-19 risk: a cohort study. *J Infect.* 2020;81(6):923-930. doi:10.1016/j.jinf.2020.10.023
148. Gombor S, Bergquist T, Pejaver V, et al. SARS-CoV-2 infection and COVID-19 severity in individuals with prior seasonal coronavirus infection. *Diagn Microbiol Infect Dis.* 2021;100(2):115338. doi:10.1016/j.diagmicrobio.2021.115338
149. Antunes RS, Pallikkuth S, Williams E, et al. Differential T cell reactivity to endemic coronaviruses and SARS-CoV-2 in community and health care workers. *J Infect Dis.* 2021;224(1):jjab176. doi:10.1093/infdis/jjab176
150. Swadling L, Diniz MO, Schmidt NM, et al. Pre-existing polymerase-specific T cells expand in abortive seronegative SARS-CoV-2. *Nature.* 2021;601(7891):110-117. doi:10.1038/s41586-021-04186-8
151. Kundu R, Narean JS, Wang L, et al. Cross-reactive memory T cells associate with protection against SARS-CoV-2 infection in COVID-19 contacts. *Nat Commun.* 2022;13(1):80. doi:10.1038/s41467-021-27674-x
152. Lederer K, Bettini E, Parvathaneni K, et al. Germinal center responses to SARS-CoV-2 mRNA vaccines in healthy and immunocompromised individuals. *Cell.* 2022;185(6):1008-1024.e15. doi:10.1016/j.cell.2022.01.027
153. Mudd PA, Minervina AA, Pogorelyy MV, et al. SARS-CoV-2 mRNA vaccination elicits a robust and persistent T follicular helper cell response in humans. *Cell.* 2022;185(4):603-613.e15. doi:10.1016/j.cell.2021.12.026
154. Ewer KJ, Barrett JR, Belij-Rammerstorfer S, et al. T cell and antibody responses induced by a single dose of ChAdOx1 nCoV-19 (AZD1222) vaccine in a phase 1/2 clinical trial. *Nat Med.* 2021;27(2):270-278. doi:10.1038/s41591-020-01194-5
155. Folegatti PM, Ewer KJ, Aley PK, et al. Safety and immunogenicity of the ChAdOx1 nCoV-19 vaccine against SARS-CoV-2: a preliminary report of a phase 1/2, single-blind, randomised controlled trial. *Lancet.* 2020;396(10249):467-478. doi:10.1016/s0140-6736(20)31604-4
156. Vikkurthi R, Ansari A, Pai AR, et al. Inactivated virus vaccine BBV152/Covaxin elicits robust cellular immune memory to SARS-CoV-2 and variants of concern. *medRxiv.* 2021;2021.11.14.21266294. doi:10.1101/2021.11.14.21266294
157. Stamatatos L, Czartoski J, Wan YH, et al. mRNA vaccination boosts cross-variant neutralizing antibodies elicited by SARS-CoV-2 infection. *Science.* 2021;372(6549):1413-1418. doi:10.1126/science.abg9175
158. Crotty S, Felgner P, Davies H, Glidewell J, Villarreal L, Ahmed R. Cutting Edge: Long-Term B Cell Memory in Humans after Smallpox Vaccination. *J Immunol.* 2003;171(10):4969-4973. doi:10.4049/jimmunol.171.10.4969
159. Yu X, Tsibane T, McGraw PA, et al. Neutralizing antibodies derived from the B cells of 1918 influenza pandemic survivors. *Nature.* 2008;455(7212):532-536. doi:10.1038/nature07231
160. Li GM, Chiu C, Wrarmert J, et al. Pandemic H1N1 influenza vaccine induces a recall response in humans that favors broadly cross-reactive memory B cells. *Proc Natl Acad Sci USA.* 2012;109(23):9047-9052. doi:10.1073/pnas.1118979109
161. Purtha WE, Tedder TF, Johnson S, Bhattacharya D, Diamond MS. Memory B cells, but not long-lived plasma cells, possess antigen specificities for viral escape mutants. *J Exp Med.* 2011;208(13):2599-2606. doi:10.1084/jem.20110740
162. Wang Z, Muecksch F, Schaefer-Babajew D, et al. Naturally enhanced neutralizing breadth against SARS-CoV-2 one year after infection. *Nature.* 2021;595(7867):426-431. doi:10.1038/s41586-021-03696-9
163. Marcotte H, Piralla A, Zuo F, et al. Immunity to SARS-CoV-2 up to 15 months after infection. *Science.* 2022;25(2):103743. doi:10.1016/j.isci.2022.103743
164. Sokal A, Chappert P, Barba-Spaeth G, et al. Maturation and persistence of the anti-SARS-CoV-2 memory B cell response. *Cell.* 2021;184(5):1201-1213.e14. doi:10.1016/j.cell.2021.01.050
165. Sakharkar M, Rappazzo CG, Wieland-Alter WF, et al. Prolonged evolution of the human B cell response to SARS-CoV-2 infection. *Sci Immunol.* 2021;6(56):eabg6916. doi:10.1126/sciimmunol.abg6916
166. Kaneko N, Kuo HH, Boucay J, et al. Loss of Bcl-6-expressing T follicular helper cells and germinal centers in COVID-19. *Cell.* 2020;183(1):143-157.e13. doi:10.1016/j.cell.2020.08.025
167. Röltgen K, Nielsen SCA, Silva O, et al. Immune imprinting, breadth of variant recognition, and germinal center response in human SARS-CoV-2 infection and vaccination. *Cell.* 2022;185(6):1025-1040.e14. doi:10.1016/j.cell.2022.01.018
168. Piccoli L, Park YJ, Tortorici MA, et al. Mapping neutralizing and immunodominant sites on the SARS-CoV-2 spike receptor-binding domain by structure-guided high-resolution serology. *Cell.* 2020;183(4):1024-1042.e21. doi:10.1016/j.cell.2020.09.037
169. Wec AZ, Haslwanter D, Abdiche YN, et al. Longitudinal dynamics of the human B cell response to the yellow fever 17D vaccine. *Proc Natl Acad Sci USA.* 2020;117(12):6675-6685. doi:10.1073/pnas.1921388117
170. Davis CW, Jackson KJL, McElroy AK, et al. Longitudinal analysis of the human B cell response to Ebola virus infection. *Cell.* 2019;177(6):1566-1582.e17. doi:10.1016/j.cell.2019.04.036
171. Allie SR, Bradley JE, Mudunuru U, et al. The establishment of resident memory B cells in the lung requires local antigen encounter. *Nat Immunol.* 2019;20(1):97-108. doi:10.1038/s41590-018-0260-6
172. Weisel NM, Weisel FJ, Farber DL, et al. Comprehensive analyses of B-cell compartments across the human body reveal novel subsets and a gut-resident memory phenotype. *Blood.* 2020;136(24):2774-2785. doi:10.1182/blood.2019002782

173. Cho A, Muecksch F, Schaefer-Babajew D, et al. Anti-SARS-CoV-2 receptor-binding domain antibody evolution after mRNA vaccination. *Nature*. 2021;600(7889):517-522. doi:10.1038/s41586-021-04060-7
174. Pauthner M, Havenar-Daughton C, Sok D, et al. Elicitation of robust tier 2 neutralizing antibody responses in nonhuman primates by HIV envelope trimer immunization using optimized approaches. *Immunity*. 2017;46(6):1073-1088.e6. doi:10.1016/j.immuni.2017.05.007
175. Havenar-Daughton C, Lee JH, Crotty S. Tfh cells and HIV bnAbs, an immunodominance model of the HIV neutralizing antibody generation problem. *Immunol Rev*. 2017;275(1):49-61. doi:10.1111/imr.12512
176. Havenar-Daughton C, Carnathan DG, Peña AT de la, et al. Direct probing of germinal center responses reveals immunological features and bottlenecks for neutralizing antibody responses to HIV Env trimer. *Cell Rep*. 2016;17(9):2195-2209. doi:10.1016/j.celrep.2016.10.085
177. Cirelli KM, Carnathan DG, Nogal B, et al. Slow delivery immunization enhances HIV neutralizing antibody and germinal center responses via modulation of immunodominance. *Cell*. 2019;177(5):1153-1171.e28. doi:10.1016/j.cell.2019.04.012
178. Lee JH, Sutton H, Cottrell CA, et al. Long-lasting germinal center responses to a priming immunization with continuous proliferation and somatic mutation. *bioRxiv*. 2021;2021.12.20.473537. doi:10.1101/2021.12.20.473537
179. Kim W, Zhou JQ, Horvath SC, et al. Germinal centre-driven maturation of B cell response to mRNA vaccination. *Nature*. 2022;604:141-145. doi:10.1038/s41586-022-04527-1
180. Turner JS, O'Halloran JA, Kalaidina E, et al. SARS-CoV-2 mRNA vaccines induce persistent human germinal centre responses. *Nature*. 2021;596(7870):109-113. doi:10.1038/s41586-021-03738-2
181. Sokal A, Barba-Spaeth G, Fernández I, et al. mRNA vaccination of naive and COVID-19-recovered individuals elicits potent memory B cells that recognize SARS-CoV-2 variants. *Immunity*. 2021;54:2893-2907.e5. doi:10.1016/j.immuni.2021.09.011
182. Gudbjartsson DF, Norddahl GL, Melsted P, et al. Humoral immune response to SARS-CoV-2 in Iceland. *N Engl J Med*. 2020;383(18):1724-1734. doi:10.1056/nejmoa2026116
183. Wajnberg A, Amanat F, Firpo A, et al. Robust neutralizing antibodies to SARS-CoV-2 infection persist for months. *Science*. 2020;370(6521):1227-1230. doi:10.1126/science.abd7728
184. Qaseem A, Yost J, Etzandia-Ikobaltzeta I, et al. What is the antibody response and role in conferring natural immunity after SARS-CoV-2 infection? Rapid, living practice points from the american college of physicians (Version 2). *Ann Intern Med*. 2022;175:556-565. doi:10.7326/m21-3272
185. Mandavilli A. Immunity to the coronavirus may last years, new data hint. *The New York Times*. 2020. <https://www.nytimes.com/2020/11/17/health/coronavirus-immunity.html>. Accessed January 1, 2022.
186. Li C, Yu D, Wu X, et al. Twelve-month specific IgG response to SARS-CoV-2 receptor-binding domain among COVID-19 convalescent plasma donors in Wuhan. *Nat Commun*. 2021;12(1):4144. doi:10.1038/s41467-021-24230-5
187. Gallais F, Gantner P, Bruel T, et al. Evolution of antibody responses up to 13 months after SARS-CoV-2 infection and risk of reinfection. *EBioMedicine*. 2021;71:103561. doi:10.1016/j.ebiom.2021.103561
188. Kannenberg J, Trawinski H, Henschler R, Buhmann R, Hönemann M, Jassoy C. Antibody course and memory B-cell response in the first year after SARS-CoV-2 infection. *J Infect Dis*. 2022;jiac034. doi:10.1093/infdis/jiac034
189. Turner JS, Kim W, Kalaidina E, et al. SARS-CoV-2 infection induces long-lived bone marrow plasma cells in humans. *Nature*. 2021;595(7867):421-425. doi:10.1038/s41586-021-03647-4
190. Isho B, Abe KT, Zuo M, et al. Persistence of serum and saliva antibody responses to SARS-CoV-2 spike antigens in COVID-19 patients. *Sci Immunol*. 2020;5(52):eabe5511. doi:10.1126/sciimmunol.abe5511
191. Yang Y, Yang M, Peng Y, et al. Longitudinal analysis of antibody dynamics in COVID-19 convalescents reveals neutralizing responses up to 16 months after infection. *Nat Microbiol*. 2022;7(3):423-433. doi:10.1038/s41564-021-01051-2
192. Wei J, Matthews PC, Stoesser N, et al. Anti-spike antibody response to natural SARS-CoV-2 infection in the general population. *Nat Commun*. 2021;12(1):6250. doi:10.1038/s41467-021-26479-2
193. Crotty S, Ahmed R. Immunological memory in humans. *Semin Immunol*. 2004;16(3):197-203. doi:10.1016/j.smim.2004.02.008
194. Amanna IJ, Carlson NE, Slifka MK. Duration of humoral immunity to common viral and vaccine antigens. *N Engl J Med*. 2007;357(19):1903-1915. doi:10.1056/nejmoa060692
195. Faria NR, Mellan TA, Whittaker C, et al. Genomics and epidemiology of the P.1 SARS-CoV-2 lineage in Manaus, Brazil. *Science*. 2021;372:815-821. doi:10.1126/science.abh2644
196. Edridge AWD, Kaczorowska J, Hoste ACR, et al. Seasonal coronavirus protective immunity is short-lasting. *Nat Med*. 2020;26(11):1691-1693. doi:10.1038/s41591-020-1083-1
197. Dobaño C, Alonso S, Vidal M, et al. Multiplex antibody analysis of IgM, IgA and IgG to SARS-CoV-2 in saliva and serum from infected children and their close contacts. *Front Immunol*. 2022;13:751705. doi:10.3389/fimmu.2022.751705
198. Steensels D, Pierlet N, Penders J, Mesotten D, Heylen L. Comparison of SARS-CoV-2 antibody response following vaccination with BNT162b2 and mRNA-1273. *JAMA*. 2021;326(15):1533-1535. doi:10.1001/jama.2021.15125
199. Israel A, Shenhar Y, Green I, et al. Large-scale study of antibody titer decay following BNT162b2 mRNA vaccine or SARS-CoV-2 infection. *Vaccines (Basel)*. 2021;10(1):64. doi:10.3390/vaccines10010064
200. Doria-Rose N, Suthar MS, Makowski M, et al. Antibody persistence through 6 months after the second dose of mRNA-1273 vaccine for Covid-19. *N Engl J Med*. 2021;384:2259-2261. doi:10.1056/nejmc2103916
201. Pegu A, O'Connell SE, Schmidt SD, et al. Durability of mRNA-1273 vaccine-induced antibodies against SARS-CoV-2 variants. *Science*. 2021;373(6561):eabj4176. doi:10.1126/science.abj4176
202. Pajon R, Doria-Rose NA, Shen X, et al. SARS-CoV-2 omicron variant neutralization after mRNA-1273 booster vaccination. *N Engl J Med*. 2022;386:1088-1091. doi:10.1056/nejmc2119912
203. Choi A, Koch M, Wu K, et al. Safety and immunogenicity of SARS-CoV-2 variant mRNA vaccine boosters in healthy adults: an interim analysis. *Nat Med*. 2021;27(11):2025-2031. doi:10.1038/s41591-021-01527-y
204. Falsey AR, Frenck RW, Walsh EE, et al. SARS-CoV-2 neutralization with BNT162b2 vaccine dose 3. *N Engl J Med*. 2021;385(17):NEJMc2113468. doi:10.1056/nejmc2113468
205. Xia H, Zou J, Kurhade C, et al. Neutralization of omicron SARS-CoV-2 by 2 or 3 doses of BNT162b2 vaccine. *bioRxiv*. 2022;2022.01.21.476344. doi:10.1101/2022.01.21.476344
206. Regev-Yochay G, Gonen T, Gilboa M, et al. 4th dose COVID mRNA vaccines' immunogenicity & efficacy against omicron VOC. *medRxiv*. 2022;2022.02.15.22270948. doi:10.1101/2022.02.15.22270948
207. Self WH, Tenforde MW, Rhoads JP, et al. Comparative effectiveness of Moderna, Pfizer-BioNTech, and Janssen (Johnson & Johnson) vaccines in preventing COVID-19 hospitalizations among adults without immunocompromising conditions – United States, March–August 2021. *MMWR Morb Mortal Wkly Rep*. 2021;70(38):1337-1343. doi:10.15585/mmwr.mm7038e1
208. Marks P. FDA Ad26.COV2.S COVID-19 vaccine emergency use authorization amendment review memorandum. 2021. <https://www.fda.gov/media/153439/download>. Accessed December 1, 2021.

209. Shen X, Tang H, Pajon R, et al. Neutralization of SARS-CoV-2 variants B.1.429 and B.1.351. *New Engl J Med*. 2021;384:2352-2354. doi:[10.1056/nejmc2103740](https://doi.org/10.1056/nejmc2103740)
210. Keech C, Albert G, Cho I, et al. Phase 1-2 Trial of a SARS-CoV-2 Recombinant Spike Protein Nanoparticle Vaccine. *New Engl J Med*. 2020;383(24):2320-2332. doi:[10.1056/nejmoa2026920](https://doi.org/10.1056/nejmoa2026920)
211. Sheikh-Mohamed S, Isho B, Chao GYC, et al. Systemic and mucosal IgA responses are variably induced in response to SARS-CoV-2 mRNA vaccination and are associated with protection against subsequent infection. *medRxiv*. 2022;2021.08.01.21261297. doi:[10.1101/2021.08.01.21261297](https://doi.org/10.1101/2021.08.01.21261297)
212. Collier AY, Brown CM, McMahan K, et al. Immune responses in fully vaccinated individuals following breakthrough infection with the SARS-CoV-2 delta variant in Provincetown, Massachusetts. *medRxiv*. 2021;2021.10.18.21265113. doi:[10.1101/2021.10.18.21265113](https://doi.org/10.1101/2021.10.18.21265113)
213. Corbett KS, Nason MC, Flach B, et al. Immune correlates of protection by mRNA-1273 vaccine against SARS-CoV-2 in nonhuman primates. *Science*. 2021;373(6561):eabj0299. doi:[10.1126/science.abj0299](https://doi.org/10.1126/science.abj0299)
214. Lund FE, Randall TD. Scent of a vaccine. *Science*. 2021;373(6553):397-399. doi:[10.1126/science.abg9857](https://doi.org/10.1126/science.abg9857)
215. Gaebler C, Wang Z, Lorenzi JCC, et al. Evolution of antibody immunity to SARS-CoV-2. *Nature*. 2021;591(7851):639-644. doi:[10.1038/s41586-021-03207-w](https://doi.org/10.1038/s41586-021-03207-w)
216. Muecksch F, Wang Z, Cho A, et al. Increased potency and breadth of SARS-CoV-2 neutralizing antibodies after a third mRNA vaccine dose. *bioRxiv*. 2022;2022.02.14.480394. doi:[10.1101/2022.02.14.480394](https://doi.org/10.1101/2022.02.14.480394)
217. Andreano E, Paciello I, Piccini G, et al. Hybrid immunity improves B cells and antibodies against SARS-CoV-2 variants. *Nature*. 2021;600(7889):530-535. doi:[10.1038/s41586-021-04117-7](https://doi.org/10.1038/s41586-021-04117-7)
218. Planas D, Saunders N, Maes P, et al. Considerable escape of SARS-CoV-2 Omicron to antibody neutralization. *Nature*. 2021;602:671-675. doi:[10.1038/s41586-021-04389-z](https://doi.org/10.1038/s41586-021-04389-z)
219. Saadat S, Tehrani ZR, Logue J, et al. Binding and neutralization antibody titers after a single vaccine dose in health care workers previously infected with SARS-CoV-2. *Jama*. 2021;325(14):1467-1469. doi:[10.1001/jama.2021.3341](https://doi.org/10.1001/jama.2021.3341)
220. Richardson SI, Madzorera VS, Spencer H, et al. SARS-CoV-2 Omicron triggers cross-reactive neutralization and Fc effector functions in previously vaccinated, but not unvaccinated individuals. *medRxiv*. 2022;2022.02.10.22270789. doi:[10.1101/2022.02.10.22270789](https://doi.org/10.1101/2022.02.10.22270789)
221. Gaebler C, DaSilva J, Bednarski E, et al. SARS-CoV-2 neutralization after mRNA vaccination and variant breakthrough infection. *medRxiv*. 2022;2022.02.09.22270692. doi:[10.1101/2022.02.09.22270692](https://doi.org/10.1101/2022.02.09.22270692)
222. Suryawanshi RK, Chen IP, Ma T, et al. Limited cross-variant immunity after infection with the SARS-CoV-2 omicron variant without vaccination. *medRxiv*. 2022;2022.01.13.22269243. doi:[10.1101/2022.01.13.22269243](https://doi.org/10.1101/2022.01.13.22269243)
223. Kotaki R, Adachi Y, Moriyama S, et al. SARS-CoV-2 Omicron-neutralizing memory B-cells are elicited by two doses of BNT162b2 mRNA vaccine. *Sci Immunol*. 2022;7:eabn8590. doi:[10.1126/sciimmunol.abn8590](https://doi.org/10.1126/sciimmunol.abn8590)
224. Janssen EM, Lemmens EE, Wolfe T, Christen U, van Herrath MG, Schoenberger SP. CD4+ T cells are required for secondary expansion and memory in CD8+ T lymphocytes. *Nature*. 2003;421(6925):852-856. doi:[10.1038/nature01441](https://doi.org/10.1038/nature01441)
225. Apostolidis SA, Kakara M, Painter MM, et al. Cellular and humoral immune responses following SARS-CoV-2 mRNA vaccination in patients with multiple sclerosis on anti-CD20 therapy. *Nat Med*. 2021;27(11):1990-2001. doi:[10.1038/s41591-021-01507-2](https://doi.org/10.1038/s41591-021-01507-2)
226. Tortorella C, Aiello A, Gasperini C, et al. Humoral- and T-cell-specific immune responses to SARS-CoV-2 mRNA vaccination in patients with MS using different disease-modifying therapies. *Neurology*. 2022;98(5):e541-e554. doi:[10.1212/wnl.00000000000013108](https://doi.org/10.1212/wnl.00000000000013108)
227. Bates TA, McBride SK, Leier HC, et al. Vaccination before or after SARS-CoV-2 infection leads to robust humoral response and antibodies that effectively neutralize variants. *Sci Immunol*. 2022;7:eabn8014. doi:[10.1126/sciimmunol.abn8014](https://doi.org/10.1126/sciimmunol.abn8014)
228. Cele S, Jackson L, Khoury DS, et al. Omicron extensively but incompletely escapes Pfizer BNT162b2 neutralization. *Nature*. 2021;602:654-656. doi:[10.1038/s41586-021-04387-1](https://doi.org/10.1038/s41586-021-04387-1)
229. Goel RR, Apostolidis SA, Painter MM, et al. Distinct antibody and memory B cell responses in SARS-CoV-2 naïve and re-covered individuals following mRNA vaccination. *Sci Immunol*. 2021;6(58):eabi6950. doi:[10.1126/sciimmunol.abi6950](https://doi.org/10.1126/sciimmunol.abi6950)
230. Rodda LB, Morawski PA, Pruner KB, et al. Imprinted SARS-CoV-2-specific memory lymphocytes define hybrid immunity. *medRxiv*. 2022;2022.01.12.22269192. doi:[10.1101/2022.01.12.22269192](https://doi.org/10.1101/2022.01.12.22269192)
231. Brouillette M. Your immune system evolves to fight coronavirus variants - scientific American. *Scientific American*. 2021. <https://www.scientificamerican.com/article/your-immune-system-evolves-to-fight-coronavirus-variants/>. Accessed December 1, 2021.
232. Tarke A, Coelho CH, Zhang Z, et al. SARS-CoV-2 vaccination induces immunological memory able to cross-recognize variants from Alpha to Omicron. *bioRxiv*. 2021;2021.12.28.474333. doi:[10.1101/2021.12.28.474333](https://doi.org/10.1101/2021.12.28.474333)
233. Laidlaw BJ, Ellebedy AH. The germinal centre B cell response to SARS-CoV-2. *Nat Rev Immunol*. 2021;22:7-18. doi:[10.1038/s41577-021-00657-1](https://doi.org/10.1038/s41577-021-00657-1)
234. Kim W, Zhou JQ, Sturtz AJ, et al. Germinal centre-driven maturation of B cell response to SARS-CoV-2 vaccination. *bioRxiv*. 2021;2021.10.31.466651. doi:[10.1101/2021.10.31.466651](https://doi.org/10.1101/2021.10.31.466651)
235. Lederer K, Parvathaneni K, Painter MM, et al. Germinal center responses to SARS-CoV-2 mRNA vaccines in healthy and immunocompromised individuals. *medRxiv*. 2021;2021.09.16.21263686. doi:[10.1101/2021.09.16.21263686](https://doi.org/10.1101/2021.09.16.21263686)
236. Muecksch F, Weisblum Y, Barnes CO, et al. Affinity maturation of SARS-CoV-2 neutralizing antibodies confers potency, breadth, and resilience to viral escape mutations. *Immunity*. 2021;54:1853-1868.e7. doi:[10.1016/j.immuni.2021.07.008](https://doi.org/10.1016/j.immuni.2021.07.008)

How to cite this article: Sette A, Crotty S. Immunological memory to SARS-CoV-2 infection and COVID-19 vaccines. *Immunol Rev*. 2022;00:1-20. doi: [10.1111/imr.13089](https://doi.org/10.1111/imr.13089)