

BMJ Open Effects of remote learning during the COVID-19 lockdown on children's visual health: a systematic review

María Camila Cortés-Albornoz, Sofía Ramírez-Guerrero , William Rojas-Carabali , Alejandra de-la-Torre , Claudia Talero-Gutiérrez 

To cite: Cortés-Albornoz MC, Ramírez-Guerrero S, Rojas-Carabali W, *et al*. Effects of remote learning during the COVID-19 lockdown on children's visual health: a systematic review. *BMJ Open* 2022;**12**:e062388. doi:10.1136/bmjopen-2022-062388

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2022-062388>).

Received 01 March 2022
Accepted 19 July 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

Neuroscience Research Group (NeURos), NeuroVitae Center for Neuroscience, School of Medicine and Health Sciences, Universidad del Rosario, Bogota D.C, Colombia

Correspondence to

Dr Claudia Talero-Gutiérrez;
claudia.talero@urosario.edu.co

ABSTRACT

Objectives Increased exposure to digital devices as part of online classes increases susceptibility to visual impairments, particularly among school students taught using e-learning strategies. This study aimed to identify the impact of remote learning during the COVID-19 lockdown on children's visual health.

Design Systematic review using the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines.

Data sources Scopus, PubMed and ScienceDirect databases from the year 2020 onwards.

Eligibility criteria We included cross-sectional, case-control, cohort studies, case series and case reports, published in English, Spanish or French, that approached the effects of remote learning during the COVID-19 lockdown on visual health in neurotypical children.

Data extraction and synthesis We included a total of 21 articles with previous quality assessments using the Joanna Briggs checklist. Risk of bias assessment was applied using the National Institutes of Health quality assessment tool for before-and-after studies with no control group; the tool developed by Hoy *et al* to assess cross-sectional studies; the Murad *et al* tool to evaluate the methodological quality of case reports and case series; and the Newcastle-Ottawa Scale for cohort studies.

Results All but one study reported a deleterious impact of the COVID-19 lockdown on visual health in children. Overall, the most frequently identified ocular effects were refractive errors, accommodation disturbances and visual symptoms such as dry eye and asthenopia.

Conclusions Increased dependence on digital devices for online classes has either induced or exacerbated visual disturbances, such as rapid progression of myopia, dry eye and visual fatigue symptoms, and vergence and accommodation disturbances, in children who engaged in remote learning during the COVID-19 lockdown.

PROSPERO registration number CRD42022307107.

INTRODUCTION

Since the WHO declared a global pandemic in March 2020, COVID-19 has become the focus of governmental decisions aimed at protecting the public and limiting the death toll. Schools, universities and businesses have been forced to close to prevent the spread of the virus, limiting in-person relationships and

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ A systematic review was conducted in three different databases, studies were filtered following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines.
- ⇒ Analysed studies approached the effects of remote learning during the COVID-19 lockdown on visual health in children.
- ⇒ To facilitate comparison, eligible studies were clustered according to the main ocular effects evaluated, including refractive errors (myopia), accommodation disturbances (esotropia) and visual symptoms (dry eye and fatigue).
- ⇒ We used quality assessment guidelines and specific risk of bias assessment tools for each study design included.
- ⇒ Heterogeneous methods used in each study, including both subjective and objective measures, limit precise comparisons between them.

substantially enhancing our digital dependence. The lifestyle and behavioural modifications that have emerged in response to the lockdowns have affected approximately 80% of the world's student population.^{1 2}

The establishment of in-house quarantine led to a significant decrease in the amount of time spent engaged in outdoor activities, reduction in exposure to sunlight and increase in time spent doing near work. These factors can enhance the risk of visual impairments, especially among school and university students encouraged to adopt a digital learning approach.³ A growing dependence on e-learning and electronic devices has increased the incidence of visual fatigue, the onset and progression of myopia, dry eye, irregular astigmatism and acute concomitant esotropia among other ocular pathologies.⁴

Even before the COVID-19 pandemic, an estimated 22.9% of the global population had myopia.⁵ During the COVID-19 lockdown, the increased need for electronic devices, digital screens and virtual classrooms

might have caused previously healthy students to develop myopia, and faster progression in those who already had impaired vision. Obligatory confinement, intensive near work activities and decreased exposure to sunlight can lead to visual fatigue, and may also enhance the risk of myopia, the most prevalent ocular condition.⁴

Digital screen use is considered a common risk factor for dry eye, characterised by the deterioration of tear film quality. The risk of dry eye and symptom severity can be exacerbated by increased digital screen time.^{6–8} Myopia and dry eye are potential visual health consequences associated with the increasing demand for children to engage in e-learning, which often starts at a very young age. To address this in the present systematic review, we sought to identify the impact of remote learning during the COVID-19 pandemic on visual health in school-age children.

METHODS

Search strategy and selection criteria

In January 2022, we conducted a systematic review using three online databases. We used the following terms in PubMed: (<https://pubmed.ncbi.nlm.nih.gov/advanced/>) (((((vision) OR (visual impairment)) OR (myopia [MeSH Terms])) AND (COVID-19)) AND (lockdown)) AND (screen time); ScienceDirect: ([sciencedirect.com/search\) \(\(vision\) OR \(visual impairment\) OR \(myopia\)\) AND \(\(COVID-19 lockdown\)\) AND \(screen time\)\); and Scopus: \(<https://www.scopus.com>\) ALL \(vision OR \('visual' AND 'impairment'\) OR myopia AND \('COVID-19' AND 'lockdown'\) AND \('screen' AND 'time'\)\) AND \(LIMIT-TO \(SUBJAREA, 'MEDI'\) OR LIMIT-TO \(SUBJAREA, 'COMP'\) OR LIMIT-TO \(SUBJAREA, 'NEUR'\) OR LIMIT-TO \(SUBJAREA, 'NURS'\) OR LIMIT-TO \(SUBJAREA, 'HEAL'\)\). The ID CRD42022307107 was generated in the International Prospective Register of Systematic Reviews \(PROSPERO\).](https://www.</p>
</div>
<div data-bbox=)

Data collection

A total of 326 articles were initially retrieved. Duplicates were removed, and the remaining articles were filtered by title and abstract following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines (figure 1 and online supplemental table 1). Five researchers divided into two groups screened all of the articles, and 28 were selected for study inclusion. At weekly meetings, the authors analysed the studies, debated disagreements and double-checked all of the articles according to the inclusion and exclusion criteria. Articles were included if they described studies on the effects of remote learning during the COVID-19 lockdown on visual health in neurotypical children. They were excluded if they (1) were published before 2020; (2) studied the




Figure 1 Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram (adapted from Moher *et al.*⁵²)

effects of remote learning during the COVID-19 lockdown on visual health in adults or university students; (3) assessed children with genetic syndromes or visual disabilities; (4) were book chapters, editorials or opinion pieces; and (5) were published in languages other than Spanish, English and French. Following this procedure, a total of 21 articles were included. These were evaluated using Joanna Briggs checklist to guarantee study quality. Additionally, we conducted a risk of bias assessment using several tools. First, we used the National Institutes of Health quality assessment tool for before-and-after (pre-post) studies with no control group.⁹ This instrument evaluates 12 major components with response options of yes/no/not applicable/cannot determine/not reported and gives a final quality rating of good, poor or fair depending on the overall item response.⁹ Second, we used the tool developed by Hoy *et al* to assess cross-sectional studies by categorising the article bias as low, moderate or high risk according to responses to 10 questions.^{10,11} Third, we used the tool proposed by Murad *et al* to evaluate the methodological quality of case reports and case series. This tool appraises the selection, ascertainment, causality and reporting bias of each article and makes an overall judgement about the methodology based on the responses to eight questions.¹² Finally, we used the Newcastle-Ottawa Scale for cohort studies to assess the selection, comparability and outcome bias of the article by applying a qualitative star scale.⁹ All domains evaluated using these tools can be found in online supplemental table 2.

Finally, we extracted data to obtain the following information: title, authors, digital object identifier number, objective, type of study, country in which the study was conducted, population (age and sample), presence of control group (age and sample), implemented test or evaluation methodology, main visual outcome, results, conclusion and answers to the question 'Did the COVID-19 lockdown impact visual health (improvement, deterioration, no change)?' All information was synthesised using qualitative and quantitative synthesis (see the Results section). Considering the heterogeneity among studies, we created subgroups for analysis, for example, studies regarding dry eye, refractive errors, clinical symptoms and other clusters. All investigators participated in the data collection and synthesis.

Patient and public involvement

This research was done without patient or public involvement. However, the findings will be shared at conferences attended by paediatric ophthalmologists and patients with myopia who access ophthalmological services.

RESULTS

We grouped the articles included in the review based on the main visual outcome associated with vision status and changes in vision in children during the COVID-19 lockdown. Overall, the main ocular effects observed were refractive errors (myopia), accommodation disturbances

(esotropia) and visual symptoms (dry eye and fatigue) (table 1). Among the studies, 16 were conducted in Asia,^{13–28} 2 in Europe^{29,30} and 3 in America.^{31,32} The risk of bias assessment revealed that all of the cross-sectional studies and case series had a low risk of bias. Three of the before-and-after studies had fair quality, and one had good quality.

We identified 11 articles that examined refractive errors related to virtual learning during the COVID-19 lockdown. Most of these examined myopia progression as the main visual outcome. Eight studies reported that myopia worsened throughout the COVID-19 lockdown in children and teenagers between 5 and 18 years old.^{15,17,19,21–24,27} One study reported a significant decrease in spherical equivalent refraction (SER) in children with hyperopia and emmetropia (see table 2, Glossary).³⁰ Interestingly, a study evaluating axial length in myopic children undergoing orthokeratology (see table 2, Glossary) did not find any change in myopia progression after lockdown.²¹ Furthermore, one study focused on risk factors and behavioural changes during the COVID-19 lockdown in terms of myopia found that all children had changes in near work time, electronic device use and outdoor time. However, myopic children had a significantly lower level of daily light exposure compared with non-myopic children.³² The monthly extent of myopia progression during the COVID-19 lockdown was reported to be -0.074 D/month, which corresponds to an annual progression in 2020 of -0.71 ± 0.46 D.^{15,20} Furthermore, rapid myopia progression was reported in a sample of 133 school students. Specifically, the percentage of children with reported annual progression for whom progression was rapid increased from 10.5% before to 45.9% during the pandemic.²⁷ SER was estimated in several studies. In 2020, the mean SER in myopic children and teenagers was between -1.94 ± 2.13 D and -2.7 ± 1.21 D, and this was significantly lower than in 2019 (-1.64 ± 5.49 D and -1.99 ± 1.04 D, $p < 0.001$).^{19,20} Similarly, there was a significant decrease in the mean SER of hyperopic and emmetropic children from 2019 to 2020, that is, 0.66 ± 2.03 D (2019) and 0.48 ± 1.81 D (2020), respectively, $p \leq 0.001$.³⁰ Finally, studies examining virtual learning during the COVID-19 lockdown as an exposure risk factor found a higher incidence of myopia in children who engaged in virtual learning ($p < 0.01$).^{22–24}

Four studies reported accommodation and vergence dysfunction (see table 2, Glossary) secondary to near work and increased screen use time.^{13,26,29,33} Two studies focused on binocular accommodation in a sample of 156 children aged 10–17 years and reported a significant increase in Convergence Insufficiency Symptom Survey (CISS) scores after exposure to longer screen time during online classes.^{11,29} The other two were case series of children who developed acquired concomitant esotropia and vergence abnormalities secondary to the excessive use of digital devices.^{27,29}

Emerging visual symptoms were identified in six studies with populations ranging from 8 to 20 years old. The

Table 1 Articles related to visual outcomes and the impact of remote learning during the COVID-19 pandemic

Title	Authors	Year	Type of study	Country of the study	Results	Is there an effect of COVID-19 lockdown on visual health?
Comparison of myopic progression before, during, and after COVID-19 lockdown	Chang <i>et al</i> ¹⁵	2021	Before-and-after study	China	Proportions of myopia and high myopia: Round 1: 48.0% and 1.3% Round 2: 53.2% and 1.9% Round 3: 73.7% and 2.8% Round 4: 67.9% and 2.7%	Worsen
Impact of COVID-19 home confinement in children's refractive errors	Alvarez-Peregrina <i>et al</i> ³⁰	2021	Cross-sectional	Spain	Mean rDSER: Period 1: -0.030 D/month (95% CI e0.031 to e0.029) Period 2 (lockdown): -0.074 D/month (95% CI e0.075 to e0.074) Period 3: 0.016 D/month (95% CI 0.015 to e0.018) Spherical equivalent: Average value in 2019: +0.66±2.03 D Average value in 2020: +0.48±1.81 D Children's lifestyle during confinement: 56% changed the amount of time spent outdoors (95% CI 53% to 58%). 47% (95% CI 45% to 50%; p<0.001) of the cases decreased this time. Children's near vision during confinement: 49% changed the amount of time they spent doing near distance activities (95% CI 47% to 52%). 44% (95% CI 41% to 46%; p<0.001) of the children increased near distance activities. 42% of the children changed the amount of time they spent with electronic devices (95% CI 40% to 45%). 39% (95% CI 37% to 42%; p<0.001) of cases increase electronic use.	Worsen
Objective and subjective behavioral measures in myopic and non-myopic children during the COVID-19 pandemic	Mirhajianmoghadam <i>et al</i> ³²	2021	Cross-sectional	USA	Children who spent more time outdoors had higher SE in preconfinement and postconfinement (p<0.001 and p=0.049). 1. Subjective measures: - Significant interaction between session and day of the week. - Time outdoors: 2 hours less during the 2020 summer of COVID-19 compared with typical summer before COVID-19. No significant differences between refractive error groups (p=0.20). - Daily electronic device use: increased on weekdays and weekends during COVID-19 (7.3±0.6 and 7.9±0.7 hours) compared with a typical summer (4.9±0.5 and 6.1±0.5 hours, p<0.001) and with a typical school period (3.4±0.3 and 5.4±0.5 hours, p<0.001). 2. Objective measures: - During COVID-19: myopic children had lower daily light exposure (183.6±39.3 lux) than non-myopic children (279.5±23.5 lux, p=0.04) (p=0.09).	Unclear

Continued

Table 1 Continued

Title	Authors	Year	Type of study	Country of the study	Results	Is there an effect of COVID-19 lockdown on visual health?
Progression of myopia in school-aged children after COVID-19 home confinement	Wang <i>et al</i> ¹⁷	2021	Prospective cross-sectional	China	Mean SER: ▲ Annual screenings from 2015 to 2019: stable for all age groups. ▲ SER decreased in 2020 compared with 2015–2019 in children aged 6 (–0.32 D), 7 (–0.28 D) and 8 (–0.29 D) years. Prevalence of myopia: ▲ 2020: 21.5% at 6 years, 26.2% at 7 years, 37.2% at 8 years. ▲ 2015–2019: 5.7% at 6 years in 2019, 16.2% at 7 years in 2018, 27.7% at 8 years in 2018.	Worsen
Survey on the progression of myopia in children and adolescents in Chongqing during COVID-19 pandemic	Wang <i>et al</i> ¹⁹	2021	Before-and-after study	China	Myopia prevalence among teenagers: ▲ 2019: 44.62%. ▲ 2020: 55.02%. Average progression rate: 10.49% Spherical equivalent: ▲ 2019: –1.64±5.49 D. ▲ 2020: –1.94±2.13 D. Myopia percentage was 84.89% in high school, 73.39% in junior school and 39.27% in primary school.	Worsen
The effect of home education on myopia progression in children during the COVID-19 pandemic	Aslan and Sahinoglu- Keskek ²⁰	2022	Before-and-after study	Turkey	Mean duration spent in front of the screen was 5.77±1.34 hours/day. The mean SER were: ▲ 2016: –1.14±0.66 D. ▲ 2017: –1.47±0.82 D. ▲ 2018: –0.45±0.91 D. ▲ 2019: –1.99±1.04 D. ▲ 2020: –2.7±1.21 D. The mean myopic progression in 2020: ▲ Overall: 0.71±0.46 D. ▲ In children who spent time outside in the daylight for 2 hours/day: 0.55±0.42 D. ▲ In children with less outside time: 0.82±0.45 D (p=0.003). The myopia progression in 2020 was slow (0.31±0.2 D) in 42.6% of the subjects, moderate (0.82±0.14 D) in 39.1% and rapid (1.42±0.29 D) in 18.3%. No correlation was found between the 2020 progression and the daily digital device use.	Worsen
The impact of COVID-19 home confinement on axial length in myopic children undergoing orthokeratology	Lv <i>et al</i> ²¹	2022	Before-and-after study	China	Monthly axial length growth: ▲ After confinement: 0.023±0.019 mm/month. ▲ During confinement: 0.018±0.021 mm/month, negatively related (p=0.002). ▲ Before confinement: 0.014±0.016 mm/month. The monthly axial length growth after and before confinement was not significantly different (p=0.333).	Remains the same

Continued

Table 1 Continued

Title	Authors	Year	Type of study	Country of the study	Results	Is there an effect of COVID-19 lockdown on visual health?
The impact of study-at-home during the COVID-19 pandemic on myopia progression in Chinese children	Ma <i>et al</i> ²²	2021	Cohort	China	Myopia progression: $p < 0.001$ ▲ In exposed group: -0.83 ± 0.56 D. ▲ In control group: -0.28 ± 0.54 D. In the exposed group, children had a larger change in myopia progression in the follow-up period (-0.83 ± 0.56 D) compared with the baseline period (-0.33 ± 0.46 D; $p < 0.001$). Increment on near work time from 2.96 ± 1.05 to 4.33 ± 1.04 hours/day ($p < 0.001$) during COVID-19. Decrease on outdoor activities from 1.84 ± 1.43 to 0.98 ± 1.01 hours/day ($p < 0.001$) during COVID-19.	Worsen
COVID-19 quarantine reveals that behavioral changes have an effect on myopia progression	Xu <i>et al</i> ²³	2021	Before-and-after study	China	Myopia prevalence: ▲ June 2019: 52.89% (95% CI 52.79% to 52.99%). ▲ December 2019: 53.9% (95% CI 53.79% to 54.01%). ▲ June 2020: 59.35% (95% CI 59.24% to 59.46%). Increase in myopia prevalence: ▲ Grades 1–6: 8.54%. ▲ Grades 7–12: 4.32%. Half-year incidence rate of myopia: ▲ Before COVID-19: 8.5%. ▲ After COVID-19: 13.62% ($p < 0.001$).	Worsen
Rates of myopia development in young Chinese schoolchildren during the outbreak of COVID-19	Hu <i>et al</i> ²⁴	2021	Cohort	China	The mean AL was 0.11 mm (95% CI 0.05 to 0.16). Exposed group: ▲ Myopic shift of SER: 0.36 D (95% CI 0.32 to 0.41; $p < 0.001$) and 0.08 mm (95% CI 0.06 to 0.10; $p < 0.001$) greater AL elongation. ▲ Incidence of myopia: 7.9% (95% CI 5.1% to 10.6%; $p < 0.001$) higher. ▲ Prevalence of myopia: 219 of 1054 students (20.8%) were 7.5% (95% CI 4.3% to 10.7%) higher than in the non-exposure group (141 of 1060 students (13.3%)).	Worsen
Impact of online classes and home confinement on myopia progression in children during COVID-19 pandemic: Digital Eye Strain among Kids (DESK study 4)	Mohan <i>et al</i> ²⁸	2022	Cross-sectional	India	Myopia progression report: ▲ Before COVID-19: 45.9% of participants. ▲ During the COVID-19: 62.5% of participants.	Worsen
Acute acquired concomitant esotropia from excessive application of near vision during the COVID-19 lockdown	Vagge <i>et al</i> ²⁹	2020	Case series	Italy	▲ A 4-year-old girl with ACE of 35 prism dioptres managed with glasses. She used the tablet around 8 hours/day. ▲ A 16-year-old boy with ACE of 30 prism dioptres managed with Fresnel prism. Computer 8 hours/day. ▲ A 16-year-old boy with ACE of 20 prism dioptres managed with Fresnel prism. Computer 10 hours/day. ▲ An 8-year-old girl with ACE of 25 prism dioptres managed conservatively. She used the tablet around 8 hours/day.	Worsen

Continued

Table 1 Continued

Title	Authors	Year	Type of study	Country of the study	Results	Is there an effect of COVID-19 lockdown on visual health?
Binocular accommodation and vergence dysfunction in children attending online classes during the COVID-19 pandemic: Digital Eye Strain in Kids (DESK study-2)	Mohan <i>et al</i> ¹³	2021	Cross-sectional	India	<p>Mean CISS scores:</p> <ul style="list-style-type: none"> ▲ In children using devices less than 4 hours/day: 21.73±12.81. ▲ In children using digital devices for 4 hours/day or more: 30.34±13.0 (p=0.019). <p>Mean values of near exophoria (p=0.03), NFV (p=0.02), NRA (p=0.057) and AA (p=0.002).</p> <p>Spearman correlation between CISS score and the duration of online classes: weak linear association (coefficient rs=0.39, p=0.007).</p>	Worsen
The visual consequences of virtual school: acute eye symptoms in healthy children	Hamburger <i>et al</i> ³³	2022	Cross-sectional	USA	<p>CISS score:</p> <ul style="list-style-type: none"> ▲ Before school: mean of 5.17 and median of 4. ▲ After school: mean of 9.82 and median of 7.5 (mean change 4.65; median change 2; p<0.001). ▲ Linear regression analysis of change in total CISS score from before to after school versus hours spent in virtual school: score increase of 1.243 per hour of virtual school (p=0.0282). <p>The asthenopia score:</p> <ul style="list-style-type: none"> ▲ Before school: mean of 1.58 and total median of 1. ▲ After school: mean of 2.74 and median of 2 (mean change 1.15, median change 1; p<0.001). ▲ Linear regression analysis of change in total asthenopia score from before to after school versus actual hours spent in virtual school: score increase of 0.280 per hour of virtual school (p=0.0807). 	Worsen
Series of cases of acute acquired comitant esotropia in children associated with excessive online classes on smartphone during COVID-19 pandemic; Digital Eye Strain among Kids (DESK study-3)	Mohan <i>et al</i> ²⁷	2021	Case series	India	<ul style="list-style-type: none"> ▲ 5/8 subjects were emmetropic, 1 myopic, 1 pseudomyopic, 1 mild hyperopic. ▲ Average use of device: 4.6±0.7 hours/day at an average of 5.5 inches from the screen. ▲ Average angle deviation for near vision with corrected vision in esotropia: 48.1±16.4 PD. ▲ 7/8 children reported horizontal diplopia. 	Worsen
Relationship between screen time and dry eye symptoms in pediatric population during the COVID-19 pandemic	Elhousseiny <i>et al</i> ¹⁸	2021	Cross-sectional	Egypt	<p>m-SPEED questionnaire score:</p> <ul style="list-style-type: none"> ▲ Pre-COVID-19: 0.83±2.04 (p<0.001). ▲ COVID-19 2020: 3.9±4.53 (higher in urban areas (4.68±4.87) versus rural areas (2.97±3.69) (p<0.001)). <p>Screen time:</p> <ul style="list-style-type: none"> ▲ Male sex: associated with greater screen time (mean difference of 0.6±0.31 hours/day (p=0.047)). ▲ Development of DED: in association with prolonged screen time. 	Worsen

Continued

Table 1 Continued

Title	Authors	Year	Type of study	Country of the study	Results	Is there an effect of COVID-19 lockdown on visual health?
New indicator of children's excessive electronic screen use and factors in meibomian gland atrophy	Creemers <i>et al</i> ³¹	2021	Cross-sectional	USA	CHESUD: <ul style="list-style-type: none"> ▲ 4 or more hours: 86% of MGA cases. ▲ 8 or more hours: 50%. Severe MGA related to: <ul style="list-style-type: none"> ▲ Less outdoor time. ▲ Higher meibography scores ($p < 0.01$). CHESUD's positive association with the increase in combined meibography scores (OR=2.81; 95% CI 1.66 to 4.77).	Worsen
Prevalence and risk factor assessment of digital eye strain among children using online e-learning during the COVID-19 pandemic: Digital Eye Strain among Kids (DESK study-1)	Mohan <i>et al</i> ¹⁴	2021	Cross-sectional	India	During the COVID-19: <ul style="list-style-type: none"> ▲ Mean duration of digital device use: 3.9 ± 1.9 hours. ▲ Percentage of children using devices for >5 hours: 36.9%. Before COVID-19: <ul style="list-style-type: none"> ▲ Mean duration of digital device use: 1.9 ± 1.1 hours. ▲ Percentage of children using devices for >5 hours: 1.8%. Survey report: <ul style="list-style-type: none"> ▲ Eyesight worsened because of the online classes: 49.76%. ▲ Overall prevalence of DES: 50.23%. 	Worsen
Impact of e-schooling on digital eye strain in coronavirus disease era: a survey of 654 students	Gupta <i>et al</i> ¹⁶	2021	Cross-sectional	India	Asthenopia and dry eye symptoms: <ul style="list-style-type: none"> ▲ Heaviness of eyelids: 79.7%, eye redness: 69.1%, eye strain: 68.2%, blinking: 57.8%, blurred vision: 56.9%, light sensitivity: 56%, stinging: 47.1%, burning: 46.3%. Digital device use: <ul style="list-style-type: none"> <4 hours: 30 (12.5%) children, 5–6 hours: 13 (6.6%) children, >6 hours: 4 (2.3%) children, had not reported any AS ($p = 0.001$). Computer vision syndrome score: CVS score was statistically significantly lower in the group aged 5–7 years versus the group aged 11–16 years ($p = 0.004$).	Worsen
Prevalence of self-reported symptoms of computer vision syndrome and associated risk factors among school students in China during the COVID-19 pandemic	Li <i>et al</i> ⁶	2021	Cross-sectional	China	53% used glasses, 47% myopia, 2.9% hyperopia, 2.1% astigmatism, 0.8% amblyopia, 0.7% anisometropia, 1.5% strabismus, 2.5% conjunctivitis, 1.7% previous eye surgery. Mean screen time 4.6 hours/day, 1.2 hours/day on outdoor activities, 9.9 hours/day on sleep. Self-reported symptoms: 13% double vision, 48% itching, eye dryness. <ul style="list-style-type: none"> ▲ Mild intensity: 1.4% coloured halos around objects, 8.8% eye dryness, 0.3% double vision, 3.2% neck or shoulder pain. ▲ Moderate intensity: 1.3% coloured halos around objects, 7.8% eye dryness. ▲ Severe intensity: 0.3% double vision, 2.7% neck or shoulder pain or feeling sight worsening. 	Worsen

Continued

Table 1 Continued

Title	Authors	Year	Type of study	Country of the study	Results	Is there an effect of COVID-19 lockdown on visual health?
Contribution of total screen/online-course time to asthenopia in children during COVID-19 pandemic via influencing psychological stress	Li et al ²⁵	2021	Cross-sectional	China	63.1% had myopia, 36% had astigmatism and 12.1% reported asthenopia. Students with asthenopia had longer screen/online-course time and less daily rest time. A 100-hour increment was associated with an increased risk of asthenopia at 9% (OR=1.09) and 11% (OR=1.11).	Worsen
AA, accommodative amplitude; AL, axial length; AS, asthenopic symptoms; CHESUD, cumulative hours of electronic screen use per day; CISS, Convergence Insufficiency Symptom Survey; CVS, computer vision syndrome; DED, Dry eye disease; DES, Digital eye strain; MGA, Meibomian gland atrophy; m-SPEED, modified Standardized Patient Evaluation of Eye Dryness questionnaire; NfV, negative fusional weakness; NRA, negative relative accommodation; PD, prism dioptre; rDSER, rate of SER change; SER, spherical equivalent refraction.						

Table 2 Glossary

Term	Definition
Accommodation	Contraction of the ciliary muscle resulting in a change of lens shape. ⁵³
Asthenopia	Subjective symptoms of ocular fatigue or eye strain. ⁵³
Astigmatism	Type of refractive error due to imperfection in the curvature of the eye that causes blurred distance and near vision. ⁵⁴
Cycloplegic refraction	A technique used to calculate the complete refractive error by temporarily paralyzing the ciliary muscle of the eye that aids in focusing. ⁵³
Diplopia	Disorder of vision in which two images of a single object are seen. ⁵³
Dry eye	Alteration of ocular surface homeostasis characterised by an alteration of the tear film.
Emmetropia	Refractive state of an eye in which parallel rays of light entering the eye are focused on the retina, creating an image that is perceived as crisp and in focus. ⁵⁵
Esotropia	Eye misalignment in which one eye is deviated inward, or nasally. ⁵⁴
Hyperopia	Ocular condition in which the refracting power of the eye causes light rays entering the eye to have a focal point that is posterior to the retina while accommodation is maintained in a state of relaxation. ⁵⁴
Myopia	Ocular condition in which the refracting power of the eye causes light rays entering the eye to have a focal point that is anterior to the retina while accommodation is maintained in a state of relaxation. ⁵⁴
Orthokeratology	Use of specially designed and fitted contact lenses to temporarily reshape the cornea to improve vision. ⁵⁶
Refractive errors	Type of vision problem that makes it hard to see clearly and happens when the shape of your eye keeps light from focusing correctly on your retina. ⁵⁵
Spherical equivalent refraction	Estimate of the eyes' refractive error, calculated independently for each eye. It is calculated by merging the spherical (near-sightedness or far-sightedness) and cylindrical (astigmatism) refractive error components. ⁵⁴
Vergence	The turning motion of the eyeballs towards (convergence) or away (divergence) from each other. ⁵³

studies reported worsening of visual symptoms such as vision impairment, asthenopia, dryness, scratchiness, headache, eye redness, eye strain and light sensitivity, among others.^{14 16 18 25 26 33}

Overall, the results of qualitative data syntheses showed a negative effect of the COVID-19 lockdown on visual health in children. Only one of the articles included did not report a deleterious impact of the lockdown on vision.²¹

DISCUSSION

Most of the studies included in this systematic review showed some degree of worsening in visual health in children exposed to virtual learning strategies during the COVID-19 lockdown. The majority of the articles focused on myopia development and progression, and reported a faster onset and progression following the beginning of the lockdown. Also, prolonged exposure to screens was associated with worsened ocular symptoms such as eye strain, blurred vision and redness, as well as an increase in the rate of dry eye, which is traditionally considered to be uncommon in the paediatric population.

Refractive errors

The COVID-19 lockdown impacted the behaviour and daily life of children and teenagers, resulting in increased digital time, near work and decreased outdoor time.³⁴ It is estimated that close to 1.37 billion students worldwide switched to a digital or e-learning school modality during the lockdown.³⁴ These changes have been related to an increase in myopia incidence and progression.³⁴ First, the relationship between near work, especially near reading, and myopia was well established before the COVID-19 pandemic, as stated in the Collaborative Longitudinal Evaluation of Ethnicity and Refractive Error Study.^{34 35} Second, several studies have focused on screen time and its association with myopia development.^{34 36 37} Third, outdoor time has been considered a protective factor against myopia onset. He *et al* showed a 23% reduction in myopia incidence after 40 min of outdoor time daily.^{34 38}

During the COVID-19 pandemic in 2020, Mirhajianmoghdam *et al* assessed subjective and objective measures in 14 myopic and 39 non-myopic children in the USA.³² Initially, parents completed the University of Houston Near Work, Environment, Activity, and Refraction survey in three sessions. The first session included questions related to summer 2020, which was during the COVID-19 pandemic. The second session served to collect data about a typical school period before the COVID-19 pandemic, and the goal of the third session was to collect data about a typical summer period before the pandemic. Later, the investigators used an actigraph device to measure physical activity, sleep and ambient illumination exposure (time spent outdoors) in children for 10 days. The results indicated that all of the children spent less time outdoors during the summer of the pandemic (2020) compared with before the lockdown and showed an increase in daily electronic device use. Furthermore, myopic children had less daily light exposure (183.6 ± 39.3 lux) and spent less time outdoors (0.2 hours/day) during COVID-19 compared with non-myopic children (279.5 ± 23.5 lux, $p=0.04$).³²

The authors of several previous studies have proposed that increased time spent using digital devices is associated with decreased time spent outdoors and impaired retinal dopamine release, which is normally stimulated by daylight exposure. This suppresses axial expansion of the eye, preventing myopia progression.^{39 40} For

instance, Wu *et al* reported that children who spent more than 11 hours/week outdoors had a 53% decrease in myopia progression,⁴¹ and Ip *et al* reported an increased incidence of progression in children living in apartment buildings compared with those living in detached houses.⁴² Additionally, Xu *et al* found that the amount of time spent online was significantly positively associated with an increased incidence of myopia and progression in students.²³ However, not all studies have shown this correlation.²⁰ Aslan and Sahinoglu-Keskek reported that myopia advancement in 2020 was mainly slow (0.31 ± 0.2 D) in most of the children evaluated (49 subjects), followed by moderate progression in 45 children (0.82 ± 0.14 D). The authors found no correlation between myopia progression and digital device time or glasses use.²⁰ Thus, the relationship between myopia progression and digital device use requires further investigation.

The studies by Mirhajianmoghdam *et al* and Aslan and Sahinoglu-Keskek support the findings of myopia progression during the COVID-19 lockdown. For example, Chang *et al* compared myopic progression before, during and after the COVID-19 lockdown in 44 187 students in China by assessing non-cycloplegic autorefractometry and the SER.¹⁵ Four evaluation rounds separated by 6 months during 2019 and 2020 indicated a transitory period of accelerated myopic progression in children that reversed after the lockdown. The mean SER during the pre-pandemic assessment was -0.030 D/month, shortly after the lockdown was -0.074 D/month and later during the lockdown was 0.016 D/month. The proportion of myopic participants was 48% before the lockdown, 45.2% at a second assessment before the lockdown, 73.7% shortly after the lockdown and 67.9% later after the lockdown during rounds 1, 2, 3 and 4, respectively. The authors considered the influence of accommodative spasms and structural changes related to restricted outdoor time, increased screen time and limited indoor space to be the leading cause of the progression. Moreover, they found that younger children were at a higher risk of myopic progression during the lockdown because their lifestyle changes were strongly associated with reduced light exposure, and accordingly, reduced retinal dopamine levels.¹⁵

This is concordant with the findings of Wang *et al*, who reported a substantial decrease in the SER after COVID-19 home confinement, especially for children aged 6 (-0.32 D), 7 (-0.28 D) and 8 (-0.29 D) years, $p < 0.05$.¹⁷ Furthermore, they found myopia development to occur earlier in girls than boys. The prevalence of myopia appeared to be approximately 3 times higher in 2020 than in other years for children aged 6 years, 2 times higher for children aged 7 years and 1.4 times higher for those aged 8 years. This led the authors to hypothesise that younger children are more sensitive to environmental changes than older children.¹⁷ Furthermore, Wang *et al* reported a prevalence of myopia of 39.27% in primary school students, 73.39% in junior school students and 84.89% in high school students, identifying an increase in the rate

of myopia among teenagers in 2020 (55.02%) compared with that in 2019 (44.64%).¹⁹

Ly *et al* investigated the potential impacts of home confinement on myopia progression from the perspective of axial growth length in children undergoing orthokeratology treatment.²¹ They found a monthly axial growth length of 0.023 ± 0.019 mm/month, 0.018 ± 0.021 mm/month and 0.014 ± 0.016 mm/month before, during and after home confinement, respectively. However, the monthly axial growth length before confinement was not significantly different from that after confinement ($p=0.333$), although age was negatively associated with the axial length growth rate during confinement in myopic children.²¹ This coincides with the findings of a previous meta-analysis that suggested that orthokeratology decreases the rate of myopia progression in children.⁴³

In contrast, Alvarez-Peregrina *et al* did not find an increase in the prevalence of myopia among children between 2019 and 2020.³⁰ However, they observed that the percentage of hyperopes decreased, and the percentage of emmetropes increased ($p<0.001$). The average SE value in 2019 was $+0.66\pm 2.03$ D, compared with $+0.48\pm 1.81$ D in 2020 ($p\leq 0.001$). This decrease was significant in children aged 5 years. Additionally, 47% (95% CI 45% to 50%) of children spent less time outdoors in 2020 vs 2019 ($p<0.001$). Children who spent more time outdoors had higher SE values both pre-confinement and post-confinement ($p<0.001$ and $p=0.049$).²⁶ Even though Alvarez-Peregrina *et al* did not demonstrate myopia progression, a reduction in SER is a strong predictive factor for myopia in emmetropic and hyperopic children, as indicated by the Wenzhou Medical University Essilor Progression and Onset of Myopia study.⁴⁴

Accommodation and vergence disturbances

A longer duration of digital device use requires more accommodative effort, and consequently increases the chance of asthenopia symptoms and dysfunctional accommodation and vergence (see table 2, Glossary). Mohan *et al* studied the effects of online classes during the COVID-19 pandemic, and considered the time spent in online classes and using digital devices such as television, video game systems and smartphones. According to the CISS survey, followed by evaluations by an optometrist and paediatric ophthalmologist, 36 out of 46 examined children had symptoms of convergence insufficiency. However, children who attended online classes for less than 4 hours/day exhibited fewer symptoms than those who attended online classes for more than 4 hours/day. Furthermore, near exophoria, near point convergence, positive fusional weakness and accommodation excess were more frequent in children exposed to longer online classes.¹³

Similarly, Hamburger *et al* evaluated ocular symptoms in 110 children who attended virtual school during the COVID-19 pandemic. They found that 61% of the children reported a significant increase in convergence

insufficiency, as evidenced by a higher CISS score after attending online classes.³³

Vagge *et al* reported four cases of children between 4 and 16 years old who developed acute acquired concomitant esotropia after intense digital device use during the COVID-19 lockdown.²⁹ All of the children experienced acute-onset diplopia (see table 2, Glossary) after more than 8 hours/day spent looking at digital screens. Ophthalmological examination reported manifest esotropia from 20 to 35 prism dioptres at far and near distances in all four patients. Two out of the four children presented bilaterally cycloplegic refraction of +1.00 to +2.00 dioptre sphere. One of them presented cycloplegic refraction of -2.50 in the right eye and -2.25 in the left eye, and another presented -0.5 bilaterally.²⁹ Some studies have suggested that digital device-induced esotropia is associated with excessive application of near vision, as well as dynamic activation of the medial rectus muscles when exposed to longer periods of digital screen time. This may affect the near vision triad, that is, the accommodation-convergence reflex: convergence of both eyes, contraction of the ciliary muscle resulting in a change of lens shape (accommodation) and pupillary constriction.^{29 45 46}

Visual symptoms

The increase in digital device use associated with the COVID-19 lockdown and remote learning has precipitated a rise in dry eye symptoms and asthenopia. Hamburger *et al* reported a significant increase in asthenopia symptoms after online classes with discomfort, fatigue and impaired vision as dominant symptoms. Moreover, an increased asthenopia score was identified after online classes in more than half of the children evaluated.³³ Likewise, Li *et al* identified a positive association between screen time and the risk of asthenopia in approximately 25 000 students aged 8–20 years, and attributed a higher risk of asthenopia to conditions such as myopia, astigmatism and mechanical factors like distance from the screen.²⁵

Elhousseiny *et al* reported a significant increase in symptoms such as eye dryness, grittiness and scratchiness associated with prolonged exposure to digital screens for education and leisure purposes in 403 children aged 10–18 years.¹⁸ Similarly, Mohan *et al* identified longer screen time during the COVID-19 lockdown compared with the pre-COVID era in 217 children, of which almost half attended online classes.¹⁴ More than a third of the evaluated children used digital devices for over 5 hours/day, and 50.23% manifested dry eye with itching and headache as predominant symptoms.

Gupta *et al* evaluated 654 students between 5 and 18 years old using the Rasch-based Computer Vision Symptom Scale.¹⁶ The authors reported a significant increase in average digital device exposure during confinement, particularly smartphone, which was greater than 5 hours/day. Visual symptoms in the children were eye redness, eye strain, blurred vision, light sensitivity and heaviness of eyelids.¹⁶ Furthermore, Li *et al* identified a

higher risk of computer vision syndrome in children with myopia with and without correction, astigmatism, fewer outdoor activities and prolonged screen time.²⁶

The relationship between digital screen time and dry eye has already been described in both adults and children, as well as before the global COVID-19 pandemic.^{47–50} Changes in blinking dynamics and ocular surface abnormalities are some of the consequences that arise from intense screen exposure. Regarding ocular surface measures, longer screen time can decrease blinking frequency and completeness, resulting in reduced tear break-up time and tear volume, as well as changes in tear lipid composition.^{6,51} This means that a longer exposure to digital devices can enhance the deterioration of tear film quality, and thus increase the risk of developing dry eye symptoms.⁶

A main limitation of this study is the inclusion of articles with different study designs, as it is difficult to compare them quantitatively and qualitatively. Moreover, the evidence reported in the selected studies was obtained using distinct evaluation methods, from symptom surveys to detailed ophthalmological examinations, influencing the objectiveness of the conclusions obtained. Given that most of the studies were developed specifically in Asian countries, extrapolations to other parts of the world should be made with caution.

CONCLUSIONS

The changes in habits and lifestyles as a result of the COVID-19 pandemic have severely impacted eye health in children. Children attending classes as part of a remote learning strategy had more rapid myopia progression, increased frequency of dry eye and visual fatigue symptoms, and exhibited signs of vergence and accommodation disturbances such as acute acquired concomitant esotropia and convergence insufficiency. Ophthalmologists, paediatricians and general physicians should make themselves aware of the effect of virtual learning on the paediatric population to enable early identification and management of these conditions. In addition, countries around the world must implement public health strategies to mitigate the impacts of a more screen-focused life, especially with respect to conditions as common and costly as myopia. Further studies are required to evaluate the long-term impacts of such changes associated with the COVID-19 pandemic.

Twitter William Rojas-Carabali @warc97

Contributors Conceptualisation: MCC-A, SR-G, WR-C, Ad-I-T, CT-G. Methodology: MCC-A, SR-G, WR-C, CT-G. Investigation: MCC-A, SR-G, WR-C, Ad-I-T, CT-G. Resources: MCC-A, SR-G, WR-C, Ad-I-T, CT-G. Data curation: MCC-A, SR-G, CT-G. Writing—original draft preparation: MCC-A, SR-G, WR-C, Ad-I-T, CT-G. Writing—review and editing: MCC-A, SR-G, WR-C, Ad-I-T, CT-G. Supervision: Ad-I-T, CT-G. Guarantor: CT-G. All authors have read and agreed to the published version of the manuscript.

Funding The review was supported by the Universidad del Rosario. We thank Sydney Koke, MFA, from Edanz (<https://www.edanz.com/ac>) for editing a draft of this manuscript.

Disclaimer The sponsors had no role in the design, data collection or analysis of the study.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available in a public, open access repository.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Sofía Ramírez-Guerrero <http://orcid.org/0000-0001-8927-8284>

William Rojas-Carabali <http://orcid.org/0000-0002-9976-8989>

Alejandra de-la-Torre <http://orcid.org/0000-0003-0684-1989>

Claudia Talero-Gutiérrez <http://orcid.org/0000-0003-1601-5015>

REFERENCES

- Arora T, Grey I. Health behaviour changes during COVID-19 and the potential consequences: a mini-review. *J Health Psychol* 2020;25:1155–63.
- Almarzooq ZI, Lopes M, Kochar A. Virtual learning during the COVID-19 pandemic. *J Am Coll Cardiol* 2020;75:2635–8.
- Enthoven CA, Tideman JW, Polling JR, et al. The impact of computer use on myopia development in childhood: the generation R study. *Prev Med* 2020;132:105988.
- Fan Q, Wang H, Kong W, et al. Online learning-related visual function impairment during and after the COVID-19 pandemic. *Front Public Health* 2021;9:645971.
- Holden BA, Fricke TR, Wilson DA, et al. Global prevalence of myopia and high myopia and temporal trends from 2000 through 2050. *Ophthalmology* 2016;123:1036–42.
- Al-Mohtaseb Z, Schachter S, Shen Lee B, et al. The relationship between dry eye disease and digital screen use. *Clin Ophthalmol* 2021;15:3811–20.
- Napoli PE, Nioi M, Fossarello M. The ‘quarantine dry eye’: the lockdown for coronavirus disease 2019 and its implications for ocular surface health. *Risk Manag Healthc Policy* 2021;14:1629–36.
- Inomata T, Iwagami M, Nakamura M, et al. Characteristics and risk factors associated with diagnosed and undiagnosed symptomatic dry eye using a smartphone application. *JAMA Ophthalmol* 2020;138:58–68.
- Ma L-L, Wang Y-Y, Yang Z-H. Methodological quality (risk of bias) assessment tools for primary and secondary medical studies: what are they and which is better? *Mil Med Res* 2020;7.
- Hoy D, Brooks P, Woolf A, et al. Assessing risk of bias in prevalence studies: modification of an existing tool and evidence of interrater agreement. *J Clin Epidemiol* 2012;65:934–9.
- Simo LP, AgborNdi E, et al. Prevalence and determinants of anaemia in children aged 6–59 months in Africa: a protocol for systematic review and meta-analysis. *BMJ Open* 2020;10:e032042.
- Murad MH, Sultan S, Haffar S, et al. Methodological quality and synthesis of case series and case reports. *BMJ Evid Based Med* 2018;23:60–3.
- Mohan A, Sen P, Shah C, et al. Binocular accommodation and vergence dysfunction in children attending online classes during the COVID-19 pandemic: digital eye strain in kids (DESK study-2). *J Pediatr Ophthalmol Strabismus* 2021;58:224–31.

- 14 Mohan A, Sen P, Shah C, *et al.* Prevalence and risk factor assessment of digital eye strain among children using online e-learning during the COVID-19 pandemic: digital eye strain among kids (DESK study-1). *Indian J Ophthalmol* 2021;69:140–4.
- 15 Chang P, Zhang B, Lin L, *et al.* Comparison of myopic progression before, during, and after COVID-19 lockdown. *Ophthalmology* 2021;128:1655–7.
- 16 Gupta R, Chauhan L, Varshney A. Impact of e-schooling on digital eye strain in coronavirus disease era: a survey of 654 students. *J Curr Ophthalmol* 2021;33:158–64.
- 17 Wang J, Li Y, Musch DC, *et al.* Progression of myopia in school-aged children after COVID-19 home confinement. *JAMA Ophthalmol* 2021;39:293–300.
- 18 Elhousseiny AM, Eleiwa TK, Yacoub MS, *et al.* Relationship between screen time and dry eye symptoms in pediatric population during the COVID-19 pandemic. *Ocul Surf* 2021;22:117–9.
- 19 Wang W, Zhu L, Zheng S, *et al.* Survey on the progression of myopia in children and adolescents in Chongqing during COVID-19 pandemic. *Front Public Health* 2021;9:646770.
- 20 Aslan F, Sahinoglu-Keskek N. The effect of home education on myopia progression in children during the COVID-19 pandemic. *Eye* 2022;36:1427–32.
- 21 Lv H, Wang Y, Sun S, *et al.* The impact of COVID-19 home confinement on axial length in myopic children undergoing orthokeratology. *Clin Exp Optom* 2022;1–5.
- 22 Ma D, Wei S, Li S-M, *et al.* The impact of study-at-home during the COVID-19 pandemic on myopia progression in Chinese children. *Front Public Health* 2021;9:720514.
- 23 Xu L, Ma Y, Yuan J, *et al.* COVID-19 quarantine reveals that behavioral changes have an effect on myopia progression. *Ophthalmology* 2021;128:1652–4.
- 24 Hu Y, Zhao F, Ding X, *et al.* Rates of myopia development in young Chinese schoolchildren during the outbreak of COVID-19. *JAMA Ophthalmol* 2021;139:1115–21.
- 25 Li L, Zhang J, Chen M, *et al.* Contribution of total screen/online-course time to asthenopia in children during COVID-19 pandemic via influencing psychological stress. *Front Public Health* 2021;9:736617.
- 26 Li R, Ying B, Qian Y, *et al.* Prevalence of self-reported symptoms of computer vision syndrome and associated risk factors among school students in China during the COVID-19 pandemic. *Ophthalmic Epidemiol* 2021;1–11.
- 27 Mohan A, Sen P, Mujumdar D, *et al.* Series of cases of acute acquired comitant esotropia in children associated with excessive online classes on smartphone during COVID-19 pandemic; digital eye strain among kids (DESK study-3). *Strabismus* 2021;29:163–7.
- 28 Mohan A, Sen P, Peeush P, *et al.* Impact of online classes and home confinement on myopia progression in children during COVID-19 pandemic: digital eye strain among kids (DESK study 4). *Indian J Ophthalmol* 2022;70:241.
- 29 Vagge A, Giannaccare G, Scarinci F, *et al.* Acute acquired concomitant esotropia from excessive application of near vision during the COVID-19 lockdown. *J Pediatr Ophthalmol Strabismus* 2020;57:e88–91.
- 30 Alvarez-Peregrina C, Martinez-Perez C, Villa-Collar C. Impact of COVID-19 home confinement in children's refractive errors. *Int J Environ Res Public Health* 2021;18.
- 31 Cremers SL, Khan ARG, Ahn J, *et al.* New indicator of children's excessive electronic screen use and factors in meibomian gland atrophy. *Am J Ophthalmol* 2021;229:63–70.
- 32 Mirhajianmoghadam H, Piña A, Ostrin LA. Objective and subjective behavioral measures in myopic and non-myopic children during the COVID-19 pandemic. *Transl Vis Sci Technol* 2021;10:4.
- 33 Hamburger JL, Lavrich JB, Rusakevich AM, *et al.* The visual consequences of virtual school: acute eye symptoms in healthy children. *J AAPOS* 2022;26:2.e1–2.e5.
- 34 Wong CW, Tsai A, Jonas JB, *et al.* Digital screen time during the COVID-19 pandemic: risk for a further myopia boom? *Am J Ophthalmol* 2021;223:333–7.
- 35 Jones-Jordan LA, Mitchell GL, Cotter SA, *et al.* Visual activity before and after the onset of juvenile myopia. *Invest Ophthalmol Vis Sci* 2011;52:1841–50.
- 36 Mccrann S, Loughman J, Butler JS, *et al.* Smartphone use as a possible risk factor for myopia. *Clin Exp Optom* 2021;104:35–41.
- 37 Lanca C, Saw S-M. The association between digital screen time and myopia: a systematic review. *Ophthalmic Physiol Opt* 2020;40:216–29.
- 38 He M, Xiang F, Zeng Y, *et al.* Effect of time spent outdoors at school on the development of myopia among children in China: a randomized clinical trial. *JAMA* 2015;314:1142–8.
- 39 McCarthy CS, Megaw P, Devadas M, *et al.* Dopaminergic agents affect the ability of brief periods of normal vision to prevent form-deprivation myopia. *Exp Eye Res* 2007;84:100–7.
- 40 Feldkaemper M, Schaeffel F. An updated view on the role of dopamine in myopia. *Exp Eye Res* 2013;114:106–19.
- 41 Wu P-C, Chen C-T, Lin K-K, *et al.* Myopia prevention and outdoor light intensity in a school-based cluster randomized trial. *Ophthalmology* 2018;125:1239–50.
- 42 Ip JM, Saw S-M, Rose KA, *et al.* Role of near work in myopia: findings in a sample of Australian school children. *Invest Ophthalmol Vis Sci* 2008;49:2903–10.
- 43 Si J-K, Tang K, Bi H-S, *et al.* Orthokeratology for myopia control: a meta-analysis. *Optom Vis Sci* 2015;92:252–7.
- 44 Wong YL, Yuan Y, Su B, *et al.* Prediction of myopia onset with refractive error measured using non-cycloplegic subjective refraction: the WEPrOM study. *BMJ Open Ophthalmol* 2021;6:e000628.
- 45 Lee HS, Park SW, Heo H. Acute acquired comitant esotropia related to excessive smartphone use. *BMC Ophthalmol* 2016;16:37.
- 46 Motlagh M, Geetha R. *Physiology, accommodation*. Treasure Island, FL: StatPearls Publishing, 2022.
- 47 Uchino M, Yokoi N, Uchino Y, *et al.* Prevalence of dry eye disease and its risk factors in visual display terminal users: the Osaka study. *Am J Ophthalmol* 2013;156:759–66.
- 48 Hanyuda A, Sawada N, Uchino M, *et al.* Physical inactivity, prolonged sedentary behaviors, and use of visual display terminals as potential risk factors for dry eye disease: JPHC-NEXT study. *Ocul Surf* 2020;18:56–63.
- 49 Bhattacharya S, Saleem SM, Singh A. Digital eye strain in the era of COVID-19 pandemic: an emerging public health threat. *Indian J Ophthalmol* 2020;68:1709–10.
- 50 Rojas-Carabali W, Uribe-Reina P, Muñoz-Ortiz J, *et al.* High prevalence of abnormal ocular surface tests in a healthy pediatric population. *Clin Ophthalmol* 2020;14:3427–38.
- 51 Rossi GCM, Scudeller L, Bettio F, *et al.* Prevalence of dry eye in video display terminal users: a cross-sectional Caucasian study in Italy. *Int Ophthalmol* 2019;39:1315–22.
- 52 Moher D, Liberati A, Tetzlaff J, *et al.* Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6:e1000097.
- 53 Merriam-Webster. Medical dictionary, 2022. Available: <https://www.merriam-webster.com/>
- 54 EyeWiki. American Academy of ophthalmology (online). Available: https://eyewiki.aao.org/Main_Page
- 55 Palay D, Krachmer J. *Primary care ophthalmology*. Mosby, 2005: 1–23.
- 56 Mukamal R. What is orthokeratology (online), 2018. Available: <https://www.aao.org/eye-health/glasses-contacts/what-is-orthokeratology>