



Deciphering the myth of icariin and synthetic derivatives in improving erectile function from a molecular biology perspective: a narrative review

Yuanjie Niu^{1,2#}, Guiting Lin^{3#}, Jiancheng Pan^{1,2#}, Jihong Liu^{4,5}, Yongde Xu⁶, Qiliang Cai^{1,2}, Tao Wang^{4,5}, Yang Luan^{4,5}, Yegang Chen^{1,2}, Yuhong Feng^{1,2}, Xiaoqing Yang², Wenjie Tian⁷, Wong Jin Bae⁷, Ruili Guan⁸, Zhongcheng Xin^{1,2,8}

¹Department of Urology, Tianjin Institute of Urology, The Second Hospital of Tianjin Medical University, Tianjin, China; ²China-Korea Joint Research Center for Male Reproductive and Sexual Medicine, Institute of Urology, Tianjin, China; ³Knappe Molecular Urology Laboratory, Department of Urology, School of Medicine, University of California, San Francisco, CA, USA; ⁴Department of Urology, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China; ⁵Institute of Urology, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China; ⁶Department of Urology, Beijing Friendship Hospital, Capital Medical University, Beijing, China; ⁷Department of Urology, Seoul St. Mary's Hospital, the Catholic University of Korea, Seoul, Korea; ⁸Andrology Center, Peking University First Hospital, Peking University, Beijing, China

Contributions: (I) Conception and design: Z Xin, R Guan; (II) Administrative support: Z Xin, Y Niu, G Lin, J Liu, WJ Bae; (III) Provision of study materials or patients: J Pan, Y Xu, Q Cai, T Wang; (IV) Collection and assembly of data: J Pan, Y Feng; (V) Data analysis and interpretation: J Pan, Y Chen, Y Lun, X Yang, W Tian; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

[#]These authors contributed equally to this work.

Correspondence to: Ruili Guan. Molecular Biology Laboratory of Andrology Center, Peking University First Hospital, Peking University, Beijing, China. Email: guanruili@bjmu.edu.cn; Zhongcheng Xin. Male reproductive and sexual medicine, department of urology, the second hospital of tianjin medical university, Tianjin, China; China-Korea Joint Research Center for Male Reproductive and Sexual Medicine, Institute of Urology, Tianjin, China; Molecular Biology Laboratory of Andrology Center, Peking University First Hospital, Peking University, Beijing, China. Email: xinzc@bjmu.edu.cn.

Background and Objective: Although epimedium herb (EH) has been widely used in ancient Chinese medicine to enhance sexual activity, its pharmacological mechanism is not clear. Modern studies have shown that epimedium herb is rich in icariin (ICA, a flavonoid compound), and 91.2% of icariin is converted to icariside II (ICA II) by hydrolytic enzymes in intestinal bacteria after oral administration. YS-10 is a synthetic derivative of icariside II. The aim of this review was to summarize the contemporary evidence regarding the pharmacokinetics, therapeutic properties, and molecular biological mechanisms of ICA and some ICA derivatives for erectile dysfunction therapy.

Methods: A detailed search was conducted in the PubMed database using keywords and phrases, such as “icariin” AND “erectile dysfunction”, “icariside II” AND “erectile dysfunction”. The publication time is limited to last 20 years. Articles had to be published in peer reviewed journals.

Key Content and Findings: ICA and its some derivatives showed the specific inhibition on phosphodiesterase type 5 (PDE5) and the promotion of testosterone synthesis. In addition, by regulating various reliable evidence of signaling pathways such as PI3K/AKT, TGFβ1/Smad2, p38/MAPK, Wnt and secretion of various cytokines, ICA and ICA derivatives can activate endogenous stem cells (ESCs) leading to endothelial cell and smooth muscle cell proliferation, nerve regeneration and fibrosis inhibition, repair pathological changes in penile tissue and improve erectile function.

Conclusions: ICA and some of its derivatives could be a potential treatment for restoring spontaneous erections. In addition ICA and his derivatives may also be valuable as a regenerative medicine approach for other diseases, but more clinical and basic researches with high quality and large samples are recommended.

Keywords: Epimedium herb (EH); icariin; erectile dysfunction; endogenous stem cells; molecular mechanism

Submitted Mar 28, 2022. Accepted for publication Jun 02, 2022.

doi: 10.21037/tau-22-232

View this article at: <https://dx.doi.org/10.21037/tau-22-232>

Introduction

Erectile dysfunction (ED) is defined as the inability to obtain and/or maintain a sufficient erection for satisfying sex over three months. Currently, oral administration of type 5 phosphodiesterase inhibitors (PDE5Is) given before intercourse are the first-line treatment for ED patients (1). Second-line treatments for ED, including the vacuum devices and intracavernose injection therapy (ICI), present numerous complications like penile pain, priapism, and fibrosis. The overall efficiency of both was about 70%, however, these treatments were only used to relieve ED symptoms. The implantation of a penile prosthesis might be the last option, but is highly cost with the risks of infection, erosion and device failure (2). Recent researches on ED regenerative therapies including gene therapy and stem cell therapy have shown significant benefits in repairing pathological changes of penile tissue of various ED animal models and are important attempts to restore natural erections. However, their clinical application is limited by ethical and potential safety issues (3). Anyhow, a natural erection restoration is always the primary pursuit.

Traditional Chinese medicine has a long history in exploring ED therapy. Many studies have demonstrated the effectiveness of Epimedium for the treatment of ED. The history of Epimedium in ED therapy can be traced back to the ancient Chinese North and South Dynasties (420–589 AD). Tao Hongjing, a famous medical scientist, learned from the shepherds that male sheep consumed a plant that significantly increased the times of penile erections and mating. Tao believed this plant could enhance the “YANG” energy (in Chinese, “Yin Yang Huo”, in English, Epimedium Herb, EH). It has also been widely used in East Asian countries for centuries, with different names such as Epimedium Brevicornum Maxim, E. Sagittatum Maxim, E. Pubescens Maxim, and E. Koreanu Nakai.

Flavonoids have been reported to be the main bioactive components of Epimedium (4). Icariin (ICA) is one of the most abundant flavonoids in Epimedium and is often used as a marker for quality control in Epimedium herbal preparation and chemical taxonomy (5,6). ICA has a similar

structure to PDE5Is. In addition, it has a wide range of other pharmacological effects, including anticancer activity (7), anti-osteoporotic activity (8,9), antidepressant activity (10,11), and aphrodisiac (12). ICA is a disaccharide, and several studies have shown that flavonoids in the form of glycosides have low bioactivity because of their low intrinsic absorption permeability (13–15). Both *in vivo* and *in vitro* metabolic studies demonstrated that the intestinal microbiota can convert ICA into icariside I, icariside II (ICAI) by hydrolyzing C-3-O-rhamnoside (R1) and C-7-O-glucopyranoside (R2) molecule, respectively (16–19). Pharmacokinetic studies have shown that the intestinal bacterial metabolites of ICA have better biological activity (20,21). Previous studies found that after oral administration, 91.2% of ICA was converted to ICA II. The maximum blood concentration (C_{max}) and the degree of absorption (AUC_{0-t}) that occurred after ICA II administration were 3.8 and 13.0 times higher than those of ICA, respectively. Therefore, ICA II, which lacks a glucose group at C-7, is the major metabolite of ICA and has higher activity than ICA (22).

By administering different doses of ICA or ICA II treatment, it was found that appropriate doses (generally low dose, ranging from 10–200 mg/kg) significantly improved erectile function in various ED animal models. Moreover, ICA and ICAII are able to reverse the injury of penile corpus cavernosum, such as restoring the content of cavernous endothelium (23) and smooth muscle (24), regenerating damaged cavernous nerves (25), inhibiting cavernous fibrosis (26), restoring the normal level of testosterone and the ratio of extracellular matrix (27,28). Therefore, ICA/ICAII may have the potential to rehabilitate the pathological injury and achieve spontaneous erection.

In current review, we summarized the advancements in terms of the biological effect of ICA and ICA derivatives on type 5 phosphodiesterase (PDE5) expression and enzymatic function, the modulation of tissue resident stem cells to regenerate penile damaged tissues, and the restoration of testosterone level. We present the following article in accordance with the Narrative Review reporting

Table 1 The search strategy summary

Items	Specification
Date of search (specified to date, month and year)	January 1, 2003 to March 1, 2022
Databases and other sources searched	PubMed
Search terms used (including MeSH and free text search terms and filters) (Note: please use an independent supplement table to present detailed search strategy of one database as an example)	As shown in Table S1
Timeframe	The publication time is limited to last 20 years
Inclusion and exclusion criteria (study type, language restrictions etc.)	<ol style="list-style-type: none"> 1. Articles had to be published in peer reviewed journals 2. Articles without experimental data related to the search terms and data that did not support relevant conclusions were excluded, as well as articles with low quality 3. Written by English
Selection process (who conducted the selection, whether it was conducted independently, how consensus was obtained, etc.)	Two experimenters were searched and screened separately. Discussion of the discrepancy literature and invitation to the other experimenter to participate
Any additional considerations, if applicable	None

checklist (available at <https://tau.amegroups.com/article/view/10.21037/tau-22-232/rc>).

Methods

We conducted a literature search for PUBMED in March 2022 following the *Table 1* search strategy. The publication time is limited to last 20 years. Articles had to be published in peer reviewed journals. Keyword searches including “erectile dysfunction” AND “icariin”, “erectile dysfunction” AND “icariside II”, “erectile dysfunction” AND “icariin derivative”, “PDE5” AND “icariin”, “PDE5” AND “icariside II”, “PDE5” AND “icariin derivative”, “testis” AND “testosterone” AND “icariin”, and “testis” AND “testosterone” AND “icariside II”, “testis” AND “testosterone” AND “icariin derivative”. Articles without experimental data related to the search terms and data that did not support relevant conclusions were excluded, as well as articles with low quality. The data were extracted from each study by two investigators (JCP and YHF) independently. After the search, summarized the literature and discussed the different literature. Another researcher (ZCX) was invited to discuss and identify the literature on which consensus could not be reached ([Table S1](#)). Our literature search covered English-language clinical or basic research articles published from January 1, 2003 to March 1,

2022, as well as related reviews and meta analyses.

Discussions

Modulating PDE5 expression and function

Numerous studies have shown that ICA and ICA derivatives affected the enzymatic function of PDE5 mainly through competitive inhibition, while the possible regulatory role of PDE5 expression needs to be further investigated (*Table 2*). Protein binding of ICA/ICA derivatives and PDE5.

The catalytic domain of PDE5 can and the active site, which realizes the role of bind and hydrolyze cyclic guanosine monophosphate (cGMP), while the H-loop structure can modulate the enzyme affinity. Crystal structure of ICAII binding to PDE5 shows that the 7-o-glucose structure is located near the entrance of the active binding site and forms hydrogen bonds with ser668 residues on the flexible H-loop. In addition, the 3-o-rhamnose structure is located in the hydrophobic region that may affect the catalytic efficiency of the enzyme (29,37). In 2019, Chau *et al.* (30) demonstrated that different functional groups at the 3-o and 7-o positions of the ICA derivative backbone play a role in modulating the ability of ICA to inhibit PDE5. The substitution of the 3-o and 7-o positions using different hydrophilic and hydrophobic groups revealed that the hydrophobic group at the 3-o position had a more

Table 2 PDE5 inhibition by ICA /ICA derivatives and mechanism

ICA and ICA derivative	Year of publication	Study type	Mechanism of PDE5 inhibition	The inhibition Effect for PDE5	Key reference
ICAI	2006	<i>In vitro</i> (crystal structure and enzyme kinetics)	Bonding with PDE5A1 flexible H-loop	Selectively inhibiting PDE5A1 with an IC50 of 2M	(29)
ICA derivative-Compounds 3	2019	<i>In vitro</i> (crystal structure and enzyme kinetics)	The hydrophobic group at the 3-o position more significantly inhibiting PDE5	Inhibiting PDE5 with an IC50 of 0.083±0.01 µM	(30)
ICA	2003	<i>In vitro</i> (enzyme kinetics)	–	Inhibiting PDE5 with an IC50 of 0.432 µM	(31)
ICA	2006	<i>In vitro</i> (rat cavernous smooth muscle cells and enzyme kinetics)	–	Inhibiting PDE5A1, A2, and A3 with an IC50 value of 1.0, 0.75, and 1.1 M	(32)
ICAI	2012	<i>In vitro</i> (rat corpus cavernosum tissue and enzyme kinetics)	–	about 50% of Sildenafil	(33)
3,7-bis(2-hydroxyethyl) Icaritin	2008	<i>In vitro</i> (enzyme kinetics)	–	With a similar IC50 to that of sildenafil (IC50 75 vs. 74 nM)	(34)
ICA	2006	<i>In vitro</i> (rat corpus cavernosum tissue and enzyme kinetics)	Inhibit PDE5 mRNA expression	Inhibiting PDE5 with EC50 was 4.62 micromol/L	(35)
ICA	2014	<i>In vivo</i> (rat corpus cavernosum tissue and enzyme kinetics)	Inhibit PDE5 expression	–	(36)

PDE5, phosphodiesterase type 5; ICA, icariin; ICAI, icariside II; IC50, the half maximal inhibitory concentration; EC50, concentration for 50% of maximal effect.

pronounced effect on the inhibition of PDE5.

Enzymatic inhibition of PDE5 function by ICA and ICA derivatives

A previous study showed that ICA exhibited a dose-dependent inhibition of PDE5 activity (31). The half-maximal inhibitory concentrations (IC50) of ICA for PDE5 and phosphodiesterase-4 (PDE4) were 0.432 mmol/L and 73.50 mmol/L, respectively. Ning and colleagues (32) also reported that ICA inhibited all three PDE5 isomers with similar IC50 values. However, the inhibitory effect of ICA on PDE5 is about one-tenth that of sildenafil, while the effect of ICA II is significantly higher than ICA, about 50% of sildenafil (33).

In addition, Dell'Agli *et al.* (34) showed that 3,7-bis(2-hydroxyethyl) icaritin has a strong inhibitory effect on PDE5A1 with a similar IC50 to that of sildenafil (IC50 75 vs. 74 nM) and 80 times than ICA. Therefore, chemical modifications of the ICA molecular structure have been investigated with the aim of obtaining more specific PDE5 inhibitory activity. Recently, Chau *et al.* (30) developed several novel semi-synthetic ICA derivatives, among

which compounds 3 and 7 synthesized by modification with hydroxyethyl substitution of C-3-O-rhamnoside showed specific PDE5 inhibitory activity close to that of commercially available PDE5 inhibitors. Additionally, compound 3 showed less phosphodiesterase-6C (PDE6C) inhibitory effect compared to sildenafil. Compared to naturally derived ICA, synthetic and chemically modified ICA derivatives have good PDE5 inhibition and specificity, and may be promising for further development as better PDE5Is candidates. However, studies on the efficacy of ICA chemically modified compounds are limited to *in vitro* experiments, while safety needs to be verified in more animal experiments and even clinical trials.

ICA regulates PDE5 expression

In contrast to commercial PDE5Is such as sildenafil, a previous study suggested that ICA may have the ability to inhibit PDE5 expression. Jiang *et al.* (35) compared the differences of cGMP, cyclic adenosine monophosphate (cAMP) and PDE5 mRNA by exposing isolated rabbit penile corpus cavernosum to ICA solution and sildenafil solution. The results showed an increase in tissue cGMP

levels in both solutions, with no significant change in cAMP. The term-half maximal effective concentration (EC₅₀) was 4.62 (ICA) and 0.42 (Sildenafil) $\mu\text{mol/L}$ respectively. More importantly, ICA treatment also differentially inhibited PDE5A1 and PDE5A2 mRNA levels in rat penile corpus cavernosum compared with sildenafil. Another study in which ICA was administered to adult rats showed that ICA significantly reduced PDE5 levels in penile tissues, and these results may be related to the reduction of Rho signaling pathway associated coiled-coil containing protein kinase 1 and 2 (ROCK1 and ROCK2) levels by ICA (36). Thus, ICA treatment is considered to have a longer-term PDE5 inhibitory effect than commercially available PDE5Is. This study suggests another possible direction for future research. However, there are only a few studies on the regulation of PDE5 expression by ICA, and definite conclusions need to be confirmed by clear mechanistic studies and experiments in large sample to exclude distortion of results due to individual differences.

Repair of penile tissue mainly through regulation of tissue-resident stem cells

In fact, the majority of ED patients are not satisfied with one-time symptom improvement and want new therapies that are safer and more effective in restoring erectile function by repairing the pathological changes in ED. Numerous studies have demonstrated the effectiveness of ICA and its derivatives in repairing injured penile tissue. The above-mentioned effect through the regulation of tissue-resident stem cell proliferation and differentiation is one of the most important mechanisms currently being investigated for the treatment of ED (Table 3). Therefore, ICA and its derivatives are considered to be regenerative medicine therapies with the potential to restore spontaneous erections in patients with ED.

Tissue resident stem cells activation

Stem cells can self-replicate and differentiate into multiple cell types, which are widely used in regenerative medicine research. Recently, many studies have shown that ICA and ICAII can regulate the biological behaviors of a variety of stem cells, including bone marrow mesenchymal stem cells (BMSCs), adipose-derived stem cells (ADSCs), neural stem cells (NSC), endothelial progenitor cells, human umbilical cord mesenchymal stem cells (53-56). Previous studies have demonstrated that ICA could induce BMSC proliferation, differentiation, and ameliorate prednisolone-induced BMSC

apoptosis *in vitro* and *in vivo* (57,58). Ye *et al.* (54) reported that diabetic ED rats treated with ADSCs and ICA showed significant improvement in the intracranial pressure (ICP) and ICP/mean arterial pressure (MAP) values. Besides, ICA application increased the survival rate of transplanted ADSCs and repaired the damaged structures of the corpus cavernosum (59). Exogenous stem cell transplantation combined with ICA is effective in treating ED methods, but the escape of stem cells after transplantation and proliferation can lead to side effects such as pulmonary embolism, which has hindered the clinical application (60).

In contrast, endogenous stem cells residing in tissues are safer. Endogenous stem cells (ESCs) or progenitor cells (EPCs), also known as tissue resident stem cells, are present in many organs. In recent years, Scholars are gradually turning their attention to endogenous stem cell research in the penis, and some breakthroughs have been achieved. The label-retaining cell (LRC) strategy is a commonly used technique for identification of stem cells in tissue (61,62). Its mechanism is based on the principle that the rapidly proliferating cells will lose the cell label in a short period of time, while quiescent cells and slow-cycling cells will retain the label for a longer period. A recent research had identified potential stem/progenitor cells in the penis by using a co-localization strategy of 5-ethynyl-2-deoxyuridine (EDU)-LRC and cell differentiation markers (63). Treatment with ICAII in obese and cavernous nerve-injured rats showed that penile stem cell mitosis was significantly increased and multidirectional differentiation occurred to repair histopathological changes in the penile corpus cavernosum (64,65). Researches on the molecular mechanisms involved has increased significantly in recent years. Wingless-type MMTV integration site family (Wnt)/ β -catenin, p38 mitogen-activated protein kinase (p38 MAPK), transforming growth factor- β (TGF- β)/Smad, phosphatidylinositol 3-kinase (PI3K)/serine/threonine-specific protein kinase Akt (Akt) and mammalian target of rapamycin (mTOR) signaling pathways are well-documented signaling pathways in the study of ICA-regulated stem cells (Figure 1) (38-41,66-70). Therefore, ICAII may have a role in regulating penile stem cells to repair pathological damage to the corpus cavernosum and restore erectile function.

It should first be stated that the pathophysiology of ED is often interconnected and coexistent, including loss of smooth muscle and endothelial cells, abnormal collagen ratios, and reduction of neural nitric oxide synthase (nNOS)-positive nerves, and that the role of ICA in

Table 3 ICA /ICA derivatives repair injured penile tissue and modulate tissue endogenous stem cells

ICA and ICA derivative	Year of publication	Study design	Study type	Stem cell type	Treatment results	Molecular Mechanisms	Effective dose	STAIR list score (full 7 points)	Key reference
ICAll	2013	Streptozotocin-induced diabetic ED rats	<i>In vivo</i>	SMCs	Increased SMC proliferation and decreased the numbers of autophagosomes	Up-regulated NO-cGMP and downregulated mTOR pathway	10 mg/kg/day	4	(24)
ICA	2017	Male rats with bilateral cavernous nerves injury	<i>In vivo</i>	NSCs	Repaired the damaged neural pathway for erection; Promoted differentiation of endogenous stem cells to Schwann cells	-	1.5 mg/kg/day	5	(25)
ICA	2011	Streptozotocin-induced diabetic ED rats	<i>In vivo</i>	-	Promoted smooth muscle/collagen ratio and endothelial cell content in the corpora cavernosa	up-regulated vWF and PECAM; down-regulated TGFβ1/Smad2 signaling pathway	1, 5, and 10 mg/kg/day	5	(26)
ICAll	2018	Zucker Fatty (ZUC-Lep ^{fa} 185; ZF) male rats	<i>In vivo</i>	-	Prevented penile smooth muscle atrophy, endothelial dysfunction, and lipid accumulation Activated more penile stem cells to proliferate and differentiate	-	1.5 mg/kg/day	4	(38)
ICAll	2014	Male rats with bilateral cavernous nerves injury	<i>In vivo</i>	-	Prevented distortion of normal neural anatomy, smooth muscle atrophy, and collagen deposition of penile Promoted differentiation of penile endogenous stem cells	-	0.5, 1.5 or 4.5 mg/kg/day	4	(39)
ICAll	2016	Streptozotocin-induced diabetic ED rats	<i>In vivo</i>	-	Increased density of dorsal nerve bundle of penile	Increased NGF expression	5 mg/kg/day	5	(40)
YS 10	2021	Male rats with bilateral cavernous nerves injury	<i>In vivo</i>	-	Repaired corpora cavernous nerves injury	Increased β-Catenin and cyclin D1 expression	2.5 mg/kg/day	5	(41)
ICA	2018	NSCs isolated from rats and EdU labeled	<i>In vitro</i>	NSCs	Increased the number of stem cell spheres	Increased mRNA and protein expression of cell cycle genes cyclin D1 and p21	50 & 100 μmol/L	-	(42)
ICA	2014	NSCs isolated from mice embryos and EdU labeled	<i>In vitro</i>	NSCs	Increased neurosphere formation and NSCs with EdU	Activated ERK/MAPK pathway	100 μmol/L	-	(43)
ICA	2016	NSCs isolated from 16-20-week human fetuses	<i>In vitro</i>	NSCs	Enhanced NSCs proliferation and neurosphere formation	Upregulated Frizzled 7, DVL3, FGFR1 and down-regulated GSK-3β	10 μmol/L	-	(44)

Table 3 (continued)

Table 3 (continued)

ICA and ICA derivative	Year of publication	Study design	Study type	Stem cell type	Treatment results	Molecular Mechanisms	Effective dose	STAIR list score (full 7 points)	Key reference
ICA	2020	Aβ25-35-treated hippocampal neural stem cells of rats	<i>In vitro</i>	NSCs	promoted the viability and differentiation into neurons and astrocytes	Activated BDNF-TrkB-ERK/Akt signaling pathway	20, 40, 80 μmol/L	-	(45)
ICA	2022	Chronic unpredictable mild stress for depression rat; Corticosterone treated NSCs from rats embryonic hippocampi	<i>In vivo</i> & <i>In vitro</i>	NSCs	Alleviated dysfunctional neurogenesis and neuronal loss Promoted neuronal proliferation and differentiation	Down-regulated Rps4x, Rps12, Rps14, Rps19, Hsp90b1, and Hsp90aa1 and up-regulated Htra1	120 mg/kg/day <i>in vivo</i> and 20 μmol/L <i>s in vitro</i>	5	(46)
ICAll	2012	Streptozotocin-induced diabetic ED rats	<i>In vivo</i>	-	Promoted corpus cavernosum smooth muscle/collagen ratio and endothelial cell content	Down-regulated TGFβ1/Smad2/CTGF and up-regulated NO-cGMP	1, 5, and 10 mg/kg/day	4	(47)
ICA	2019	High glucose-induced rats bone marrow derived EPCs	<i>In vitro</i>	EPCs	Partially restored EPCs migration and tube formation	Inhibited p38/CREB pathway and activated Akt/eNOS/NO pathway	1 μmol/L	-	(48)
ICA	2015	H2O2-induced rats bone marrow derived EPCs	<i>In vitro</i>	EPCs	Promoted cell migration and capillary tube formation, abrogated apoptotic and autophagic programmed cell death	Reduced ROS levels and restored ΔYm; activated rapamycin / p70S6K/4EBP1 and increased ATF2 and ERK1/2 protein levels	7.5, 15, and 30 μM	-	(49)
ICAll	2020	Streptozotocin-induced diabetic ED rats	<i>In vivo</i>	SMCs	Increased smooth muscle cell/collagen fibril proportions, decreased mitochondrial autophagy, and AGE concentrations	-	10 mg/kg/day	4	(50)
ICA	2020	Monocrotaline-Induced Pulmonary Arterial rats	<i>In vivo</i>	-	Decreased right ventricular systolic pressure (RVSP) and the right ventricular hypertrophy index (RI)	Inhibited TGF-β1, Smad2/3, P-Smad2/3, and MMP2 expressions	50 or 100 mg/kg/day	4	(51)
ICAll	2018	Hemorrhage injection model of subarachnoid hemorrhage rats	<i>In vivo</i>	-	Inhibited subarachnoid fibrosis, attenuated ventriculomegaly, and chronic hydrocephalus	Inhibited TGF-β1/Smad/CTGF signaling pathway	1, 5, 10 mg/kg/day	4	(52)

ICA, icarilin; ICAll, icarilide II; NSCs, neural stem cell; EPCs, endothelial progenitor cells; SMCs, smooth muscle cells; TGF-β1, transforming growth factor β1; Smad2, SMAD family member 2; CREB, cAMP response element binding protein; AGE, advanced glycation end products; ERK, extracellular signal-regulated kinase; Akt, protein kinase B; MAPK, mitogen-activated protein kinase; NO, nitric oxide; eNOS, nitric oxide synthase; Frizzled 7, frizzled class receptor 7; DVL3, dishevelled segment polarity protein 3; FGFR1, fibroblast growth factor receptor 1; GSK-3β, Glycogen synthase kinase-3β; ΔYm, mitochondrial membrane potential; ox-LDL, oxidized low-density lipoprotein; EdU, 5-ethynyl-2-deoxyuridine; STAIR list, initial stroke therapy academic industry roundtable (evaluation tools commonly used in quality assessment of animal experiments); BDNF, brain-derived neurotrophic factor; cGMP, cyclic guanosine monophosphate; vWF, von Willebrand factor; PECAM, platelet endothelial cell adhesion molecule; NGF, nerve growth factor; ERK, extracellular regulated protein kinases; BDNF, brain derived neurotrophic factor; TrkB, tyrosine kinase receptor B;CTGF, connective tissue growth factor; p70S6K, ribosomal protein S6 kinase, 70kDa; 4EBP1, 4E-binding protein 1; ATF2, activating transcription factor-2; MMP2, matrix metalloproteinase; Rps4x, ribosomal protein S4 X; Rps12, ribosomal protein 12; Rps14, ribosomal protein 14; Rps19, ribosomal protein 19; Hsp90b1, heat shock protein 90 beta family member 1; and Hsp90aa1, heat shock protein 90 alpha family class A member 1; Htra1, high-temperature requirement protein A1.




Figure 1 ICA and ICA derivatives regulate PI3K/AKT, Wnt/ β -catenin and TGF- β /Smad signaling pathways to regulate stem cell biological behavior. ICA and ICA derivatives promote p-AKT expression, which in turn promotes downstream eNOS/NO expression, activate mTORC1/p70S6K/4EBP1 to promote cell proliferation, inhibits Bcl-2 and GSK3 expression to inhibit apoptosis, and inhibits transcription factor CREB expression. ICA and ICA derivatives regulate the TGF- β /Smad and p38 MAPK pathways mainly by inhibiting Smad2/3 and p38 MAPK phosphorylation, while regulation of the Wnt pathway may be achieved by increasing frizzled class receptor expression and β -catenin phosphorylation. The regulation of these pathways by ICA and ICA derivatives ultimately leads to increased proliferation and differentiation of stem cells and inhibition of apoptosis and fibrosis. ICA, icariin; PI3K, phosphoinositide 3-kinases; Akt, threonine-specific protein kinase Akt; mTORC1, mechanistic target of rapamycin complex 1; ROS, reactive oxygen species; CREB, cAMP-response element binding protein; GSK3 β , glycogen synthase kinase 3 β ; Wnt, wingless-Type MMTV Integration Site Family; AGE, advanced glycation end products; TGF β 1, transforming growth factor- β 1; Smad2/3, mothers against decapentaplegic homolog 2/3; p70S6K, ribosomal protein S6 kinase, 70kDa; 4EBP1, 4E-binding protein 1; PIP2, phosphatidylinositol-3,4-bisphosphate; PIP3, phosphatidylinositol-3,4,5-trisphosphate; ATF-2, activating transcription factor-2; DVL, dishevelled segment polarity protein; Bcl-2, B-cell lymphoma-2; NO, nitric oxide; eNOS, endothelial nitric oxide synthase; PRAS60, rolin-rich Akt substrate of 60 kD; CK1 α , casein kinase 1 α .

repairing penile injured and restoring erectile function should be studied as a whole. In order to more clearly illustrate the therapeutic effect of ICA, it was classified according to the study type of animal model and stem cell.

Cavernous nerve regeneration

Erection is a complex physiological process. After the brain receives sexual stimulation, nerves transmit it to the target organ and secrete nitric oxide synthase (nNOS), which coordinates three downstream hemodynamic events: smooth muscle relaxation, arterial dilation and venous restriction, culminating in penile erection. The cavernous nerve (CN) originating from the pelvic ganglion (PG), with sympathetic and parasympathetic fibers, is capable of releasing nitric oxide (NO) and nNOS and activating endothelial cell NOS

(eNOS), resulting in a rise in NO content in the relaxed smooth muscle of the cavernous tissue.

ICAI restored nerve density and microstructure of pelvic floor ganglia in the penile corpus cavernosum of ED rats with diabetes-induced nerve damage, and *in vitro* ICAII resulted in longer synapses and more branching of ganglion tissue (42). In addition, YS-10, a new flavonoid based on the ICA II structure, treated rats with bilateral cavernous nerve injury (BCNI) and showed a significant reduction in smooth muscle atrophy, collagen deposition, endothelial and neurological dysfunction (43). In order to assess the efficacy and mechanisms of ICA treated BCNI rat. Newborn male rats were injected with 5-ethynyl-2-deoxyuridine (EdU, 50 mg/kg) to track endogenous stem cells in penis. Adult rats underwent bilateral cavernous nerve injury in the

penis, and the experimental group was treated with ICA (1.5 mg/kg/d) by gavage for 8 weeks. Nerve structures at the ICA-treated injury were restored to normal, and the number of EDU and S100 (a nervous specificity protein)-positive co-expressing cells and nNOS levels were positively correlated and significantly increased (25). It is suggested that promoting endogenous stem cell differentiation is an important mechanism for ICA to repair damaged cavernous nerves. Further studies confirmed the role of ICA to stimulate NSCs self-renewal, proliferation and differentiation *in vitro* (44,45).

Previous study found that ICAII significantly increased the levels of S100 and NGF in the penile tissue of rats after cavernous nerve injury and prevented distortion of neuroanatomical structures and neurotransmission dysfunction. Meanwhile, the expression trend of p38 MAPK in the penis was basically the same as that of S100, suggesting that the p38 MAPK signaling pathway may be involved in this process (69). *In vitro* experiments reveal more mechanisms of ICA regulation of NSCs. Yang *et al.* (46) found that ICA (10 μ M for 7 days) enhanced NSCs proliferation and neurosphere formation, by upregulating the expression of frizzled class receptor 7 (the key proteins of Wnt pathway), dishevelled segment polarity protein 3 (the key proteins of Wnt pathway), bFGF receptor 1, and downregulating glycogen synthase kinase-3 β (the Wnt pathway inhibitor) in NSCs. In another study suggested that ICA promoted the proliferation and differentiation of NSCs through the brain-derived neurotrophic factor (BDNF)-tyrosine kinase receptor B (TrkB)-extracellular regulated protein kinases (ERK)/Akt signaling pathway. By treating β -amyloid protein (A β)-damaged NSCs cells, the scholars found that ICA reversed the reduction of BDNF and TrkB expression and ERK/Akt phosphorylation caused by A β toxicity, and promoted NSCs proliferation and differentiation (47). In addition, the PI3K/Akt and IL-17 pathways were found to play a role in the regulation of functional proteins such as Rps14, Hsp90b1 and Htra1 in the cerebrospinal fluid by ICA (48).

However, it should be cautioned that mechanistic studies are often more advantageous in *in vitro* experiments, but due to technical limitations, penile stem cells and cavernous nerve stem cells are difficult to isolate. With the improvement of technology various tissue stem cell isolation will also be an important research direction.

Endothelium regeneration

After receiving sexual stimulation from cavernous nerve, the

cavernous endothelial cells secrete NO to maintain smooth muscle relaxation. Therefore, normal function and number of endothelial cells are important for erectile function of the penis. Studies found that the content of smooth muscle and endothelial cells were significantly reduced in diabetic rats. The expression of platelet endothelial cell adhesion molecule-1 (PECAM-1), eNOS and Von Willebrand factor (vWF) in the cavernous sinus endothelium were also reduced. ICA and ICAII treatments reversed these changes and increased the number of endothelial cells and smooth muscle cells (SMCs), possibly associated with downregulation of the TGF β 1/Smad2 signaling pathway (26,49).

Chen *et al.* (71) used high glucose to inhibit endothelial progenitor cell viability in a dose-dependent manner, while 1 μ M ICA treatment partially restored glucose-induced impairment of EPCs migration and tube formation. One μ M ICA significantly inhibited high glucose-induced phosphorylation of p38 and CREB (cAMP-response element binding protein) and increased Akt and eNOS activity in endothelial progenitor cells. Activation of Akt/eNOS could increase NO expression to regulate migration of endothelial progenitor cells. Tang *et al.* (50) found that ICA had the ability to activate rapamycin/70 kDa ribosomal protein S6 kinase (p70S6K)/eukaryotic translation initiation factor 4E (eIF4E)-binding protein 1 (4EBP1) and increased ATF2 and ERK1/2 protein levels to inhibit oxidative stress-induced damage to EPCs and promoted proliferation and differentiation of EPCs, and augmented capillary tube formation. Similar findings on the effects of ICA treatment on EPCs across different causes of injury further added credibility to the protection and activation effects of ICA for aforementioned signaling pathways on ECP.

Cavernous smooth muscle regeneration

Smooth muscle relaxation leads to rapid filling of the cavernous sinus with arterial blood, resulting in penile erection. Vascular smooth muscle has multiple sources, including mainly the proliferation of smooth muscle itself settled in the vasculature, transdifferentiation of circulating hematopoietic stem cells, and transdifferentiation of stem cells in the outer stromal layer of the vasculature.

Previous studies have demonstrated that ICA and ICAII treatment may reverse the reduction of penile SMCs caused by diabetes through downregulation of the TGF- β 1 signaling and upregulate α -smooth muscle actin (α -SMA) (26,49). Furthermore, Ruan *et al.* (64) found that ICA II (1.5 mg/kg/day for 4 weeks) improved erectile function and smooth muscle pathological changes through activation

of endogenous stem cells in obesity-related ED rats. Zhang *et al.* (24) demonstrated that ICA II (10 mg/kg for 8 weeks) ameliorated SMCs injury and restored smooth muscle to collagen ratio in diabetic rats. In this study, the percentage of SMCs in S phase (DNA synthesis phase), proliferation index (PI) were higher in the ICAII-treated group compared to the diabetic ED group. These effects may be related to the upregulation of the NO-cGMP pathway and the inhibition of excessive SMC autophagy and advanced glycation end products (AGE) deposition by ICAII. However, Hu *et al.* (72) showed that ICA dose-dependently (10 and 40 μ M) inhibited oxidized low-density lipoprotein (ox-LDL)-induced vascular SMC proliferation and suppressed ERK1/2 pathway and PCNA expression. The findings of this study diverge from those mentioned above but seem to be explicable. The complex processes and mechanisms such as oxidative stress in the penis of diabetic and obese rats ultimately lead to reduced damage to SMCs, and ox-LDL *in vitro* may be a stronger stimulus to activate the ERK1/2 pathway and promote SMC proliferation. Inhibition of oxidative stress by ICAII may be responsible for the discrepancy in the above findings. However, this study has less experimental data and lack of *in vitro* mechanism studies. Further experimental validation is necessary.

Modulation extracellular matrix and inhibition of fibrosis

Recent studies reported that diabetic rats had a lower ratio of type I to type III collagen, fragmented elastic fibers, and decreased elastic fiber content (27,73). After treating with ICA II (10 mg/kg/day) for 12 weeks, the α -SMA content and the ratio of collagen I to III were significantly higher in penis of diabetic rat compared to the untreated diabetic group. The results also showed a dramatic change in elastic fibers (71), which might be related to the fact that ICA II has the ability to improve lipid metabolism, reduce AGE concentration and mitochondrial autophagy.

There are at least 60 related genes that are downstream targets of TGF- β 1 (74). TGF- β 1 has been shown to increase collagen synthesis in cultured human corpus cavernosa SMCs *in vitro* (75). TGF- β 1 activated Smad2 and Smad3, leading to fibrosis-related changes (76,77). Connective tissue growth factor (CTGF), which plays an important role in connective tissue homeostasis, fibroblast proliferation, migration, and adhesion, is another downstream target of TGF- β 1 (78). The expression of TGF- β 1, total Smad2 and phospho-Smad2 was significantly higher in arteries,

vena cava and penile corpus cavernosum of diabetic rats. In addition, higher CTGF expression was shown in fibroblasts, endothelial cells and SMCs in the penile tissue of diabetic rats (51,52,79). In contrast, ICA II (1, 5, 10 mg/kg) treatment significantly reduced the expression of TGF- β 1/Smad/CTGF signaling pathway members in brain and penile tissues of diabetic rats and inhibited the fibrotic process in penile tissues (80).

Regulating endogenous testosterone (Table 4)

ICA and testosterone production

In the absence of testosterone, men may have symptoms including decreased libido, erectile dysfunction, reduced muscle mass and bone density, depression, and anemia. Although testosterone supplementation therapy (TST) has been reported to be potentially effective for these conditions, TST has potential adverse effects (81). The effect of ICA and ICAII in increasing testosterone levels could be related to their ability to mitigate the effects of Leydig cell damage in a variety of testicular damaging environmental exposures (e.g., diabetes, toxic substances) (82,84,85). More importantly, ICA and ICAII directly increase the level of steroids (the raw material required for testosterone production) and the expression of several key enzymes in testosterone synthesis (86-88).

Biological effects and molecular mechanisms of ICA in alleviating Leydig cell injury

Approximately 90% of testosterone is derived from the conversion of cholesterol by testicular Leydig cells and its level is regulated by follicle stimulating hormone (FSH) and luteinizing hormone (LH). Sun and his colleagues showed the protective effect of ICA (1 μ m/mL) reduced diethylhexyl phthalate (DEHP) injury-induced reactive oxygen species (ROS) levels of Leydig cells *In vitro* experiments, and increased mitochondrial membrane potential ($\Delta\Psi$ m) (82). Further study Sun *et al.* found that estrogen receptor 1 (Esr1)/Src family kinases (Src)/Akt/CREB/steroidogenic factor-1 (Sf-1) pathway may play an important role in ICA to protect Leydig cells from DEHP damage (83).

In addition, ICAII treatment (1.5 or 4.5 mg/kg/d for 28 days) increased superoxide dismutase (SOD), glutathione peroxidase (GPx) activity and inhibited malondialdehyde (MDA) activity, thereby attenuating diabetes, nicotine and aging on rat testicular Leydig cell damage (28,85).

Oxidative stress is one of the most important pathogenesis of testicular and penile tissue damage. Current study found

Table 4 Effect of ICA and ICA II repair on testosterone production of animals with testicular injury

ICA/ICA II	Year of publication	Study design	Study type	Treatment results	Molecular mechanisms	Effective dose	STAIR list score (full 7 points)	Key reference
ICA	2020	Mice with nicotine	<i>In vivo</i>	Improved sperm density, hormone levels and antioxidant enzyme activity	Activated antioxidant enzymes	75 mg/kg/day	5	(28)
ICA	2019	Mouse and Leydig cells with (2-Ethylhexyl) Phthalate	<i>In vivo</i> and <i>in vitro</i>	Promoted cell proliferation, and testosterone levels; Inhibited reactive oxygen species levels, mitochondrial membrane potential	Increased SF-1 and steroidogenic enzymes (CYP11, 3 β -HSD and 17 β -HSD)	50, 100 or 150 mg/kg/day <i>in vivo</i> ; 1 μ g/mL, and 5 μ g/mL <i>in vitro</i>	5	(80)
ICA	2021	Rat with high fat diet and streptozotocin	<i>In vivo</i>	recovered the number of spermatogonia, primary spermatocytes and Sertoli cells	upregulated the expression of PCNA, activated SRIT1-HIF-1 α signaling pathway; Up-regulated the expression of Bcl-2 and down-regulated the expression of Bax and caspase 3	80 mg/kg/day	4	(81)
ICAI	2014	Rat with streptozotocin	<i>In vivo</i>	Increased epididymal sperm parameters and testicular Johnsen's scores	Increased antioxidant enzyme activities and the expression of Sertoli cell Vimentin filaments, and	0.5, 1.5 or 4.5 mg/kg/day	4	(82)
ICA	2022	Mice and Leydig cells with (2-Ethylhexyl) Phthalate	<i>In vivo</i> and <i>in vitro</i>	Promotes testosterone synthesis	Activated Esr1/Src/Akt/Creb/Sf-1 signaling pathway	100 mg/kg/day <i>in vivo</i> ; 5 μ g/mL <i>in vitro</i>	5	(83)

ICA, icariin; ICAII, icariside II; ROS, reactive oxygen species; PBR, peripheral-type benzodiazepine receptor; SF-1, Steroidogenic factor-1; CYP11, Cytochrome P450 Family 11; 3 β -HSD, 3-beta (β)-hydroxysteroid dehydrogenase; 17 β -HSD, 3-beta (β)-hydroxysteroid dehydrogenase; STAIR list, The Initial Stroke Therapy Academic Industry Roundtable (Evaluation tools commonly used in quality assessment of animal experiments); PCNA, proliferating cell nuclear antigen; SRIT1, sirtuin 1; HIF-1 α , hypoxia-inducible factor-1; Bcl-2, B-cell lymphoma-2; Bax, Bcl-2-associated X; Esr1, estrogen receptor 1; Src, Src family kinases; Akt, threonine-specific protein kinase Akt; Creb, cAMP response element binding protein.

that nuclear factor- κ B-related factor 2 (Nrf2) could prevent ROS overexpression and accumulation (86,87). ICA can upregulate the expression of Nrf2 and its downstream heme oxygenase-1 (HO-1), nicotinamide adenine dinucleotide phosphate (NADPH), quinone oxidoreductase-1 (NQO-1) in high glucose-stimulated TM4 cells, which can increase SOD activity, decrease MDA content, and inhibit the production of ROS by high glucose stimulation. The above effects of ICA were diminished after Nrf2 knockdown treatment. These results suggest that the Nrf2 pathway is an important

molecular to mediate excessive oxidative stress inhibition activity of ICA. Further study revealed that the activation of Nrf2 pathway by ICA was mainly mediated by G protein-coupled estrogen receptor (GPER), which further promoted the dissociation of Nrf2/keap1 complex and the translocation of Nrf2 to nucleus (88).

Molecular mechanism of ICA on testosterone production

In addition to protecting testicular Leydig cells, ICA

plays an important regulatory role in testosterone production. Steroids are the precursors of testosterone, which are transported to the mitochondria of Leydig cells by acute regulatory protein (StAR) and then converted to testosterone through a series of cleavage and dehydrogenation reactions by enzymes including 3 β -Hydroxysteroid Dehydrogenase (3 β -HSD) and 17 β -Hydroxysteroid Dehydrogenase (17 β -HSD) (89). ICA (1 μ g/mL) treatment can reverse the DEPH-induced decrease of steroid levels and expression of StAR, 3 β -HSD, 17 β -HSD, and SF-1 in Leydig cells, enhancing testosterone synthesis (82).

In addition, the cGMP/PKG signaling pathway in Leydig cells is involved in the regulation of steroidogenic activity. It has been demonstrated that cGMP and protein kinase G (PKG) promote phosphorylation of StAR protein involved in testosterone synthesis, which can be inhibited by PDE5 (90). Therefore, the activation of cGMP and PKG by ICA and ICA derivatives and the inhibition of PDE5 may also be a mechanism for the regulation of testosterone production.

Summary

Researches in recent years have demonstrated the therapeutic effects of ICA, ICAII (an ICA metabolite isolated from epimedium herb) and its synthetic derivative YS-10 on ED, including inhibition of PDE5 enzymatic activity, promotion of testosterone production, and modulation of endogenous stem cells to promote regeneration of damaged penile tissues. As novel candidate agents for regenerative medicine on ED, the ICA, ICAII and YS-10 displayed bright application and research prospects. There are also several aspects that need to be studied in future research: (I) Flavonoids are generally insoluble in water, and experiments usually require the use of toxic organic solvents such as DMSO, so chemical modifications are also needed to increase the solubility of ICA and its derivatives; (II) The safety of the chemically modified compounds and the possible metabolic pathways need to be investigated; (III) ICA and ICA derivatives increase the number of penile stem cells and may also be the result of recruitment of stem cells from other sources, which warrants further investigation; (IV) ICA has been used for efficacy and mechanism studies in a variety of diseases, but no human data is currently available. This is likely to be the most important direction for future research. High quality and large sample of clinical and evidence-based medicine

studies are recommended while ensuring safety.

Acknowledgments

Funding: This work was supported and funded by the National Natural Science Foundation of China (grant No. 81971379).

Footnote

Reporting Checklist: The authors have completed the Narrative Review reporting checklist. Available at <https://tau.amegroups.com/article/view/10.21037/tau-22-232/rc>

Peer Review File: Available at <https://tau.amegroups.com/article/view/10.21037/tau-22-232/prf>

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <https://tau.amegroups.com/article/view/10.21037/tau-22-232/coif>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Lei H, Liu J, Li H, et al. Low-intensity shock wave therapy and its application to erectile dysfunction. *World J Mens Health* 2013;31:208-14.
2. Krakhotkin DV, Ralph DJ, Blecher GA, et al. Management of residual penile curvature after penile prosthesis placement. *Asian J Androl* 2021;23:129-34.
3. Rota C, Morigi M, Imberti B. Stem Cell Therapies in Kidney Diseases: Progress and Challenges. *Int J Mol Sci* 2019;20:2790.

4. Ma H, He X, Yang Y, et al. The genus Epimedium: an ethnopharmacological and phytochemical review. *J Ethnopharmacol* 2011;134:519-41.
5. Zhang HF, Yang TS, Li ZZ, et al. Simultaneous extraction of epimedin A, B, C and icariin from Herba Epimedii by ultrasonic technique. *Ultrason Sonochem* 2008;15:376-85.
6. Zhang DW, Cheng Y, Wang NL, et al. Effects of total flavonoids and flavonol glycosides from Epimedium koreanum Nakai on the proliferation and differentiation of primary osteoblasts. *Phytomedicine* 2008;15:55-61.
7. Zhang DC, Liu JL, Ding YB, et al. Icariin potentiates the antitumor activity of gemcitabine in gallbladder cancer by suppressing NF- κ B. *Acta Pharmacol Sin* 2013;34:301-8.
8. Kapoor S. Icariin and its emerging role in the treatment of osteoporosis. *Chin Med J (Engl)* 2013;126:400.
9. Liu M, Zhong C, He RX, et al. Icariin associated with exercise therapy is an effective treatment for postmenopausal osteoporosis. *Chin Med J (Engl)* 2012;125:1784-9.
10. Pan Y, Wang FM, Qiang LQ, et al. Icariin attenuates chronic mild stress-induced dysregulation of the LHPA stress circuit in rats. *Psychoneuroendocrinology* 2010;35:272-83.
11. Pan Y, Kong LD, Li YC, et al. Icariin from Epimedium brevicornum attenuates chronic mild stress-induced behavioral and neuroendocrinological alterations in male Wistar rats. *Pharmacol Biochem Behav* 2007;87:130-40.
12. Liao H, Jacob R. Chinese herbal drugs for erectile dysfunction through NO-cGMP-PDE5 signaling pathway. *Zhonghua Nan Ke Xue* 2012;18:260-5.
13. Chen Y, Zhao YH, Jia XB, et al. Intestinal absorption mechanisms of prenylated flavonoids present in the heat-processed Epimedium koreanum Nakai (Yin Yanghuo). *Pharm Res* 2008;25:2190-9.
14. Cao YF, He RR, Cao J, et al. Drug-Drug Interactions Potential of Icariin and Its Intestinal Metabolites via Inhibition of Intestinal UDP-Glucuronosyltransferases. *Evid Based Complement Alternat Med* 2012;2012:395912.
15. Cai WJ, Huang JH, Zhang SQ, et al. Icariin and its derivative icariside II extend healthspan via insulin/IGF-1 pathway in *C. elegans*. *PLoS One* 2011;6:e28835.
16. Cheng T, Zhang Y, Zhang T, et al. Comparative Pharmacokinetics Study of Icariin and Icariside II in Rats. *Molecules* 2015;20:21274-86.
17. Wu H, Kim M, Han J. Icariin Metabolism by Human Intestinal Microflora. *Molecules* 2016;21:1158.
18. Xu W, Zhang Y, Yang M, et al. LC-MS/MS method for the simultaneous determination of icariin and its major metabolites in rat plasma. *J Pharm Biomed Anal* 2007;45:667-72.
19. Liu M, Liu H, Lu X, et al. Simultaneous determination of icariin, icariside II and osthole in rat plasma after oral administration of the extract of Gushudan (a Chinese compound formulation) by LC-MS/MS. *J Chromatogr B Analyt Technol Biomed Life Sci* 2007;860:113-20.
20. Qian Q, Li SL, Sun E, et al. Metabolite profiles of icariin in rat plasma by ultra-fast liquid chromatography coupled to triple-quadrupole/time-of-flight mass spectrometry. *J Pharm Biomed Anal* 2012;66:392-8.
21. Chang Q, Wang GN, Li Y, et al. Oral absorption and excretion of icaritin, an aglycone and also active metabolite of prenylflavonoids from the Chinese medicine Herba Epimedii in rats. *Phytomedicine* 2012;19:1024-8.
22. Wong SP, Shen P, Lee L, et al. Pharmacokinetics of prenylflavonoids and correlations with the dynamics of estrogen action in sera following ingestion of a standardized Epimedium extract. *J Pharm Biomed Anal* 2009;50:216-23.
23. Ahn GJ, Sohn YS, Kang KK, et al. The effect of PDE5 inhibition on the erectile function in streptozotocin-induced diabetic rats. *Int J Impot Res* 2005;17:134-41.
24. Zhang J, Li AM, Liu BX, et al. Effect of icarisid II on diabetic rats with erectile dysfunction and its potential mechanism via assessment of AGEs, autophagy, mTOR and the NO-cGMP pathway. *Asian J Androl* 2013;15:143-8.
25. Xu Y, Xin H, Wu Y, et al. Effect of icariin in combination with daily sildenafil on penile atrophy and erectile dysfunction in a rat model of bilateral cavernous nerves injury. *Andrology* 2017;5:598-605.
26. Liu T, Xin H, Li WR, et al. Effects of icariin on improving erectile function in streptozotocin-induced diabetic rats. *J Sex Med* 2011;8:2761-72.
27. Zhang J, Li S, Zhang S, et al. Effect of Icariside II and Metformin on Penile Erectile Function, Histological Structure, Mitochondrial Autophagy, Glucose-Lipid Metabolism, Angiotensin II and Sex Hormone in Type 2 Diabetic Rats With Erectile Dysfunction. *Sex Med* 2020;8:168-77.
28. Ni G, Zhang X, Afedo SY, et al. Evaluation of the protective effects of icariin on nicotine-induced reproductive toxicity in male mouse -a pilot study. *Reprod Biol Endocrinol* 2020;18:65.
29. Wang H, Liu Y, Huai Q, et al. Multiple conformations of phosphodiesterase-5: implications for enzyme function and drug development. *J Biol Chem* 2006;281:21469-79.

30. Chau Y, Li FS, Levsh O, et al. Exploration of icariin analog structure space reveals key features driving potent inhibition of human phosphodiesterase-5. *PLoS One* 2019;14:e0222803.
31. Xin ZC, Kim EK, Lin CS, et al. Effects of icariin on cGMP-specific PDE5 and cAMP-specific PDE4 activities. *Asian J Androl* 2003;5:15-8.
32. Ning H, Xin ZC, Lin G, et al. Effects of icariin on phosphodiesterase-5 activity in vitro and cyclic guanosine monophosphate level in cavernous smooth muscle cells. *Urology* 2006;68:1350-4.
33. Zhang J, Wang YB, Ma CG, et al. Icarisid II, a PDE5 inhibitor from *Epimedium wanshanense*, increases cellular cGMP by enhancing NOS in diabetic ED rats corpus cavernosum tissue. *Andrologia* 2012;44 Suppl 1:87-93.
34. Dell'Agli M, Galli GV, Dal Cero E, et al. Potent inhibition of human phosphodiesterase-5 by icariin derivatives. *J Nat Prod* 2008;71:1513-7.
35. Jiang Z, Hu B, Wang J, et al. Effect of icariin on cyclic GMP levels and on the mRNA expression of cGMP-binding cGMP-specific phosphodiesterase (PDE5) in penile cavernosum. *J Huazhong Univ Sci Technolog Med Sci* 2006;26:460-2.
36. Li Y, Jiang J, He Y, et al. Icariin combined with breviscapine improves the erectile function of spontaneously hypertensive rats. *J Sex Med* 2014;11:2143-52.
37. Li WK, Zhang RY, Xiao PG. Flavonoids from *Epimedium wanshanense*. *Phytochemistry* 1996;43:527-30.
38. Fan JJ, Cao LG, Wu T, et al. The dose-effect of icariin on the proliferation and osteogenic differentiation of human bone mesenchymal stem cells. *Molecules* 2011;16:10123-33.
39. Zhang S, Feng P, Mo G, et al. Icariin influences adipogenic differentiation of stem cells affected by osteoblast-osteoclast co-culture and clinical research adipogenic. *Biomed Pharmacother* 2017;88:436-42.
40. Jones NC, Tyner KJ, Nibarger L, et al. The p38alpha/beta MAPK functions as a molecular switch to activate the quiescent satellite cell. *J Cell Biol* 2005;169:105-16.
41. Zhai YK, Guo XY, Ge BF, et al. Icariin stimulates the osteogenic differentiation of rat bone marrow stromal cells via activating the PI3K-AKT-eNOS-NO-cGMP-PKG. *Bone* 2014;66:189-98.
42. Bai GY, Zhou F, Hui Y, et al. Effects of Icariside II on corpus cavernosum and major pelvic ganglion neuropathy in streptozotocin-induced diabetic rats. *Int J Mol Sci* 2014;15:23294-306.
43. Gu SJ, Li M, Yuan YM, et al. A novel flavonoid derivative of icariside II improves erectile dysfunction in a rat model of cavernous nerve injury. *Andrology* 2021;9:1893-901.
44. Fu X, Li S, Zhou S, et al. Stimulatory effect of icariin on the proliferation of neural stem cells from rat hippocampus. *BMC Complement Altern Med* 2018;18:34.
45. Huang JH, Cai WJ, Zhang XM, et al. Icariin promotes self-renewal of neural stem cells: an involvement of extracellular regulated kinase signaling pathway. *Chin J Integr Med* 2014;20:107-15.
46. Yang P, Guan YQ, Li YL, et al. Icariin promotes cell proliferation and regulates gene expression in human neural stem cells in vitro. *Mol Med Rep* 2016;14:1316-22.
47. Lu Q, Zhu H, Liu X, et al. Icariin sustains the proliferation and differentiation of Aβ25-35-treated hippocampal neural stem cells via the BDNF-TrkB-ERK/Akt signaling pathway. *Neurol Res* 2020;42:936-45.
48. Zeng NX, Li HZ, Wang HZ, et al. Exploration of the mechanism by which icariin modulates hippocampal neurogenesis in a rat model of depression. *Neural Regen Res* 2022;17:632-42.
49. Zhou F, Xin H, Liu T, et al. Effects of icariside II on improving erectile function in rats with streptozotocin-induced diabetes. *J Androl* 2012;33:832-44.
50. Tang Y, Jacobi A, Vater C, et al. Icariin promotes angiogenic differentiation and prevents oxidative stress-induced autophagy in endothelial progenitor cells. *Stem Cells* 2015;33:1863-77.
51. Castañares C, Redondo-Horcajo M, Magán-Marchal N, et al. Signaling by ALK5 mediates TGF-beta-induced ET-1 expression in endothelial cells: a role for migration and proliferation. *J Cell Sci* 2007;120:1256-66.
52. Hyman KM, Seghezzi G, Pintucci G, et al. Transforming growth factor-beta1 induces apoptosis in vascular endothelial cells by activation of mitogen-activated protein kinase. *Surgery* 2002;132:173-9.
53. Yang A, Yu C, Lu Q, et al. Mechanism of Action of Icariin in Bone Marrow Mesenchymal Stem Cells. *Stem Cells Int* 2019;2019:5747298.
54. Ye Y, Jing X, Li N, et al. Icariin promotes proliferation and osteogenic differentiation of rat adipose-derived stem cells by activating the RhoA-TAZ signaling pathway. *Biomed Pharmacother* 2017;88:384-94.
55. Li X, Yang HF, Chen Y, et al. Effect of the icariin on endothelial microparticles, endothelial progenitor cells, platelets, and erectile function in spontaneously hypertensive rats. *Andrology* 2022;10:576-84.
56. Cui H, Liu Z, Wang L, et al. Icariin-treated human umbilical cord mesenchymal stem cells decrease chronic

- liver injury in mice. *Cytotechnology* 2017;69:19-29.
57. Luo Z, Liu M, Sun L, et al. Icaritin recovers the osteogenic differentiation and bone formation of bone marrow stromal cells from a rat model of estrogen deficiency-induced osteoporosis. *Mol Med Rep* 2015;12:382-8.
 58. Sun ZB, Wang JW, Xiao H, et al. Icaritin may benefit the mesenchymal stem cells of patients with steroid-associated osteonecrosis by ABCB1-promoter demethylation: a preliminary study. *Osteoporos Int* 2015;26:187-97.
 59. Wang X, Liu C, Xu Y, et al. Combination of mesenchymal stem cell injection with icaritin for the treatment of diabetes-associated erectile dysfunction. *PLoS One* 2017;12:e0174145.
 60. Xu Y, Yang Y, Zheng H, et al. Intracavernous injection of size-specific stem cell spheroids for neurogenic erectile dysfunction: Efficacy and risk versus single cells. *EBioMedicine* 2020;52:102656.
 61. Kurzrock EA, Lieu DK, Degraffenried LA, et al. Label-retaining cells of the bladder: candidate urothelial stem cells. *Am J Physiol Renal Physiol* 2008;294:F1415-21.
 62. Huang YL, Tao X, Xia J, et al. Distribution and quantity of label-retaining cells in rat oral epithelia. *J Oral Pathol Med* 2009;38:663-7.
 63. Song J, Shu L, Zhang Z, et al. Reactive oxygen species-mediated mitochondrial pathway is involved in Baohuoside I-induced apoptosis in human non-small cell lung cancer. *Chem Biol Interact* 2012;199:9-17.
 64. Ruan Y, Lin G, Kang N, et al. In Situ Activation and Preservation of Penile Progenitor Cells Using Icariside II in an Obesity-Associated Erectile Dysfunction Rat Model. *Stem Cells Dev* 2018;27:207-15.
 65. Xu Y, Guan R, Lei H, et al. Implications for differentiation of endogenous stem cells: therapeutic effect from icariside II on a rat model of postprostatectomy erectile dysfunction. *Stem Cells Dev* 2015;24:747-55.
 66. Ding L, Liang XG, Zhu DY, et al. Icaritin promotes expression of PGC-1alpha, PPARalpha, and NRF-1 during cardiomyocyte differentiation of murine embryonic stem cells in vitro. *Acta Pharmacol Sin* 2007;28:1541-9.
 67. Qin S, Zhou W, Liu S, et al. Icaritin stimulates the proliferation of rat bone mesenchymal stem cells via ERK and p38 MAPK signaling. *Int J Clin Exp Med* 2015;8:7125-33.
 68. Fu S, Yang L, Hong H, et al. Wnt/ β -catenin signaling is involved in the icaritin induced proliferation of bone marrow mesenchymal stem cells. *Journal of Traditional Chinese Medicine* 2016;36:360-8.
 69. Luo G, Xu B, Huang Y. Icariside II promotes the osteogenic differentiation of canine bone marrow mesenchymal stem cells via the PI3K/AKT/mTOR/S6K1 signaling pathways. *Am J Transl Res* 2017;9:2077-87.
 70. Lin G, Alwaal A, Zhang X, et al. Presence of stem/progenitor cells in the rat penis. *Stem Cells Dev* 2015;24:264-70.
 71. Chen S, Wang Z, Zhou H, et al. Icaritin reduces high glucose-induced endothelial progenitor cell dysfunction via inhibiting the p38/CREB pathway and activating the Akt/eNOS/NO pathway. *Exp Ther Med* 2019;18:4774-80.
 72. Hu Y, Liu K, Yan M, et al. Icaritin inhibits oxidized low-density lipoprotein-induced proliferation of vascular smooth muscle cells by suppressing activation of extracellular signal-regulated kinase 1/2 and expression of proliferating cell nuclear antigen. *Mol Med Rep* 2016;13:2899-903.
 73. Hsu GL, Brock G, von Heyden B, et al. The distribution of elastic fibrous elements within the human penis. *Br J Urol* 1994;73:566-71.
 74. Verrecchia F, Chu ML, Mauviel A. Identification of novel TGF-beta/Smad gene targets in dermal fibroblasts using a combined cDNA microarray/promoter transactivation approach. *J Biol Chem* 2001;276:17058-62.
 75. Moreland RB. Is there a role of hypoxemia in penile fibrosis: a viewpoint presented to the Society for the Study of Impotence. *Int J Impot Res* 1998;10:113-20.
 76. Verrecchia F, Mauviel A. Control of connective tissue gene expression by TGF beta: role of Smad proteins in fibrosis. *Curr Rheumatol Rep* 2002;4:143-9.
 77. McNairn AJ, Brusadelli M, Guasch G. Signaling moderation: TGF-beta in exocrine gland development, maintenance, and regulation. *Eur J Dermatol* 2013. doi: 10.1684/ejd.2013.1979.
 78. Yamagishi S, Inagaki Y, Okamoto T, et al. Advanced glycation end products inhibit de novo protein synthesis and induce TGF-beta overexpression in proximal tubular cells. *Kidney Int* 2003;63:464-73.
 79. Xiang Y, Cai C, Wu Y, et al. Icaritin Attenuates Monocrotaline-Induced Pulmonary Arterial Hypertension via the Inhibition of TGF- β 1/Smads Pathway in Rats. *Evid Based Complement Alternat Med* 2020;2020:9238428.
 80. Dong C, Ming X, Ye Z, et al. Icariside II Attenuates Chronic Hydrocephalus in an Experimental Subarachnoid Hemorrhage Rat Model. *J Pharm Pharm Sci* 2018;21:318-25.
 81. Lang PO, Samaras D, Samaras N. Testosterone replacement therapy in reversing "andropause": what is the proof-of-principle?. *Rejuvenation Res* 2012;15:453-65.

82. Sun J, Wang D, Lin J, et al. Icariin protects mouse Leydig cell testosterone synthesis from the adverse effects of di(2-ethylhexyl) phthalate. *Toxicol Appl Pharmacol* 2019;378:114612.
83. Sun J, Xu W, Zheng S, et al. Icariin promotes mouse Leydig cell testosterone synthesis via the Esr1/Src/Akt/Creb/Sf-1 pathway. *Toxicol Appl Pharmacol* 2022;441:115969.
84. He W, Liu H, Hu L, et al. Icariin improves testicular dysfunction via enhancing proliferation and inhibiting mitochondria-dependent apoptosis pathway in high-fat diet and streptozotocin-induced diabetic rats. *Reprod Biol Endocrinol* 2021;19:168.
85. Xu Y, Lei H, Guan R, et al. Prophylactic protective effects and its potential mechanisms of IcarisideII on streptozotocin induced spermatogenic dysfunction. *Int J Mol Sci* 2014;15:16100-13.
86. Loboda A, Damulewicz M, Pyza E, et al. Role of Nrf2/HO-1 system in development, oxidative stress response and diseases: an evolutionarily conserved mechanism. *Cell Mol Life Sci* 2016;73:3221-47.
87. Jha JC, Banal C, Chow BS, et al. Diabetes and Kidney Disease: Role of Oxidative Stress. *Antioxid Redox Signal* 2016;25:657-84.
88. Wang K, Zheng X, Pan Z, et al. Icariin Prevents Extracellular Matrix Accumulation and Ameliorates Experimental Diabetic Kidney Disease by Inhibiting Oxidative Stress via GPER Mediated p62-Dependent Keap1 Degradation and Nrf2 Activation. *Front Cell Dev Biol* 2020;8:559.
89. Chen X, Liu YN, Zhou QH, et al. Effects of low concentrations of di-(2-ethylhexyl) and mono-(2-ethylhexyl) phthalate on steroidogenesis pathways and apoptosis in the murine leydig tumor cell line MLTC-1. *Biomed Environ Sci* 2013;26:986-9.
90. Andric SA, Janjic MM, Stojkov NJ, et al. Testosterone-induced modulation of nitric oxide-cGMP signaling pathway and androgenesis in the rat Leydig cells. *Biol Reprod* 2010;83:434-42.

Cite this article as: Niu Y, Lin G, Pan J, Liu J, Xu Y, Cai Q, Wang T, Luan Y, Chen Y, Feng Y, Yang X, Tian W, Bae WJ, Guan R, Xin Z. Deciphering the myth of icariin and synthetic derivatives in improving erectile function from a molecular biology perspective: a narrative review. *Transl Androl Urol* 2022;11(7):1007-1022. doi: 10.21037/tau-22-232