



The carbon footprint of respiratory treatments in Europe and Canada: an observational study from the CARBON programme

Copyright ©The authors 2022.

This version is distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0. For commercial reproduction rights and permissions contact permissions@ersnet.org

Received: 20 Oct 2021
Accepted: 5 June 2022

To the Editor:

Climate change represents a global challenge and nations are increasingly looking to decarbonise their economies by developing roadmaps for reducing greenhouse gas (GHG) emissions in accordance with international treaties, such as the Paris Agreement [1]. As the healthcare sector remains a key contributor to GHG emissions [2], an examination of the global carbon footprint of its operations and treatment pathways is essential to identify targets for decarbonisation.

In respiratory treatment, the environmental impact of controller inhalers has received considerable attention due to the hydrofluorocarbon propellants used in metered-dose inhalers (MDIs), which have global warming potential [3]. In the United Kingdom (UK), where MDIs represented ~3% of health and social care system carbon emissions [4] and 13.1% of emissions related to the delivery of care in 2019 [2], targets are in place to reduce total emissions by 80% by 2036–2039, including those from MDIs [5]. However, the focus on controller inhalers omits other contributors, such as the impact of short-acting β_2 -agonists (SABAs), presenting an incomplete picture of the carbon footprint of respiratory treatments for both asthma and COPD.

Patients with mild asthma, who represent approximately half of the European asthma population [6], are commonly prescribed SABA-only treatment [7], placing them at increased risk of poor outcomes [8]. Additionally, findings from the real-world SABA use in Asthma (SABINA) programme revealed that approximately one-third of patients with asthma across Europe overuse SABA (prescription/dispensing of three or more canisters per year) [9], which is associated with an increased risk of exacerbations and healthcare resource use [10, 11]. Moreover, increased healthcare resource use in respiratory treatment, whether associated with poor disease control of asthma [12] or progression of COPD [13], carries an additional carbon burden [14].

We considered that suboptimal care of patients with asthma or COPD may drive a higher, yet potentially modifiable, contribution to global GHG emissions, and developed the Healthcare-based Carbon Cost of Treatment (CARBON) programme, an evolving healthcare sustainability scheme to better understand the carbon footprint of respiratory disease control and progression [15]. As part of the CARBON programme, the SABA CARBON Europe and Canada observational cohort study quantified the carbon footprint associated with 1) the use of both reliever and controller inhalers in 20 European countries and in Canada and 2) SABA overuse (prescription/dispensing of three or more canisters per year) in five European countries and two Canadian provinces (Alberta and Nova Scotia) from the SABINA programme.

Inhaler sales data, as a surrogate for inhaler use, for SABA and controller medications (MDIs and dry powder inhalers (DPIs)), across all respiratory uses, were obtained from the IQVIA quarterly MIDAS database Q3 2019 (September 2018 to September 2019), accessed and analysed *via* the AstraZeneca in-house STAR system. This analysis included patients treated for any respiratory condition. Controller treatments included inhaled corticosteroid-containing drugs, long-acting β_2 -agonists (LABA), long-acting muscarinic antagonists (LAMA) and LAMA/LABA combinations.



Shareable abstract (@ERSpublications)

Relievers account for the majority of inhaler use and associated GHG emissions. Implementing treatment guidelines can reduce the unmet need in respiratory care by improving disease control and reducing reliever overuse and the overall carbon footprint. <https://bit.ly/3zh3c2B>

Cite this article as: Janson C, Maslova E, Wilkinson A, *et al.* The carbon footprint of respiratory treatments in Europe and Canada: an observational study from the CARBON programme. *Eur Respir J* 2022; 60: 2102760 [DOI: 10.1183/13993003.02760-2021].

SABA overuse in patients with asthma (aged ≥ 12 years) of any severity was assessed using prescription/dispensing data from the SABINA programme (2006–2019) [10, 16–19].

Data were compared by dose, preventing confounding from inhaler actuation count differences. A 1:1 equivalence of actuation and dose was assumed for SABAs and controller medication delivered *via* a DPI, whereas a 2:1 ratio for actuation to dose was assumed for controller medication delivered *via* a MDI. Both analyses were descriptive in nature and annual GHG emissions were expressed as carbon dioxide equivalent (CO₂e) and quantified using published data [3, 20] and internal AstraZeneca estimates. Per capita inhaler usage and associated carbon footprint were calculated using the national population for IQVIA sales data and using the study populations for the SABINA data.

SABA use was common across 20 countries in Europe and in Canada (table 1). Although SABAs were mostly administered *via* MDIs, this varied across countries, ranging from 28.2% in Sweden to 100% in Italy. Per capita SABA use ranged from 98 770 (Poland) to 1 033 535 (UK) doses per 10 000 persons per year. Compared with SABA use, per capita controller medication use was lower, ranging from 58 506 (Romania) to 437 945 (UK) doses per 10 000 persons per year.

As a proportion of total inhaler use, SABA inhalers ranged from 33% (Belgium) to 71% (Canada and Ireland), with SABA GHG emissions ranging from 47% (Netherlands and Sweden) to 80% (Romania). Compared with per capita GHG emissions from SABA, which ranged from 12 (Sweden) to 134 (UK) tonnes CO₂e per 10 000 persons per year, per capita GHG emissions from controller medication use were lower and ranged from 4 (Romania) to 65 (UK) tonnes CO₂e per 10 000 persons per year. Total GHG emissions from the use of SABA and controller medications were approximately 2 and 1 million tonnes CO₂e, respectively, with SABA use accounting for 66% of the total GHG emissions from inhalers.

Across the seven SABINA datasets comprising 1 131 416 patients with asthma (table 1), most SABA prescriptions were received by patients who were overusing SABA (three or more canisters per year), ranging from 69% (Italy and Sweden) to 94% (Canada (Nova Scotia)). SABA overuse contributed to excess per capita GHG emissions, ranging from 78 (Sweden) to 864 (Canada (Nova Scotia)) tonnes CO₂e per 10 000 persons per year. Per capita GHG emissions from SABA overuse in Canada (Nova Scotia) were 1.6- to 11.1-fold higher *versus* the UK and Sweden, respectively. As an example, SABA overuse when scaled to the national asthma population of ~ 5.4 million in the UK translated to an excess carbon footprint of 293 227 tonnes CO₂e. Across the two Canadian provinces, both SABA overuse and the associated per capita emissions were higher in Nova Scotia compared with Alberta.

Although inhaler sales and prescriptions/dispensing data may not reflect actual medication use and final disposal (together accounting for $\sim 90\%$ of the GHG emissions) [3], this study provides an understanding of how high SABA use, a marker of poor disease control and suboptimal disease management [8], drives the associated carbon footprint of respiratory treatment.

Overall, our findings reveal that suboptimal respiratory treatment, in the form of high SABA use across Europe and Canada, remains widespread, representing approximately two-thirds of total GHG emissions. These findings highlight the importance of assessing the contribution of SABAs to the carbon footprint of respiratory treatment, which in many countries were commonly used and administered by MDIs, thereby explaining higher GHG emissions associated with SABA *versus* controller inhaler use. Furthermore, an analysis of SABINA datasets demonstrated that SABA overuse, as defined by the threshold of three or more canisters per year [8], drives the majority of SABA prescriptions/dispensing in asthma, suggesting suboptimal disease management in a high proportion of patients who are at increased risk of asthma exacerbations and exacerbation-related healthcare resource use, thereby further contributing to the total carbon footprint.

In most countries, SABAs represented the majority of respiratory-related inhaler use, indicating suboptimal disease control in these populations. The highest per capita use of both SABA and controller inhalers was observed in the UK. SABA overuse in asthma was prevalent despite the different healthcare and reimbursement policies of each country, a finding consistent with previous studies [9–11, 17, 18, 21, 22]. Across all SABINA datasets, SABA prescribing/dispensing was primarily driven by patients who were potentially overusing SABA relievers. However, these findings should be interpreted in light of diverse asthma management practices and differences in healthcare delivery systems and socioeconomic status across the individual datasets, particularly in relation to access to medications [19]. For example, in Germany and Sweden, SABA is a prescription-only medicine [9], while in Italy, SABA is available without a prescription [17]. Thus, actual SABA use in Italy may have been higher than observed.

TABLE 1 Greenhouse gas (GHG) emissions related to short-acting β_2 -agonist (SABAs) *versus* controller medication in 21 countries, and SABA overuse among patients with asthma from the seven SABA use in Asthma (SABINA) datasets and associated GHG emissions

	Per capita SABA use, doses per 10 000 persons per year	Per capita controller medication use, doses per 10 000 persons per year	SABA <i>versus</i> total inhaler use, %	GHG emissions from SABA, tonnes CO ₂ e	GHG emissions from controller medication, tonnes CO ₂ e	SABA <i>versus</i> total inhaler GHG emissions, %	Per capita GHG emissions from SABA, tonnes CO ₂ e per 10 000 persons per year	Per capita GHG emissions from controller medication, tonnes CO ₂ e per 10 000 persons per year	Total patients in SABINA database (GINA equivalent steps 1–2–3–5)	Volume of SABA prescriptions (GINA equivalent steps 1–2–3–5)	SABA prescriptions received by patients potentially overusing SABA reliever (GINA equivalent steps 1–2–3–5), %	Total GHG emissions from SABA overuse, tonnes CO ₂ e within this cohort	Per capita GHG emission from SABA overuse, tonnes CO ₂ e per 10 000 persons
Belgium	131 047	267 567	33	19 672	17 422	53	17	15					
Bulgaria	125 630	141 248	47	11 959	6949	63	17	10					
Canada [#]	543 796	224 999	71	193 356	73 898	72	55	21					
Alberta									107 444	274 175	81	4303	401
Nova Scotia									8034	34 677	94	694	864
Croatia	126 186	111 056	53	7253	4175	64	17	10					
Czech Republic	125 088	210 556	37	17 563	18 485	49	17	17					
Denmark	281 377	284 622	50	10 889	8017	58	19	14					
Finland	278 298	337 388	45	10 963	11 563	49	20	21					
France	383 001	229 696	63	334 716	126 131	73	50	19					
Germany	275 899	234 476	54	293 638	144 077	67	36	18	13 030	39 643	74	540	415
Greece	237 510	288 218	45	34 223	20 988	62	32	19					
Hungary	182 449	144 174	56	22 597	12 560	64	23	13					
Ireland	743 424	300 928	71	48 986	16 782	75	98	34					
Italy	125 516	144 659	46	104 503	86 040	55	17	14	22 102	17 866	69	240	108
The Netherlands	256 961	288 783	47	46 559	52 922	47	27	31					
Norway	285 983	279 637	51	14 776	10 188	59	28	19					
Poland [¶]	98 770	166 863	37	49 893	43 738	53	13	11	98 876	135 424	74	1928	195
Romania	126 896	58 506	68	36 749	9371	80	17	4					
Spain	318 751	222 311	59	195 771	86 977	69	40	18					
Sweden ⁺	238 512	349 211	41	11 632	12 895	47	12	13	365 324	794 589	69	2844	78
Switzerland	154 456	150 331	51	16 067	6174	72	20	8					
UK	1 033 535	437 945	70	862 685	415 345	68	134	65	516 606	1 753 804	85	28 052	543

GHG emissions associated with medication use were quantified using SimaPro life cycle assessment software modelling resource and energy consumption data, in addition to Ecoinvent datasets and certified published studies. Per capita GHG emissions were calculated to allow comparisons across countries/datasets. CO₂e: carbon dioxide equivalent; GINA: Global Initiative for Asthma. #: data from Alberta and Nova Scotia from the SABINA datasets were analysed separately to compare SABA overuse and associated emissions across the two provinces; ¶: patients with zero SABA use could not be categorised across GINA steps; +: 8061 patients from Sweden could not be categorised according to GINA steps.

SABA prescribing/dispensing patterns in Poland may be attributable to underfunding of the healthcare system, leading to relatively high out-of-pocket spending [23]. Variations in SABA overuse between the two Canadian provinces may be due, in part, to differences in socioeconomic status [24] that may have influenced access to recommended asthma medications. However, further research is needed to verify these findings.

Although global asthma guidelines no longer recommend as-needed SABA use alone due to safety concerns [8] and an inconsistent evidence base, asthma management practices have not yet caught up with current evidence-based recommendations, and SABA overuse therefore continues to drive the majority of GHG emissions across countries. Consequently, implementation of clinical guidelines, adherence to asthma action plans, and delivery of personalised care along with a focus on the management of modifiable risk factors for poor disease control, such as SABA overuse, poor medication adherence and incorrect inhaler technique should be prioritised to improve patient outcomes [8]. This approach will subsequently reduce SABA reliever use and additional healthcare resource use, thereby benefiting patients and realising carbon savings that go beyond the reduction in SABA use alone. As suboptimal disease management continues to be an unacceptable unmet need in respiratory treatment, this is a call to action for healthcare professionals and policymakers to ensure that treatment-related decisions are guided by current evidence-based recommendations and tailored to patient needs, thereby reducing SABA use and associated carbon emissions in respiratory treatment, without risking improvements in patient outcomes or causing harm.

Christer Janson ¹, **Ekaterina Maslova**², **Alexander Wilkinson** ³, **Erika Penz**^{4,5}, **Alberto Papi** ⁶, **Nigel Budgen**⁷, **Claus F. Vogelmeier**⁸, **Maciej Kupczyk**^{9,10}, **John Bell** ² and **Andrew Menzies-Gow**¹¹

¹Department of Medical Sciences, Respiratory, Allergy and Sleep Research, Uppsala University, Uppsala, Sweden. ²BioPharmaceuticals Medical, AstraZeneca, Cambridge, UK. ³East and North Hertfordshire NHS Trust, Stevenage, UK. ⁴Department of Medicine, Division of Respiriology, Critical Care and Sleep Medicine, University of Saskatchewan, Saskatoon, SK, Canada. ⁵Respiratory Research Centre, University of Saskatchewan, Saskatoon, SK, Canada. ⁶Respiratory Medicine, Department of Translational Medicine, University of Ferrara, Ferrara, Italy. ⁷Global Sustainability, AstraZeneca, Macclesfield, UK. ⁸Department of Medicine, Pulmonary and Critical Care Medicine, University Medical Center Giessen and Marburg, Philipps-University Marburg, Member of the German Center for Lung Research (DZL), Marburg, Germany. ⁹Department of Internal Medicine, Asthma and Allergy, Barlicki University Hospital, Medical University of Lodz, Lodz, Poland. ¹⁰Center for Allergy Research, IMM, Karolinska Institutet, Stockholm, Sweden. ¹¹Lung Division, Royal Brompton Hospital, London, UK.

Corresponding author: Christer Janson (christer.janson@medsci.uu.se)

Acknowledgements: The authors thank Dorota Brzostek (AstraZeneca, Poland), Gunilla Telg (AstraZeneca, Sweden), Silvia Boarino (AstraZeneca, Italy), Ingo Mokros (AstraZeneca, Germany) and Stephen Noorduy (AstraZeneca, Canada) for their assistance in coordinating data collection. Writing and editorial support was provided by Praveen Kaul and Frances Gambling, of Cactus Life Sciences (part of Cactus Communications, Mumbai, India) in accordance with Good Publication Practice (GPP3) guidelines (<http://www.ismpp.org/gpp3>) and fully funded by AstraZeneca.

IQVIA did not provide any support for the analysis or interpretation of the data.

Author contributions: J. Bell and E. Maslova conceptualised the study design. E. Maslova, N. Budgen and J. Bell performed the data analysis. All authors contributed to data collection, data interpretation and writing.

Conflict of interest: C. Janson reports personal fees from AstraZeneca, Boehringer Ingelheim, Chiesi, GlaxoSmithKline PLC, Novartis and Teva, outside the submitted work. A. Wilkinson is a member of the Montreal protocol Medical and Chemical Technical Options Committee and has made unpaid contributions to publications on the carbon footprint of inhalers and respiratory treatment which were sponsored by GlaxoSmithKline and AstraZeneca. E. Penz has received honoraria and consulting fees from AstraZeneca, GlaxoSmithKline, Sanofi Genzyme, International Centre for Evidence-Based Medicine in Canada and Boehringer Ingelheim. A. Papi reports grants and personal fees from GlaxoSmithKline, AstraZeneca, Boehringer Ingelheim, Chiesi Farmaceutici, Menarini and Sanofi/Regeneron; personal fees from Mundipharma, Zambon, Novartis, Edmond Pharma and Roche; and grants from Fondazione Maugeri and Fondazione Chiesi. C.F. Vogelmeier has delivered presentations at symposia and/or served on scientific advisory boards sponsored by Aerogen, AstraZeneca, Boehringer Ingelheim, CSL

Behring, Chiesi, GlaxoSmithKline, Grifols, Menarini, Novartis, Nuaira and MedUpdate. M. Kupczyk reports grants from AstraZeneca and personal fees from AstraZeneca, Chiesi, GlaxoSmithKline, Novartis, Lekam, Alvogen, Emma, Nexter and Berlin Chemie. E. Maslova, N. Budgen and J. Bell are employees of AstraZeneca. A. Menzies-Gow has attended advisory boards for GlaxoSmithKline, Novartis, AstraZeneca, Sanofi and Teva; received speaker fees from Novartis, AstraZeneca, Vectura, Teva and Roche; participated in research with AstraZeneca; attended international conferences with Teva; and has consultancy agreements with AstraZeneca, Vectura and Sanofi.

Support statement: AstraZeneca funded the CARBON and SABINA studies and was involved in designing the study, developing the study protocol, conducting the study and performing the analyses. AstraZeneca was given the opportunity to review the manuscript before submission. Funding information for this article has been deposited with the Crossref Funder Registry.

References

- 1 United Nations. The Paris Agreement. 2015. <https://unfccc.int/process-and-meetings/the-paris-agreement/the-paris-agreement> Date last accessed: 8 March 2022. Date last updated: June 2020.
- 2 Tennison I, Roschnik S, Ashby B, *et al.* Health care's response to climate change: a carbon footprint assessment of the NHS in England. *Lancet Planet Health* 2021; 5: e84–e92.
- 3 Janson C, Henderson R, Löfdahl M, *et al.* Carbon footprint impact of the choice of inhalers for asthma and COPD. *Thorax* 2020; 75: 82–84.
- 4 Public Health England, NHS England. Reducing the Use of Natural Resources in Health and Social Care. 2018. https://networks.sustainablehealthcare.org.uk/sites/default/files/resources/20180912_Health_and_Social_Care_NRF_web.pdf Date last accessed: 8 March 2022. Date last updated: October 2020.
- 5 National Health Service. Delivering a 'Net Zero' National Health Service. www.england.nhs.uk/greenernhs/wp-content/uploads/sites/51/2020/10/delivering-a-net-zero-national-health-service.pdf Date last accessed: 8 March 2022. Date last updated: October 2020.
- 6 Tran TN, King E, Sarkar R, *et al.* Oral corticosteroid prescription patterns for asthma in France, Germany, Italy and the UK. *Eur Respir J* 2020; 55: 1902363.
- 7 Ding B, Small M. Disease burden of mild asthma: findings from a cross-sectional real-world survey. *Adv Ther* 2017; 34: 1109–1127.
- 8 Global Initiative for Asthma. Global Strategy for Asthma Management and Prevention. 2021. Available from: <http://ginasthma.org/> Date last accessed: 8 March 2022. Date last updated: June 2021.
- 9 Janson C, Menzies-Gow A, Nan C, *et al.* SABINA: an overview of short-acting β_2 -agonist use in asthma in European countries. *Adv Ther* 2020; 37: 1124–1135.
- 10 Bloom CI, Cabrera C, Arnetorp S, *et al.* Asthma-related health outcomes associated with short-acting β_2 -agonist inhaler use: an observational UK study as part of the SABINA global program. *Adv Ther* 2020; 37: 4190–4208.
- 11 Nwaru BI, Ekström M, Hasvold P, *et al.* Overuse of short-acting β_2 -agonists in asthma is associated with increased risk of exacerbation and mortality: a nationwide cohort study of the global SABINA programme. *Eur Respir J* 2020; 55: 1901872.
- 12 Pavord ID, Mathieson N, Scowcroft A, *et al.* The impact of poor asthma control among asthma patients treated with inhaled corticosteroids plus long-acting β_2 -agonists in the United Kingdom: a cross-sectional analysis. *NPJ Prim Care Respir Med* 2017; 27: 17.
- 13 Anzueto A. Impact of exacerbations on COPD. *Eur Respir Rev* 2010; 19: 113–118.
- 14 Sustainable Healthcare Coalition. Care Pathway Carbon Calculator. Available from: <https://shcpathways.org/> Date last accessed: 8 March 2022.
- 15 Wilkinson A, Maslova E, Janson C, *et al.* Environmental sustainability in respiratory care: an overview of the healthCARE-Based environmental Cost of Treatment (CARBON) programme. *Adv Ther* 2022; 39: 2270–2280.
- 16 Cabrera CS, Nan C, Lindarck N, *et al.* SABINA: global programme to evaluate prescriptions and clinical outcomes related to short-acting β_2 -agonist use in asthma. *Eur Respir J* 2020; 55: 1901858.
- 17 Di Marco F, D'Amato M, Lombardo FP, *et al.* The burden of short-acting β_2 -agonist use in asthma: is there an Italian case? An update from SABINA Program. *Adv Ther* 2021; 38: 3816–3830.
- 18 Worth H, Crieé CP, Vogelmeier CF, *et al.* Prevalence of overuse of short-acting beta-2 agonists (SABA) and associated factors among patients with asthma in Germany. *Respir Res* 2021; 22: 108.
- 19 Quint JK, Arnetorp S, Kocks JWH, *et al.* Short-acting β_2 -agonist exposure and severe asthma exacerbations: SABINA findings from Europe and North America. *J Allergy Clin Immunol Pract* 2022; in press [<https://doi.org/10.1016/j.jaip.2022.02.047>].
- 20 Jeswani HK, Azapagic A. Life cycle environmental impacts of inhalers. *J Cleaner Prod* 2019; 237: 117733.
- 21 FitzGerald JM, Tavakoli H, Lynd LD, *et al.* The impact of inappropriate use of short acting beta agonists in asthma. *Respir Med* 2017; 131: 135–140.
- 22 Kupczyk M, Barg W, Bochenek G, *et al.* Overprescription of short-acting beta2-agonists in asthma management? Pharmacy reports from 91,673 patients in Poland. *Eur Respir J* 2019; 54: Suppl. 63, OA2107.

- 23 European Commission. State of Health in the EU: Poland, Country Health Profile. 2017. Available from: www.euro.who.int/__data/assets/pdf_file/0006/355992/Health-Profile-Poland-Eng.pdf?ua=1 Date last accessed: 8 March 2022.
- 24 Statistics Canada, Government of Canada. Socio-economic Status in Canadian Provinces. www150.statcan.gc.ca/n1/pub/81-590-x/2007001/tables/5002494-eng.htm Date last accessed: 8 March 2022. Date last updated: November 2008.