

An assessment of preventive care offered to orthodontic patients by oral health therapists in NSW Australia

Angela V. Masoe¹, Anthony S. Blinkhorn², Jane Taylor¹ and Fiona A. Blinkhorn¹

¹School of Health Sciences, Faculty of Health and Medicine, Oral Health, University of Newcastle, Ourimbah, NSW, Australia; ²Department of Population Oral Health, Faculty of Dentistry, University of Sydney, Westmead, NSW, Australia.

Objective: The aim of this study was to record preventive oral health care planned by dental therapists and oral health therapists (therapists) for patients with poor oral hygiene undergoing orthodontic treatment. **Materials and methods:** A cross-sectional survey, using a clinical vignette of a patient with poor oral hygiene undergoing therapy with a fixed appliance, was undertaken to record the preventive care offered to this individual by therapists working across 15 Local Health Districts (LHDs). This orthodontic vignette was inserted between two dental caries-related vignettes. Data were coded and descriptive statistics were used to report the findings. **Results:** One-hundred and seventeen therapists returned questionnaires (giving a response rate of 64.6%), of whom 82.0% ($n = 95$) completed the orthodontic vignette. Adopting motivational interviewing techniques to facilitate communication with the patient and their parent was recommended by 88.4% ($n = 84$) respondents, 98.0% ($n = 93$) offered oral-hygiene instruction, 70.5% ($n = 67$) recorded plaque levels and used disclosing solution and 60.0% ($n = 57$) offered dietary advice. Products recommended for use at home included fluoride toothpaste [1,450 ppm F (80.0%; $n = 76$) and 5,000 ppm F (59.0%; $n = 24$)] and casein phosphopeptide amorphous phosphates plus fluoride (CPP-ACPF) paste (33.3%; $n = 32$). Less than 20% offered fissure sealants. **Conclusion:** Preventive advice and care was offered inconsistently by therapists in this study. To ensure that all therapists adopt a scientifically based approach to prevention, LHD clinical directors should implement continuous professional education programmes for therapists to improve patient's health outcomes.

Key words: Orthodontic adolescent patient, public oral health, preventive care

INTRODUCTION

Orthodontic care is available to disadvantaged eligible adolescents under 18 years of age through the New South Wales (NSW) public oral health services, providing that they or their parents are holders of a government Medicare health-care card^{1–3}. The policy criteria for referral of patients to specialist orthodontists states that 'eligible patients who have active dental caries, chronic marginal gingivitis or whose oral hygiene is not at an optimal level should not be offered orthodontic treatment'¹. Additionally, the policy clearly stipulates that if the patient is unable to maintain acceptable oral hygiene standards during treatment and does not respond to an improvement

programme, orthodontic treatment should be discontinued¹.

Orthodontic treatment often commences during adolescence, which is a significant period for behaviour, personality and self-image development⁴. Orthodontic brackets, wires, ligatures and other parts of the appliance create areas that encourage accumulation of plaque biofilm and food stagnation, which present challenges for adolescents regarding maintaining a daily optimal level of oral hygiene during the treatment period^{4,5}.

Enamel demineralisation [white spot lesions (WSLs)] is an adverse complication associated with therapy with fixed appliances, as is chronic hyperplastic gingivitis with increased pocket depths, with slight, yet significant, loss of periodontal support associated with retention of plaque biofilm^{5–8}. Increased gingivitis and gingival hyperplasia are reported as problems during orthodontic treatment; however, these rarely lead to periodontitis⁹. Salivary flow is altered by an

The authors are responsible for the content of this study and do not reflect the views of the NSW Ministry of Health or the funding Agency.

orthodontic fixed appliance, which interrupts the saliva's 'self-cleansing' action of eliminating food waste, leading to demineralisation of the dental enamel^{5,10}.

Several factors are associated with increased risk of developing WSLs, dental caries and gingivitis:

- The adolescent's ability to maintain motivation for an effective daily oral hygiene regime
- Build up of the plaque biofilm, which increases oral bacterial activity and decreases salivary pH
- Adolescents tend not to follow advice or do not see themselves as vulnerable to health problems
- A high consumption of carbonated drinks and sugar-containing snacks^{4,5}.

The NSW public health system offers all adolescents under 18 years of age free oral health care and preventive advice, which is provided, in the majority of cases, by dental therapists and oral health therapists (therapists)³. These clinicians have a fundamental role and responsibility for the prevention of oral disease, particularly dental caries and periodontal disease^{10–12}. Thus, they are able to assist an orthodontic patient's treatment pathway by providing much needed regular preventive care and advice^{5,6,10}. However, researchers found that therapists¹¹ reported that therapists in rural areas offered less preventive care to their patients because of a greater demand for emergency dental care and urgent treatment compared with metropolitan areas. A study investigating the provision of dental care to more than 29,000 adolescent patients in NSW over a 1-year period reported that the offer of preventive care and advice varied from 32% to 55% of therapists' clinical activity¹³.

The development of WSLs and gingivitis, leading to the need for periodontal treatment during orthodontic treatment, is preventable¹⁰. Researchers recommend that dentists (and therapists) should assess each patient's risk factors at the initial visit and throughout their course of care and offer appropriate preventive agents and therapies, such as fluoride treatments, antimicrobials, xylitol gum, casein phosphopeptide amorphous calcium phosphate (CPP-ACP) paste, dietary counselling and oral hygiene instruction^{5,6,10}. There is currently a dearth of information on the clinical preventive practice of therapists, particularly for orthodontic patients accessing the NSW public oral health service. Therefore, the purpose of this study was to use a clinical vignette to record what preventive oral health care therapists would offer to an adolescent patient undergoing orthodontic treatment.

METHODS

Clinical vignettes have been defined as re-creations of actual clinical scenarios that can be used to elicit participants' knowledge, attitudes and perceptions in accordance with their clinical practice in their

natural milieu^{14–16}. Vignettes may be distinct and standardised, enabling all participants to respond to the same stimulus¹⁷. A cross-sectional self-administered survey using a clinical vignette for therapists working within all 16 Local Health Districts (LHDs) of NSW Health was developed. The survey also encompassed demographic information about the participants.

Based on research literature and academic curriculum teaching, clinical problems commonly seen in adolescents were chosen by an advisory team made up of two paediatric dental specialists, an academic clinical curriculum convenor and two experienced therapists. Three vignettes based on these problems were created, and photographs were used to help the therapists visualise the clinical issues. The orthodontic case scenario for this study was inserted between the two dental caries vignettes that are not reported in this paper. The clinical intra-oral photograph was provided, with full written consent, by the patient and guardian for research use, with personal details and location de-identified.

The orthodontic vignette was designed using the classic clinical dental presentation, with a focus on: (i) history of the chief complaint; (ii) overall dental history; (iii) clinical examination; and (iv) diagnostic tests. These were used by the therapists to develop an assessment and management plan. The orthodontic vignette described a 14-year-old male patient (TJ) who presented with his mother because of concerns regarding a halitosis problem (*Figure 1*). Therapists were requested to use the scenario description, photograph and charting provided for the vignette, aligned with their clinical practice protocols¹⁸, to respond to the following questions:

- 1 What treatment, if any, would you carry out for TJ today?
- 2 Would you bring TJ back to the clinic for treatment, if yes, what?

The vignette was pilot tested with five therapists who were working in the Australian Capital Territory, and minor amendments were made before starting the main survey.

The names and contact details for all therapists working within the NSW public oral health services were obtained by contacting directors of each of the 15 LHDs. An information document outlining the research and inviting participants to consent voluntarily to participate by completing and returning the survey was developed. One-hundred and ninety-two potential participants were identified. Survey questionnaires, an information document and return postage-paid envelopes were mailed and reminder letters were posted out 2 weeks later. Further reminders to non-respondents were undertaken 1, 2 and 3 months after the initial mailing.

TJ is a 14 year old male patient. His mother says that all of the family are complaining of TJ's bad breath. TJ has not noticed anything of concern, except the braces are hard to keep clean. His mother is more concerned about his teeth than he is. TJ is a healthy young lad with no history of any serious illness. TJ is a regular dental attender and has accessed public health services all his life.

He is under the orthodontist for treatment of a Class II Division I malocclusion and his next appointment with the orthodontist is in 4 weeks' time. He had 14 & 24 extracted under LA in order to commence orthodontic treatment 12 months ago. He has never required restorations in primary or permanent dentitions.

TJ is the youngest of 3 children in the family. He lives in Sydney. The parents are both employed intermittently with local commercial companies. He lives close to the dental clinic. TJ has a good diet according to mum. He drinks tap water and soft drinks are only bought occasionally; sweets are available in the form of biscuits, cakes and muesli bars for school lunches.

Clinical Observation:

Extra-oral: No abnormality detected.

Intra-oral: Oral hygiene: poor. Plaque present on most teeth. No calculus present.

Gingivae: Red and inflamed – chronic marginal gingivitis, bleeding on gentle probing.

Occlusion: Skeletal Class 1, Dental Class 1

Using the information, photograph and dental chart you have been provided with:

1. What treatment, if any, would you carry out for TJ today?
2. Would you bring TJ back to the clinic for treatment, if yes, what?




Figure 1. Adolescent orthodontic presentation.

A coding index system was constructed from the first 35 written responses, guided by The Australian National Dental Schedule System¹⁹. These codes were reviewed, amended and confirmed in consultation with the advisory team, with subsequent responses coded and entered into a Microsoft Excel database; these were later collated into key clinical preventive categories. Respondent's narratives to clarify clinical decisions were also uploaded and analysed. To ensure rigor, two independent non-clinician oral health professionals were recruited to review and confirm data entry, data coding and narrative upload before data analysis. The advisory group systematically reviewed and verified data-analysis processes within specific time frames.

Ethics approval for the study was obtained from the Hunter New England Local Health District Lead Health and Research Ethics Committee (HREC) Reference No. 12/02/15/5.04 and all 15 LHDs. This research was conducted in full accordance with the World Medical Association Declaration of Helsinki.

RESULTS

Following the initial mail out, further information was received on therapist numbers. The original sample of 192 was reduced by 11 because of retirements and job changes, giving a final sample of 181, of whom 117 (64.6%) responded. Respondents were: (i) dental therapists (79.1%; $n = 91$); (ii) dental hygienists (1.7%; $n = 2$); and (iii) oral health therapists (20.5%; $n = 24$). Most (61.5%; $n = 72$) respondents worked in rural LHDs compared to metropolitan LHDs (38.4%; $n = 45$). The mean time \pm standard deviation since completion of their academic qualification was 21.9 ± 12.7 years.

Ninety-five (82.0%) of the 117 respondents completed the orthodontic vignette reported in this paper. Therapists noted that this patient had halitosis because he was having difficulties cleaning his teeth as a result of his fixed orthodontic appliance. The types of preventive oral health care recommended for the patient by the therapists are shown in *Table 1*.

Table 1 Therapists' record of immediate oral health treatment for the orthodontic patient ($n = 95$)

Treatment option	<i>n</i>	%
Comprehensive oral examination	28	27.5
Bitewing radiographs	15	15.8
Intra-oral photographs	6	6.3
Plaque Index (PI) or Periodontal Screening Record (PSR)	7	7.4
Plaque disclosing	67	70.5
Oral hygiene, including tongue cleaning instruction	93	98.0
Use of a hand mirror as oral hygiene instruction patient-learning tool	78	82.2
Use Motivational Interviewing (coaching technique) for TJ (<i>and parent</i>)	84	88.4
Dietary advice	57	60.0
Super Floss and Piksters	60	63.1
Professional cleaning (plaque and calculus removal)	65	68.4
Topical fluoride applications (varnish and gels)	45	47.3
Fluoride toothpaste (Colgate Total antibacterial, 1,450 ppm F)	76	80.0
Recommend use of Neutra Fluor 5000 (5,000 ppm F)	24	25.3
Recommend use of mouth rinse (chlorhexidine, antibacterial agents, saline rinses and Neutra Fluor 900 ppm F)	23	24.2
Recommend use of Tooth Mousse plus fluoride (CPP-ACPF)	32	33.7
Fissure sealants	17	17.9
Issue oral health products and relevant brochures	15	15.8
Seek medical practitioner advice (if oral health practices improve and halitosis persists as it may be caused by other underlying health issues)	25	26.3

CPP-ACPF, casein phosphopeptide amorphous phosphates plus fluoride.

Immediate care (Question 1) consisted of an oral examination (27.5%; $n = 28$), bitewing radiographs (15.8%; $n = 15$), plaque disclosing ($n = 67$; 70.5%), recording the Plaque Index or Periodontal Screening Record (7.4%; $n = 7$) and providing oral hygiene instruction and advice on tongue cleaning (98.0%; $n = 93$) (Table 1). As the chief complaint was halitosis, the majority of respondents (82.2%; $n = 78$) focused on toothbrushing instruction, which included asking the patient whether he cleaned his tongue. Education regarding the importance of tongue cleaning to reduce the bacterial load, which might be contributing to the halitosis, was part of the care plan.

The use of super-floss and piksters as tools to improve cleaning between teeth was recorded by 63.1% ($n = 60$) of the respondents. Motivational interviewing techniques, such as having the patient demonstrate how he currently brushed his teeth, followed by the therapist using a hand mirror to show where he needed to improve, was suggested by 88.4% ($n = 84$) of the participants. The majority (68.4%; $n = 65$) reported that after offering oral hygiene

instruction, they would undertake a professional clean (removal of plaque and calculus). Sixty per cent ($n = 57$) provided dietary advice with a focus on sugar consumption and its impact on bacterial growth as a contributor to halitosis. Placement of fissure sealants was recorded by 17.9% ($n = 17$) of the respondents.

Eighty per cent ($n = 76$) stated that they would offer advice on fluoride toothpaste (1,000–1,450 ppm F), including use of Neutra Fluor 5000 (25.3%, $n = 24$) and Tooth Mousse (33.7%, $n = 32$), for the management of demineralised enamel areas. Mouth rinses were also recommended (24.2%, $n = 23$), which included the use of antibacterial agents, such as chlorhexidine, Neutra Fluor 900 mouthwash (900 ppm F, once weekly) and saline rinses in conjunction with the oral hygiene regime.

When asked if they would bring the patient back for further treatment (Question 2), 44.2% ($n = 42$) stated they would bring the patient back weekly until the condition had stabilised and to monitor the oral hygiene home practices and gingival health (Table 2). Approximately one-third (28.4%, $n = 27$) stated that they would give him fortnightly appointments, and if there was no improvement or there was evidence of further deterioration of gingival health they would inform TJ that an early referral back to the orthodontist for de-banding was almost inevitable to prevent further oral health issues. Less than 5% ($n = 3$) recommended bringing the patient back in 1 month;

Table 2 Therapists' management plan for follow-up of the orthodontic patient ($n = 95$)

Would you bring TJ back, if yes, what treatment would you provide?	<i>n</i>	%
Weekly follow-up	42	44.2
Review patient oral hygiene and preventive home-care practices		
Provide oral hygiene support where indicated		
Seek medical practitioner advice if halitosis has not improved		
Fortnightly follow-up	27	28.4
Review patient oral hygiene and preventive home care practices		
Communicate with orthodontist if condition has not improved with consideration for de-band (therapist caution TJ and parent)		
Seek medical practitioner advice if halitosis has not improved		
One-month follow-up	3	3.1
Recall appointment, general review of patient's oral hygiene practices		
Repeat above oral hygiene and home-care advice		
Seek medical practitioner advice if halitosis has not improved		
6-month recall appointment	15	15.8
Review and undertake general oral health care examination		
12-month general oral health recall	5	5.3
Regular oral health review		

15.8% ($n = 15$) suggested a 6-month review and 5.3% ($n = 5$) offered re-appointment at 12 months (Table 2). Just over one-quarter (26.3%; $n = 25$) stated that they would discuss other possible reasons for halitosis, suggesting that TJ and his mother should seek medical advice if the problem did not improve (Table 1).

DISCUSSION

The objective of this study was to record therapists' planned preventive oral health care for an orthodontic adolescent patient using a clinical vignette. Most adolescents accessing public oral health systems are from disadvantaged backgrounds and the working poor²⁰. There is a dearth of research in the area of preventive clinical practices to support public orthodontic patients; thus, this study, utilising a clinical vignette to record therapists' preventive care plans, has provided valuable information. Furthermore, there is a major flaw in the NSW public oral health system for tracking referral of eligible adolescent patients to orthodontic specialist services and follow-up care. Thus, there is scope for future clinical research into the referral and feedback processes to ensure continuous quality care of patients.

NSW Health, which commissions the public dental service, does not have a policy or protocol specifically for preventive care for patients under orthodontic treatment. However, there are general preventive oral health care policies on the use of pit-and-fissure sealants, topical fluorides and smoking-cessation advice^{21–24}. Therapists should apply these same principles to all orthodontic patients.

This study found fairly adequate levels of motivational interviewing to facilitate communication with the patient to enhance his oral hygiene instruction, including the use of a hand mirror as an educational tool for demonstrating an interactive learning session. However, the infrequent use of the plaque index and periodontal screening recorded by the therapists is a major concern considering the patient's very poor gingival health and halitosis. Furthermore, the use of tri-plaque disclosing solutions, a relatively simple procedure, should have been recommended by all therapists to monitor the patient's current and future oral hygiene practices^{5,25}.

Hadler-Olsen *et al.*'s⁵ public health study in Norway, of adolescents under 16 years of age, reported that assessment of plaque levels using plaque-disclosing tablets in conjunction with oral hygiene instructions was important. Adolescents were provided with an oral health kit containing an orthodontic toothbrush, interdental brushes, plaque-disclosing tablets, fluoride toothpaste and mouth rinse to facilitate the preventive regime⁵. Those ($n = 9$) who complied with

the comprehensive oral hygiene regime developed, on average, one new WSL, patients with moderate compliance ($n = 27$) developed 1.4 WSLs and those ($n = 4$) with poor compliance developed 3.3 WSLs⁵. Instituting a comprehensive oral hygiene regimen for orthodontic patients was reported as challenging by Hadler-Olsen *et al.*⁵, but it illustrates the importance of providing rigorous preventive measures for orthodontic patients.

A study by Derk *et al.*²⁶ reviewed the use, by orthodontic practices, of measurements to prevent decalcification during fixed appliance treatment, and comparison of these measures with the available scientific evidence found that many orthodontists failed to implement procedures in their dental establishments to prevent enamel demineralisation. Thus, these authors²⁶ recommended the development of practice guidelines for the prevention of enamel demineralisation. If this strategy is to be used by public oral health services, then clinical quality-improvement mechanisms would have to be adopted to monitor the implementation and compliance of clinicians to ensure improved patient health outcomes, as part of clinical governance²⁷.

Dietary advice was offered by 60% of the respondents, which is disappointing considering the critical role that sugar plays in plaque formation and in the aetiology of dental caries. Some form of advice on restricting sugary foods and drinks between meals, and healthy alternatives for snacks, should have been part of the immediate care option^{6,10}.

Only a small proportion (17.9%) suggested that fissure sealants should be offered at this visit, which shows that the majority of respondents had focused on the main clinical problems of gingivitis and halitosis. The patient was described as caries free, so sealants were not an urgent requirement.

Application of topical fluoride varnish (and gels) recorded in this study was low (47.3%), and as the patient is described as having no previous dental caries, it is not an immediate and urgent issue to discuss fluoride and other remineralising products. This is better left to future visits because too much information at the initial visit will confuse both the mother and the patient. However, because of the increased caries risk in fixed appliance therapy it is vital that the patient is given this information in an early follow-up appointment.

The majority of respondents did recommend the use of 1,450 ppm F toothpaste (80%); 25.3% recommended 5,000 ppm F and 33.7% offered CPP-ACPF. A study conducted by Sonesson *et al.*⁸ to establish the efficacy of daily toothbrushing with 5,000 ppm F toothpaste on enamel demineralisation, found that the prevalence of WSLs was significantly lower in the group using the high-fluoride toothpaste ($P = 0.04$). There was an 18.1% incidence of WSL in the high-fluoride

toothpaste group compared with an incidence of WSL of 26.6% in the reference group⁸. Therefore, an orthodontic patient's WSL risk should be assessed, and use of a suite of remineralising agents, such as 5,000 ppm F toothpaste, in conjunction with CPP-ACP agents, should be discussed and advice offered at future visits, to ensure that optimal levels of calcium, phosphate and fluoride ions are present in the saliva to support enamel remineralisation during orthodontic treatment¹⁰.

Gingivitis, leading to gingival enlargement (gingival hyperplasia), can be controlled by adopting high standards of oral hygiene. However, Zachrisson and Zachrisson's longitudinal study of 49 patients, 11-13 years of age at the commencement of treatment, reported that despite good oral hygiene and sodium fluoride rinsing performed twice weekly throughout the study, most patients developed generalised moderate gingival hyperplasia. The authors reported that gingival health improvement was noted after the first month of orthodontic band removal. Conversely, the review by Blinkhorn *et al.*²⁸, relating to the effectiveness, safe delivery and use, by patients, of triclosan/copolymer toothpaste, found strong support for its positive medicinal effect on preventing biofilm formation and promoting gingival health with twice-daily use to control plaque and slow progression of periodontal disease. Thus, LHDs in NSW should review the scientific efficacy of oral health products regularly to ensure their appropriate prescription to assist patients' oral health home regimes.

Of concern, this study found inconsistencies in the follow-up time frames used to monitor the patient's oral hygiene status. Considering the presenting clinical oral health status of the patient, 20% of the respondents recorded a follow-up time frame of between 6 and 12 months, which is deemed far too long to offer support to the patient ensuring improved oral health outcomes. A study by Bardal *et al.*²² reported findings over a period of 6 months; they found that monitoring oral hygiene at 6, 12 and 24 weeks gave positive results in terms of good gingival health for orthodontic patients. Therefore, a review of time frames used by public health systems for vulnerable adolescents undergoing fixed orthodontic treatment should be developed and publicised.

It was somewhat surprising to note that 15.8% of the respondents would take bitewing radiographs at the first visit. The diagnostic yield will be compromised by the orthodontic brackets and the patient is a low caries risk. The heavy deposits of plaque will predispose the patient to smooth surface lesions, which will not be shown on a bitewing. The US Food and Drug Administration, in collaboration with the American Dental Association²⁹, urges dental professionals to minimise radiation exposure. A patient who is receiving orthodontic care will have a full

treatment-planning schedule of radiographs; therefore, it is most unwise to prescribe further radiographs that will be of little diagnostic value.

A potential limitation of this study was capturing and reporting the multifaceted preventive activities during the communication interplay between the clinician and the patient. Similar difficulties in recording clinical-preventive activities have been previously reported by Tickle *et al.*³⁰. Therefore, caution should be exercised in the generalisation of this study's findings.

Nonetheless, this study, utilising a vignette as a way of replicating a real event to elicit therapists' clinical preventive care planning according to how they would behave in clinical practice, has yielded new information to assist the public oral health services of NSW to develop clinical preventive care quality improvement programmes.

CONCLUSION

Preventive oral health strategies reported by respondents for the clinical management of a patient undergoing fixed orthodontic treatment varied markedly. It is recommended that rigorous preventive care and clinical treatment for adolescents should be embedded in the clinical practice of therapists for disadvantaged and high-risk patients. Clinical directors should provide therapists with ongoing scientific professional education on the management of dental caries and periodontal disease, including the resourcing of relevant oral health products for offering to patients, in order to ensure good clinical outcomes. Additionally, evaluation mechanisms to monitor implementation and compliance to NSW Health preventive policies and protocols should be a component of annual clinical governance processes.

Acknowledgements

This study was funded by NSW Ministry of Health Rural and Remote Allied Health Professional Scholarship Scheme and Centre for Oral Health Strategy. We are grateful to Australian Capital Territory therapists and hygienists who participated in pilot testing the survey questionnaire; Rachael Moir research officer, University of Newcastle for research data assistance; NSW LHDs clinical directors, health service managers and NSW Oral Health Promotion Network members for supporting the research activities.

Author contributions

All authors contributed to the design of the study. AVM, ASB and FAB participated in analysing the data and drafting the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

REFERENCES

1. Centre for Oral Health Strategy NSW. *Oral Health Specialist Referral Protocol*. In: NSW COHS. Sydney: NSW Health; 2011. Available from: http://www0.health.nsw.gov.au/policies/pd/2011/pdf/PD2011_071.pdf. Accessed 03 December 2014.
2. Centre for Oral Health Strategy NSW. Oral health – eligibility of persons for public oral health care in NSW, 2013. Available from: http://www0.health.nsw.gov.au/policies/pd/2009/pdf/PD2009_074.pdf. Accessed 12 March 2014.
3. NSW Ministry of Health. *Oral Health 2020: A Strategic Framework for Dental Health in NSW*. Sydney: NSW Health; 2013. Available from: <http://www.health.nsw.gov.au/oralhealth/Publications/Oral-Health-2020.pdf>. Accessed 03 December 2014.
4. Al-Jewair TS, Suri S, Tompson BD. Predictors of adolescent compliance with oral hygiene instructions during two-arched multibracket fixed orthodontic treatment. *Angle Orthod* 2011 81: 525–531.
5. Hadler-Olsen S, Sandvik K, El-Agroudi MA *et al.* The incidence of caries and white spot lesions in orthodontically treated adolescents with a comprehensive caries prophylactic regimen-a prospective study. *Eur J Orthod* 2012 34: 633–639.
6. Chang HS, Walsh LJ, Freer TJ. Enamel demineralisation during orthodontic treatment. Aetiology and prevention. *Aust Dent J* 1997 42: 322–327.
7. Chapman JA, Roberts WE, Eckert GJ *et al.* Risk factors for incidence and severity of white spot lesions during treatment with fixed orthodontic appliances. *Am J Orthod Dentofac Orthop* 2010 138: 188–194.
8. Sonesson M, Twetman S, Bondemark L. Effectiveness of high-fluoride toothpaste on enamel demineralisation during orthodontic treatment-a multicentre randomized controlled trial. *Eur J Orthod* 2013: 1–5. Available from: <http://ejo.oxfordjournals.org/content/early/2013/12/27/ejo.cjt096>. Accessed 11 January 2015
9. Zachrisson S, Zachrisson BU. Gingival condition associated with orthodontic treatment. *Angle Orthod* 1972 42: 26–34.
10. Sudjalim TR, Woods MG, Manton DJ. Prevention of white spot lesions in orthodontic practice: a contemporary review. *Aust Dent J* 2006 51: 284–289.
11. Satur J, Gussy M, Marino R *et al.* Patterns of dental therapists' scope of practice and employment in Victoria, Australia. *J Dent Educ* 2009 73: 416–425.
12. Nash DA. Envisioning an oral healthcare workforce for the future. *Community Dent Health* 2012 2: 141–147.
13. Masoe AV, Blinkhorn AS, Taylor J *et al.* Preventive and clinical care provided to adolescents attending public oral health services New South Wales, Australia: a retrospective study. *BMC Oral Health* 2014 14: 1–9.
14. Gould D. Using vignettes to collect data for nursing research studies: how valid are the findings. *J Clin Nurs* 1996 5: 207–212.
15. Wilson J, While AE. Methodological issues surrounding the use of vignettes in qualitative research. *J Interprof Care* 1998 12: 79–86.
16. International Training and Education Centre. Structured clinical vignettes: what are they and how are they used? Available from: <http://www.go2itech.org/HTML/CM08/toolkit/tools/vignettes.html>. Accessed 26 August 2014.
17. Peabody JW, Luck J, Glassman P *et al.* Measuring the quality of physician practice by using clinical vignettes: a prospective validation study. *Annals of International Medicine* 2004 141: 771–780.
18. Centre for Oral Health Strategy NSW. *Oral Health Record Protocols*. NSW Ministry of Health: NSW Health Australia; 2008. Available from: http://www0.health.nsw.gov.au/policies/pd/2008/pdf/PD2008_024.pdf. Accessed 12 March 2014.
19. Australian Dental Association Inc. *The Australian Schedule of Dental Services and Glossary*, 10th ed. Australian Dental Association Inc.; 2013. Available from: http://www.ada.org.au/app_cmslib/media/lib/1303/m538110_v1_10th%20edition%20schedule%20and%20glossary_web.pdf. Accessed 03 December 2014.
20. Australia's National Oral Health Plan 2015-2024(Draft). Available from: <http://oralhealthplan.com.au/australias-oral-health>. Accessed 12 October 2014.
21. Centre for Oral Health Strategy NSW. Pit and fissure sealants: use of in oral health services. Available from: http://www0.health.nsw.gov.au/policies/pd/2013/pdf/PD2013_025.pdf. Accessed 30 January 2013.
22. Centre for Oral Health Strategy NSW. Fluorides – use of in NSW. Available from: <http://www.health.nsw.gov.au/environment/water/Documents/PD2006-076-Use-of-Fluorides-in-NSW.pdf>. Accessed 12 December 2014.
23. Centre for Oral Health Strategy NSW. Smoking cessation brief intervention at the chairside: role of public oral health/dental service, 2009. Available from: http://www0.health.nsw.gov.au/policies/pd/2009/pdf/PD2009_046.pdf. Accessed 12 December 2013.
24. Davies RM, Blinkhorn AS. Preventing Dental Caries: Part 1 the scientific rationale for preventive advice. *Dental Update* 2013 22: 24–26.
25. Worthington HV, Hill KB, Mooney J *et al.* A cluster randomized controlled trial of a dental health education program for 10-year-old children. *J Public Health Dent* 2001 61: 22–27.
26. Derks A, Kuijpers-Jagtman AM, Frencken JE *et al.* Caries preventive measures used in orthodontic practices: an evidence-based decision? *Am J Orthod Dentofac Orthop* 2007 132: 165–170.
27. NSW Health. NSW State Health Plan: Towards 2021, Sydney. Available from: <http://www.health.nsw.gov.au/statehealthplan/Pages/NSW-State-Health-Plan-Towards-2021.aspx>. Accessed 09 October 2014.
28. Blinkhorn AS, Bartold PM, Cullinan MP *et al.* Is there a role for triclosan/copolymer toothpaste in the management of periodontal disease? *Br Dent J* 2009 207: 117–125.
29. US Food and Drug Administration. *The Selection of Patients for Dental Radiographic Examination*. American Dental Association; 2012. Available from: <http://www.fda.gov/RadiationEmittingProducts/RadiationEmittingProductsandProcedures/MedicalImaging/MedicalX-Rays/ucm116504.htm>. Accessed 25 January 2015
30. Tickle M, Milsom KM, King D *et al.* The influences on preventive care provided to children who frequently attend the UK General Dental Service. *Br Dent J* 2003 194: 329–332.

Correspondence to:

Angela V. Masoe,
Faculty of Health and Medicine,
School of Health Sciences, Oral Health,
University of Newcastle,
PO Box 729,
Queanbeyan,
NSW 2620,
Australia.

Email: Angela.Masoe@gsahs.health.nsw.gov.au