

CORRECTION

Open Access



Correction to: Shp2 positively regulates cigarette smoke-induced epithelial mesenchymal transition by mediating MMP-9 production

Ya-nan Liu^{1,2,3,4†}, Yan Guan^{5†}, Jian Shen^{2,6}, Yong-liang Jia^{2,6}, Jian-cang Zhou⁵, Yun Sun^{3,4}, Jun-xia Jiang², Hui-juan Shen², Qiang Shu¹, Qiang-min Xie^{1,2*}  and Yicheng Xie^{1*}

Correction to: *Respiratory Research* (2020) 21:161

<https://doi.org/10.1186/s12931-020-01426-9>

Following publication of the original article [1], the authors identified an error in Fig. 6B. During the preparation of the figures in the above article, the authors regret that an error occurred during the assembly of Fig. 6B. Erroneous duplication of E-cadherin images were

mistakenly assembled for MMP-9⁺ PPHS-1⁺ and MMP⁺ Shp2(siRNA)⁺ group.

The authors apologize for any inconvenience caused, and have confirmed that the conclusions were not affected.

The correct Fig. 6B is given in this correction article.

The original article can be found online at <https://doi.org/10.1186/s12931-020-01426-9>.

[†]Ya-nan Liu and Yan Guan contributed equally to this work*Correspondence: xieqm@zju.edu.cn; ycxie@zju.edu.cn

¹The Children's Hospital, Zhejiang University School of Medicine, National Clinical Research Center for Child Health, Zhejiang 310052, Hangzhou, China Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.




Fig. 6 MMP-9 inhibition, Shp2 inhibition or Shp2 knockdown suppresses the expression of EMT-related factors induced by recombinant MMP-9 in NCI-H292 cells. **a** NCI-H292 cells with no treatment exhibit a pebble-like shape and display cell-cell contacts consistent with an epithelial morphology. The cells treated with human recombinant MMP-9 (2 µg/ml, 48 h) exhibit a fibroblast-like morphology with cellular elongation and reduction of cell-cell contacts. SB-3CT (1 µM) prevents the MMP-9-induced cellular changes and preserves normal epithelial morphology. The cells treated with recombinant MMP-9 for 48 h exhibit weaker expression of E-cadherin and stronger expression of α-SMA, compared with control. MMP-9 inhibition by SB-3CT (1 µM) alleviates the recombinant MMP-9 induced changes of E-cadherin and α-SMA expression. Scale bar = 100 µm. **b** Pharmacological inhibition or Shp2 knock down reverses the recombinant MMP-9 induced changes of E-cadherin and α-SMA expression. Scale bar = 100 µm. **c** MMP-9 inhibition by SB-3CT (1 µM) prevents the recombinant MMP-9 (2 µg/ml) induced decreases in E-cadherin expression and increases in α-SMA mRNA expression assessed by real-time PCR. $n = 3$ per group. $^{\#}p < 0.05$ compared with control (no treatment); $^{*}p < 0.05$ compared with cells treated with recombinant MMP-9. **d** Shp2 inhibition by PHPS1 (10 µM) or knock down by siRNA prevents the recombinant MMP-9 (2 µg/ml) induced decreases in E-cadherin expression and increases in α-SMA protein expression assessed by western blot. Data are expressed as mean \pm SEM of three independent experiments. $n = 3$ per group. $^{\#}p < 0.05$ compared with control (no treatment); $^{*}p < 0.05$ compared with cells treated with recombinant MMP-9. At least three independent experiments were completed for each group assessment. Data are presented as the mean \pm SEM. Statistical significance is determined by one-way ANOVA followed by the Student-Newman-Keuls test

Author details

¹The Children's Hospital, Zhejiang University School of Medicine, National Clinical Research Center for Child Health, Zhejiang 310052, Hangzhou, China. ²Zhejiang Respiratory Drugs Research Laboratory of Food and Drug Administration of China, Zhejiang University School of Medicine, Zhejiang 310058, Hangzhou, China. ³The First People's Hospital of Yancheng, Yancheng 224001, Jiangsu, China. ⁴Medical College of Yangzhou University, 11 Huaihai Road, Yangzhou 225001, Jiangsu, China. ⁵Sir Run Run Shaw Hospital, Zhejiang University School of Medicine, Zhejiang 310000, Hangzhou, China. ⁶Breath Smooth Biotech Hangzhou Co, LTD., Zhejiang 310012, Hangzhou, China.

Published online: 24 August 2022

Reference

- Liu Y, Guan Y, Shen J, Jia Y, Zhou J, Sun Y, Jiang J, Shen H, Shu Q, Xie Q, Xie Y. Shp2 positively regulates cigarette smoke-induced epithelial mesenchymal transition by mediating MMP-9 production. *Respir Res*. 2020;21(1):161. <https://doi.org/10.1186/s12931-020-01426-9>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

