




# The Brazilian Journal of INFECTIOUS DISEASES

[www.elsevier.com/locate/bjid](http://www.elsevier.com/locate/bjid)



## Original article

# Mutations in the S gene and in the overlapping reverse transcriptase region in chronic hepatitis B Chinese patients with coexistence of HBsAg and anti-HBs



Feng Ding<sup>a</sup>, Xi-Li Miao<sup>c</sup>, Yan-Xia Li<sup>a</sup>, Jin-Fen Dai<sup>a</sup>, Hong-Gang Yu<sup>a,b,\*</sup>

<sup>a</sup> Department of Gastroenterology, Renmin Hospital of Wuhan University, Hubei Province, PR China

<sup>b</sup> Institute for Gastroenterology and Hepatology, Wuhan University Medical School, Hubei Province, PR China

<sup>c</sup> Department of Clinical Laboratory, Puai Hospital Affiliated to Tongji Medical College of Huazhong University of Science & Technology, Hubei Province, PR China

## ARTICLE INFO

### Article history:

Received 28 May 2015

Accepted 31 August 2015

Available online 21 November 2015

### Keywords:

Chronic hepatitis B

Coexistence of HBsAg and anti-HBs

S gene

Reverse transcriptase

## ABSTRACT

**Background:** The mechanism underlying the coexistence of hepatitis B surface antigen and antibodies to HBsAg in chronic hepatitis B patients remains unknown.

**Aims:** This research aimed to determine the clinical and virological features of the rare pattern.

**Methods:** A total of 32 chronic hepatitis B patients infected by HBV genotype C were included: 15 carrying both HBsAg and anti-HBs (group I) and 17 solely positive for HBsAg (group II). S gene and reverse transcriptase region sequences were amplified, sequenced and compared with the reference sequences.

**Results:** The amino acid variability within major hydrophilic region, especially the "a" determinant region, and within reverse transcriptase for regions overlapping the major hydrophilic region in group I is significantly higher than those in group II. Mutation s126S/T within the "a" determinant was the most frequent change, and only patients from group I had the sQ129R, sG130N, sF134I, sG145R amino acid changes, which are known to alter immunogenicity.

**Conclusions:** In chronic patients, the concurrent HBsAg/anti-HBs serological profile is associated with an increased aa variability in several key areas of HBV genome. Additional research on these genetic mutants are needed to clarify their biological significance for viral persistence.

© 2015 Elsevier Editora Ltda. All rights reserved.

\* Corresponding author.

E-mail address: [yuhonggang@yahoo.com](mailto:yuhonggang@yahoo.com) (H.-G. Yu).

<http://dx.doi.org/10.1016/j.bjid.2015.08.014>

1413-8670/© 2015 Elsevier Editora Ltda. All rights reserved.

## Introduction

Hepatitis B virus (HBV) infection is a serious global public health problem, and its prevalence varies considerably from region to region. Worldwide, more than 350 million people are chronic carriers of HBV, and have a risk of dying from development of cirrhosis and hepatocellular carcinoma.<sup>1</sup> Clinically, the appearance of circulating hepatitis B surface antigen (HBsAg) heralds HBV infection, whereas the presence of the antibody to HBsAg (anti-HBs) usually indicates resolution of infection and is considered indicative of immunity to HBV infection.<sup>2</sup> It is generally believed that anti-HBs can neutralize and clear HBsAg. Therefore, the concomitant presence in the same serum of HBsAg and anti-HBs could be possible. However, the persistence of HBsAg associated with anti-HBs in patients with chronic HBV infection has been reported in previous studies.<sup>3-5</sup> So far, the mechanism underlying the concurrent detection of HBsAg and anti-HBs remains largely controversial.

HBV genome contains four partially overlapping open reading frames (ORF), and the "a" determinant is located at codon positions 124–147 within the major hydrophilic region (MHR) of the HBsAg. This determinant is the main target of recognition of HBsAg by anti-HBs and immune response cells during the course of the initial immune response in acute hepatitis B. HBV has been classified into eight genotypes, designated as A–H, and the most prevalent genotypes in China are genotypes B and C.<sup>5</sup> Notably, previous reports documented that, in comparison to genotype B, genotype C takes a more aggressive disease course and has a lower response rate to antiviral therapy.<sup>6-9</sup>

Several reports showed the paradoxical coexistence of HBsAg and anti-HBs might be associated with the selection of HBV immune escape mutants during chronic carriage.<sup>10-12</sup> These studies documented increased amino acid (aa) mutations in and around the "a" determinant, which is responsible for the stability and immunogenicity of HBsAg.<sup>13</sup> Nonetheless, a later report rejected this hypothesis and instead suggested that the pattern of simultaneous appearance of HBsAg and anti-HBs was related to the weak binding of anti-HBs to HBsAg.<sup>12</sup> As the reverse transcriptase (RT) region of HBV polymerase gene overlaps the HBsAg ORF, mutations within HBsAg gene might result in structural and functional alterations in the HBV reverse transcriptase with potential influence on viral replication capacity and efficacy of antiviral drugs.<sup>14,15</sup>

Therefore, the aims of this study were to investigate the prevalence of the coexistence of anti-HBs in HBsAg-positive CHB patients infected with genotype C HBV, and explore the relationship between the variability of the HBV S gene and the paradoxical serological profile.

## Materials and methods

### Patients

From January to December 2013, 1194 patients with CHB recruited from Wuhan, Huangshi, and Yichang were enrolled in the study. The inclusion criteria were HBsAg carriage

for more than six months, HBV DNA concentration of  $>1 \times 10^4$  IU/mL, serum alanine aminotransferase (ALT)  $>2$  ULN (upper limit of normal), and associated symptoms. Exclusion criteria included having received passive immunization, previous antiviral therapy, coinfection with other hepatitis viruses, or HIV. Sera were collected and stored at  $-70^{\circ}\text{C}$ . All individuals provided written informed consent before entering the study.

### Serologic testing

HBV serological markers were determined by using the chemiluminescent microparticle immunoassay (CMIA) technique with an Architect-i2000 automatic analyzer (Abbott Laboratories, USA). Commercially available kits were purchased from Abbott Laboratories. The analytical threshold of anti-HBs was defined as 10 mIU/mL. Serum was diluted according to the Manual Dilution Procedure if HBsAg concentration value  $>250,00$  IU/mL. ALT was assayed with an ADVIA automatic biochemical analyzer (Siemens, Germany).

### HBV DNA quantification

HBV DNA was quantified in serum using a commercially available real time fluorescence quantitative kit (Da An, China) with a lower detection threshold of 500 copies/mL.

### Serum DNA extraction

Viral DNA was extracted from 200  $\mu\text{L}$  of serum using QIAamp DNA blood mini kit (Qiagen, Germany) according to the manufacturer's instructions.

### PCR amplification and sequencing of PCR fragment

Amplification and sequencing of the full-length S gene and the overlapping RT regions were performed using the protocol as follows. The primer sequences were synthesized by Beijing SBS Biotechnology Company. PCR was carried out in a 50- $\mu\text{L}$  reaction mixture containing 10  $\mu\text{L}$  of HBV DNA template, 0.2  $\mu\text{M}$  each primer, 0.2 mM of each dNTP, 2 mM MgCl<sub>2</sub>, and 1  $\mu\text{L}$  of high-fidelity Taq polymerase (Takara, China). Besides, PCR was performed with hot start and denaturation at  $94^{\circ}\text{C}$  for 5 min, 35 cycles at  $94^{\circ}\text{C}$  for 1 min, at  $50^{\circ}\text{C}$  for 1 min, and at  $72^{\circ}\text{C}$  for 2 min, then incubated at  $72^{\circ}\text{C}$  for 5 min. In order to obtain a complete description of HBV quasispecies, sequence analysis was performed with the powerful ultra-deep sequencing (UDS) approach, using the primers according to the reference sequence X04615 (the upstream primer: 5'-GTCACCATATTCTGGGAAC-3' nt 2818-2837; the downstream primer: 5'-CATATCCCATGAAGTTAAGG-3' nt 888-869).<sup>12</sup>

### Sequence analysis

Genomic sequences obtained for the S gene and the overlapping RT regions were translated into aa sequences and compared with HBV reference sequences used on the NCBI Website (<http://www.ncbi.nih.gov/projects/genotyping/view.cgi?db=2>). For analysis, the full-length S protein was divided into three regions corresponding to structural and/or functional domains: the N terminal region (aa 1-99), the MHR

**Table 1 – Summary of clinical features of patients with concurrent HBsAg and anti-HBs.**

Patient	Gender/age	HBsAg (IU/mL)	Anti-HBs (mIU/mL)	ALT (IU/L)	HBeAg	HBV-DNA
P1	F/23	325.6	1326.5	224	Pos	8.63E+07
P2	M/40	103.19	51.21	95	Neg	5.35E+05
P3	M/27	102.28	156.8	168	Pos	5.36E+06
P4	F/21	179.88	36.9	147	Pos	5.20E+06
P5	M/30	167.17	23.6	103	Neg	2.36E+04
P6	M/29	122.47	111.25	115	Pos	3.12E+06
P7	F/31	85.72	62.36	98	Pos	6.45E+06
P8	F/22	669.81	20.65	214	Pos	2.80E+08
P9	M/29	92.28	18.52	136	Pos	6.35E+06
P10	M/31	116.6	26.32	94	Neg	3.68E+05
P11	F/25	522.53	19.62	197	Pos	5.61E+07
P12	M/35	93.18	85.26	115	Pos	3.26E+05
P13	M/28	72.01	35.15	94	Pos	3.72E+04
P14	F/31	126.07	25.36	97	Neg	4.35E+05
P15	F/33	718.63	120.56	319	Pos	3.16E+09

M, male; F, female; Pos, positive; Neg, negative; ALT, alanine aminotransferase.

(aa 100–169), which contains the “a” determinant (aa 124–147), and the C terminal region (aa 170–226).

### Statistical analysis

Data were stored using Microsoft Excel and analyzed using SPSS 17.0 software (Chicago, IL). Quantitative data were compared using Mann–Whitney or t test, and mutation rates were analyzed with the Chi-square test. Statistical significance was determined at a *p*-value <0.05.

## Results

### Characteristics of patients

Out of the 1194 CHB patients enrolled in this study, in 36 (3.0%) HBsAg and anti-HBs were concomitantly detected. In order to elucidate the mechanisms of coexistence of HBsAg and anti-HBs in patients infected with HBV of the genotype C, two groups were set up on the basis of their serological profile.

Fifteen HBsAg+/anti-HBs+ CHB patients infected with genotype C HBV were enrolled as the experimental group (group I). Meanwhile, 17 HBsAg+/anti-HBs– CHB patients infected with genotype C HBV were selected as the control group (group II). None of the patients had neither received vaccination against HBV nor antiviral therapy.

**Table 1** shows the clinical characteristics of group I. There were no significant differences between group I and group II with regard to age, gender, serum ALT level, HBsAg level, HBeAg positivity rate, HBV DNA concentration (*p*=0.127, 0.755, 0.692, 0.610, 0.388, 0.234, respectively) (**Table 2**). Importantly, the patients with anti-HBs had rather higher anti-HBs level, with a median titer of 36.9 mIU/mL (range, 18.52–1326.5).

### Sequencing of S gene in HBV

The entire S encoding gene sequences for patients with and without anti-HBs were available by PCR amplification and sequencing, and subsequently compared to reference sequences of the same genotype (**Figs. 1 and 2** and **Table 3**). Thirteen patients from group I had single or multiple aa substitutions within the S protein.

The distributions of aa substitutions were heterogeneous along the S protein (**Fig. 2**). Apparently, the highest frequency of aa substitutions was observed within the “a” determinant for both groups. Besides, aa variability within the MHR was markedly higher than that in the N-terminal region and the C-terminal region.

Furthermore, when considering the full-length S coding region, a marked aa substitution difference was observed between group I and group II (1.45 vs. 0.55, for substitutions per 100 aa, the same below, *p*=0.000) (**Table 3**). Moreover, the percentage of aa substitutions in the MHR for group I was

**Table 2 – Comparative analysis of clinical features of patients between group I and group II.**

	Group I (n = 15)	Group II (n = 17)	<i>p</i>
Age, median years (range)	29 (21–40)	30 (18–43)	0.127
Gender, female:male	7:8	7:10	0.755
HBsAg titer, median IU/mL (range)	122.47 (72.01–718.63)	119.37 (68.25–625.53)	0.610
HBeAg+ (%)	73.3	58.8	0.388
Serum ALT level, median IU/L (range)	115 (94–319)	147 (92–325)	0.692
Serum HBV DNA level, median (log 10 copies/mL)	6.71 (4.37–9.50)	5.57 (4.42–8.71)	0.234

ALT, alanine aminotransferase.



**Table 3 – Number of substitutions per 100 amino acid positions within HBsAg and overlapping regions of reverse transcriptase.**

Region of HBsAg	HBsAg			HBsAg overlapping regions of reverse transcriptase		
	Group I	Group II	p	Group I	Group II	p
Complete HBsAg	1.45	0.55	0.000			
Complete RT				0.71	0.26	0.005
N-terminal Region	1.01	0.42	0.044	0.34	0.18	0.595
MHR	2.67	0.67	0.000	1.14	0.17	0.003
a-Determinant	4.12	1.47	0.022	2.22	0.49	0.073
C-terminal Region	0.70	0.62	0.828	0.82	0.52	0.425

group I compared to 18% of HBV sequences from group II. In addition, only patients from group I had aa changes sQ129R, sG130N, sF134I, and sG145R, which have been proven to alter immunogenicity.<sup>4,16</sup> However, substitution sG145R, which was popular in previous study, was only found in one patient from group I.

#### HBV reverse transcriptase sequences

As the RT regions overlap the S gene, mutations within S gene might lead to aa substitutions in RT regions. Theoretically, the highest aa variability regions within the RT should be those overlapping the MHR, and especially the “a” determinant located in a structural and functional “finger” subdomain of the enzyme (Table 3).<sup>17</sup> It was natural that the RT overlapping region the MHR, the number of aa substitutions of group I patients was markedly increased compared to those of group II patients (1.14 vs. 0.17, p = 0.003). Surprisingly, in the RT overlapping region the “a” determinant was not significantly different in terms of frequencies of aa substitutions between groups I and II (2.22 vs. 0.49, p = 0.073). When considering the region of RT overlapping the C-terminal region, which stands within the structural and functional “palm” subdomain of the enzyme,<sup>17</sup> the difference of aa substitutions was not significant between the two groups (0.82 vs. 0.52, p = 0.425). Furthermore, substitutions rtE134V, rtR138K, rtN139Q, corresponding to sI126S, sG130N, sT131N, were more frequent in patients from group I than in patients from group II.

#### Discussion

The present study described the prevalence of coexistence of anti-HBs in HBsAg-positive CHB patients infected with genotype C HBV and analyzed the clinical and virological features, including aa substitutions in S gene and the overlapping RT regions. The presence of both HBsAg and anti-HBs was documented in 3.0% of 1194 CHB patients, which was similar to several previous reports,<sup>12,17-20</sup> but lower compared to others.<sup>4,21</sup> The prevalence diversity might be explained by differences in demographic, ethnic, geographical diversion, and diagnosis criteria. In this study, only an anti-HBs titer >10 mIU/mL was considered positive according to the Chinese consensus criteria.<sup>22</sup>

Genotypes B and C are the most common HBV genotypes in China, accounting for approximately 95% of HBV-infected patients.<sup>23</sup> Previous reports indicated that HBV genotype C takes more aggressive clinical course than genotype

B.<sup>7,24</sup> In addition, some studies suggested that concomitant HBsAg/anti-HBs is associated with more advanced liver diseases, such as fibrosis, cirrhosis, and hepatocellular carcinoma.<sup>21,25,26</sup> Therefore, it is necessary to explore the mechanism underlying the coexistence of HBsAg and anti-HBs in genotype C CHB patients.

The mechanism underlying the simultaneous HBsAg/anti-HBs positivity is still unknown, but one explanation might be the selection of HBsAg immune escape variants. Several studies reported that, for the MHR, especially for the “a” determinant region, genetic variability in patients carrying concurrent HBsAg/anti-HBs is much higher than that in patients carrying HBsAg only.<sup>4,16,18,19,27</sup> It has been reported that the accumulation of aa substitutions in and around the “a” determinant region may change the conformational structure and immunogenicity of HBsAg and T-cell epitope structure, leading to escape from recognition by the host immune system.<sup>13,14</sup> This hypothesis is strengthened by the results of our study.

In the present study, a significantly higher aa variability was also observed within the MHR and the “a” determinant region of HBV from HBsAg+/anti-HBs+ patients than from controls, which is in line with previous reports.<sup>4,16,18,19,27</sup> Whether and how these mutations affect the immunogenicity of HBsAg, either alone or in combination, still needs much more in-depth studies. However, regarding the specific aa substitutions, mutations sI126S/T, sQ129R, sG130N, sF134I, and sG145R found in this study were associated with alteration of HBsAg antigenicity, as reported previously.<sup>4,16,18,19,27</sup> Interestingly, the most frequent substitution sG145R reported in previous studies was only found in one patient from group I.<sup>4,19</sup> This result is similar to several previous reports from China.<sup>16,18,27</sup> Diversities in genotypes may be an important reason for the preferred mutations in the “a” determinant. All patients enrolled in this study and most of those enrolled in other studies from China were infected with genotype C HBV, in contrast with genotypes A and D HBV prevalent in Europe and the US.

In this study, we also observed significantly higher aa variability within the RT region of HBV from group I than from group II. Furthermore, we specifically observed significantly higher aa variability in the RT region overlapping the MHR region of HBsAg. However, in the RT region overlapping the C terminal region of HBsAg, which is associated with the emergence of viral resistance, there was no significant difference in the percentage of aa substitutions between group I and group II. This result is in line with one report,<sup>16</sup> but contradicts another study.<sup>19</sup> Although the virological and

clinical implications of our findings remain unknown, some mutations in RT region have already been confirmed to be associated with specific clinical outcomes previously. Mutation rtM204V, observed in one patient in our study, was previously found to partially lead to the production and intracellular retention of truncated HBsAg forms.<sup>28</sup> In addition, previous study has demonstrated that mutations rtL180M and rtM204V confer Lamivudine resistance.<sup>29</sup> Furthermore, we observed that the substitution rtS78T, which was found in group I, was previously found to decrease RT binding affinity of adefovir and potentially contributes by itself to adefovir resistance.<sup>30</sup> Besides, mutation rtS78T, corresponding to a stop-codon at the HBsAg-position 69, might potentially affect HBV pathogenicity and oncogenic potential.<sup>31</sup> Therefore, further studies deserve careful follow-up of these patients once antiviral treatments are introduced.

Notably, though most of previous reports supported the relationship of the HBsAg sequence variations and the concurrent HBsAg/anti-HBs serological profile, another study objected. In that study, similar aa variability was found in the HBsAg sequences derived from patients with and without anti-HBs.<sup>12</sup> They thought the exposure of patients to HBV infection of different subtypes might be a possible explanation.<sup>12</sup>

In summary, our study has shown the prevalence of anti-HBs in HBsAg-positive CHB patients infected with genotype C HBV. We observed a higher aa variability within the N-terminal region and the MHR, especially the "a" determinant region, and within RT for regions overlapping the MHR in patients with concurrent HBsAg/anti-HBs serological profile. Additional research on these genetic mutants is needed to clarify their biological significance for viral persistence. Further studies are needed to determine whether these simultaneous HBV mutants might positively influence the clinical course and alter nucleos(t)ide analogs efficacy.

## Conflicts of interest

The authors declare no conflicts of interest.

## Acknowledgment

This study was supported by the National Natural Science Foundation of China (81272693).

## REFERENCES

- Lok AS, McMahon BJ. Chronic hepatitis B: update 2009. *Hepatology*. 2009;50:661-2.
- Pungpapong S, Kim WR, Poterucha JJ. Natural history of hepatitis B virus infection: an update for clinicians. *Mayo Clin Proc*. 2007;82:967-75.
- Tabor E, Gerety RJ, Smallwood LA, et al. Coincident hepatitis B surface antigen and antibodies of different subtypes in human serum. *J Immunol*. 1977;118:369-70.
- Lada O, Benhamou Y, Poynard T, et al. Coexistence of hepatitis B surface antigen (HBs Ag) and anti-HBs antibodies in chronic hepatitis B virus carriers: influence of "a" determinant variants. *J Virol*. 2006;80:2968-75.
- Zeng G, Wang Z, Wen S, et al. Geographic distribution, virologic and clinical characteristics of hepatitis B virus genotypes in China. *J Viral Hepat*. 2005;12:609-17.
- Choi MS, Kim DY, Lee DH, et al. Clinical significance of pre-S mutations in patients with genotype C hepatitis B virus infection. *J Viral Hepat*. 2007;14:161-8.
- You J, Sriplung H, Chongsuvivatwong V, et al. Profile, spectrum and significance of hepatitis B virus genotypes in chronic HBV-infected patients in Yunnan, China. *Hepatobiliary Pancreat Dis Int*. 2008;7:271-9.
- Liu CJ, Kao JH. Genetic variability of hepatitis B virus and response to antiviral therapy. *Antivir Ther*. 2008;13:613-24.
- Zeng AZ, Deng H, Yang C, et al. Hepatitis B virus genotype-associated variability in antiviral response to adefovir dipivoxil therapy in Chinese Han population. *Tohoku J Exp Med*. 2008;216:205-11.
- Mathet VL, Feld M, Espinola L, et al. Hepatitis B virus S gene mutants in a patient with chronic active hepatitis with circulating anti-HBs antibodies. *J Med Virol*. 2003;69:18-26.
- Margeridon S, Lachaux A, Trepo C, et al. A quasi-monoclonal anti-HBs response can lead to immune escape of 'wild-type' hepatitis B virus. *J Gen Virol*. 2005;86:1687-93.
- Zhang JM, Xu Y, Wang XY, et al. Coexistence of Hepatitis B surface antigen (HBsAg) and heterologous subtype-specific antibodies to HBsAg among patients with chronic hepatitis B virus infection. *Clin Infect Dis*. 2007;44:1161-9.
- Weber B. The diagnostic and clinical impact of the genetic variability of the S gene of hepatitis B virus. *J Lab Med*. 2004;28:56-69.
- Torresi J. The virological and clinical significance of mutations in the overlapping envelope and polymerase genes of hepatitis B virus. *J Clin Virol*. 2002;25:97-106.
- Hussain M, Lok AS. Mutations in the hepatitis B virus polymerase gene associated with antiviral treatment for hepatitis B. *J Viral Hepat*. 1999;6:183-94.
- Liu WW, Hu T, Wang XY, et al. Coexistence of hepatitis B surface antigen and anti-HBs in Chinese chronic hepatitis B virus patients relating to genotype C and mutations in the S and P gene reverse transcriptase region. *Arch Virol*. 2012;157:627-34.
- Lin X, Yuan ZH, Wu L, Ding JP, Wen YM. A single amino acid in the reverse transcriptase domain of hepatitis B virus affects virus replication efficiency. *J Virol*. 2001;75:11827-33.
- Chen Y, Qian FC, Yuan Q, et al. Mutations in hepatitis B virus DNA from patients with coexisting HBsAg and anti-HBs. *J Clin Virol*. 2011;52:198-203.
- Colson P, Borentain P, Motte A, et al. Clinical and virological significance of the co-existence of HBsAg and anti-HBs antibodies in hepatitis B chronic carriers. *Virology*. 2007;367:30-40.
- Lee BS, Cho YK, Jeong SH, et al. Nationwide seroepidemiology of hepatitis B virus infection in South Korea in 2009 emphasizes the coexistence of HBsAg and anti-HBs. *J Med Virol*. 2013;85:1327-33.
- Seo SI, Choi HS, Choi BY, et al. Coexistence of hepatitis B surface antigen and antibody to hepatitis B surface may increase the risk of hepatocellular carcinoma in chronic hepatitis B virus infection: a Retrospective Cohort Study. *J Med Virol*. 2014;86:124-30.
- Chinese Society of Hepatology, Chinese Medical Association, Chinese Society of Infectious Diseases, Chinese Medical Association. Guideline on prevention and treatment of chronic hepatitis B in China (2005). *Chin Med J (Engl)*. 2007;120:2159-73.
- Wang Z, Huang Y, Wen S, et al. Hepatitis B virus genotypes and subgenotypes in China. *Hepatol Res*. 2007;37:S36-41.
- Chan HL, Wong GL, Tse CH, et al. Hepatitis B virus genotype C is associated with more severe liver fibrosis than genotype B. *Clin Gastroenterol Hepatol*. 2009;7:1361-6.

25. Shiels MT, Taswell HF, Czaja AJ, et al. Frequency and significance of concurrent hepatitis B surface antigen and antibody in acute and chronic hepatitis B. *Gastroenterology*. 1987;93:675-80.
26. Jang JS, Kim HS, Kim HJ, et al. Association of concurrent hepatitis B surface antigen and antibody to hepatitis B surface antigen with hepatocellular carcinoma in chronic hepatitis B virus infection. *J Med Virol*. 2009;81:1531-8.
27. Wang L, Liu H, Ning XX, et al. Sequence analysis of the S gene region in HBV DNA from patients positive for both HBsAg and HBsAb tests. *Hepatol Res*. 2010;40:1212-8.
28. Warner N, Locarnini S. The antiviral drug selected hepatitis B virus rtA181T/sW172\* mutant has a dominant negative secretion defect and alters the typical profile of viral rebound. *Hepatology*. 2008;48:88-98.
29. Seigneres B, Pichoud C, Martin P, et al. Inhibitory activity of dioxolane purine analogs on wild-type and lamivudine-resistant mutants of hepadnaviruses. *Hepatology*. 2002;36:710-22.
30. Cento V, Van Hemert F, Neumann-Fraune M, et al. Anti-HBV treatment induces novel reverse transcriptase mutations with reflective effect on HBV S antigen. *J Infect*. 2013;67:303-12.
31. Svicher V, Cento V, Salpini R, et al. Role of hepatitis B virus genetic barrier in drug-resistance and immune-escape development. *Dig Liver Dis*. 2011;43:975-83.