



The Brazilian Journal of INFECTIOUS DISEASES

www.elsevier.com/locate/bjid



Original article

Gestational toxoplasmosis in Paraná State, Brazil: prevalence of IgG antibodies and associated risk factors

Fabiana Maria Ruiz Lopes-Mori^{a,*}, Regina Mitsuka-Breganó^b,
Laura Helena França de Barros Bittencourt^a, Renata Cristina Ferreira Dias^a,
Daniela Dib Gonçalves^a, Jaqueline Dario Capobiango^b, Edna Maria Vissoci Reiche^b,
Helena Kaminami Morimoto^b, Roberta Lemos Freire^b, Italmir Teodorico Navarro^b

^a Postgraduate Program in Animal Science, Department of Veterinary Medicine, Universidade Estadual de Londrina (UEL), Londrina, Paraná, Brazil

^b Professor of UEL, Londrina, Paraná, Brazil

ARTICLE INFO

Article history:

Received 23 August 2012

Accepted 5 December 2012

Available online 9 May 2013

Keywords:

Seroepidemiology

Prevalence

Toxoplasmosis

Pregnancy

ABSTRACT

The aim of the present study was to verify the association between seropositivity for IgG anti-*Toxoplasma gondii* antibodies and social, economic and environmental variables of pregnant women attending the public health centers of Paraná, Brazil. From January 2007 to July 2010, 2226 pregnant women were interviewed and detection of anti-*T. gondii* specific IgG and IgM antibodies was performed by chemiluminescence test. Seropositivity for anti-*T. gondii* IgG was observed in 1151 (51.7%) pregnant women, 29 of which (1.3%) presented IgM reagent with IgG of high avidity. The variables associated with the presence of IgG were residency in the rural area, more than one pregnancy, less than or equal to eight years schooling, low per capita income, age group, raw or poorly cooked meat ingestion, and contact with the soil. There was neither association with raw fruit and vegetable ingestion nor with the presence of cats in the residencies.

© 2013 Elsevier Editora Ltda. Este é um artigo Open Access sob a licença de [CC BY-NC-ND](#)

Introduction

Congenital toxoplasmosis occurs when tachyzoites from *Toxoplasma gondii* reaches the fetus via the placenta.¹ Toxoplasmosis prevalence varies with region, due to climatic and, especially, cultural differences in the population. Serological investigations of IgG antibodies in pregnant women from several regions of the world have shown a variation ranging from

9.1 (Kent, United Kingdom) to 90.0% (Panamá).^{2,3} In Brazil, this level of variation occurs among the different states and populations investigated. The prevalence in pregnant women in Caxias do Sul, in Rio Grande do Sul, was 31.0%, and in the state of Mato Grosso do Sul it was 91.6%.^{4,5} Still in Brazil, the estimated maternal seroconversion rate ranges from 0.6% to 8.6%^{6,7} and congenital toxoplasmosis incidence from 0.3 per 1000 to 8 per 1000 live births.^{8,9}

* Corresponding author at: Pós-graduação em Ciência Animal, Universidade Estadual de Londrina, Rodovia Celso Garcia Cid, Pr 445, Km 380, CP 6001, Londrina, PR, 86051-980, Brazil.

E-mail addresses: fabiuel@yahoo.com.br, fabiana.mori@unifil.br (F.M.R. Lopes-Mori).

1413-8670 © 2013 Elsevier Editora Ltda. Este é um artigo Open Access sob a licença de [CC BY-NC-ND](#)

<http://dx.doi.org/10.1016/j.bjid.2012.12.003>

Health promotion strategies should be based on knowledge of the risk factors for infection that differ from region to region. Cook et al.¹⁰ demonstrated that between 3% and 63% of the infections were attributed to ingestion of poorly cooked or cured meat and from 6.0% to 17.0% from contact with the soil, but they found no association with contact with cats or their feces. In China an association between serum prevalence and raw or poorly cooked meat ingestion, unwashed vegetable and fruit ingestion, contact with cats, residency in the rural zone and low educational level has been reported.¹¹ On the other hand, a study carried out in the Democratic Republic of São Tomé and Príncipe found no association between seropositivity and the ingestion of raw meat, contact with animals, ingestion of poorly washed vegetables or consumption of unboiled water.¹²

In Brazil, a study conducted in the southern region demonstrated that contact with the soil was the major factor associated with serum prevalence,¹³ while in the northeast of the country, direct contact with cats and/or seropositive persons was highly associated with the infection.¹⁴ Other factors such as low educational level, low social and economic level and limited knowledge of the disease were also associated with the infection.

The objective of the present study was to determine the prevalence of anti-*T. gondii* IgG and IgM antibodies and the social and epidemiological factors associated with the infection in pregnant women.

Materials and methods

The study protocol was approved by the Ethics in Research Committee of Londrina State University under number 182/07. The purpose and procedures of the study were explained to all participants, and a written informed consent was obtained from all of them.

Pregnant women were included in the research if they received antenatal care in public health care facilities, in five municipalities of the State of Paraná, Brazil (Londrina, Rolândia, Cambe, Cascavel and Palotina) after their free and informed consent. In these municipalities a toxoplasmosis health surveillance programme was implemented based on prenatal screening with serological monitoring of seronegative pregnant women quarterly.¹⁵

The ideal sample size for each city was determined using the expected prevalence of 50%, standard error of 5% and a significance level of 5%. In Londrina, the ideal number of pregnant women was estimated to be 357, in Rolândia 254, in Cambé 278, in Cascavel 341, and in Palotina 196. In all cities, the number of pregnant women who participated in the research was higher than the calculated sample size.

Quantitative assessments of anti-*T. gondii* IgG and IgM antibodies were performed by chemiluminescence technique (LIAISON®, DiaSorin, Salluggia, Italy) following the manufacturer's recommendations. Positive anti-*T. gondii* IgG and IgM samples were submitted to IgG-avidity test (LIAISON®, DiaSorin, Salluggia, Italy).

To detect the variables associated with infection by *T. gondii*, each pregnant woman responded to a questionnaire collecting information about their pregnancy, eating and hygiene habits,

Table 1 – Prevalence of anti-*Toxoplasma gondii* IgG and IgM antibodies from pregnant women attending public health care facilities in Paraná, Brazil, 2007–2010.

Antibodies anti- <i>T. gondii</i> ^a	Prevalence n (%)	95% CI
IgG (+) IgM (–)	1122 (50.4)	[48.3, 52.5]
IgG (+) IgM (+)	29 (1.3)	[0.9, 1.8]
Total = IgG (+)	1151 (51.7)	[49.6, 53.7]
IgG (–) IgM (–)	1075 (48.3)	
Total	2226 (100.0)	

^a IgG and IgM antibodies against *T. gondii* were evaluated by chemiluminescence reaction (LIAISON®, DiaSorin, Salluggia, Italy). (+), positive; (–), negative.

as well as social and economic variables. The information was analyzed by the EpiInfo 3.5.1 software.¹⁶ Chi Square test and odds ratio were computed to assess the association between *T. gondii* infection and the variables researched. The 5% level of significance was used.

Results

The mean age of the pregnant women was 25 years (± 6.3 years). 65.4% initiated antenatal care in the Basic Health Units (BHU) in the first quarter, 26.4% in the second quarter, and 8.1% in the third quarter of pregnancy.

The prevalence of anti-*T. gondii* IgG antibodies was 51.7% (CI 95%: 49.6, 53.7%) (Table 1). Twenty nine of the pregnant women (1.3%, CI 95%: 0.9, 1.8%) presented positive IgM and IgG of high avidity.

Data analysis of the social and demographic variables associated with the serological results of anti-*T. gondii* IgG antibodies are shown in Tables 2 and 3, respectively.

Table 2 shows that the seropositivity for anti-*T. gondii* IgG antibodies were associated with residency in the rural area, low per capita income, age group, low educational level and more than one pregnancy. The variables related to eating habits and behavior presented significant association only with contact with the soil. There was no association with water consumption from the public system, raw or poorly cooked meat ingestion, raw fruit and vegetable ingestion and the presence of cats in the residencies (Table 3).

Discussion

The prevalence of anti-*T. gondii* IgG antibodies observed in the present study (51.7%) was within the limits of seropositivity reported in pregnant women in Brazil ranging from 31% to 91.6%, depending on the region and populations studied.^{4,5,17,18} Prevalence studies carried out in different parts of the world have also shown great variation that can be explained by environmental and cultural factors of the population.¹⁹

Another relevant aspect detected in the present study was that almost half the pregnant women attending the public health centers in the State of Paraná (48.3%), Brazil, were susceptible to primo-infection by *T. gondii* and they were at risk for fetal transmission. This result underscores the importance

Table 2 – Analysis of sociodemographic characteristics of pregnant women attending public health care facilities in Paraná, Brazil, 2007–2010, according to seropositivity for anti-*Toxoplasma gondii* IgG antibodies.

Sociodemographic characteristics	Positive anti- <i>T. gondii</i> IgG antibodies/total (%)	Odds ratio [95% CI]	p-value ^a
Residence			
Urban area	1046/2060 (50.8)	1.74 [1.22, 2.49]	0.001
Rural area	99/154 (64.3)		
Age group (years)			
<20	287/611 (47.0)	1.0 reference	0.001
21–30	600/1164 (51.5)	1.20 [0.98, 1.47]	
> 31	260/445 (58.4)	1.59 [1.23, 2.05]	
Per capita income^b			
<U\$ 77.00	288/452 (63.7)	1.0 reference	<0.001
U\$ 78.00–179.00	491/956 (51.3)	0.60 [0.47, 0.76]	
>U\$ 180.00	338/747 (45.2)	0.47 [0.37, 0.60]	
Education level (years in school)			
≤8	526/898 (58.5)	1.60 [1.34, 1.90]	<0.001
>8	620/1321 (47.0)		
Parity			
Primipara	445/979 (45.5)	0.64 [0.54, 0.76]	<0.001
Multipara	698/1232 (56.6)		
Gestational age at prenatal initiation			
1st quarter	697/1361 (51.2)	1.0 reference	0.585
2nd quarter	290/550 (52.7)	1.06 [0.87, 1.30]	
3rd quarter	93/169 (55.0)	1.17 [0.84, 1.63]	

CI, confidence interval.
^a Chi Square test for trend.
^b R\$ Brazilian currency (R\$ 1.00 = U\$ 1.95, approximately).

Table 3 – Analysis of eating habits and behavior associated with seropositivity for anti-*Toxoplasma gondii* IgG antibodies in pregnant women in Paraná, Brazil, 2007–2010.

Variable	Positive anti- <i>T. gondii</i> IgG antibodies/total (%)	Odds ratio [95% CI]	p-value ^a
Water consumption from the public system			
No	78/131 (59.5)	0.71 [0.49, 1.04]	0.078
Yes	1072/2093 (51.2)		
Consumption of raw or rare meat			
Yes	364/688 (52.9)	1.04 [0.87, 1.26]	0.678
No	759/1464 (51.8)		
Consumption of raw vegetables			
Yes	1077/2081 (51.7)	1.06 [0.74, 1.51]	0.817
No	70/139 (50.3)		
Presence of vegetable garden at home			
Yes	240/439 (54.7)	1.16 [0.93, 1.43]	0.193
No	908/1778 (51.0)		
Presence of cats at home			
Yes	722/1376 (52.4)	1.10 [0.92, 1.32]	0.318
No	388/774 (50.1)		
Presence of dogs at home			
Yes	714/1388 (51.4)	0.97 [0.81, 1.16]	0.753
No	421/806 (52.2)		
Habit of manipulating dirt or sand			
Yes	169/275 (61.4)	1.58 [1.21, 2.06]	<0.001
No	976/1941 (50.2)		

CI, confidence interval.
^a Chi Square test for trend.

of prenatal screening to identify the pregnant women's risk to acquire primo-infection during pregnancy with subsequent transmission to the fetus.

The occurrence of simultaneous positivity of IgG and IgM antibodies in 29 pregnant women (1.3%) was similar to that reported in the Midwestern region of Brazil,¹⁷ but lower rate was found in Caxias do Sul, Rio Grande do Sul, Brazil (5.5%).⁴ IgG and IgM positivity in pregnant women with high avidity IgG antibodies in the first quarter of pregnancy is indicative of infection prior to pregnancy and therefore with no risk of fetal infection.²⁰ However, when detected in the second quarter, as it was the case in one pregnant woman, high avidity IgG after the 16th week of pregnancy cannot rule out infection at the beginning of pregnancy.²¹

The pregnant woman with negative IgG and positive IgM had a second serology 15 days later and maintained the same result. Thus she was considered a false-positive or nonspecific IgM and this pregnant woman was considered susceptible to primo-infection by *T. gondii*. This fact demonstrated the importance of confirming the results. IgM results should be interpreted carefully to prevent the exposure of the mother and fetus to unnecessary procedures such as treatment and amniocentesis, which are often harmful to both. False positive diagnosis also causes anxiety in the mother and family and reduces the positive expectations regarding the new child.²²

Toxoplasmosis prevalence was greater among pregnant women from the rural area than among those from the urban area, this difference was significant and similar to that reported by other authors meaning that pregnant women living in the rural areas are in greater contact with the parasite. Studies involving a larger number of pregnant women should be carried out to identify the risk factors in this population.^{11,23} According to Frenkel²⁴ the higher the age, the higher the percentage of individuals seropositive for toxoplasmosis. In pregnant women of the present study, this finding was confirmed as the highest frequency of positivity was observed in women aged over 31 years. A significant association between age (<20 years) and toxoplasmosis, as well as the results found in pregnant women in Rio Grande do Sul, Brazil²⁵ and in pregnant women in Rio Grande do Norte, Brazil.¹⁴

The fact that women with low per capita income and low educational level presented greater chance of being infected by *T. gondii* underscores the importance of investments in education as a means of promoting health in the population. Similar results were reported in pregnant women from Goiania and Londrina, Brazil.^{6,18}

Another variable associated with infection by *T. gondii* was the number of pregnancies. Having more than one pregnancy was associated with greater prevalence. This result was similar to that observed by other authors.²⁶ In a cohort study carried out on women at fertile age, it was observed that pregnant women had 2.2 times greater chance of acquiring the infection,²⁷ showing that pregnancy itself may be a risk factor for the infection, probably because of the immune depression that occurs during pregnancy.^{28,29} However, maternal age can be a confusing factor because increase in pregnancies is associated with increase in age which is explained by the increased exposure to the source of infection during life.²⁷

The variable habit of manipulating earth and sand also showed statistical significance with toxoplasmosis in this

study ($p < 0.001$) and this finding is similar to that reported by Spalding et al.¹³ in pregnant women from Rio Grande do Sul, Brazil.

An important risk factor for infection is ingestion of raw or badly cooked fresh or cured meat,^{10,11} but in the present study this was not associated with toxoplasmosis. This finding may be justified by the reduction in the prevalence of toxoplasmosis in pigs in the northern region of Paraná, where the municipality of Londrina is located.³⁰ The reduction in this rate probably occurred because more technology and sanitary improvements have been adopted by swine farmers in recent years.

Similarly, the presence of cats in the residence was not associated with infection by *T. gondii*, the same result was reported by others authors.^{10,12} The association between cats and human toxoplasmosis is difficult to be ascertained by epidemiological surveys. The oocysts do not adhere to cat fur as do helminth eggs, and furthermore cats remove the oocysts from their fur before they become infectant.³¹

Regarding other factors studied such as raw fruit and/or vegetable ingestion, type of water consumed, presence of dogs, and presence of a vegetable garden in the residency were not significant either. However, the various studies carried out in Brazil and in the world have shown variation in these results, which can be explained by the climactic, cultural and hygiene differences of the various populations and also by the difference in methodologies employed in the studies, making it difficult to compare the results.^{11,12,27,28}

Due to the low incidence of acute toxoplasmosis during pregnancy and the high cost of cohort studies, most of the studies that aim to determine toxoplasmosis risk factors are on prevalence and not on incidence. Reactivity for IgG is used as a marker of infection by *T. gondii*. As anti-*T. gondii* IgG antibodies persist for many years, many infections may have been acquired years before and the environment and behavior patterns may have changed and the risk factors associated with infection may no longer be present. This is a limitation of the present study, but the knowledge of the risk factors in each region is important to define toxoplasmosis control programmes, especially in pregnancy. Thus in the impossibility of carrying out longitudinal studies, the studies based on prevalence have become important tools for epidemiological characterization of the infection.

In conclusion, in the population studied, 48.3% of the pregnant women were anti-*T. gondii* IgG and IgM antibody negative, showing the importance of implementing specific primary prevention programmes to guide pregnant women on the ways they can prevent the infection. Serum monitoring throughout pregnancy, so as to detect cases of maternal seroconversion allowing for early maternal treatment is also crucial. Such intervention could decrease the rate of fetal transmission or even reduce the sequelae in cases where congenital transmission has already occurred.

Financial support

SETI-PR, SESA-PR, Fundação Araucária-PR, CNPq and PROPPG-UEL.

Conflict of interest

All authors declare to have no conflict of interest

Acknowledgements

To the Municipal Health Office of Londrina (Basic Health Units and CENTROLAB) and to the University Hospital from Universidade Estadual de Londrina.

REFERENCES

- Dubey JP. Sources of *Toxoplasma gondii* infection in pregnancy. Until rates of congenital toxoplasmosis fall, control measures are essential. *Br Med J*. 2000;321:127-8.
- Nash JQ, Chissel S, Jones J, Warburton F, Verlander NQ. Risk factors for toxoplasmosis in pregnant women in Kent, United Kingdom. *Epidemiol Infect*. 2005;133:475-83.
- Souza OE, Sanze RE, Frenkel JK. Toxoplasmosis in Panamá: a 10-years study. *Am J Trop Med Hyg*. 1988;38:315-22.
- Detanico L, Basso RMC. Toxoplasmose: perfil sorológico de mulheres em idade fértil e gestantes. *Rev Bras Anál Clín*. 2006;38:15-8.
- Figueiró-Filho EA, Lopes AHA, Senefonte FRA, et al. Toxoplasmose aguda: estudo da frequência, taxa de transmissão vertical e relação entre os testes diagnósticos materno-fetais em gestantes em Estado da Região Centro-Oeste do Brasil. *Rev Bras Ginec Obst*. 2005;27:442-9.
- Avelino MM, Campos-Júnior D, Parada JB, Castro AM. Risk factors for *Toxoplasma gondii* infection in women of childbearing age. *Braz J Infect Dis*. 2004;8:164-74.
- Nóbrega OT, Karnikowski MGO. An estimation of the frequency of gestational toxoplasmosis in the Brazilian Federal District. *Rev Soc Bras Med Trop*. 2005;38:358-60.
- Neto EC, Anele E, Rubin R, et al. High prevalence of congenital toxoplasmosis in Brazil estimated in a 3-year prospective neonatal screening study. *Int J Epidemiol*. 2000;29:941-7.
- Segundo GRS, Silva DAO, Mineo JR, Ferreira MS. Congenital toxoplasmosis in Uberlândia, MG, Brazil. *J Trop Pediatr*. 2004;50:50-3.
- Cook AJC, Gilbert RE, Buffolano W, et al. Sources of *Toxoplasma* infection in pregnant women: European multicentre case-control study. *Br Med J*. 2000;321:142-7.
- Liu Q, Wei F, Gao S, et al. *Toxoplasma gondii* infection in pregnant women in China. *Trans R Soc Trop Med Hyg*. 2009;103:162-6.
- Hung CC, Fan C-K, Su K-E, et al. Serological screening and toxoplasmosis exposure factors among pregnant women in the Democratic Republic of Sao Tome and Principe. *Trans R Soc Trop Med Hyg*. 2007;101:134-9.
- Spalding SM, Amendoeira MRR, Klein CH, Ribeiro LC. Serological screening and toxoplasmosis exposure factors among pregnant women in South of Brazil. *Rev Soc Bras Med Trop*. 2005;38:173-7.
- Barbosa IR, Holanda CMCX, Andrade-Neto VF. Toxoplasmosis screening and risk factors amongst pregnant females in Natal, northeastern Brazil. *Trans R Soc Trop Med Hyg*. 2009;103:377-82.
- Lopes-Mori FMR, Mitsuka-Breganó R, Capobiango JD, et al. Programs for control of congenital toxoplasmosis. *Rev Assoc Med Bras*. 2011;57:581-6.
- Dean AG, Dean JA, Coulomerier D, et al. Epi Info, version 6: a word processing, data bases, and statistic program for epidemiology on microcomputers. Atlanta, GA, USA: Center for Diseases Control and Prevention; 1994.
- Lago EG, Carvalho RL, Jungblut R, Silva VB, Fiori RM. Screening for *Toxoplasma gondii* antibodies in 2513 consecutive parturient women and evaluation of newborn infants at risk for congenital toxoplasmosis. *Sci Med*. 2009;19:27-34.
- Lopes FMR, Mitsuka-Breganó R, Gonçalves DD, et al. Factors associated with seropositivity for anti-*Toxoplasma gondii* antibodies in pregnant women of Londrina, Paraná, Brazil. *Mem Inst Oswaldo Cruz*. 2009;104:378-82.
- Dubey JP. *Toxoplasma*, *Hammondia*, *Besnoitia*, *Sarcocystis* and others tissue cyst-forming coccidia of man and animals. In: Kreier JP, editor. *Parasitic protozoa*. New York: Academic Press; 1977.
- Jones JL, Lopez A, Wilson M. Congenital toxoplasmosis. *Am Fam Physician*. 2003;67:2131-8.
- Montoya JG, Liesenfeld O, Kinney S, Press C, Remington JS. VIDAS test for avidity of *Toxoplasma gondii*-specific immunoglobulin G for confirmatory testing of pregnant women. *J Clin Microbiol*. 2002;40:2504-8.
- Khoshnood B, De Vigan C, Goffinet F, Leroy V. Prenatal screening and diagnosis of congenital toxoplasmosis: a review of safety issues and psychological consequences for women who undergo screening. *Prenat Diagn*. 2007;27:395-403.
- Mandai ON, Lopes FMR, Mitsuka-Breganó R. Prevalência de anticorpos IgG e IgM anti-*Toxoplasma gondii* em gestantes atendidas nas unidades básicas de saúde do município de Londrina - Paraná, no período de 2003 e 2004. *Rev Bras Anál Clín*. 2007;39:247-9.
- Frenkel JK. Prevention of *Toxoplasma* infection in pregnant women and their fetuses. *Clin Infect Dis*. 1995;20:727-9.
- Cademartori BG, Farias NAR, Brod CS. Soroprevalência e fatores de risco à infecção por *Toxoplasma gondii* em gestantes de Pelotas. Sul do Brasil. *Rev Pan Infect*. 2008;10:30-5.
- Birgisdottir A, Asbjörnsdóttir H, Cook E, et al. Seroprevalence of *Toxoplasma gondii* in Sweden, Estonia and Iceland. *Scand J Infect Dis*. 2006;38:625-31.
- Avelino MM, Campos Junior D, Parada JCB, Castro AM. Pregnancy as a risk factor for acute toxoplasmosis seroconversion. *Eur J Obstet Gynecol Reprod Biol*. 2003;108:19-24.
- Crouch SP, Crocker IP, Fletcher J. The effect of pregnancy on polymorphonuclear leukocyte function. *J Immunol*. 1995;155:5436-43.
- Feinberg BB, Gonik B. General precepts of the immunology of pregnancy. *Clin Obstet Gynecol*. 1991;34:3-16.
- Carletti RT, Freire RL, Shimada MT, et al. Prevalência da infecção por *Toxoplasma gondii* em suínos abatidos no Estado do Paraná, Brasil. *Semin Ciên Agrár*. 2005;26:563-8.
- Dubey JP. Duration of immunity to shedding of *Toxoplasma gondii* oocysts by cats. *J Parasitol*. 1995;81:410-5.