



The Brazilian Journal of INFECTIOUS DISEASES

www.elsevier.com/locate/bjid



Original article

Assessment of antioxidants status and superoxide dismutase activity in HIV-infected children



Camila Pugliese^a, Rose Vega Patin^a, Cecilia Zanin Palchetti^a,
Cristiane Chiantelli Claudio^a, Aída de Fátima Thomé Barbosa Gouvêa^b,
Regina Célia de Menezes Succi^b, Olga Maria Silverio Amancio^a,
Sílvia Maria Franciscato Cozzolino^c, Fernanda Luisa Ceragioli Oliveira^{a,*}

^a Division of Nutrology, Department of Pediatrics, Universidade Federal de São Paulo (UNIFESP/EPM), São Paulo, SP, Brazil

^b Division of Pediatric Infectious Diseases, Department of Pediatrics, Universidade Federal de São Paulo (UNIFESP/EPM), São Paulo, SP, Brazil

^c Department of Food and Experimental Nutrition, Faculdade de Ciências Farmacêuticas, Universidade de São Paulo (USP), São Paulo, SP, Brazil

ARTICLE INFO

Article history:

Received 24 June 2013

Accepted 4 February 2014

Available online 26 April 2014

Keywords:

Antioxidants

HIV and nutrition

Oxidative stress

Trace elements

ABSTRACT

Objective: This study aims to assess the nutritional status of selenium, copper and zinc; and also the erythrocyte superoxide dismutase activity of HIV-infected children compared to a control group.

Methods: A cross-sectional study was carried out with prepubertal HIV-infected children ($n = 51$) and their healthy siblings ($n = 32$). All biochemical measurements including plasma selenium, serum copper levels, serum and erythrocyte zinc levels and erythrocyte superoxide dismutase activity were evaluated according to dietary, clinical and biochemical parameters.

Results: Compared to the control group, the HIV-infected children had lower z-score values for height-for-age ($p = 0.0006$), higher prevalence of stunting (11.8%) ($p = 0.047$), lower selenium levels ($p = 0.0006$) and higher copper levels ($p = 0.019$). No difference was found concerning superoxide dismutase activity ($p > 0.05$). The HIV-infected group presented a higher proportion (45.1%) of children with zinc intakes below the estimated average requirement ($p = 0.014$); however, no association with zinc biochemical parameters was found.

Conclusion: HIV-infected children have an inadequate selenium and copper nutritional status, which could influence the progression to AIDS. An adequate micronutrient status could improve the clinical conditions in these patients and minimize free radical production and cellular oxidative stress.

© 2014 Elsevier Editora Ltda. Este é um artigo Open Access sob a licença de [CC BY-NC-ND](http://creativecommons.org/licenses/by-nc-nd/4.0/)

* Corresponding author.

E-mail addresses: fernandalco.dped@epm.br, fernandalco@gmail.com (F.L.C. Oliveira).

<http://dx.doi.org/10.1016/j.bjid.2014.02.003>

1413-8670/© 2014 Elsevier Editora Ltda. Este é um artigo Open Access sob a licença de [CC BY-NC-ND](http://creativecommons.org/licenses/by-nc-nd/4.0/)

Introduction

Oxidative stress, which is caused by an increased production of free radicals and/or an inadequate antioxidant protection mechanism, can be worsened by HIV infection, thereby compromising the immune system.^{1,2}

Antioxidants protect cells and tissues against inflammatory and infectious processes and inhibit lipid peroxidation, acting as anti-atherogenic agents.³ Selenium and copper may influence the progression from HIV to AIDS.^{4,5} Zinc and copper are components of superoxide dismutase (CuZnSOD), an important antioxidant enzyme; therefore, deficiencies in these minerals may affect this enzyme activity.⁶ Inadequate intake of antioxidants, such as vitamins and minerals, can lead to impaired immune system and increase in lipid peroxidation, predisposing the individual to infections and aggravating the malnutrition problem.³

In addition to antioxidant nutrients, antioxidant enzymes (e.g., superoxide dismutase, catalase, and glutathione peroxidase) play crucial roles in eliminating free radicals and minimizing the effects of cellular oxidative stress.⁷

HIV infection itself and its treatment can determine metabolic changes that may increase the mortality risk for HIV-infected children. Nutritional status and biochemical parameters should be evaluated in HIV-infected individuals with low intakes of antioxidant nutrients to minimize adverse health effects.

The objectives of this study were to assess the trace elements antioxidant nutritional status and to determine the erythrocyte superoxide dismutase (SOD) activity in pre-pubertal HIV-infected individuals compared to the control group.

Materials/subjects and methods

Subjects

A cross-sectional study, which included 83 prepubertal children from both sex, followed up at the Care Center of the Division of Pediatric Infectious Diseases, Department of Pediatrics (CEADIPe), Universidade Federal de São Paulo – UNIFESP/EPM, São Paulo, Brazil. Data were collected from October 2009 to October 2011. This study was approved by the Ethics Committee of Universidade Federal de São Paulo – UNIFESP/EPM and written informed consent was obtained from all of the subjects.

Subjects were distributed into two groups:

- I. The HIV-infected group consisted of 51 children to whom HIV was transmitted vertically.
- II. The control group consisted of 32 healthy siblings who were born before the diagnosis of HIV infection in the mother. Mothers were not infected yet with HIV by the time of their pregnancy and/or delivery and it was confirmed by laboratory tests.

Only children who had a Tanner stage 1 pubertal development⁸ were enrolled in the study. Children with

other chronic diseases that could interfere with micronutrient nutritional status or who were receiving vitamin or mineral supplement and also those that were hospitalized or presented severe clinical complications in the year before entering the study were not included.

Clinical and immunological assessment

HIV infection was classified according to the clinical and immunological parameters described by the CDC guidelines (1994).⁹ Plasma HIV-1 RNA (viral load) was measured by polymerase chain reaction using the Amplicor HIV-1 Monitor (Amplicor Roche Diagnostics), which has a detection limit of 400 copies/mL; values were expressed as log₁₀. The CD4⁺ cell count was determined by flow cytometry and was expressed as percentage (%).⁹

Anthropometry assessment and dietary intake

Weight (kg) and height (cm) measurements were used to calculate height-for-age (H/A) z-score and body mass index (BMI) z-score, according to WHO Child Growth Standards (2007).¹⁰

Energy and macro and micronutrient intake were obtained from the average of four 24-h dietary recalls.¹¹ Data were analyzed using DietWin[®] software – Porto Alegre, Rio Grande do Sul, Brazil. Estimated energy requirement (EER) equations were used to assess the daily dietary energy requirements. For the macronutrients, we used the acceptable macronutrient distribution range (AMDRs) and the intake was expressed as a percentage of the total energy value (TE).¹² The estimated average requirement (EAR) was used for the micronutrients evaluation, values below the cut-off points indicated inadequate nutrient intake.¹³

Laboratory analyses

Biochemical parameters were measured in the absence of infectious processes. Following a 12-h fast, 10 mL of blood was collected by venipuncture.

Plasma selenium levels were determined by hydride generation atomic absorption spectrometry (HITACHI brand, model Z-5000) coupled to a quartz cell (HGQTAAS).¹⁴ Values were considered low or high when they were below the 2.5 percentile or above the 9.7 percentile, respectively, for age and gender.¹⁵ Serum copper measurement were analyzed by atomic absorption spectrophotometry (PERKIN ELMER brand, model 5.100).¹⁶ The reference values used were 70–140 µg/dL for males and 80–155 µg/dL for females.¹⁷ Serum and erythrocyte zinc concentrations were analyzed by atomic absorption spectrophotometry (PERKIN ELMER brand, model 5.100).¹⁶ The reference values are 40–44 µg Zn/g Hb for erythrocyte zinc,¹⁸ 65–120 µg/dL for serum zinc in males, and 60–120 µg/dL for serum zinc in females.¹⁷ SOD activity in erythrocytes was measured by spectrophotometry and adequate values range from 4.7 to 5.7 U/mg of hemoglobin (Hb).¹⁹

The Advia Chemistry System 1650 (Bayer[®]) analyzer and a turbidimetric assay were used to evaluate the level of C-reactive protein (CRP) in the subjects.

Table 1 – Anthropometric characteristics, nutritional status, and virological and immunological assessment of HIV-infected (n = 51) and control (n = 32) groups.

	HIV-infected	Control	p value
Age (years) ^{a,b}	8.3 (7.4–9.4)	9.1 (8.4–9.7)	0.0110^b
Male ^{c,d}	31 (60.8)	18 (56.2)	0.6830
BMI ^c (kg/m ²) ^{a,b}	15.5 (14.7–16.6)	16.2 (14.9–17.6)	0.1536
BMI z-score ^{b,e,f}	–0.18 (–0.91 to 0.39)	–0.08 (–0.76 to 0.54)	0.4402
Height for age z-score ^{b,f}	–1.05 (–1.55 to –0.23)	–0.24 (–0.80 to 0.72)	0.0006
Nutritional status ^f			
Stunting ^{f,g}	6 (11.8)	0 (0)	0.0470
Eutrophic ^{f,g}	45 (88.2)	28 (87.5)	0.4280
Overweight + obesity ^f	4 (7.8)	4 (12.5)	
Undernutrition ^f	2 (3.9)	0 (0)	
Immunological parameters			
Viral load (log ₁₀) ^a	3.483 (0–4.500)	NA	
CD4 (%) ^a	31.3 (24.9–38.1)		
Antiretroviral therapy			
Age of ART initiation (months) ^a	21 (11–36)	NA	
Duration of ART (months) ^a	100.8 (64.9–85.5)	NA	

NA, not applicable; ART, antiretroviral therapy.

^a Median (interquartile range 25–75).

^b Mann–Whitney *U*-test.

^c *n* (%)

^d Chi square test.

^e BMI (body mass index).

^f WHO (2007).

^g Fisher's exact test.

Statistical analyses

Data were analyzed using Epi info 6.04 (Centers for Diseases Control and Prevention, Atlanta, GA, USA) and STATA 10.0 (Stata Corporation, College Station, TX, USA) statistical software.

Differences in the anthropometric characteristics, dietary intake and biochemical parameters among the children were assessed using Student's *t*-test (for normal distributions) or a Mann–Whitney *U*-test (for non-normal distributions). Chi-square test and Fisher's exact test were applied to assess the association between the categorical variables according to pre-established cut-off points. Pearson's and Spearman's tests were used to study the correlations between the dietary intake and the biochemical parameters. The statistical significance level was set at *p*-value <0.05.

Results

There were 83 prepubertal children who participated in the study, 51 were HIV-infected children and 32 were HIV-uninfected children. Table 1 shows the anthropometric characteristics and the nutritional status of both groups and also the virological and immunological parameters of HIV-infected children. The groups were comparable in terms of gender, BMI and proportion of individuals with normal weight. The median age of the HIV-infected group was lower than that of the control group. The proportion of stunted individuals (smaller height-for-age z-score) was higher in the HIV-infected group than in the control group (*p* = 0.0006). No significant correlation between zinc levels and short stature was found.

The macro- and micronutrient intakes are shown in Table 2. There was higher percentage of subjects (45.1%) in the HIV-infected group with dietary zinc intake below the EAR (*p* = 0.014) and no statistically significant differences between groups with respect to macronutrients (g), selenium and copper consumption. HIV-infected group presented lower plasma selenium (*p* = 0.0006) and higher copper levels when compared to the control group (*p* = 0.019). Moreover, no difference was found concerning SOD activity or erythrocyte and serum zinc levels (Table 3). There were no significant differences in the concentration of trace elements according to different clinical and immunological categories of AIDS.

Discussion

Progression of HIV infection and adverse health effects of antiretroviral medication lead to a chronic inflammatory condition and to impaired immune system in the child. It is therefore crucial to evaluate and monitor the antioxidant defense mechanisms that might influence the progression of HIV infection.^{20,21}

In this study, the HIV-infected group had a higher proportion of stunting (11.8%) than the control group. Inadequate growth in HIV-infected children can be directly caused by viral infection or indirectly by adverse environmental effects or by a combination. The groups did neither differ in energy nor in macronutrients intake.

Children frequently consume monotonous diets characterized by high intakes of simple carbohydrates, saturated and trans-fats (e.g., sugar, artificial juices, sweets, sausages and fries). Compared to the control group, the HIV-infected children had the highest proportion of subjects with zinc intakes

Table 2 – Dietary assessment of energy, macronutrients and micronutrients of HIV-infected (n = 51) and control (n = 32) groups.

Dietary assessment	HIV-infected	Control	p value
TE ^a (Kcal) ^{b,c}	1892.18 (411.97)	2002.66 (460.13)	0.2591
EER ^{b,d}	1593 (172)	1788 (319)	
Carbohydrates (g) ^{b,c}	250.44 (61.04)	261.01 (57.70)	0.4356
Proteins (g) ^{b,c}	66.78 (16.65)	69.42 (18.58)	0.5032
Total fats (g) ^{b,c}	69.12 (15.74)	75.63 (21.37)	0.1143
Selenium (μg) ^{b,c}	51.15 (16.16)	49.08 (15.25)	0.5635
Selenium < EAR ^{e,f,g,h}	5 (9.8)	4 (12.5)	0.4823
Zinc (mg) ^{i,j}	8.8 (7.4–10.7)	9.0 (7.4–10.7)	0.7013
Zinc < EAR ^{e,f,g,k}	23 (45.1)	6 (18.8)	0.0140
Copper (mg) ^{i,j}	0.96 (0.84–1.18)	1.10 (0.84–1.18)	0.3212
Copper < EAR ^{e,f,g,h}	3 (5.9)	2 (6.2)	0.6437

^a TE (total energetic value).
^b Mean (standard deviation).
^c Student's t-test.
^d EER (estimated energy requirement).
^e EAR (estimated average requirement).
^f Altered values < EAR.
^g n (%).
^h Fisher's exact test.
ⁱ Median (interquartile range 25–75).
^j Mann-Whitney U-test.
^k Chi square test.

Table 3 – Biochemical parameters of HIV-infected and control groups.

Biochemical parameter	n	HIV group	n	Control group	p value
Plasma selenium (μg/L) ^{a,b}	50	31.45 (8.91)	32	38.80 (9.49)	0.0006
Low < 2.5 percentile ^{c,d,e}		48 (96.0)		27.0 (84.4)	0.0780
Erythrocyte zinc (μg/gHb) ^{a,f}	51	25.0 (7.9)	32	25.0 (10.14)	0.8223
Low (<40 μg/gHb) ^{d,e,g}		48 (94.12)		30 (93.8)	0.644
Serum zinc (mg/dL) ^{a,f}	51	52.3 (26.5)	32	69.4 (42.4)	0.0830
Low < 65 mg/dL (M) < 60 mg/dL (F) ^{d,h,i}		33 (64.7)		16 (50.0)	0.185
Serum copper (μg/dL) ^{a,f}	51	120.83 (48.93)	32	105.78 (37.73)	0.0190
High > 140 μg/dL (M) > 155 μg/dL (F) ^{d,e,h}		16 (31.4)		7 (21.9)	0.9200
SOD activity (U/mg Hb) ^{f,j}	51	6.0 (5.3–6.5)	31	6.0 (5.4–6.6)	0.5856
High (>5.7 U/mg Hb) ^{d,i}		30 (58.8)		28 (96.6)	0.8366
CRP (mg/dL) ^{j,k}	48	0.01 (0.01–0.22)	29	0.05 (0.01–0.07)	0.5022
Low (<0.8 mg/dL) ^{a,e}		7 (14.6)		1 (3.4)	0.1194

F, female; M, male.
^a Mean (standard deviation).
^b Student's t-test.
^c Bates et al.¹⁵
^d n (%).
^e Fisher's exact test.
^f Mann-Whitney U-test.
^g Guthrie and Picciano.¹⁸
^h Gibson.¹⁷
ⁱ Chi square test.
^j Median (interquartile range 25–75).

below the EAR; however no association with zinc biochemical parameters was found. This result can be explained by the low consumption of protein that is rich in zinc.

Lower selenium levels were found in the HIV infected group. These results suggest that the low plasma selenium of the HIV-infected group could be associated with increased

requirement for antioxidants to neutralize oxidative stress. Plasma selenium level is a sensitive indicator of disease progression and risk of mortality in HIV-infected children.^{4,22} It maintains a healthy immune system, acts as an antioxidant and might regulate viral expression.⁴ Selenium deficiency was associated with higher mortality risk regardless of the CD4⁺ T

cell count⁴ and may be related to protein-energy malnutrition commonly found in HIV-infected patients.¹

A study found no selenium deficiency in healthy and HIV-infected adolescents; however, immune function assessed by plasma neopterin concentration was negatively correlated with plasma selenium levels.²³ Poor selenium intakes and malabsorption may contribute to selenium deficiency, which affects the immune system and the cardiac muscle of HIV-infected subjects.¹

The HIV-infected group had higher serum copper level. Moreno et al. found higher serum copper concentrations in HIV-infected subjects regardless of gender when compared to the healthy controls. These results suggest that copper is an acute phase reactant and its serum concentration can be used as a surrogate marker for HIV progression, as well as in other chronic infections and inflammatory diseases.²⁴ Serum copper levels are not good markers of the inflammatory response because the concentration of this micronutrient appears to reflect interactions among different pathological processes.²⁵ Because of its participation in LDL-c oxidation within artery walls, under and overconsumption of copper can result in atherogenesis.^{26,27} However, copper is a constituent of SOD and ceruloplasmin, which are involved in preventing oxidative stress.^{26,27}

The CRP level was not significantly different between the groups; therefore, it was not possible to characterize the acute inflammatory process in the children. This result may be explained by the fact that, despite having moderate or severe clinical and immunological categories, the children were not hospitalized or presented severe clinical complications in the year prior to entering the study.

Our study had a small number of subjects in the HIV-infected and HIV-uninfected groups, which limits the extrapolation of our findings to other HIV-infected children. However, our HIV-infected and control populations were from the same families; therefore they had similar food access.

HIV-infected children can be exposed to numerous risk factors that impair their immune function, including inadequate nutritional status for trace elements, which could influence the progression of HIV infection. Assessing dietary intake and biochemical parameters are essential in the clinical practice in order to identify and treat early micronutrients deficiency minimizing clinical and metabolic alterations.

Contributions

Camila Pugliese – this article is a result of MSc dissertation – contributed to conception and design of the study, acquisition of data, micronutrient assessment, analysis and interpretation of data, drafting the article, and final approval of the version to be published.

Rose Vega Patin contributed to conception and design of the study, acquisition of data, analysis and interpretation of data, drafting the article, important intellectual content, and final approval of the version to be published.

Cecilia Zanin Palchetti contributed to conception and design of the study, acquisition of data, interpretation of data, drafting the article, important intellectual content, and final approval of the version to be published.

Cristiane Chiantelli Claudio contributed to conception and design of the study, acquisition of data, interpretation of data, drafting the article, and final approval of the version to be published.

Aida de Fátima Thomé Barbosa Gouvêa contributed to acquisition of medical data, important intellectual content, and final approval of the version to be published.

Regina Célia de Menezes Succi contributed to conception and design of the study, interpretation of data, drafting the article, important intellectual content, and final approval of the version to be published.

Olga Maria Silverio Amancio contributed to micronutrient assessment, interpretation of data, important intellectual content, and final approval of the version to be published.

Silvia Maria Franciscato Cozzolino contributed to micronutrient assessment, interpretation of data, important intellectual content and final approval of the version to be published.

Fernanda Luisa Ceragioli Oliveira contributed to conception and design of the study, analysis and interpretation of data, drafting the article, important intellectual content, and final approval of the version to be published.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgments

We are grateful to the families and the children for participating in this study and to the research teams of the Division of Nutrology, Department of Pediatrics, Universidade Federal de São Paulo – UNIFESP/EPM, Brazil and of the Nutrition and Minerals Laboratory, Faculdade de Ciências Farmacêuticas, Universidade de São Paulo, USP, Brazil.

REFERENCES

1. Dworkin BM. Selenium deficiency in HIV infection and the acquired immunodeficiency syndrome (AIDS). *Chem Biol Interact.* 1994;91:181–6.
2. Seifried HE, Anderson DE, Sorkin BC, Costello RB. Free radicals: the pros and cons of antioxidants. *American Society for Nutritional Sciences. J Nutr.* 2004;134:3143–63.
3. Maggini S, Wintergerst ES, Beveridge S, Hornig DH. Selected vitamins and trace elements support immune function by strengthening epithelial barriers and cellular and humoral immune responses. *Br J Nutr.* 2007;98:29–35.
4. Campa A, Shor-Posner G, Indacochea F, et al. Mortality risk in selenium-deficient HIV-positive children. *J Acquir Immune Defic Syndr Hum Retrovirol.* 1999;20:508–13.
5. Kassu A, Yabutani T, Mohammad A, et al. Alterations in serum levels of trace elements in tuberculosis and HIV infections. *Eur J Clin Nutr.* 2006;60:580–6.
6. Ishigami N, Isoda K, Adachi T, et al. Deficiency of CuZn superoxide dismutase promotes inflammation and alters medial structure following vascular injury. *J Atheroscler Thromb.* 2011;18:1009–17.
7. Catania AS, Barros CR, Ferreira SRG. Vitamins and minerals with antioxidant properties and cardiometabolic risk:

- controversies and perspectives. *Arq Bras Endocrinol Metab.* 2009;53:550-9.
8. Marshall EA, Tanner JM. Growth and physiological development during adolescence. *Ann Rev Med.* 1975;19:283-300.
 9. Centers for Disease Control Prevention. Revised classification system for human immunodeficiency virus infection in children less than 13 years of age. *MMWR Recomm Rep.* 1994;43:1-10.
 10. WHO. The WHO child growth standards. Growth reference, 5-19y. Geneva, Switzerland: World Health Organization; 2007.
 11. Smit E, Tang A. Nutritional assessment in intravenous drug users with HIV/AIDS. *J Acquir Immune Defic Syndr.* 2000;25:62-9.
 12. Institute of Medicine. Dietary reference intakes for energy, carbohydrate, fiber, fat, protein and amino acids (macronutrients). Washington, DC: National Academy Press; 2002.
 13. Institute of Medicine. Dietary reference intakes: vitamin A, vitamin K, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium and zinc. Washington, DC: National Academy Press; 2001.
 14. Hao D, Xie G, Zhang Y, Tian G. Determination of serum selenium by hydride generation flame atomic absorption spectrometry. *Talanta.* 1996;43:595-600.
 15. Bates CJ, Thane CW, Prentice A, Delves HT, Gregory J. Selenium status and associated factors in a British National Diet and Nutrition Survey: young people aged 4-18 y. *Eur J Clin Nutr.* 2002;56:873-81.
 16. Whitehouse R, Prasad AS, Rabbani PI, Cossak ZT. Zinc in plasma, neutrophils, lymphocytes and erythrocytes as determined by flameless atomic absorption spectrophotometry. *Clin Chem.* 1982;28:475-80.
 17. Gibson RS. Assessment of trace-element status. In: Gibson RS, editor. Principles of nutrition assessment. 2nd ed. New York: Oxford University Press; 1990.
 18. Guthrie HA, Picciano M. Micronutrient mineral. In: Human nutrition. St Louis: Mosby; 1995.
 19. McCord JM, Fridovich I. Superoxide dismutase. An enzymic function for erythrocyte (hemocuprein). *J Biol Chem.* 1969;244:6049-55.
 20. Grinspoon SK, Grunfeld C, Kotler DP, et al. State of the science conference: initiative to decrease cardiovascular risk and increase quality of care for patients living with HIV/AIDS: executive summary. *Circulation.* 2008;118:198-210.
 21. Allard JP, Aghdassi E, Chau J, Salit I, Walmsley S. Oxidative stress and plasma antioxidant micronutrients in humans with HIV infection. *Am J Clin Nutr.* 1998;67:143-7.
 22. Look MP, Rockstroh JK, Rao GS, et al. Serum selenium, plasma glutathione (GSH) and erythrocyte glutathione peroxidase (GSH-Px)-levels in asymptomatic versus symptomatic human immunodeficiency virus-1 (HIV-1)-infection. *Eur J Clin Nutr.* 1997;51:266-72.
 23. Stephensen CB, Marquis GS, Douglas SD, Kruzich LA, Wilson CM. Glutathione, glutathione peroxidase, and selenium status in HIV-positive and HIV-negative adolescents and young adults. *Am J Clin Nutr.* 2007;85:173-81.
 24. Moreno T, Artacho R, Navarro M, Pérez A, Ruiz-López MD. Serum copper concentration in HIV-infection patients and relationships with other biochemical indices. *Sci Total Environ.* 1998;217:21-6.
 25. Ucar SK, Coker M, Sözmen E, Simsek DG, Darcan S. An association among iron, copper, zinc, and selenium, and antioxidative status in dyslipidemic pediatric patients with glycogen storage disease types IA and III. *J Trace Elem Med Biol.* 2010;24:42-5.
 26. Ferns GA, Lamb DJ, Taylor A. The possible role of copper ions in atherogenesis: the Blue Janus. *Atherosclerosis.* 1997;133:139-52.
 27. Mazur A, Gueux E, Bureau I, Feillet-Coudray C, Rock E, Rayssiquier Y. Copper deficiency and lipoprotein oxidation. *Atherosclerosis.* 1998;137:443-5.