
CORRIGENDUM

WILEY

In Jinli Ding et al., the published article contains errors in Figure 3C, D. The wound healing picture from the group of Co-culture + PD in Figure 3C and the image of invasion capacity for G-CSF+LY in Figure 3D are incorrect in the original publication due to technical error during image preparation. The corrected Figure 3 is shown below. The authors confirm that all the results and conclusions of this article remain unchanged.



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. Journal of Cellular and Molecular Medicine published by Foundation for Cellular and Molecular Medicine and John Wiley & Sons Ltd.

FIGURE 3 M2 macrophage-derived G-CSF promotes EMT, migration and invasion of trophoblasts via activating PI3K/Akt/Erk1/2 pathway. A, Western blot analysis of HTR-8 alone, G-CSF-stimulated HTR-8, macrophage-co-cultured HTR-8 and G-CSF depleted macrophage-co-cultured HTR-8. B, Western blot analysis of HTR-8 alone, G-CSF-stimulated HTR-8 and macrophage-co-cultured HTR-8 in the presence or absence of LY29004 (20 μ M) or PD98059 (15 μ M). C and D, Cell migration and invasion capacity in HTR-8 alone, G-CSFstimulated HTR-8 and macrophage-co-cultured HTR-8 in the presence or absence of LY29004 (20 μ M) or PD98059 (15 μ M) were determined by wound healing assay and transwell system, respectively. Representative photographs of migratory or invasive cells (magnification, ×200) are shown. Notes: compared with control group, *p < 0.05, **p < 0.01; **p < 0.001; compared with G-CSF-stimulated HTR-8 group, *p < 0.05, ##p<0.01; compared with co-culture group, &p<0.05, &&p<0.01

REFERENCE

1. Ding J, Yang C, Zhang Y, et al. M2 macrophage-derived G-CSF promotes trophoblasts EMT, invasion and migration via activating PI3K/Akt/Erk1/2 pathway to mediate normal pregnancy. J Cell Mol Med. 2021;25:2136-2147.