

Erratum

KLF10 inhibits cell growth by regulating PTTG1 in multiple myeloma under the regulation of microRNA-106b-5p: Erratum

Mimi Zhou¹, Jinqiu Chen², Hui Zhang², Hailing Liu², Huan Yao², Xiaman Wang², Wanggang Zhang², Yingren Zhao¹, Nan Yang¹✉

1. Department of Infectious Diseases, the First Affiliated Hospital of Xi'an Jiaotong University, Yanta West Road No. 277, Xi'an 710061, China
2. Department of Hematology, the Second Affiliated Hospital of Xi'an Jiaotong University, West Five Road No. 157, Xi'an 710004, China

✉ Corresponding author: Nan Yang, Phone number: +8613 572 227 780, E-mail: nan_yang@xjtu.edu.cn.

© The author(s). This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>). See <http://ivyspring.com/terms> for full terms and conditions.

Published: 2022.08.20

Corrected article: *Int J Biol Sci* 2020; 16(12): 2063-2071. doi: 10.7150/ijbs.45999

In our paper, the authors noted one error in Fig. 3A. The subcutaneous xenograft for tumor formation picture was misused. The authors carefully checked the original data and found that the images were unintentionally included from another project (ongoing in the same lab) due to the similarity of appearance and file name. All the original pictures from this experiment were taken on the same day. We confirmed that this mistake in Figure 3A did not affect the research results and conclusion of this article. All authors have agreed to the Erratum, and we apologize for the negligence in our work and hope to get the opportunity to correct this mistake. Figure 3A should be corrected as follows.




Figure 3. KLF10 inhibits tumor growth and promotes apoptosis in vivo. (A) Representative pictures of MM xenografts from RPMI8226-KLF10 and RPMI8226-control. (B) Tumor growth curve revealed that KLF10 overexpression significantly inhibited tumor growth in vivo. Tumor nodules were subjected to immunohistochemical staining for Ki-67 (C) and TUNEL (D) assays and quantitative analysis. Representative immunostaining and TUNEL assays revealed that KLF10 overexpression significantly decreased the number of Ki-67 positive cells and increased the number of apoptotic cells. *P<0.05.