

Croat Med J. 2022;63: 335-42

<https://doi.org/10.3325/cmj.2022.63.335>

# Survival after hospital discharge in patients hospitalized for acute coronavirus disease 2019: data on 2586 patients from a tertiary center registry

**Aim** To assess the long-term survival after hospital discharge of patients hospitalized due to coronavirus disease 2019 (COVID-19).

**Methods** We retrospectively reviewed data on post-discharge survival of 2586 COVID-19 patients hospitalized in our tertiary hospital from March 2020 to March 2021.

**Results** Among 2586 patients, 1446 (55.9%) were men. The median age was 70 years, interquartile range (IQR, 60-80). The median Charlson comorbidity index was 4 points, IQR (2-5). The median length of hospital stay was 10 days, IQR (7-16). During a median follow-up of 4 months, 192 (7.4%) patients died. The median survival time after hospital discharge was not reached, and 3-month, 6-month, and 12-month survival rates were 93%, 92%, and 91%, respectively. In a multivariate analysis, mutually independent predictors of worse mortality after hospital discharge were age >75 years, Eastern Cooperative Oncology Group status 4, white blood cell count  $>7 \times 10^9/L$ , red cell distribution width >14%, urea on admission  $>10.5 \text{ mmol/L}$ , mechanical ventilation during hospital stay, readmission after discharge, absence of obesity, presence of chronic obstructive pulmonary disease, dementia, and metastatic malignancy ( $P < 0.05$  for all).

**Conclusion** Substantial risk of death persists after hospital admission due to COVID-19. Factors related to an increased risk are older age, higher functional impairment, need for mechanical ventilation during hospital admission, parameters indicating more pronounced inflammation, impaired renal function, and particular comorbidities. Interventions aimed at improving patients' functional capacity may be needed.

Marko Lucijanić<sup>1,2,3</sup>, Nevenka Piskač Živković<sup>2,4</sup>, Marko Zelenika<sup>2,5</sup>, Mislav Barišić-Jaman<sup>2,5</sup>, Ivana Jurin<sup>2,6</sup>, Ana Matijaca<sup>2,7</sup>, Nikola Zagorec<sup>2,8</sup>, Marko Lagančić<sup>2,9</sup>, Besa Osmani<sup>2,8</sup>, Iva Bušić<sup>3</sup>, Sara Šakota<sup>2,9</sup>, Ivan Vukoja<sup>10,11</sup>, Ivica Lukšić<sup>2,3,12</sup>, Bruno Baršić<sup>2,3</sup>

<sup>1</sup>Hematology Department, Dubrava University Hospital, Zagreb, Croatia

<sup>2</sup>Primary Respiratory and Intensive Care Center, Dubrava University Hospital, Zagreb, Croatia

<sup>3</sup>University of Zagreb, School of Medicine, Zagreb, Croatia

<sup>4</sup>Pulmonology Department, Dubrava University Hospital, Zagreb, Croatia

<sup>5</sup>Department of Gastroenterology, Hepatology and Clinical Nutrition, Dubrava University Hospital, Zagreb, Croatia

<sup>6</sup>Cardiology Department, Dubrava University Hospital, Zagreb, Croatia

<sup>7</sup>Endocrinology Department, Dubrava University Hospital, Zagreb, Croatia

<sup>8</sup>Nephrology Department, Dubrava University Hospital, Zagreb, Croatia

<sup>9</sup>Department of Emergency and Intensive Care Medicine, Dubrava University Hospital, Zagreb, Croatia

<sup>10</sup>Gastroenterology and Nephrology Department, General County Hospital Požega, Požega, Croatia

<sup>11</sup>Faculty of Medicine Osijek, Josip Juraj Strossmayer University of Osijek, Osijek, Croatia

<sup>12</sup>Department of Maxillofacial Surgery, Dubrava University Hospital, Zagreb, Croatia

Received: March 25, 2022

Accepted: June 10, 2022

**Correspondence to:**

Marko Lucijanić  
Hematology Department  
University Hospital Dubrava  
Av. Gojka Šuška 6  
10000 Zagreb, Croatia  
[markolucijanic@yahoo.com](mailto:markolucijanic@yahoo.com)

Coronavirus disease 2019 (COVID-19) affects multiple organ systems, but its clinical presentation is dominated by respiratory symptoms and high rates of respiratory deterioration (1). The disease strains health care systems and affects both hospital and ambulatory care. Up to 15%-20% of pre-vaccination patients were hospitalized due to respiratory insufficiency or active comorbidities that required hospital care (2). Many people still remain unvaccinated, presenting a high risk group for short-term mortality (3). Elderly patients and those burdened with comorbidities are especially susceptible to unfavorable clinical course during hospital stay (4). Data on survival subsequent to hospital discharge after overcoming respiratory insufficiency are scarce. The main concerns affecting these patients are death, rehospitalizations, and reduced functioning (5,6). Due to the importance of the period after hospital discharge for patients' rehabilitation and social re-inclusion, health care providers and caregivers need to be informed on the risks associated with this period. Thus, we aimed to assess the mortality and risk factors for unfavorable outcome after hospital discharge among COVID-19 patients treated in our institution.

## PATIENTS AND METHODS

The presented data are part of the Registry project encompassing clinical and laboratory data as well as the outcomes of all hospitalized COVID-19 patients treated in Dubrava University Hospital during the COVID-19 pandemic. Dubrava University Hospital was repurposed into a regional tertiary COVID-19 center. Of 4102 hospitalizations and 4014 consecutive COVID-19 patients treated from March 2020 to March 2021, 2586 patients were discharged and are included in the analysis. All patients were whites. All had positive polymerase chain reaction (PCR) or antigen COVID-19 test before hospital admission. During hospital stay, patients were treated according to the contemporary guidelines with various exposure to low molecular weight heparin, corticosteroids, and remdesivir. Post-discharge follow-up and survival were assessed through a review of patients' electronic records of post-discharge follow-up visits and through telephone check-up. COVID-19 disease severity on admission was graded based on the World Health Organization (WHO) recommendations. The study was approved by the Institutional Review Board of Dubrava University Hospital (2021/2503-04).

### Statistical analysis

The normality of distribution of numerical variables was tested with the Kolmogorov-Smirnov test. Numerical

variables are presented as median and interquartile range. Categorical variables are presented as frequencies and percentages. The receiver operating characteristic (ROC) curve analysis was used to define optimal cut-off levels for numerical variables regarding survival prediction. The presented survival curves and estimates are based on the Kaplan-Meier method. Univariate and multivariate survival associations were assessed with the Cox regression analysis. Variables were selected with the backward approach, with  $P < 0.05$  and  $P > 0.1$  inclusion and exclusion criteria, respectively.  $P$  values  $< 0.05$  were considered statistically significant. All analyses were performed with the MedCalc statistical software, version 20.006 (MedCalc Software Ltd, Ostend, Belgium).

## RESULTS

### Patients' characteristics

We enrolled 2586 patients who were discharged from hospital after index admission due to COVID-19. The median age was 70 years, IQR (60-80). A total of 1446 (55.9%) patients were men. The median Charlson comorbidity index was 4, IQR (2-5). Severe or critical COVID-19 symptoms on admission were present in 1956 (75.6%) patients. The median time from the start of COVID-19 symptoms to admission was 5 days, IQR (1-10), and the median length of hospital admission was 10 days, IQR (7-16).

During the median follow-up of 4 months, 192 (7.4%) patients died. The median survival time after hospital discharge was not reached. Three-month, 6-month, and 12-month survival rates were 93%, 92%, and 91%, respectively.

### Associations of demographic parameters and medical history with post-discharge survival

Higher mortality after hospital discharge was significantly associated with older age (hazard ratio [HR] 1.07,  $P < 0.001$ ) (Figure 1A). According to the ROC curve analysis, the optimal cut-off for survival discrimination was  $> 75$  years, with older patients having a higher risk of post-discharge mortality (HR 4.51, 95% confidence interval [CI] 3.3-6.14;  $P < 0.001$ ). Sex was not significantly associated with post-discharge survival ( $P = 0.166$ ).

Charlson comorbidity index was significantly associated with higher post-discharge mortality (HR 1.34;  $P < 0.001$ ). The ROC curve analysis yielded the cut-off level of  $> 4$  points, above which the patients experienced a higher risk

of death (HR 5.29, 95% CI 3.79-7.37;  $P < 0.001$ ) (Figure 1B). The frequency of individual comorbidities and their relationship with post-discharge survival are shown in Table 1.


#### Relationship of COVID-19 disease severity, clinical status on admission, and hospital stay parameters with post-discharge survival

The indication for hospital admission and the referral origin were significantly associated with post-discharge mortality. Patients with asymptomatic COVID-19 (admitted due to non-medical and social reasons, HR 2.2;  $P = 0.008$ ), patients admitted due to a neurologic indication (HR 2.08;  $P = 0.004$ ), and nursing-home residents (HR 3.95;  $P < 0.001$ ) experienced a worse post-discharge survival.

A worse post-discharge survival was significantly associated with a shorter disease duration on admission (HR 0.94;

$P < 0.001$ ), worse Eastern Cooperative Oncology Group (ECOG) status on admission (HR 1.88;  $P < 0.001$ ), higher symptom severity on presentation (HR 1.2;  $P = 0.046$ ), other infection on admission (HR 2.09;  $P < 0.001$ ), higher white blood cell count (WBC; HR 1.03;  $P < 0.001$ ), lower hemoglobin (HR 0.98;  $P < 0.001$ ), higher red blood cell distribution width (RDW; HR 1.13;  $P < 0.001$ ), lower estimated glomerular filtration rate (eGFR; HR 0.98;  $P < 0.001$ ), higher urea (HR 1.04;  $P < 0.001$ ), and higher D-dimer on admission (HR 1.28;  $P < 0.001$ ). Post-discharge survival stratified according to the admission ECOG status and WHO disease severity categories are shown in Figure 1C and 1D, respectively.

Hospital stay length negatively affected post-discharge prognosis (HR 1.03;  $P < 0.001$ ), and patients requiring  $> 11$  days of hospital stay experienced higher post-discharge mortality (HR 2.05, 95% CI 1.54-2.73;  $P < 0.001$ ). Mechanical ventilation survivors (HR 3.72;  $P < 0.001$ ) and patients



**FIGURE 1.** Post-discharge survival stratified according to the **A)** age categories (patients  $< 70$  years of age and  $\geq 90$  years of age were grouped together due to similar outcomes in younger patients and small number of patients/events in the elderly), **B)** Charlson comorbidity index quartiles, **C)** Eastern cooperative oncology group (ECOG) status on admission, and **D)** World Health Organization (WHO) disease severity categories.

who experienced prolonged immobilization during hospital stay ( $\geq 7$  days without bathroom privileges HR 2.56;  $P < 0.001$ ) had a worse post-discharge survival. Higher mortality was also observed in patients requiring hospital readmission (HR 4.92;  $P < 0.001$ ). Pneumonia, need for oxygen therapy, high-flow oxygen therapy, or intensive care unit stay were not significantly associated with post-discharge survival (Table 2).

### Multivariate analysis of factors affecting post-discharge survival

The variables significantly univariately associated with post-discharge mortality (excluding Charlson comorbidity index and including specific comorbidities) were included in the Cox regression model building process by using the backward approach. Numerical variables were dichotomized at optimal cut-off levels determined by using the ROC curve analysis (Table 3). Mutually independent predictors of a worse post hospital discharge mortality were

age  $> 75$  years, ECOG status 4,  $WBC > 7 \times 10^9/L$ ,  $RDW > 14\%$ , urea  $> 10.5$  mmol/L on admission, mechanical ventilation during hospital stay, readmission post discharge, absence of obesity, presence of chronic obstructive pulmonary disease, dementia, and metastatic malignancy.

### DISCUSSION

Our study showed that hospital treatment for COVID-19 led to prolonged health risks. In the post-discharge period, patients experienced substantial mortality, especially those with critical presentation of disease on admission and older patients with specific comorbidities. Similar long-term sequelae have been reported in other viral illnesses (7).

In our study, patients admitted to hospital from nursing homes and referred from other institutions had worse post-discharge outcomes than patients coming from home and those admitted due to acute neurological conditions and non-medical/social reasons (thus marked as as-

**TABLE 1.** Demographic and medical history parameters and their relationship with post-discharge survival (N = 2586)\*

	n (%) <sup>†</sup>	Univariate association with survival	
		HR and 95% CI	P
Age (years); median (IQR)	70 (60-80)	1.07 (1.06-1.09)	<0.001
Male sex	1446 (55.9)	0.81 (0.62-1.09)	0.166
Charlson comorbidity index; median (IQR)	4 (2-5)	1.34 (1.28-1.41)	<0.001
Arterial hypertension	1707 (66)	1.41 (1.03-1.95)	0.032
Diabetes mellitus	727 (28.1)	0.97 (0.7-1.33)	0.833
Obesity	708 (27.4)	0.51 (0.35-0.75)	<0.001
Chronic renal disease	256 (9.9)	1.89 (1.29-2.77)	0.001
COPD	164 (6.3)	1.83 (1.16-2.92)	0.010
Congestive heart failure	318 (12.3)	2.16 (1.54-3.03)	<0.001
Atrial fibrillation	357 (13.8)	2.85 (2.09-3.89)	<0.001
Liver cirrhosis	20 (0.8)	2.22 (0.71-6.9)	0.172
Dementia	357 (13.8)	4.32 (3.23-5.77)	<0.001
Active malignant disease	233 (9)	2.37 (1.64-3.42)	<0.001
Prior or active malignancy	412 (15.9)	1.74 (1.25-2.42)	0.001
Metastatic malignant disease	141 (5.5)	2.92 (1.93-4.42)	<0.001
Autoimmune/rheumatic disease	109 (4.2)	0.23 (0.06-0.92)	0.038
History of VTE	99 (3.8)	0.79 (0.35-1.78)	0.565
History of CVI	253 (9.8)	2.61 (1.84-3.7)	<0.001
History of MI	199 (7.7)	0.88 (0.5-1.54)	0.654
History of bleeding	127 (4.9)	1.43 (0.81-2.51)	0.212
VTE during hospitalization	155 (6)	0.96 (0.51-1.82)	0.904
MI during hospitalization	28 (1.1)	1.0 (0.25-4.08)	0.996
CVI during hospitalization	57 (2.2)	2.6 (1.37-4.92)	0.003
GI tract bleeding during hosp.	66 (2.6)	2.08 (1.06-4.06)	0.032

\*Abbreviations: HR – hazard ratio, CI – confidence interval, COPD – chronic obstructive pulmonary disease, VTE – venous thromboembolism, CVI – cerebrovascular infarction, MI – myocardial infarction, GI – gastrointestinal. †if not otherwise indicated.

ymptomatic) rather than due to COVID-19 pneumonia *per se*. Patients referred from other institutions and those com-

ing from nursing homes were significantly older and more burdened with comorbidities, as well as had a significantly

**TABLE 2.** Parameters related to coronavirus disease 2019 hospital admission and their relationship with post-discharge survival (N = 2568)\*

	n (%)	Univariate association with survival HR and 95% CI	P
<b>Indication for admission</b>			
asymptomatic	85 (3.3)	2.2 (1.24-3.93)	0.008
pneumonia	1610 (62.3)	Reference category	
high temperature without pneumonia	95 (3.7)	0.73 (0.29-1.79)	0.489
acute medical condition	356 (13.8)	0.33 (0.81-1.86)	0.327
acute neurological condition	136 (5.3)	2.08 (1.27-3.44)	0.004
acute surgical condition	304 (11.8)	0.98 (0.61-1.59)	0.964
<b>Origin of referral</b>			
patient's home	1018 (39.4)	Reference category	
nursing home	223 (8.6)	3.95 (2.54-6.15)	<0.001
other hospital	1345 (52)	2.03 (1.43-2.88)	<0.001
<b>Disease duration on admission (days)</b>	5 (1-10)	0.94 (0.91-0.97)	<0.001
<b>ECOG status on admission</b>			
0	307 (11.9)	Reference category	
1	659 (25.5)	6.06 (1.43-25.6)	0.014
2	648 (25.1)	8.55 (2.05-35.6)	0.003
3	545 (21.1)	17.8 (4.36-73.35)	<0.001
4	427 (16.5)	31.6 (7.75-128.76)	<0.001
<b>MEWS score</b>	2 (1-3)	1.09 (1.01-1.19)	0.036
<b>WHO severity on admission</b>			
mild	433 (16.7)	Reference category	
moderate	197 (7.6)	1.04 (0.54-2.01)	0.914
severe	1744 (67.4)	1.19 (0.79-1.82)	0.393
critical	212 (8.2)	1.99 (1.16-3.43)	0.013
<b>Admission WBC (x 10<sup>9</sup>/L)</b>	7.4 (5.5-10.3)	1.03 (1.02-1.05)	<0.001
<b>Admission hemoglobin (g/L)</b>	129.5 (115-142)	0.98 (0.97-0.99)	<0.001
<b>Admission RDW</b>	13.8 (13.2-14.9)	1.13 (1.09-1.18)	<0.001
<b>Admission platelets (x 10<sup>9</sup>/L)</b>	230 (172-305)	0.99 (0.99-1.0)	0.169
<b>Admission CRP</b>	72.2 (27.8-131.9)	1.0 (0.99-1.0)	0.239
<b>Admission ferritin</b>	640 (347-1132)	0.99 (0.99-1.0)	0.648
<b>Admission eGFR</b>	79.2 (55.4-94.2)	0.98 (0.97-0.99)	<0.001
<b>Admission urea (mmol/L)</b>	7.1 (5.2-10.3)	1.04 (1.03-1.05)	<0.001
<b>Admission D-dimers (mg/L FEU)</b>	1.14 (0.63-2.62)	1.28 (1.14-1.43)	<0.001
<b>Pneumonia</b>	2124 (82.1)	1.37 (0.91-2.06)	0.131
<b>Other infection on admission</b>	324 (12.5)	2.09 (1.49-2.95)	<0.001
<b>Oxygen therapy</b>	1872 (72.4)	1.3 (0.93-1.82)	0.124
<b>High flow oxygen therapy</b>	192 (7.4)	1.15 (0.68-1.95)	0.598
<b>Mechanical ventilation</b>	74 (2.9)	3.72 (2.23-6.22)	<0.001
<b>Need for ICU</b>	240 (9.3)	1.21 (0.76-1.93)	0.409
<b>Length of hospitalization</b>	10 (7-16)	1.03 (1.02-1.04)	<0.001
<b>Immobilization ≥7 days</b>	739 (28.6)	2.56 (1.93-3.41)	<0.001
<b>Readmission to the hospital</b>	76 (2.9)	4.92 (3.16-7.68)	<0.001

\*Abbreviations: HR – hazard ratio, CI – confidence interval, ECOG – Eastern cooperative oncology group, MEWS – modified early warning score, WHO – World health organization, WBC – white blood cell count, RDW – red blood cell distribution width, CRP – C reactive protein, eGFR – estimated glomerular filtration rate, ICU – intensive care unit.

worse functional status on admission, than patients coming from home, which is probably the reason for our finding since these patients had less severe COVID-19 symptoms on admission (data not shown). No association with CRP on admission was found despite the association with other measures of increased inflammation, such as higher D-dimer and RDW. Nevertheless, a composite CRP-based index determined on admission has been shown to predict long-term outcomes (8). As the current study shows, the intensity of COVID-19 symptoms quantified with the modified early warning score at hospital admission and mechanical ventilation requirement have negative prognostic value in the post-discharge period. Systemic inflammatory response associated with severe acute respiratory syndrome coronavirus 2 infection might lead to a deterioration in the existing comorbidities (9) after the recovery of respiratory self-sufficiency.

Lower hemoglobin levels and higher RDW, which are recognized negative prognostic factors for acute COVID-19 infection (10,11), were also shown to affect the outcomes in the post-discharge period. A similar role was observed for parameters of renal excretion function. The quality of hematopoiesis and parameters of renal excretion function, as universal markers of overall health fitness, were shown to be prognostic in a number of chronic and malignant diseases in the pre-COVID era (12-14). Higher urea may indicate worse hydration status on admission, especially considering that it outperformed the presence of chronic kidney disease and creatinine in a multivariate prognostic model. Leukocytosis, as a non-specific parameter of physical stress, also seems to be associated with a worse long-

term prognosis. Nevertheless, platelets, which might play an important role in the protection of lung parenchyma during infection (15), were not associated with post-discharge survival.

Sex, which significantly affects the survival of patients in the hospital setting (4), was not associated with post-discharge survival, which is contrary to other reports (16). However, older age and higher comorbidity burden evidently affected the post-discharge outcomes. Post-discharge mortality could be considered as a surrogate measure of cardiovascular events, especially considering that COVID-19 might increase long-term cardiovascular risks (17). In the current study, particularly detrimental effects on post-discharge survival were evident in patients with dementia, cardiovascular, pulmonary, and malignant diseases. These comorbidities highly impair daily activities and functional independence. Indeed, the measures of functional status on admission (like ECOG scale), as well as during hospital stay (immobilization without bathroom privileges due to imposed or objective medical reasons) and the need for a prolonged hospital stay all negatively affected post-discharge survival. These findings agree with other reports that recognize impaired functional status as an important negative consequence of COVID-19 infection after hospital treatment for acute disease (5,6,16,18). Functional status is a potentially modifiable factor, and preventive measures and targeted interventions aimed at its improvement might contribute to better outcomes of patients with a higher mortality risk. The need for rehospitalization also predicted a worse post-discharge survival, which highlighted the fact that patients with long-term

**TABLE 3.** Cox regression analysis model investigating mutually independent associations of patient-related and hospital stay-related parameters with survival after hospital discharge

Variable	Multivariate association with survival	
	HR and 95% CI	P
Age>75 years	2.92 (1.9-4.48)	<0.001
ECOG status 4	1.94 (1.3-2.9)	0.001
WBC>7 ×10 <sup>9</sup> /L	1.5 (1.02-2.21)	0.039
RDW>14%	1.89 (1.26-2.84)	0.002
Urea>10.5 mmol/L	1.81 (1.24-2.64)	0.002
Mechanical ventilation	8.68 (4.71-15.97)	<0.001
Readmission post discharge	5.28 (2.92-9.54)	<0.001
Obesity	0.47 (0.29-0.78)	0.003
Chronic obstructive pulmonary disease	2.06 (1.15-3.72)	0.016
Dementia	2.19 (1.45-3.32)	<0.001
Metastatic malignancy	3.23 (1.89-5.55)	<0.001
History of cardiovascular insult	1.58 (0.99-2.51)	0.055

\*Abbreviations: HR – hazard ratio, CI – confidence interval, ECOG – Eastern cooperative oncology group, WBC – white blood cell count, RDW – red blood cell distribution width.

mortality risk are frail and might need home health services or chronic care (16).

Limitations of our study are single-center experience, retrospective study design, and the fact that it was conducted in a pandemic period characterized by low vaccination rates. The data were collected in a high-volume tertiary center treating the most severe cases of COVID-19 and patients who required urgent medical care for other conditions but also tested positive for SARS-CoV-2. The patients treated in our institution might not be directly comparable to those from other cohorts due to ethnical and regional differences, and due to differences in patient profile. Our results should be primarily interpreted as representative of Croatian population. Furthermore, the follow-up period was relatively short. Despite these limitations, our results provide valuable and unique insights into the post-discharge period risks in patients previously hospitalized for COVID-19.

In conclusion, the hospital stay for COVID-19 leads to a substantial risk of death post discharge. Patients with increased risk for death are those of older age, higher functional impairment, need for mechanical ventilation, parameters indicative of more pronounced inflammation, impaired renal function, and particular comorbidities. These patients might require interventions aimed at improvement of functional capacity.

**Acknowledgments** This study is part of the project Registar hospitalno liječenih bolesnika u Respiracijskom centru KB Dubrava/Registry of Hospitalized Patients in University Hospital Dubrava Respiratory Center.

**Funding** None.

**Ethical approval** given by the University Hospital Dubrava Review Board (2021/2503-04).

Declaration of authorship ML conceived and designed the study; ML, NPŽ, MZ, MBJ, UJ, AM, NZ, MLa, BO, IB, SS, IV acquired the data; all authors analyzed and interpreted the data; ML drafted the manuscript; all authors critically revised the manuscript for important intellectual content; all authors gave approval of the version to be submitted; all authors agree to be accountable for all aspects of the work.

**Competing interests** ML is a statistical editor in the *Croatian Medical Journal*. To ensure that any possible conflict of interest relevant to the journal has been addressed, this article was reviewed according to best practice guidelines of international editorial organizations. All authors have completed the Unified Competing Interest form at [www.icmje.org/coi\\_disclosure.pdf](http://www.icmje.org/coi_disclosure.pdf) (available on request from the corresponding author) and declare: no support from any organization for the submitted work; no financial relationships with any organizations that might have an interest in the submitted work in the previous 3 years; no other relationships or activities that could appear to have influenced the submitted work.

## References

- Attaway AH, Scheraga RG, Bhimraj A, Bieh M, Hatipoglu U. Severe covid-19 pneumonia: pathogenesis and clinical management. *BMJ*. 2021;372. [Medline:33692022](#) doi:10.1136/bmj.n436
- Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (covid-19) outbreak in China: summary of a report of 72 314 cases from the Chinese Center for Disease Control and Prevention. *JAMA*. 2020;323:1239-42. [Medline:32091533](#) doi:10.1001/jama.2020.2648
- Busic N, Lucijanic T, Barsic B, Luksic I, Busic I, Kurdija G, et al. Vaccination provides protection from respiratory deterioration and death among hospitalized COVID-19 patients: Differences between vector and mRNA vaccines. *J Med Virol*. 2022;94:2849-54. [Medline:35187697](#) doi:10.1002/jmv.27666
- Piskač Živković N, Lucijanić M, Bušić N, Jurin I, Atić A, Andrilović A, et al. The associations of age, sex, and comorbidities with survival of hospitalized patients with coronavirus disease 2019: data from 4014 patients from a tertiary-center registry. *Croat Med J*. 2022;63:36-43. [Medline:35230004](#) doi:10.3325/cmj.2022.63.36
- Chopra V, Flanders SA, O'Malley M, Malani AN, Prescott HC. Sixty-day outcomes among patients hospitalized with COVID-19. *Ann Intern Med*. 2021;174:576-8. [Medline:33175566](#) doi:10.7326/M20-5661
- Banić M, Janković Makek M, Samaržija M, Muršić D, Boras Z, Trkeš V, et al. Risk factors and severity of functional impairment in long COVID: a single-center experience in Croatia. *Croat Med J*. 2022;63:27-35. [Medline:35230003](#) doi:10.3325/cmj.2022.27
- Ahmed H, Patel K, Greenwood DC, Halpin S, Lewthwaite P, Salawu A, et al. Long-term clinical outcomes in survivors of severe acute respiratory syndrome and Middle East respiratory syndrome coronavirus outbreaks after hospitalisation or ICU admission: A systematic review and meta-analysis. *J Rehabil Med*. 2020;52:jrm00063. [Medline:32449782](#) doi:10.2340/16501977-2694
- Lucijanic M, Stojic J, Atic A, Cikara T, Osmani B, Barisic-Jaman M, et al. Clinical and prognostic significance of C-reactive protein to albumin ratio in hospitalized coronavirus disease 2019 (COVID-19) patients: Data on 2309 patients from a tertiary center and validation in an independent cohort. *Wien Klin Wochenschr*. 2022;134:377-84. [Medline:35038003](#) doi:10.1007/s00508-021-01999-5
- Guo T, Fan Y, Chen M, Wu X, Zhang L, He T, et al. Cardiovascular implications of fatal outcomes of patients with coronavirus disease 2019 (COVID-19). *JAMA Cardiol*. 2020;5:811-8. [Medline:32219356](#) doi:10.1001/jamacardio.2020.1017
- Tao Z, Xu J, Chen W, Yang Z, Xu X, Liu L, et al. Anemia is associated with severe illness in COVID-19: A retrospective cohort study. *J Med Virol*. 2021;93:1478-88. [Medline:32813298](#) doi:10.1002/jmv.26444
- Lucijanić M, Jordan A, Jurin I, Piskač Živković N, Sorić E, Hadžibegović I, et al. Red cell distribution width is a potent prognostic parameter for in-hospital and post-discharge mortality in hospitalized coronavirus disease 2019 patients: a registry-based cohort study on 3941 patients. *Croat Med J*. 2022;63:44-52. [Medline:35230005](#) doi:10.3325/cmj.2022.63.44

- 12 Kust D, Lucijanac M, Urch K, Samija I, Celap I, Kruljac I, et al. Clinical and prognostic significance of anisocytosis measured as a red cell distribution width in patients with colorectal cancer. *QJM*. 2017;110:361-7. [Medline:28069908](#) [doi:10.1093/qjmed/hcw223](#)
- 13 Lucijanac M, Krecak I, Verstovsek S, Soric E, Galusic D, Holik H, et al. Higher red blood cell distribution width predicts thrombosis risk in primary and secondary myelofibrosis. *Ann Hematol*. 2022. [Medline:34825960](#) [doi:10.1007/s00277-021-04724-4](#)
- 14 Lucijanac M, Galusic D, Krecak I, Sedinic M, Holik H, Perisa V, et al. Reduced renal function strongly affects survival and thrombosis in patients with myelofibrosis. *Ann Hematol*. 2020;99:2779-85. [Medline:32862283](#) [doi:10.1007/s00277-020-04239-4](#)
- 15 Lucijanac M, Krecak I, Soric E, Sedinic M, Sabljic A, Derek L, et al. Thrombocytosis in COVID-19 patients without myeloproliferative neoplasms is associated with better prognosis but higher rate of venous thromboembolism. *Blood Cancer J*. 2021;11:189. [Medline:34845183](#) [doi:10.1038/s41408-021-00585-2](#)
- 16 Bowles KH, McDonald M, Barrón Y, Kennedy E, O'Connor M, Mikkelsen M. Surviving COVID-19 after hospital discharge: symptom, functional, and adverse outcomes of home health recipients. *Ann Intern Med*. 2021;174:316-25. [Medline:33226861](#) [doi:10.7326/M20-5206](#)
- 17 Xie Y, Xu E, Bowe B, Al-Aly Z. Long-term cardiovascular outcomes of COVID-19. *Nat Med*. 2022;28:583-90. [Medline:35132265](#) [doi:10.1038/s41591-022-01689-3](#)
- 18 Chojnicki M, Neumann-Podczaska A, Seostianin M, Tomczak Z, Tariq H, Chudek J, et al. Long-term survival of older patients hospitalized for COVID-19. Do clinical characteristics upon admission matter? *Int J Environ Res Public Health*. 2021;18. [Medline:34682421](#) [doi:10.3390/ijerph182010671](#)