


Targetability of the neurovascular unit in inflammatory diseases of the central nervous system

Brandon C. Smith^{1,2} | Rachel A. Tinkey^{1,3} | Benjamin C. Shaw¹  | Jessica L. Williams^{1,4} 

¹Department of Neurosciences, Lerner Research Institute, Cleveland Clinic, Cleveland, Ohio, USA

²Department of Biological, Geological, and Environmental Sciences, Cleveland State University, Cleveland, Ohio, USA

³School of Biomedical Sciences, Kent State University, Kent, Ohio, USA

⁴Brain Health Research Institute, Kent State University, Kent, Ohio, USA

Correspondence

Jessica L. Williams, Lerner Research Institute, Cleveland Clinic, 9500 Euclid Avenue/NC30, Cleveland, OH 44195, USA.

Email: williaj39@ccf.org

Funding information

National Heart and Lung Institute, Grant/Award Number: T32 HL150389; National Institute of Neurological Disorders and Stroke, Grant/Award Number: K00 NS120365 and R01 NS119178

Summary

The blood–brain barrier (BBB) is a selectively permeable barrier separating the periphery from the central nervous system (CNS). The BBB restricts the flow of most material into and out of the CNS, including many drugs that could be used as potent therapies. BBB permeability is modulated by several cells that are collectively called the neurovascular unit (NVU). The NVU consists of specialized CNS endothelial cells (ECs), pericytes, astrocytes, microglia, and neurons. CNS ECs maintain a complex “seal” via tight junctions, forming the BBB; breakdown of these tight junctions leads to BBB disruption. Pericytes control the vascular flow within capillaries and help maintain the basal lamina. Astrocytes control much of the flow of material that has moved beyond the CNS EC layer and can form a secondary barrier under inflammatory conditions. Microglia survey the border of the NVU for noxious material. Neuronal activity also plays a role in the maintenance of the BBB. Since astrocytes, pericytes, microglia, and neurons are all able to modulate the permeability of the BBB, understating the complex contributions of each member of the NVU will potentially uncover novel and effective methods for delivery of neurotherapies to the CNS.

KEYWORDS

blood–brain barrier, endothelial cell, glia, neuroinflammatory disease, pericyte, tight junction

1 | INTRODUCTION

The blood–brain barrier (BBB) is a complex, highly regulated system with multiple cell types influencing its maintenance and formation. The BBB forms a privileged environment for the central nervous system (CNS), restricting entry of a wide breadth of potential hazards, including pathogens, immune cells, antibodies, and pharmaceuticals. This tight regulation protects the brain from external insults, but simultaneously prevents access of many therapeutics meant to treat neuroinflammatory and neurodegenerative disorders. As a result, multiple clinical trials have failed despite promise in preclinical studies, underscoring the need for a more complete understanding of the

BBB and its modulatory mechanisms. In this review, we discuss the composition of the neurovascular unit (NVU), known mechanisms of BBB modulation, and potential therapeutic targets for neuroinflammatory disorders.

The existence of a selectively permeable BBB was first postulated at the turn of the 20th century when it was noted that water-soluble dyes injected into the periphery did not permeate the CNS,^{1,2} and dyes injected into the CNS parenchyma did not exit to the periphery.^{2,3} Further complexity of the BBB was demonstrated by seminal evidence indicating immune privilege, wherein immune cells are excluded and unable to surveil the CNS, over 100 years ago by Shirai⁴ and later confirmed by Murphy and Sturm.⁵ However,

This article is part of a series of reviews covering Neuroimmunology appearing in Volume 311 of *Immunological Reviews*

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *Immunological Reviews* published by John Wiley & Sons Ltd.

even now, the permeability, complexity, and dynamics of the BBB are incompletely understood.^{6,7}

The BBB acts as a robust biological gateway, responsible for the flow of nearly all molecules and cells passing into and out of the CNS.^{6,8,9} This selective permeability is critical in sparing the CNS from many circulating toxins and pathogens and allowing for availability of necessary nutrients to the CNS.^{2,10,11} Instead of passing through fenestrations in endothelial cells (ECs) as it does in the peripheral vasculature, material is transported and scrutinized by one or more cells closely associated with the BBB.^{11,12} However, it is this very system, designed to protect the CNS, that also hinders therapeutic development to treat CNS disorders including multiple sclerosis (MS), Alzheimer's disease (AD), Parkinson's disease, traumatic brain injuries, and brain cancers.^{10,11,13} Several researchers have even employed artificial intelligence to determine the likelihood of small molecules to penetrate the BBB.^{13,14}

This review will examine some of the NVU cell-specific effects on BBB permeability as well as the intercellular communication that is critical for proper function of the NVU. While other complex CNS barriers, including the blood-retinal barrier, the blood-cerebrospinal fluid barrier, and the arachnoid layer,¹⁵ are also important to consider in therapeutic availability to the CNS, we will only focus on the BBB for the purposes of this review.

2 | STRUCTURE AND ORGANIZATION OF THE NVU

The NVU is the fundamental multicellular unit supporting the BBB and consists of ECs, pericytes, astrocytes, microglia, and neurons (Figure 1).^{6,7} The capillary lumen is the primary area of contact between the NVU and the periphery, and interfacing factors vary from cytokines and hormones to immune cells that have the potential to extravasate.^{16,17} Forming the vessel lumen is specialized CNS ECs, which are semipermeable under homeostatic conditions due to the presence of tight junctions.^{6,16,18–20} Perivascular mural cells, or pericytes, have recently been shown to be more complex and varied than initially assumed. Pericytes are functionally heterogeneous^{21,22} and further understanding of their diverse roles in the CNS is an active area of research. Pericytes were classically described to help govern vasoconstriction and dilation of capillaries,^{23,24} which is highly regulated in the NVU, to limit BBB permeability. Recent studies are beginning to uncover pericyte coordination with other members of the NVU, particularly ECs and astrocytes.^{21,24} Astrocytes are vital to the NVU and while the other members of the NVU have critical roles, it is largely the astrocytes that set the NVU apart from other capillary layers in the body.^{25,26} Understanding the specific mechanisms by which astrocytes regulate tight junctions in CNS ECs could prove incredibly beneficial to optimizing BBB permeability for drug delivery. Astrocytes are relatively large and abundant CNS trophic cells²⁷ and their endfeet form an additional barrier around CNS capillaries.^{6,25} The perivascular, or Virchow-Robin, space refers to the area between ECs and astrocytes and is a key checkpoint prior to entering

the CNS parenchyma.^{6,28} Microglia are a recent addition to our understanding of the NVU. Although astrocytes cover the majority of the perivascular space, there are some gaps in coverage. Recently, Kisler et al. used two-photon in vivo imaging to observe microglial processes covering many of these gaps.²⁹ How involved microglia are in the structure of the NVU is still being investigated. By area, neurons have a relatively minor role in the structure of the NVU but can modulate BBB permeability via neuronal activity.^{30,31} Due to the complex nature of the NVU, it has proven difficult to study. In vitro models have been useful; however, the heterogeneity between species, individuals, and even CNS regions have added to this investigational barrier.^{32,33} Here, we will discuss the components of the NVU and how each cell type may impact BBB permeability.

3 | ENDOTHELIAL CELLS

ECs are found throughout the body and make up the walls of arteries, veins, and capillaries. They form a tube-like structure to allow passage of various blood cells and proteins.³⁴ ECs secrete and maintain the basal lamina—the extracellular matrix on which they reside.^{35,36} The CNS ECs differ in several ways from peripheral ECs. Most notably, they lack the fenestrations found in peripheral vessels and instead create a continuous cellular barrier with significantly reduced permeability along the capillary lumen.³⁷ Instead, CNS ECs form tight junctions which prevent hydrophobic molecules from penetrating the endothelial layer unless transported through the cell. These tight junctions are responsible for much of the impermeability of the BBB. Tight junctions are composed primarily of claudin-5, occludin, and other junctional adhesion molecules³⁸ and bind to the actin cytoskeleton via ZO-1, ZO-2, or ZO-3.³⁹ Tight junctions are highly complex and can vary depending on the proteins coupled.^{40,41} Transport of nutrients, waste, and signaling molecules between the CNS and the periphery is necessary under physiological conditions and is typically achieved by a myriad of EC transporters.^{6,42} In fact, these transporters make up 10–15% of the total protein in CNS ECs and allow specific molecules, peptides, and even cells to cross into the perivascular space, bypassing tight junctions.^{33,43}

3.1 | ECs and immunity

Classically, it was thought that immune cells could not access the CNS parenchyma.^{18,44} While mostly impermeable during homeostatic conditions, CD4⁺ and CD8⁺ T cells are allowed passage for immune surveillance of the CNS, facilitated by a highly regulated, multistep process through a tricellular junction.⁴⁵ This process involves endothelial ligands vascular cell adhesion molecule (VCAM)-1 and intercellular adhesion molecule (ICAM)-1 which recognize lymphocyte function-associated antigen 1 (LFA-1) and very late antigen-4 (VLA-4) on T cells. These interactions arrest T cells on the endothelium and allow migration against the flow of blood to these tricellular junctions and ultimately extravasation.^{45,46} These




FIGURE 1 Cellular components of the neurovascular unit. The NVU is composed of a complex network of cells that are functionally diverse. ECs form the walls of blood vessels and capillaries and contribute to the formation and maintenance of the basal lamina and extracellular matrix. In addition, tight junctions formed between ECs and expression of adhesion molecules regulate BBB permeability. Pericytes reside in the capillary bed and, with regard to BBB integrity, are primarily responsible for modulation of vascular flow as well as structural changes in tight junctions and the extracellular matrix. Astrocyte endfeet cover 90%–95% of the area surrounding the BBB and contribute to a variety of processes that include, but are not limited to, osmotic homeostasis, trophic factor concentration, molecular transport into and out of the perivascular space, and formation of gap junctions under neuroinflammatory conditions. Microglia have been found to cover the remaining space around BBB ECs, respond to injury and infection, and regulate tight junction formation between ECs. Neurons predominantly communicate with astrocyte endfeet and aid in regulation of tight junctions and molecular transport.

tracellular junctions contain the proteins tricellulin and angulin-1 which direct and aid T-cell diapedesis.⁴⁵ While CNS ECs express VCAM-1 and ICAM-1 under normal conditions, the expression is sparse, limiting the number of leukocytes that can traverse triclular junctions.^{45,47}

In response to infection, autoimmunity, or injury, the permeability of the BBB is significantly enhanced and can lead to severe inflammation.⁴⁸ This is partially due to an increase in the expression of adhesion molecules on the EC surface, primarily ICAM-1 and VCAM-1, which can arrest a greater number of immune cells for extravasation.⁴⁹ Inflammation can lead to break down of the tight junctions, allowing leukocytes to invade the perivascular space.⁴⁸ Many secreted inflammatory factors, cell damage signals, and pathogen components can alter tight junction integrity including CCL2 and transforming growth factor (TGF)- β , which is known to alter expression of claudin-5, occludin, and ZO-1.^{39,50,51} Likewise, tumor necrosis factor (TNF)- α , lipopolysaccharide (LPS), and mitochondrial damage can induce BBB permeability via actin filament rearrangement.^{52–55} Prolonged dysfunction of the BBB, as in chronic inflammatory CNS diseases such as MS or AD, can lead to permanent CNS tissue damage and neuroaxonal loss.^{56–58} However, although overt and/or chronic inflammation can lead to the loss of BBB integrity

and extensive damage to the CNS, low-to-moderate amounts of inflammation can partially restore the BBB and limit peripheral immune cell infiltration into the CNS parenchyma, mitigating injury, or infection.^{56,59–61} Taking advantage of this immunological state and exploiting molecular signals used in the formation of triclular junctions may be a novel avenue for therapeutic development.

3.2 | Targeting ECs

It is possible to permeate the BBB to treat neurological disorders, but the potential bystander effects of leaky BBB during diseases such as glioblastoma, MS, AD, and others imposes significant risk. The BBB is disrupted in many neurodegenerative and psychological disorders. Additional access of immune cells, toxins, or other inflammatory mediators could enhance inflammation-mediated damage to the CNS. Additionally, BBB dysregulation disrupts homeostatic transport across the barrier, which provides trophic factors and controls osmotic regulation during physiological conditions.⁶² Bypassing the ECs of the BBB is a difficult challenge; however, there are several other members of the NVU that introduce potentially novel avenues for drug targeting.

4 | PERICYTES

Proportionally, there is a higher density of pericytes in the CNS relative to peripheral organs,²⁴ suggesting they have a critical role in the CNS. As vascular mural cells, pericytes play a large role in the dilation of capillaries in the NVU. Of note, pericytes are found in the capillary bed and not surrounding arteries or veins, which house similar, but distinct, vascular mural cells, and vascular smooth muscle cells.⁶³ In addition to vasodilation and vasoconstriction, pericytes also help form and maintain the basal lamina.^{64,65} Control of vascular flow by pericytes can indirectly impact the BBB as changes in blood flow can allow for more or less cellular contact in the capillary lumen. Increases in cellular interaction can stretch and stress CNS ECs,^{66–68} causing pericyte dysfunction, which is associated with aberrant immune cell trafficking.⁶⁹ Further, in mouse models deficient in functional pericytes, there is a loss of vascular control, dysfunctional tight junction regulation, and aberrant angiogenic sprouting, highlighting the importance of pericytes in maintaining a healthy BBB.^{69–73}

New roles for pericytes continue to be uncovered, including their ability to maintain and produce elements of the basal lamina^{74–77} and intricate regulation of tight junctions. Contrary to ECs, TGF- β and angiopoietin-1 (Ang-1) signaling in pericytes enhances the expression of occludin on CNS ECs, reducing the permeability of the BBB. TGF- β from pericytes also activates Smad4 signaling to upregulate bone morphogenic proteins (BMPs). BMPs can then ensure tight

adherence of pericytes to ECs via N-cadherins, which is upregulated by vascular endothelial growth factor (VEGF), reinforcing the communication and physical interaction between pericytes and CNS ECs.^{78–80} While basal levels of VEGF signaling can reduce BBB permeability, excessive VEGF signaling to pericytes can lead to downregulation of claudin-5 on ECs and dysregulate BBB tight junctions (Figure 2).^{81–84}

4.1 | Heterogeneity of pericytes

Similar to ECs, pericytes respond robustly to various signaling molecules. Inflammatory and non-inflammatory pericytes have been described and subdivided into Type-1 pericytes (PC1) and Type-2 pericytes (PC2). PC1s are non-inflammatory and tend to be the resting state of pericytes without injury or infection. PC2s tend to increase in frequency with and are highly responsive to inflammation.^{73,85} It is likely, as is the case with astrocytes and microglia that there is a spectrum of activation states, but as the study of pericytes is still in its infancy, this has not yet been fully elucidated. In young healthy patients, nearly all pericytes exhibit a non-inflammatory morphology.^{71,85,86} However, aging results in an increase in the population of inflammatory pericytes, which is consistent with enhanced inflammation and BBB permeability associated with age.⁸⁵ Both pericyte subtypes contribute to the




FIGURE 2 NVU interaction network. Cells within the NVU interact through intricate signaling mechanisms that allow for proper functioning of the BBB. ECs receive a variety of protective signals from other NVU cells that upregulate tight junction formation thus enhancing BBB integrity. These factors include, but are not limited to, TGF β , Ang-1, APOE, Shh, Wnts, glial-derived neurotrophic factor, insulin-like growth factor (IGF)-1, and retinoic acid. Contrastingly, ECs may also receive signals that downregulate tight junction proteins, particularly during inflammatory events, such as TNF α , NO, MMPs, endothelins, and glutamate that lead to increased BBB permeability. Importantly, ECs also maintain the ability to signal to pericytes through VEGF and N-cadherin, which are necessary for BBB maintenance. Immune checkpoint proteins, such as the PD-1/PD-L1 complex, regulate the cellular activity of microglia and neurons to modulate BBB integrity as well as dampen the inflammatory activity of infiltrating peripheral immune cells, which increases BBB permeability. Finally, neuronal activity is a critical modulator of cellular and molecular transport across the BBB.

basal lamina, but PC2s tend to produce a more irregular basal lamina, impacting both ECs and astrocytes.^{21,85} Additionally, PC2s produce less laminin-111 and laminin-211, which leads to cell hypertrophy and BBB disruption.^{87,88} Exposing pericytes to inflammatory cytokines, LPS, or reactive oxygen species induces immunoreactivity in pericytes, altering their morphology, as well as inducing their separation from the basal lamina.⁸⁷⁻⁸⁹ The dynamic morphology of pericytes suggests they have a critical role in the integrity and function of the BBB, which is altered in pathological states and in aging.

4.2 | Targeting pericytes

Examining the targetability of pericytes and its potential influence on the BBB is difficult but may prove promising. Affecting pericytes may be a subtler approach to drug penetrance rather than reducing the integrity of the EC layer itself. A deeper understanding of how pericytes communicate with CNS ECs and astrocytes could reveal potential nuanced approaches to bypass the BBB. While pericytes may be a promising target to leverage BBB permeability, much about pericyte mechanisms of BBB control remains unknown.

5 | ASTROCYTES

Astrocytes are large stellate cells with extensive processes that extend throughout the CNS.^{27,90} However, unlike CNS ECs and pericytes, astrocytes are unique in that only the ends of their processes are considered part of the NVU (Figure 1). These endfeet cover roughly 90–95% of the area surrounding the BBB and have properties that are unique from the rest of the astrocyte.⁹¹⁻⁹³ Endfeet contain aquaporin-4 and the potassium channel $K_{ir}4.1$ to modulate water and ion balance and, lacking tight junctions, astrocytic endfeet permit immune cell extravasation.^{92,94,95} Although endfeet lack the barrier that tight junctions provide, ablation of astrocytes leads to rapid and extensive BBB deterioration.^{25,93} And interestingly, early studies transplanting astrocytes outside of the CNS demonstrated that transplanted astrocytes develop a BBB-like morphology surrounding the peripheral vasculature.^{96,97} In addition to the physical barrier that astrocytes provide, during homeostasis, astrocytes are responsible for the transport of material from the perivascular space into the parenchyma.^{6,25} This function makes them excellent and critical supportive cells in the maintenance of the BBB.

Astrocytes form gap junctions between endfeet forming a much “looser” network of connections compared with that of CNS ECs.⁹⁸ Gap junctions are formed by a hexamer of proteins called connexins. The material that is transported by a given gap junction is largely dependent on the specific combination of connexins that make up the junction.^{99,100} Notable gap junctions in astrocytes are formed by connexin 30 and connexin 43, which help mediate glucose and lactate transport to distal neurons.¹⁰¹ Astrocytes use these gap junctions to communicate with other cells of the NVU including neurons,

microglia, and other astrocytes using ion gradients, electrical signals, and signaling molecules.¹⁰²

5.1 | Tight junction modulation

Astrocytes release a variety of trophic factors that help maintain a functional NVU.¹⁰³⁻¹⁰⁵ Many of the factors released under physiologic conditions increase the amount and order of tight junction proteins between BBB ECs.^{106,107} For instance, astrocytes secrete sonic hedgehog (Shh), which increases tight junctions in CNS ECs by inducing Patch1 signaling.¹⁰⁷ Other astrocytic factors that can enhance tight junctions in ECs include Wnt signaling, TGF β , and apolipoprotein E (ApoE).^{104,106,108-110} ApoE and TGF β may work indirectly through pericytes to impact EC tight junctions.^{111,112} Additionally, astrocytes produce Ang-1, which signals to EC Tie2 to further upregulate tight junction proteins and reduce adhesion molecules, reducing the likelihood of leukocyte entry.^{113,114} Further, astrocytes are incredibly responsive to inflammatory stimuli,^{115,116} and during inflammation, it has been shown that BBB ECs downregulate claudin-5, while astrocytes upregulate tight junction proteins claudin-1, claudin-4, and JAM-A.^{117,118} This suggests that astrocyte endfeet form a secondary barrier to prevent excessive immune cell entry past the perivascular space, regulating access to the CNS parenchyma.¹¹⁷ This accumulation of peripheral immune cells in the perivascular space is often referred to as perivascular cuffing and is commonly seen in MS lesions.^{117,119}

However, although astrocytes provide an impressive physical barrier, activated astrocytes can also secrete trophic factors, chemokines, and cytokines that dysregulate EC tight junctions and recruit peripheral immune cells (Figure 2).^{25,39,119,120} These factors can include VEGF, nitric oxide, MMPs, endothelins, and glutamate; all of which can downregulate endothelial tight junction proteins.^{39,121,122} Importantly, astrocytes also promote a return to homeostasis following an inflammatory event, secreting several beneficial trophic factors including Shh, astrocyte-derived Ang-1, glial-derived neurotrophic factor (GDNF), insulin-like growth factor-1, ApoE, and retinoic acid. These factors not only prevent EC apoptosis, but also stimulate tight junction formation and a return to a homeostatic state.^{116,121} Taken together, astrocytes orchestrate a complex modulation of the BBB and may represent an amenable cell type for drug manipulation.

5.2 | Astrocytes as an immunologic barrier

Since astrocytes are highly responsive to immune stimuli, they have the ability to upregulate a wide array of chemokines and adhesion molecules that can serve as an immunologic barrier to prevent excessive influx of inflammatory cells during a CNS insult (Figure 2).^{120,123} Interestingly, in the event immune cells breach the BBB, enter the perivascular space, and cross astrocyte endfeet into the parenchyma, astrocytes can express immune checkpoint

molecules, or inhibitory receptors, which can induce exhaustion and death of leukocytes,^{124,125} providing the CNS multiple layers of immunologic protection. As an example, astrocytes are known to upregulate the immune checkpoint protein programmed death ligand 1 (PD-L1) in response to inflammatory cytokines, primarily interferons.¹²⁶ PD-L1 typically signals to cells via programmed death 1 (PD-1) to reduce activation and induce apoptosis.^{127,128} While many CNS cells, including microglia and even neurons, express PD-1, it is highly enriched on immune cells, making them particularly susceptible to exhaustion.¹²⁶⁻¹²⁹ In summary, astrocytes act not only as a physical barrier, but also as a trophic and immunologic barrier to limit and control the activation state of immune cells that enter the CNS during neuroinflammation.

5.3 | Targeting astrocytes

Astrocytes have great potential as a therapeutic target for BBB maintenance. Their ability to weaken EC tight junctions while maintaining a barrier could prove beneficial in the development of drugs to bypass traditional CNS barriers. There are many examples of astrocytes guiding various cell types and molecules across the BBB. This principle—if better understood—could be leveraged to advance CNS drug permeability. Similar to pericytes, astrocytes have great influence on the maintenance of EC tight junctions. Elucidating the molecular control that astrocytes have over BBB permeability will inevitably lead to more avenues of drug delivery.

6 | MICROGLIA

Microglia are the resident immune cells of the CNS. As such, their functions are multifaceted. While they share many similarities with peripheral macrophages, they have many distinct characteristics, including their origins. Macrophages are generated in the bone marrow and circulate in the blood, while microglia migrate from the embryonic yolk sac.¹³⁰ In general, microglia are responsible for initial responses to injury and infection, clearance of waste, synaptic pruning and maintenance, as well as providing several trophic factors to other cells of the NVU.¹³⁰ Although it has been long appreciated that microglia is vital to the health of the CNS, surveying along the BBB and responding quickly to breaches, they are a relatively recent addition to the NVU and provide coverage of BBB areas not wrapped by astrocytes.^{29,131} In addition, microglia are known to signal to other members of the NVU and communicate with peripheral immune cells,⁶ attracting and/or activating them within the CNS parenchyma during injury and infection as sentinels of the CNS.¹³²⁻¹³⁵ Activated microglia and peripheral immune cells can secrete a number of factors that modulate BBB integrity (Figure 2). This is discussed in a number of reviews, but as an example, TNF- α and TGF- β can be secreted from reactive microglia under certain conditions which can either increase or decrease BBB integrity, respectively.^{123,136,137} While ablation or depletion of microglia did not result in overt BBB

breakdown,¹³⁸ recently, microglia was described to intricately associate with CNS capillaries and contribute to blood flow regulation and vasodilation.¹³⁹⁻¹⁴¹ Additionally, microglia are incredibly motile and migrate quickly in response to injury, BBB leakage, and/or inflammation.¹⁴² Finally, similar to astrocytes, microglia can upregulate PD-L1 in response to neuroinflammation,^{143,144} providing multiple mechanisms of potential control of the BBB by microglia, although there is still much to learn about the intricacies of microglia and their impact on BBB permeability.

Given their emerging role in the NVU, the mechanisms underlying microglial regulation of the BBB remain incompletely described. Targeting microglia may modify CNS capillary permeability or other NVU cells to temporarily allow access to the CNS, but there is still much to learn about this exciting new potential target.

7 | NEURONS

Neurons are the functional unit of the CNS and most other cells in the CNS support them, either directly or indirectly.²⁷ Despite their indispensable role in the CNS, as a member of the NVU, they do not directly provide a physical barrier, but instead release a number of factors that modulate other NVU cells^{27,145} using primarily their axons and dendrites.⁶ Interestingly, like astrocytes, neurons can induce a BBB-like barrier in neighboring ECs,^{96,97} suggesting neurons influence the formation and maintenance of the BBB *in vivo*.^{146,147} Astrocytes and neurons are highly communicative,^{148,149} providing a potential route of BBB control. Additionally, neural activity can influence the integrity of the BBB,¹⁵⁰⁻¹⁵² although it is unclear if neural activity directly impacts ECs or if neuronal signals are propagated through astrocytes.¹⁵³

Neuroinflammation can alter the activity of neurons, which in turn can lead to alterations in the BBB.^{150,154} As mentioned above, neurons express PD-1, to which astrocytes and microglia can signal via PD-L1 expression.¹²⁷⁻¹²⁹ Neuronal PD-1 signaling does not induce apoptosis, but instead hyperpolarizes the neuronal membrane to inhibit action potential frequency.¹²⁹ This in turn could lead to a reduction in activity-dependent transport across the BBB.¹⁵⁵ Neurons also produce Wnt ligands which reduce BBB permeability by increasing tight junction proteins in ECs.¹⁵⁶ While neurons partially modulate the BBB via activity, they largely rely on communication with other NVU members (Figure 2). While more information is necessary to determine how neurons might control the BBB, neurons represent an unlikely target for BBB permeability modulation given their moderate impact on the BBB and other essential functions.

8 | FUTURE DIRECTIONS

Each cellular component of the NVU has both unique and overlapping roles in maintaining BBB permeability. ECs are the initial barrier to the periphery, reinforced with tight junctions,

specialized to transport material between the periphery and the CNS. Pericytes, and to some extent, microglia, regulate vascular flow and, while they may also have a structural role, pericytes provide critical trophic support to ECs. Astrocytes, and potentially neurons, are critical to the formation of the BBB, aiding in tight junction formation. Microglia and neurons also potentially modulate BBB permeability through complex signaling networks, although this is an emerging field. The intricate dynamics of BBB formation, function, and maintenance has created a literal and figurative barrier when it comes to therapy development for the treatment of chronic neurologic diseases such as glioblastoma, MS, and AD. While it is possible to increase the permeability of the BBB, this typically results in detrimental off target effects. The BBB is disrupted in many neurological diseases, thus additional interventional disruption would likely prove deleterious as unwanted CNS "intruders" including immune cells and large molecules are able to enter the CNS without proper scrutiny, leading to a cascade of osmotic, trophic, and inflammatory dysregulation. Thus, optimizing targeted and temporary entry of therapeutics into the CNS with minimal BBB dysregulation is the holy grail of next-generation neurotherapeutics. Promising modalities include encapsulating drugs of interest in a form that allows them to be transported through the BBB and astrocyte endfeet or creating a transient passage that is quickly repaired. Both of these methods are active areas of research using nanoparticles and ultrasonic disruption of the BBB, respectively.^{12,157,158} Ultimately, a deeper understanding of how each of the NVU components modulate CNS drug accessibility may shed new light on actionable therapeutic modalities.

AUTHOR CONTRIBUTIONS

All authors contributed to the writing, editing, and review of the manuscript.

ACKNOWLEDGEMENTS

We apologize to authors whose work could not be referenced in this review due to scope and space limitations. Graphical illustrations were made using BioRender (<https://biorender.com/>). This work was supported by grants from the CSU CD-CAVS training grant NIH T32 HL150389 (awarded to Brandon C. Smith), NIH K00 NS120365 (awarded to Benjamin C. Shaw), NIH R01 NS119178 (awarded to Jessica L. Williams).

CONFLICT OF INTEREST

The authors declare no competing financial interests.

DATA AVAILABILITY STATEMENT

There are no data in this manuscript.

ORCID

Benjamin C. Shaw  <https://orcid.org/0000-0001-8862-3840>

Jessica L. Williams  <https://orcid.org/0000-0002-9801-9580>

REFERENCES

- Biedl A, Kraus R. Über eine bisher unbekannte toxische Wirkung der Gallensäuren auf das Zentralnervensystem. *Zentralblatt Inn Med.* 1898;19:1185-1200.
- Goldmann EE. *Vitalfärbung am Zentralnervensystem: Beitrag zur Physio-Pathologie des Plexus chorioideus und der Hirnhäute.* Königl. Akademie der Wissenschaften; 1913.
- Dreser H. Ueber den experimentellen Nachweis der Vertiefung und Verlangsamung der Athemzüge nach therapeutischen Heroingaben. *Archiv für Die Gesamte Physiologie Des Menschen Und der Tiere.* 1900;80:86-95. doi:10.1007/BF01661930
- Shirai Y. On the transplantation of the rat sarcoma in adult heterogenous animals. *Jap Med World.* 1921;1:14-15.
- Murphy JB, Sturm E. Conditions determining the transplantability of tissues in the brain. *J Exp Med.* 1923;38:183-197. doi:10.1084/jem.38.2.183
- Segarra M, Aburto MR, Acker-Palmer A. Blood-brain barrier dynamics to maintain brain homeostasis. *Trends Neurosci.* 2021;44:393-405. doi:10.1016/j.tins.2020.12.002
- Salmina AB, Komleva YK, Malinovskaya NA, et al. Blood-brain barrier breakdown in stress and neurodegeneration: biochemical mechanisms and new models for translational research. *Biochemistry (Mosc).* 2021;86:746-760. doi:10.1134/S0006297921060122
- Kolářová H, Ambrůzová B, Švihálková Šindlerová L, Klinke A, Kubala L. Modulation of endothelial glycocalyx structure under inflammatory conditions. *Mediators Inflamm.* 2014;2014:694312. doi:10.1155/2014/694312
- Kutuzov N, Flyvbjerg H, Lauritzen M. Contributions of the glycocalyx, endothelium, and extravascular compartment to the blood-brain barrier. *Proc Natl Acad Sci U S A.* 2018;115:E9429-E9438. doi:10.1073/pnas.1802155115
- Schinkel AH, Wagenaar E, Mol CA, van Deemter L. P-glycoprotein in the blood-brain barrier of mice influences the brain penetration and pharmacological activity of many drugs. *J Clin Invest.* 1996;97:2517-2524. doi:10.1172/JCI118699
- Li J, Zheng M, Shimoni O, et al. Development of novel therapeutics targeting the blood-brain barrier: from barrier to carrier. *Adv Sci (Weinh).* 2021;8:2101090. doi:10.1002/advs.202101090
- Anthony DP, Hegde M, Shetty SS, Rafic T, Mutalik S, Rao BSS. Targeting receptor-ligand chemistry for drug delivery across blood-brain barrier in brain diseases. *Life Sci.* 2021;274:119326. doi:10.1016/j.lfs.2021.119326
- Nicolazzo JA, Charman SA, Charman WN. Methods to assess drug permeability across the blood-brain barrier. *J Pharm Pharmacol.* 2006;58:281-293. doi:10.1211/jpp.58.3.0001
- Liu L, Zhang L, Feng H, et al. Prediction of the blood-brain barrier (BBB) permeability of chemicals based on machine-learning and ensemble methods. *Chem Res Toxicol.* 2021;34:1456-1467. doi:10.1021/acs.chemrestox.0c00343
- Strazielle N, Ghersi-Egea JF. Physiology of blood-brain interfaces in relation to brain disposition of small compounds and macromolecules. *Mol Pharm.* 2013;10:1473-1491. doi:10.1021/mp300518e
- Ando Y, Okada H, Takemura G, et al. Brain-specific ultrastructure of capillary endothelial glycocalyx and its possible contribution for blood brain barrier. *Sci Rep.* 2018;8:17523. doi:10.1038/s41598-018-35976-2
- Zaragozá R. Transport of amino acids across the blood-brain barrier. *Front Physiol.* 2020;11:973. doi:10.3389/fphys.2020.00973
- Pachter JS, de Vries HE, Fabry Z. The blood-brain barrier and its role in immune privilege in the central nervous system. *J Neuropathol Exp Neurol.* 2003;62:593-604. doi:10.1093/jnen/62.6.593

19. Wang W, Dentler WL, Borchardt RT. VEGF increases BMEC monolayer permeability by affecting occludin expression and tight junction assembly. *Am J Physiol Heart Circ Physiol*. 2001;280:H434-H440. doi:10.1152/ajpheart.2001.280.1.H434
20. Liu W-Y, Wang Z-B, Zhang L-C, Wei X, Li L. Tight junction in blood-brain barrier: an overview of structure, regulation, and regulator substances. *CNS Neurosci Ther*. 2012;18:609-615. doi:10.1111/j.1755-5949.2012.00340.x
21. Bohannon DG, Long D, Kim W-K. Understanding the heterogeneity of human pericyte subsets in blood-brain barrier homeostasis and neurological diseases. *Cell*. 2021;10:890. doi:10.3390/cells10040890
22. Rustenhoven J, Aalderink M, Scotter EL, et al. TGF-beta1 regulates human brain pericyte inflammatory processes involved in neurovasculature function. *J Neuroinflammation*. 2016;13:37. doi:10.1186/s12974-016-0503-0
23. Zimmermann KW. Der feinere Bau der Blutcapillaren. *Z Anat Entwicklungsgesch*. 1923;68:29-109. doi:10.1007/BF02593544
24. Hamilton N, Attwell D, Hall C. Pericyte-mediated regulation of capillary diameter: a component of neurovascular coupling in health and disease. *Front Neuroenergetics*. 2010;2:5. doi:10.3389/fnene.2010.00005
25. Heithoff BP, George KK, Phares AN, Zuidhoek IA, Munoz-Ballester C, Robel S. Astrocytes are necessary for blood-brain barrier maintenance in the adult mouse brain. *Glia*. 2021;69:436-472. doi:10.1002/glia.23908
26. Rubin LL, Hall DE, Porter S, et al. A cell culture model of the blood-brain barrier. *J Cell Biol*. 1991;115:1725-1735. doi:10.1083/jcb.115.6.1725
27. Bear M, Connors B, Paradiso MA. *Neuroscience: Exploring the Brain, Enhanced Edition*. Jones & Bartlett Learning; 2020.
28. Ueno M, Akiguchi I, Hosokawa M, Kotani H, Kanenishi K, Sakamoto H. Blood-brain barrier permeability in the periventricular areas of the normal mouse brain. *Acta Neuropathol*. 2000;99:385-392. doi:10.1007/s004010051140
29. Kisler K, Nikolakopoulou AM, Zlokovic BV. Microglia have a grip on brain microvasculature. *Nat Commun*. 2021;12:5290. doi:10.1038/s41467-021-25595-3
30. Cauli B, Tong XK, Rancillac A, et al. Cortical GABA interneurons in neurovascular coupling: relays for subcortical vasoactive pathways. *J Neurosci*. 2004;24:8940-8949. doi:10.1523/jneurosci.3065-04.2004
31. McConnell HL, Kersch CN, Woltjer RL, Neuwelt EA. The translational significance of the neurovascular unit. *J Biol Chem*. 2017;292:762-770. doi:10.1074/jbc.R116.760215
32. Bagchi S, Chhibber T, Lahooti B, Verma A, Borse V, Jayant RD. In vitro blood-brain barrier models for drug screening and permeation studies: an overview. *Drug Des Devel Ther*. 2019;13:3591-3605. doi:10.2147/DDDT.S218708
33. Wong A et al. The blood-brain barrier: an engineering perspective. *Front Neuroeng*. 2013;6:7. doi:10.3389/fneng.2013.00007
34. Jaffe EA. Cell biology of endothelial cells. *Hum Pathol*. 1987;18:234-239. doi:10.1016/S0046-8177(87)80005-9
35. Kramer RH, Bensch KG, Davison PM, Karasek MA. Basal lamina formation by cultured microvascular endothelial cells. *J Cell Biol*. 1984;99:692-698. doi:10.1083/jcb.99.2.692
36. Wolburg H, Noell S, Mack A, Wolburg-Buchholz K, Fallier-Becker P. Brain endothelial cells and the glio-vascular complex. *Cell Tissue Res*. 2009;335:75-96. doi:10.1007/s00441-008-0658-9
37. Ruck T, Bittner S, Meuth SG. Blood-brain barrier modeling: challenges and perspectives. *Neural Regen Res*. 2015;10:889-891. doi:10.4103/1673-5374.158342
38. Shayan G, Choi YS, Shusta EV, Shuler ML, Lee KH. Murine in vitro model of the blood-brain barrier for evaluating drug transport. *Eur J Pharm Sci*. 2011;42:148-155. doi:10.1016/j.ejps.2010.11.005
39. Greene C, Hanley N, Campbell M. Claudin-5: gatekeeper of neuro-logical function. *Fluids Barriers CNS*. 2019;16:3. doi:10.1186/s12987-019-0123-z
40. Branca JJV, Maresca M, Morucci G, et al. Effects of cadmium on ZO-1 tight junction integrity of the blood brain barrier. *Int J Mol Sci*. 2019;20:6010.
41. Lochhead JJ, Yang J, Ronaldson PT, Davis TP. Structure, function, and regulation of the blood-brain barrier tight junction in central nervous system disorders. *Front Physiol*. 2020;11:914. doi:10.3389/fphys.2020.00914
42. Nitta T, Hata M, Gotoh S, et al. Size-selective loosening of the blood-brain barrier in claudin-5-deficient mice. *J Cell Biol*. 2003;161:653-660. doi:10.1083/jcb.200302070
43. Enerson BE, Drewes LR. The rat blood-brain barrier transcriptome. *J Cereb Blood Flow Metab*. 2006;26:959-973. doi:10.1038/sj.jcbfm.9600249
44. Fabry Z, Sandor M, Gajewski TF, et al. Differential activation of Th1 and Th2 CD4+ cells by murine brain microvessel endothelial cells and smooth muscle/pericytes. *J Immunol*. 1993;151:38-47.
45. Castro Dias M, Odriozola Quesada A, Soldati S, et al. Brain endothelial tricellular junctions as novel sites for T cell diapedesis across the blood-brain barrier. *J Cell Sci*. 2021;134:jcs253880. doi:10.1242/jcs.253880
46. Greenwood J, Wang Y, Calder VL. Lymphocyte adhesion and transendothelial migration in the central nervous system: the role of LFA-1, ICAM-1, VLA-4 and VCAM-1. *Off. Immunology*. 1995;86:408-415.
47. Cinamon G, Shinder V, Alon R. Shear forces promote lymphocyte migration across vascular endothelium bearing apical chemokines. *Nat Immunol*. 2001;2:515-522. doi:10.1038/88710
48. Ronaldson PT, Demarco KM, Sanchez-Covarrubias L, Solinsky CM, Davis TP. Transforming growth factor-beta signaling alters substrate permeability and tight junction protein expression at the blood-brain barrier during inflammatory pain. *J Cereb Blood Flow Metab*. 2009;29:1084-1098. doi:10.1038/jcbfm.2009.32
49. Majerova P, Michalicova A, Cente M, et al. Trafficking of immune cells across the blood-brain barrier is modulated by neurofibrillary pathology in tauopathies. *PLoS One*. 2019;14:e0217216. doi:10.1371/journal.pone.0217216
50. Arima M, Nakao S, Yamaguchi M, et al. Claudin-5 redistribution induced by inflammation leads to anti-VEGF-resistant diabetic macular edema. *Diabetes*. 2020;69:981-999. doi:10.2337/db19-1121
51. Wang J, Zhang C, Zhu J, Ding J, Chen Y, Han X. Blood-brain barrier disruption and inflammation reaction in mice after chronic exposure to microcystin-LR. *Sci Total Environ*. 2019;689:662-678. doi:10.1016/j.scitotenv.2019.06.387
52. Xia H, Banks WA, Niehoff ML, Morley JE. Effect of LPS on the permeability of the blood-brain barrier to insulin. *Brain Res*. 2001;896:36-42. doi:10.1016/S0006-8993(00)03247-9
53. Hansson E. Actin filament reorganization in astrocyte networks is a key functional step in neuroinflammation resulting in persistent pain: novel findings on network restoration. *Neurochem Res*. 2015;40:372-379. doi:10.1007/s11064-014-1363-6
54. Descamps L, Coisne C, Dehouck B, Cecchelli R, Torpier G. Protective effect of glial cells against lipopolysaccharide-mediated blood-brain barrier injury. *Glia*. 2003;42:46-58. doi:10.1002/glia.10205
55. Hoshi Y, Uchida Y, Tachikawa M, Ohtsuki S, Terasaki T. Actin filament-associated protein 1 (AFAP-1) is a key mediator in inflammatory signaling-induced rapid attenuation of intrinsic P-gp function in human brain capillary endothelial cells. *J Neurochem*. 2017;141:247-262. doi:10.1111/jnc.13960
56. Van Skike CE, Jahrling JB, Olson AB, et al. Inhibition of mTOR protects the blood-brain barrier in models of Alzheimer's disease and vascular cognitive impairment. *Am J Physiol Heart Circ Physiol*. 2018;314:H693-H703. doi:10.1152/ajpheart.00570.2017

57. Kurz C, Walker L, Rauchmann B-S, Pernecky R. Dysfunction of the blood-brain barrier in Alzheimer's disease: evidence from human studies. *Neuropathol Appl Neurobiol.* 2022;48:e12782. doi:10.1111/nan.12782
58. Nishihara H, Perriot S, Gastfriend BD, et al. Intrinsic blood-brain barrier dysfunction contributes to multiple sclerosis pathogenesis. *Brain.* 2022;awac019. doi:10.1093/brain/awac019. Online ahead of print.
59. Pan W, P. Stone K, Hsuchou H, K. Manda V, Zhang Y, J. Kastin A. Cytokine signaling modulates blood-brain barrier function. *Curr Pharm Des.* 2011;17:3729-3740. doi:10.2174/138161211798220918
60. Zamudio F, Loon AR, Smeltzer S, et al. TDP-43 mediated blood-brain barrier permeability and leukocyte infiltration promote neurodegeneration in a low-grade systemic inflammation mouse model. *J Neuroinflammation.* 2020;17:283. doi:10.1186/s12974-020-01952-9
61. Saand AR, Yu F, Chen J, Chou SHY. Systemic inflammation in hemorrhagic strokes - a novel neurological sign and therapeutic target? *J Cereb Blood Flow Metab.* 2019;39:959-988. doi:10.1177/0271678X19841443
62. Villaseñor R, Lampe J, Schwaninger M, Collin L. Intracellular transport and regulation of transcytosis across the blood-brain barrier. *Cell Mol Life Sci.* 2019;76:1081-1092. doi:10.1007/s00018-018-2982-x
63. Grubb S, Lauritzen M, Aalkjær C. Brain capillary pericytes and neurovascular coupling. *Comp Biochem Physiol A Mol Integr Physiol.* 2021;254:110893. doi:10.1016/j.cbpa.2020.110893
64. Kim S, Lee S, Lim J, et al. Human bone marrow-derived mesenchymal stem cells play a role as a vascular pericyte in the reconstruction of human BBB on the angiogenesis microfluidic chip. *Biomaterials.* 2021;279:121210. doi:10.1016/j.biomaterials.2021.121210
65. Xu L, Nirwane A, Yao Y. Basement membrane and blood-brain barrier. *Stroke Vasc Neurol.* 2018;4:78-82. doi:10.1136/svn-2018-000198
66. Peppiatt CM, Howarth C, Mobbs P, Attwell D. Bidirectional control of CNS capillary diameter by pericytes. *Nature.* 2006;443:700-704. doi:10.1038/nature05193
67. Sá-Pereira I, Brites D, Brito MA. Neurovascular unit: a focus on pericytes. *Mol Neurobiol.* 2012;45:327-347. doi:10.1007/s12035-012-8244-2
68. Ferro MP, Heilshorn SC, Owens RM. Materials for blood brain barrier modeling in vitro. *Mater Sci Eng R Rep.* 2020;140:100522. doi:10.1016/j.mser.2019.100522
69. Török O, Schreiner B, Schaffenerath J, et al. Pericytes regulate vascular immune homeostasis in the CNS. *Proc Natl Acad Sci U S A.* 2021;118:e2016587118. doi:10.1073/pnas.2016587118
70. Mäe MA, He L, Nordling S, et al. Single-cell analysis of blood-brain barrier response to pericyte loss. *Circ Res.* 2021;128:e46-e62. doi:10.1161/CIRCRESAHA.120.317473
71. Dubrac A, Künzel SE, Künzel SH, et al. NCK-dependent pericyte migration promotes pathological neovascularization in ischemic retinopathy. *Nat Commun.* 2018;9:3463. doi:10.1038/s41467-018-05926-7
72. Warmke N, Platt F, Bruns AF, et al. Pericyte insulin receptors modulate retinal vascular remodeling and endothelial angiotensin signaling. *Endocrinology.* 2021;162:bqab182. doi:10.1210/endo/bqab182
73. Vazquez-Liebanas E, Nahar K, Bertuzzi G, Keller A, Betsholtz C, Mäe MA. Adult-induced genetic ablation distinguishes PDGFB roles in blood-brain barrier maintenance and development. *J Cereb Blood Flow Metab.* 2022;42:264-279. doi:10.1177/0271678X211056395
74. Rudziak P, Ellis CG, Kowalewska PM. Role and molecular mechanisms of pericytes in regulation of leukocyte diapedesis in inflamed tissues. *Mediators Inflamm.* 2019;2019:4123605. doi:10.1155/2019/4123605
75. Nag TC, Gorla S, Kumari C, Roy TS. Aging of the human choriocapillaris: evidence that early pericyte damage can trigger endothelial changes. *Exp Eye Res.* 2021;212:108771. doi:10.1016/j.exer.2021.108771
76. Kisler K, Nikolakopoulou AM, Sweeney MD, Lazic D, Zhao Z, Zlokovic BV. Acute ablation of cortical pericytes leads to rapid neurovascular uncoupling. *Front Cell Neurosci.* 2020;14:27. doi:10.3389/fncel.2020.00027
77. Nikolakopoulou AM, Montagne A, Kisler K, et al. Pericyte loss leads to circulatory failure and pleiotrophin depletion causing neuron loss. *Nat Neurosci.* 2019;22:1089-1098. doi:10.1038/s41593-019-0434-z
78. Gerhardt H, Wolburg H, Redies C. N-cadherin mediates pericytic-endothelial interaction during brain angiogenesis in the chicken. *Dev Dyn.* 2000;218:472-479.
79. Schofield CL, Rodrigo-Navarro A, Dalby MJ, Van Agtmael T, Salmeron-Sanchez M. Biochemical- and biophysical-induced Barrierogenesis in the blood-brain barrier: a review of Barrierogenic factors for use in in vitro models. *Adv NanoBiomed Res.* 2021;1:2000068. doi:10.1002/anbr.202000068
80. Tillet E, Vittet D, Féraud O, Moore R, Kemler R, Huber P. N-cadherin deficiency impairs pericyte recruitment, and not endothelial differentiation or sprouting, in embryonic stem cell-derived angiogenesis. *Exp Cell Res.* 2005;310:392-400. doi:10.1016/j.yexcr.2005.08.021
81. Kuo Y-C, Lee C-L, Rajesh R. Regulation of human brain vascular pericytes and human astrocytes in a blood-brain barrier model using human brain microvascular endothelial cells: expression of TGF- β 1, VEGF, MMP-9 and P-gp. *J Taiwan Inst Chem Eng.* 2018;86:9-17. doi:10.1016/j.jtice.2018.03.003
82. Zheng Z, Chopp M, Chen J. Multifaceted roles of pericytes in central nervous system homeostasis and disease. *J Cereb Blood Flow Metab.* 2020;40:1381-1401. doi:10.1177/0271678X20911331
83. Laredo F, Plebanski J, Tedeschi A. Pericytes: problems and promises for CNS repair. *Front Cell Neurosci.* 2019;13:546. doi:10.3389/fncel.2019.00546
84. Geranmayeh MH, Rahbarghazi R, Farhoudi M. Targeting pericytes for neurovascular regeneration. *Cell Commun Signal.* 2019;17:26. doi:10.1186/s12964-019-0340-8
85. Bohannon DG, Okhravi HR, Kim J, Kuroda MJ, Didier ES, Kim WK. A subtype of cerebrovascular pericytes is associated with blood-brain barrier disruption that develops during normal aging and simian immunodeficiency virus infection. *Neurobiol Aging.* 2020;96:128-136. doi:10.1016/j.neurobiolaging.2020.08.006
86. Escartin C, Galea E, Lakatos A, et al. Reactive astrocyte nomenclature, definitions, and future directions. *Nat Neurosci.* 2021;24:312-325. doi:10.1038/s41593-020-00783-4
87. Gautam J, Cao Y, Yao Y. Pericytic laminin maintains blood-brain barrier integrity in an age-dependent manner. *Transl Stroke Res.* 2020;11:228-242. doi:10.1007/s12975-019-00709-8
88. Yao Y, Chen Z-L, Norris EH, Strickland S. Astrocytic laminin regulates pericyte differentiation and maintains blood brain barrier integrity. *Nat Commun.* 2014;5:3413. doi:10.1038/ncomms4413
89. Yamanaka G, Takata F, Kataoka Y, et al. The neuroinflammatory role of pericytes in epilepsy. *Biomedicine.* 2021;9:759.
90. Miller SJ. Astrocyte heterogeneity in the adult central nervous system. *Front Cell Neurosci.* 2018;12:401. doi:10.3389/fncel.2018.00401
91. Mathiisen TM, Lehre KP, Danbolt NC, Ottersen OP. The perivascular astroglial sheath provides a complete covering of the brain microvessels: an electron microscopic 3D reconstruction. *Glia.* 2010;58:1094-1103. doi:10.1002/glia.20990
92. Wolburg-Buchholz K, Mack AF, Steiner E, Pfeiffer F, Engelhardt B, Wolburg H. Loss of astrocyte polarity marks blood-brain

- barrier impairment during experimental autoimmune encephalomyelitis. *Acta Neuropathol.* 2009;118:219-233. doi:10.1007/s00401-009-0558-4
93. Kubotera H, Ikeshima-Kataoka H, Hatashita Y, Allegra Mascaro AL, Pavone FS, Inoue T. Astrocytic endfeet re-cover blood vessels after removal by laser ablation. *Sci Rep.* 2019;9:1263. doi:10.1038/s41598-018-37419-4
 94. Higashi K, Fujita A, Inanobe A, et al. An inwardly rectifying K⁺ channel, Kir4.1, expressed in astrocytes surrounds synapses and blood vessels in brain. *Am J Physiol Cell Physiol.* 2001;281:C922-C931. doi:10.1152/ajpcell.2001.281.3.C922
 95. Warth A, Mittelbronn M, Wolburg H. Redistribution of the water channel protein aquaporin-4 and the K⁺ channel protein Kir4.1 differs in low- and high-grade human brain tumors. *Acta Neuropathol.* 2005;109:418-426. doi:10.1007/s00401-005-0984-x
 96. Kaiser MA, Sajja RK, Prasad S, Abhyankar VV, Liles T, Cucullo L. New experimental models of the blood-brain barrier for CNS drug discovery. *Expert Opin Drug Discovery.* 2017;12:89-103. doi:10.1080/17460441.2017.1253676
 97. Ponio JB-D, el-Ayoubi F, Glacial F, et al. Instruction of circulating endothelial progenitors in vitro towards specialized blood-brain barrier and arterial phenotypes. *PLoS One.* 2014;9:e84179. doi:10.1371/journal.pone.0084179
 98. Huang X, Su Y, Wang N, et al. Astroglial connexins in neurodegenerative diseases. *Front Mol Neurosci.* 2021;14:657514. doi:10.3389/fnmol.2021.657514
 99. Van Campenhout R, Gomes AR, TWM DG, Muyldermans S, Devoogdt N, Vinken M. Mechanisms underlying connexin hemichannel activation in disease. *Int J Mol Sci.* 2021;22:3503.
 100. Peng B, Xu C, Wang S, Zhang Y, Li W. The role of connexin hemichannels in inflammatory diseases. *Biology.* 2022;11:237.
 101. Rouach N, Koulakoff A, Abudara V, Willecke K, Giaume C. Astroglial metabolic networks sustain hippocampal synaptic transmission. *Science.* 2008;322:1551-1555. doi:10.1126/science.1164022
 102. Xing L, Yang T, Cui S, Chen G. Connexin hemichannels in astrocytes: role in CNS disorders. *Front Mol Neurosci.* 2019;12:23. doi:10.3389/fnmol.2019.00023
 103. Spampinato SF, Merlo S, Sano Y, Kanda T, Sortino MA. Astrocytes contribute to A β -induced blood-brain barrier damage through activation of endothelial MMP9. *J Neurochem.* 2017;142:464-477. doi:10.1111/jnc.14068
 104. Song S, Huang H, Guan X, et al. Activation of endothelial Wnt/ β -catenin signaling by protective astrocytes repairs BBB damage in ischemic stroke. *Prog Neurobiol.* 2021;199:101963. doi:10.1016/j.pneurobio.2020.101963
 105. Han RT, Kim RD, Molofsky AV, Liddel SA. Astrocyte-immune cell interactions in physiology and pathology. *Immunity.* 2021;54:211-224. doi:10.1016/j.immuni.2021.01.013
 106. Fu J, Li L, Huo D, et al. Astrocyte-derived TGF β 1 facilitates blood-brain barrier function via non-canonical hedgehog signaling in brain microvascular endothelial cells. *Brain Sci.* 2021;11:77.
 107. Feng S, Zou L, Wang H, He R, Liu K, Zhu H. RhoA/ROCK-2 pathway inhibition and tight junction protein upregulation by Catalpol suppresses Lipopolysaccharide-induced disruption of blood-brain barrier permeability. *Molecules.* 2018;23:2371.
 108. Liu C-C, Yamazaki Y, Heckman MG, et al. Tau and apolipoprotein E modulate cerebrovascular tight junction integrity independent of cerebral amyloid angiopathy in Alzheimer's disease. *Alzheimers Dement.* 2020;16:1372-1383. doi:10.1002/alz.12104
 109. Jackson RJ, Meltzer JC, Nguyen H, et al. APOE4 derived from astrocytes leads to blood-brain barrier impairment. *Brain.* 2021;awab478. doi:10.1093/brain/awab478. Online ahead of print.
 110. Guérit S, Fidan E, Macas J, et al. Astrocyte-derived Wnt growth factors are required for endothelial blood-brain barrier maintenance. *Prog Neurobiol.* 2021;199:101937. doi:10.1016/j.pneurobio.2020.101937
 111. Montagne A, Nation DA, Sagare AP, et al. APOE4 leads to blood-brain barrier dysfunction predicting cognitive decline. *Nature.* 2020;581:71-76. doi:10.1038/s41586-020-2247-3
 112. Yamazaki Y, Shinohara M, Yamazaki A, et al. ApoE (apolipoprotein E) in brain pericytes regulates endothelial function in an isoform-dependent manner by modulating basement membrane components. *Arterioscler Thromb Vasc Biol.* 2020;40:128-144. doi:10.1161/ATVBAHA.119.313169
 113. Zhang ZG, Zhang L, Croll SD, Chopp M. Angiotensin-1 reduces cerebral blood vessel leakage and ischemic lesion volume after focal cerebral embolic ischemia in mice. *Neuroscience.* 2002;113:683-687. doi:10.1016/S0306-4522(02)00175-6
 114. Venkat P, Yan T, Chopp M, et al. Angiotensin-1 mimetic peptide promotes neuroprotection after stroke in type 1 diabetic rats. *Cell Transplant.* 2018;27:1744-1752. doi:10.1177/0963689718791568
 115. Giovannoni F, Quintana FJ. The role of astrocytes in CNS inflammation. *Trends Immunol.* 2020;41:805-819. doi:10.1016/j.it.2020.07.007
 116. Smith BC, Sinyuk M, Jenkins JE, Psenicka MW, Williams JL. The impact of regional astrocyte interferon- γ signaling during chronic autoimmunity: a novel role for the immunoproteasome. *J Neuroinflammation.* 2020;17:184. doi:10.1186/s12974-020-01861-x
 117. Horng S, Therattil A, Moyon S, et al. Astrocytic tight junctions control inflammatory CNS lesion pathogenesis. *J Clin Invest.* 2017;127:3136-3151. doi:10.1172/JCI91301
 118. Neuhaus W, Gaiser F, Mahringer A, Franz J, Riethmüller C, Förster C. The pivotal role of astrocytes in an in vitro stroke model of the blood-brain barrier. *Front Cell Neurosci.* 2014;8:352. doi:10.3389/fncel.2014.00352
 119. McCandless EE, Wang Q, Woerner BM, Harper JM, Klein RS. CXCL12 limits inflammation by localizing mononuclear infiltrates to the perivascular space during experimental autoimmune encephalomyelitis. *J Immunol.* 2006;177:8053-8064. doi:10.4049/jimmunol.177.11.8053
 120. Williams JL, Manivasagam S, Smith BC, et al. Astrocyte-T cell crosstalk regulates region-specific neuroinflammation. *Glia.* 2020;68:1361-1374. doi:10.1002/glia.23783
 121. Michinaga S, Koyama Y. Dual roles of astrocyte-derived factors in regulation of blood-brain barrier function after brain damage. *Int J Mol Sci.* 2019;20:571.
 122. Kim GW, Gasche Y, Grzeschik S, Copin JC, Maier CM, Chan PH. Neurodegeneration in striatum induced by the mitochondrial toxin 3-nitropropionic acid: role of matrix metalloproteinase-9 in early blood-brain barrier disruption? *J Neurosci.* 2003;23:8733-8742.
 123. Psenicka MW, Smith BC, Tinkey RA, Williams JL. Connecting neuroinflammation and neurodegeneration in multiple sclerosis: are oligodendrocyte precursor cells a nexus of disease? *Front Cell Neurosci.* 2021;15:654284. doi:10.3389/fncel.2021.654284
 124. X. Gao, W. Li, F. Syed, F. Yuan, P. Li, Q. Yu. PD-L1-expressing astrocytes act as a gate-keeper for neuroinflammation in the central nervous system of mice with traumatic brain injury. *bioRxiv.* 2021. doi:10.1101/2021.11.04.467368
 125. Gao X, Li W, Syed F, Yuan F, Li P, Yu Q. PD-L1 signaling in reactive astrocytes counteracts neuroinflammation and ameliorates neuronal damage after traumatic brain injury. *J Neuroinflammation.* 2022;19:43. doi:10.1186/s12974-022-02398-x
 126. Kummer MP, Ising C, Kummer C, et al. Microglial PD-1 stimulation by astrocytic PD-L1 suppresses neuroinflammation and Alzheimer's disease pathology. *EMBO J.* 2021;40:e108662. doi:10.15252/embj.2021108662
 127. Keir ME, Butte MJ, Freeman GJ, Sharpe AH. PD-1 and its ligands in tolerance and immunity. *Annu Rev Immunol.* 2008;26:677-704. doi:10.1146/annurev.immunol.26.021607.090331

128. Latchman Y, Wood CR, Chernova T, et al. PD-L2 is a second ligand for PD-1 and inhibits T cell activation. *Nat Immunol.* 2001;2:261-268. doi:10.1038/85330
129. Chen G, Kim YH, Li H, et al. PD-L1 inhibits acute and chronic pain by suppressing nociceptive neuron activity via PD-1. *Nat Neurosci.* 2017;20:917-926. doi:10.1038/nn.4571
130. Li Q, Barres BA. Microglia and macrophages in brain homeostasis and disease. *Nat Rev Immunol.* 2018;18:225-242. doi:10.1038/nri.2017.125
131. Mondo E, Becker SC, Kautzman AG, et al. A developmental analysis of Juxtavascular microglia dynamics and interactions with the vasculature. *J Neurosci.* 2020;40:6503-6521. doi:10.1523/jneurosci.3006-19.2020
132. Shigemoto-Mogami Y, Hoshikawa K, Sato K. Activated microglia disrupt the blood-brain barrier and induce chemokines and cytokines in a rat in vitro model. *Front Cell Neurosci.* 2018;12:494. doi:10.3389/fncel.2018.00494
133. da Fonseca ACC, Matias D, Garcia C, et al. The impact of microglial activation on blood-brain barrier in brain diseases. *Front Cell Neurosci.* 2014;8:362. doi:10.3389/fncel.2014.00362
134. Halder SK, Milner R. Mild hypoxia triggers transient blood-brain barrier disruption: a fundamental protective role for microglia. *Acta Neuropathol Commun.* 2020;8:175. doi:10.1186/s40478-020-01051-z
135. Ulvestad E, Williams K, Vedeler C, et al. Reactive microglia in multiple sclerosis lesions have an increased expression of receptors for the fc part of IgG. *J Neurol Sci.* 1994;121:125-131. doi:10.1016/0022-510X(94)90340-9
136. Chen A-Q, Fang Z, Chen XL, et al. Microglia-derived TNF- α mediates endothelial necroptosis aggravating blood brain-barrier disruption after ischemic stroke. *Cell Death Dis.* 2019;10:487. doi:10.1038/s41419-019-1716-9
137. Kuroda E, Nishimura K, Kawanishi S, et al. Mouse bone marrow-derived microglia-like cells secrete transforming growth factor- β 1 and promote microglial A β phagocytosis and reduction of brain A β . *Neuroscience.* 2020;438:217-228. doi:10.1016/j.neuroscience.2020.05.004
138. Kerkhofs D, van Hagen BT, Milanova IV, et al. Pharmacological depletion of microglia and perivascular macrophages prevents vascular cognitive impairment in ang II-induced hypertension. *Theranostics.* 2020;10:9512-9527. doi:10.7150/thno.44394
139. Bisht K, Okojie KA, Sharma K, et al. Capillary-associated microglia regulate vascular structure and function through PANX1-P2RY12 coupling in mice. *Nat Commun.* 2021;12:5289. doi:10.1038/s41467-021-25590-8
140. Császár E, Lénárt N, Cserép C, et al. Microglia modulate blood flow, neurovascular coupling, and hypoperfusion via purinergic actions. *J Exp Med.* 2022;219:e20211071. doi:10.1084/jem.20211071
141. Halder SK, Milner R. A critical role for microglia in maintaining vascular integrity in the hypoxic spinal cord. *Proc Natl Acad Sci U S A.* 2019;116:26029-26037. doi:10.1073/pnas.1912178116
142. Davalos D, Kyu Ryu J, Merlini M, et al. Fibrinogen-induced perivascular microglial clustering is required for the development of axonal damage in neuroinflammation. *Nat Commun.* 2012;3:1227. doi:10.1038/ncomms2230
143. Chen Q, Xu L, du T, et al. Enhanced expression of PD-L1 on microglia after surgical brain injury exerts self-protection from inflammation and promotes neurological repair. *Neurochem Res.* 2019;44:2470-2481. doi:10.1007/s11064-019-02864-8
144. Wang L-L, Li ZH, Hu XH, Muyayalo KP, Zhang YH, Liao AH. The roles of the PD-1/PD-L1 pathway at immunologically privileged sites. *Am J Reprod Immunol.* 2017;78:e12710. doi:10.1111/aji.12710
145. Zhang Y, Chen K, Sloan SA, et al. An RNA-sequencing transcriptome and splicing database of glia, neurons, and vascular cells of the cerebral cortex. *J Neurosci.* 2014;34:11929-11947. doi:10.1523/jneurosci.1860-14.2014
146. Tontsch U, Bauer H-C. Glial cells and neurons induce blood-brain barrier related enzymes in cultured cerebral endothelial cells. *Brain Res.* 1991;539:247-253. doi:10.1016/0006-8993(91)91628-E
147. Stanness KA, Neumaier JF, Sexton TJ, et al. A new model of the blood-brain barrier: co-culture of neuronal, endothelial and glial cells under dynamic conditions. *Neuroreport.* 1999;10:3725-3731. doi:10.1097/00001756-199912160-00001
148. Barreto GE, Gonzalez J, Torres Y, Morales L. Astrocytic-neuronal crosstalk: implications for neuroprotection from brain injury. *Neurosci Res.* 2011;71:107-113. doi:10.1016/j.neures.2011.06.004
149. Mulica P, Grünewald A, Pereira SL. Astrocyte-neuron metabolic crosstalk in neurodegeneration: a mitochondrial perspective. *Front Endocrinol (Lausanne).* 2021;12:668517. doi:10.3389/fendo.2021.668517
150. Kaplan L, Chow BW, Gu C. Neuronal regulation of the blood-brain barrier and neurovascular coupling. *Nat Rev Neurosci.* 2020;21:416-432. doi:10.1038/s41583-020-0322-2
151. Yang G, Zhang Y, Ross ME, Iadecola C. Attenuation of activity-induced increases in cerebellar blood flow in mice lacking neuronal nitric oxide synthase. *Am J Physiol Heart Circ Physiol.* 2003;285:H298-H304. doi:10.1152/ajpheart.00043.2003
152. Hogan-Cann AD, Lu P, Anderson CM. Endothelial NMDA receptors mediate activity-dependent brain hemodynamic responses in mice. *Proc Natl Acad Sci U S A.* 2019;116:10229-10231. doi:10.1073/pnas.1902647116
153. Lia A, Henriques VJ, Zonta M, et al. Calcium signals in astrocyte microdomains, a decade of great advances. *Front Cell Neurosci.* 2021;15:673433. doi:10.3389/fncel.2021.673433
154. Semmler A, Hermann S, Mormann F, et al. Sepsis causes neuroinflammation and concomitant decrease of cerebral metabolism. *J Neuroinflammation.* 2008;5:38. doi:10.1186/1742-2094-5-38
155. Pulido RS, Munji RN, Chan TC, et al. Neuronal activity regulates blood-brain barrier efflux transport through endothelial circadian genes. *Neuron.* 2020;108:937-952. doi:10.1016/j.neuron.2020.09.002
156. Salinas PC, Zou Y. Wnt signaling in neural circuit assembly. *Annu Rev Neurosci.* 2008;31:339-358.
157. Gasca-Salas C, Fernández-Rodríguez B, Pineda-Pardo JA, et al. Blood-brain barrier opening with focused ultrasound in Parkinson's disease dementia. *Nat Commun.* 2021;12:779. doi:10.1038/s41467-021-21022-9
158. Zhang W, Mehta A, Tong Z, Esser L, Voelcker NH. Development of polymeric nanoparticles for blood-brain barrier transfer—strategies and challenges. *Adv Sci.* 2021;8:2003937. doi:10.1002/adv.202003937

How to cite this article: Smith BC, Tinkey RA, Shaw BC, Williams JL. Targetability of the neurovascular unit in inflammatory diseases of the central nervous system. *Immunol Rev.* 2022;311:39-49. doi: [10.1111/imr.13121](https://doi.org/10.1111/imr.13121)