

Unexpected Presence of Defective *glpR* Alleles in Various Strains of *Escherichia coli*

C. KAY HOLTMAN, RICHARD THURLKILL, AND DONALD W. PETTIGREW*

Department of Biochemistry and Biophysics and Program in Microbial Genetics and Genomics, Texas A&M University, College Station, Texas 77843-2128

Received 6 September 2000/Accepted 14 November 2000

Alleles of *glpR* associated with the same $GlpR^-$ phenotype produce substitutions in different conserved portions of the glycerol 3-phosphate repressor which are not part of the helix-turn-helix motif. Analysis of the effects on growth and enzyme expression show that glucose repression of glycerol utilization is not dependent on a functional repressor.

In the course of strain construction for studies of allosteric regulatory properties of glycerol kinase (EC 2.7.1.30; ATP-glycerol 3-phosphotransferase), we identified a novel allele for the glycerol 3-phosphate repressor. It confers a $GlpR^-$ phenotype that is indistinguishable from that seen for the *glpR2* allele, which contains an undetermined defect (5). We report here that the sequences of these two alleles are different. Phenotypes associated with these alleles reveal that a functional glycerol 3-phosphate repressor is not required for glucose inhibition of glycerol utilization.

Strain KH11 was constructed to serve as a prototrophic strain with a wild-type glycerol kinase. Characterization of KH11 showed high levels of glycerol kinase enzymatic activity when cells are grown in Luria-Bertani (LB) medium, i.e., non-inducing conditions (Table 1). The specific activities of glycerol kinase were determined in extracts of cells that were grown overnight in LB medium at 37°C, collected by centrifugation, and disrupted by sonication after being resuspended in 0.1 M triethanolamine-HCl, 2 mM glycerol, and 2 mM β -mercaptoethanol (pH 7.0). Glycerol kinase activities were determined by using an ADP-coupled continuous spectrophotometric assay at pH 7.0 and 25°C with 2 mM glycerol and 2.5 mM ATP (8). Protein concentrations were determined by using the Bio-Rad assay with bovine serum albumin as the standard. The glycerol kinase activity of strain KH11 is much higher than that of strains MC4100, MG1655, and MM130. High expression of glycerol kinase after growth in rich medium was reported for Lin strain 43 (11) and is verified as shown in Table 1. The phenotype has been termed constitutive and is associated with an undetermined defect in the *glpR2* allele (5). Table 1 shows also that the same specific activity is obtained for strains KH11, MC4100, and MG1655 after growth under normal induction conditions in minimal glycerol medium. The absence of further induction in strain KH11 under these conditions indicates that the expression is constitutive rather than an elevation of the basal level. However, the specific activity after the growth of strain KH11 in LB medium is about twofold higher, indicating that additional induction is possible in rich medium.

None of the *Escherichia coli* strains that were used in construction of KH11 were known to contain defective *glpR* alleles. Strain DG1, from which strain KH11 was derived, was constructed to provide a $GlpK^-$ background (6). DG1 is a derivative of TB1, which was chosen for efficient transformation and α -complementation (1). Table 1 shows that both TB1 and its parent, JM83 (7), display the constitutive phenotype. These results suggest that parent strain JM83 and its derivatives lack glycerol 3-phosphate repressor function. Strain JM83 was generated from strains CSH51 and 71.18 during the construction of strain JM101 (7). Table 1 also shows that JM101 displays the constitutive phenotype.

The role of the *glpR* locus in the constitutive phenotype of strain JM83 was investigated by using P1 transduction. The entry for strain KH20 in Table 1 shows that repair of the constitutive phenotype cotransduces with the *malt::Tn10* marker from strain MM130. The cotransduction frequencies were 70 and 72% in two separate experiments, a finding which agrees well with the frequency that is expected (73% [9]) from the proximity of the marker, located at 76.5 min, to the *glpEGR* locus at 76.7 min (2). Strain KH21, isolated as Tet^r from the same transduction, retains the constitutive phenotype, showing that the restored repression is not associated with the marker per se. Table 1 shows also that P1 transduction of strain KH11 restores the $GlpR^+$ phenotype (KH62). These results indicate that the constitutive phenotype is not associated with defects at the *glpFKX* locus at 88.8 min, i.e., the altered operator.

Diauxic growth in glucose-glycerol was used to evaluate the role of the repressor in glucose control of glycerol utilization and the expression of glycerol kinase. Figure 1 shows growth and glycerol kinase specific activity in cultures incubated at 37°C in minimal M9 medium with 2.5 mM glucose and 5 mM glycerol for strains MC4100, TS100, and KH44. The medium was inoculated with cells from overnight cultures in minimal glucose that were washed with minimal M9 salts. The constitutive phenotype is associated with shortening of the plateau of growth that occurs following glucose depletion and preceding glycerol utilization. However, repression of glycerol kinase by glucose does not depend on the *glpR* phenotype. This contrasts with results reported for glucose repression of the lactose operon, for which β -galactosidase increases in *lacI* strains (4). Glycerol kinase activity is repressed by glucose in the absence of glycerol in the *glpR2* strains, Lin7 and Lin43 (11). Results

* Corresponding author. Mailing address: Department of Biochemistry and Biophysics, Texas A&M University, 2128 TAMU, College Station, TX 77843-2128. Phone: (979) 845-9621. Fax: (979) 845-9274. E-mail: dpettigrew@tamu.edu.

TABLE 1. Glycerol kinase specific activities in *E. coli* strains

Strain	Genotype ^a	Source and/or reference	Growth medium ^b	Mean glycerol kinase sp act (U/mg) ± SD
JM83	F ⁻ <i>ara glpR208</i> Δ(<i>lac-proAB</i>) <i>rpsL</i> (Str ^r) [φ80 <i>dlac</i> Δ(<i>lacZ</i>)M15] <i>thi</i>	Ry Young (7)	LB	1.9 ± 0.1
JM101	<i>glpR208 supE thi</i> Δ(<i>lac-proAB</i>) [F' <i>traD36 proAB lacI</i> ^q ΔM15]	CGSC (7)	LB	1.8 ± 0.1
TB1	JM83 <i>glpR208 hsdR</i> (r _K ⁻ m _K ⁺)	Ry Young (1)	LB	1.9 ± 0.1
DG1	TB1 Δ <i>glpK202 glpR208</i>	6	LB	0.3 ± 0.02
KH11	DG1 <i>glpK</i> ⁺ <i>glpR208 pro</i> ⁺	DG1 × Hfr PK3	LB	2.5 ± 0.6
			mG	1.3 ± 0.5
KH62	KH11 <i>glpR</i> ⁺ <i>malT</i> ::Tn10	KH11 × P1.MM130	LB	0.3 ± 0.02
KH20	JM83 <i>glpR</i> ⁺ <i>malT</i> ::Tn10	JM83 × P1.MM130	LB	0.3 ± 0.02
KH21	JM83 <i>glpR208 malT</i> ::Tn10	JM83 × P1.MM130	LB	1.7 ± 0.6
Lin43	HfrC <i>jadL701 glpK22 glpR2 phoA8 pit-10 relA1 sopT1 tonA22 T₂</i> ^R	CGSC (11)	LB	2.4 ± 0.9
MC4100	F ⁻ <i>araD139</i> Δ(<i>argF-lac</i>)U169 <i>deoC1 flbB5301 ptsF25 rbsR relA1 rpsL150</i> (Str ^r)	CGSC	LB	0.4 ± 0.2
			mG	1.6 ± 0.7
TS100	MC4100 <i>glpR2</i>	Tim Larson	LB	2.5 ± 0.5
KH44	MC4100 Δ <i>glpEGR::kan</i>	MC4100 × P1.NZ42	LB	1.7 ± 0.2
NZ42	MC4100 φ(<i>glpK::lacZ</i>)(Hyb) Δ <i>glpEGR::kan</i>	Tim Larson		ND ^c
MG1655	λ ⁻ <i>rph-1</i>	CGSC	LB	0.2 ± 0.1
			mG	1.5 ± 0.2
MM130	<i>malT</i> ::Tn10	Mike Manson	LB	0.4 ± 0.02

^a The allele number *glpR208* was assigned by the *E. coli* Genetic Stock Center (CGSC) after the DNA sequence was determined and is added here to the genotypes of the relevant strains.

^b LB, LB medium; mG, minimal M9 salts with 1% (vol/vol) glycerol.

^c ND, not determined.

from diauxic growth curves and glycerol kinase expression for strains KH62 and KH11 (not shown) are the same as those shown for MC4100 and TS100, respectively. Strain KH44 (Δ*glpR*) was constructed to assess the extent of the repressor defect. Diauxic growth and glycerol kinase expression are indistinguishable for strains TS100 and KH44, indicating that the constitutive phenotype is associated with complete loss of repressor function.

Loss of repressor function could be associated with *glpEGR* promoter defects and/or the expression of a defective repressor. To determine whether these alleles encode variant repressors, the *glpR* locus from strains MC4100, MM130, Lin43, JM83, and DG1 was amplified from chromosomal DNA by PCR using *Taq* DNA polymerase and a Perkin-Elmer 2400 thermal cycler. Thermocycling conditions were as follows: 92°C for 4 min, followed by 30 cycles of 92°C for 30 s, 45°C for 1 min,




FIG. 1. Effect of *glpR* mutations on diauxic growth and the expression of glycerol kinase. Cultures were prepared as described in the text. The open symbols show the change in the A_{600} following inoculation, and the filled symbols show the glycerol kinase specific activity (S.A.). Symbols: circles, MC4100; squares, TS100; triangles, KH44.

and 72°C for 2 min, and finally, 72°C for 6 min. Both strands of the amplification products were sequenced using ABI Prism Dye Terminator Cycle Sequencing Core Kit with Big Dye premix from Perkin-Elmer, and thermocycling conditions were as recommended by ABI. Sequence products were purified using spin columns of P-30 polyacrylamide gel from Bio-Rad. The purified product was loaded onto an ABI 373 or 377 DNA Sequencing System in the Gene Technologies Laboratory of Texas A&M University. Sequence analysis was performed using Sequencher software.

The sequences of the alleles from strains MC4100 and MM130 are identical to the sequence reported for the allele from strain MC4100 (10). The sequence of the *glpR2* allele from strain 43 contains only one mutation relative to the sequence from MC4100, a G-to-C transversion at nucleotide 164 resulting in the amino acid change of glycine-55 to alanine. The sequences of the alleles from strains JM83 and DG1 are identical and contain only one mutation relative to the sequence from MC4100, i.e., a C-to-T transition at nucleotide 245 resulting in the amino acid change of alanine-82 to valine. This allele is designated *glpR208*. Thus, *glpR2* and *glpR208* are different alleles that are associated with the same phenotype. The same amino acid substitution found in *glpR2* has been identified independently in the *glpR200* allele from strain C600 (3) (Ann Flower, University of North Dakota, personal communication).

The *E. coli* glycerol 3-phosphate repressor is related to several other repressors in both gram-negative and gram-positive bacteria, as revealed by protein sequence alignments reported by Zeng et al. (10). It contains a helix-turn-helix DNA-binding motif which is located at amino acid residues 22 to 41. Thus, the substitutions that result from the *glpR2* and *glpR208* alleles do not occur in the DNA-binding motif, and direct effects on the recognition helix do not account for the loss of repressor function for these alleles. The sites of the substitutions are, however, highly conserved in the members of this repressor family. The glycine at position 55 is found in all but one member of the family, and the valine at position 82 is conserved absolutely. The extent of conservation at these two positions indicates their importance for repressor structure and function, and the complete loss of function that is associ-

ated with the variant alleles is consistent with the effects of substitutions of such highly conserved amino acids. Defective repressors associated with the substitutions found here could have effects on DNA-binding amino acids not in the helix-turn-helix motif, altered conformations that affect DNA binding, or decreased stability that prevents proper protein folding and expression.

This work was supported by a grant from the National Institutes of Health (GM-49992) and by the Texas Agricultural Experiment Station (H-6559).

We thank Donna Barker, Audra Boettcher, and Geneva Sampson for expert technical assistance and Ry Young, Mike Manson, and Tim Larson for generously providing strains.

REFERENCES

1. Baldwin, T. O., T. Berends, T. A. Bunch, T. F. Holzman, S. K. Rausch, L. Shamansky, M. L. Treat, and M. M. Ziegler. 1984. Cloning of the luciferase structural genes from *Vibrio harveyi* and expression of bioluminescence in *Escherichia coli*. *Biochemistry* **23**:3663–3667.
2. Berlyn, M. K. B., K. B. Low, and K. E. Rudd. 1996. Linkage map of *Escherichia coli* K-12, 9th ed., p. 1715–1902. In F. C. Neidhardt et al. (ed.), *Escherichia coli* and *Salmonella*: cellular and molecular biology, 2nd ed, vol. 2. ASM Press, Washington, D.C.
3. Elvin, C. M., C. M. Hardy, and H. Rosenberg. 1985. P_i exchange mediated by the GlpT-dependent sn-glycerol-3-phosphate transport system in *Escherichia coli*. *J. Bacteriol.* **161**:1054–1058.
4. Inada, T., K. Kimata, and H. Aiba. 1996. Mechanism responsible for glucose-lactose diauxie in *Escherichia coli*: challenge to the cAMP model. *Genes Cells* **1**:293–301.
5. Koch, J. P., S.-I. Hayashi, and E. C. C. Lin. 1964. The control of dissimilation of glycerol and L- α -glycerophosphate in *Escherichia coli*. *J. Biol. Chem.* **239**:3106–3108.
6. Liu, W. Z., R. Faber, M. Feese, S. J. Remington, and D. W. Pettigrew. 1994. *Escherichia coli* glycerol kinase: role of a tetramer interface in regulation by fructose 1,6-bisphosphate and phosphotransferase system regulatory protein III^{glc}. *Biochemistry* **33**:10120–10126.
7. Messing, J. 1979. A multipurpose cloning system based on the single-stranded DNA bacteriophage M13. NIH publication no. 79–99. Recombinant DNA Tech. Bull. **2**:43–48.
8. Pettigrew, D. W., G. B. Smith, K. P. Thomas, and D. C. Dodds. 1998. Conserved active site aspartates and domain-domain interactions in regulatory properties of the sugar kinase superfamily. *Arch. Biochem. Biophys.* **349**:236–245.
9. Wu, T. T. 1966. A model for three point analysis of random general transduction. *Genetics* **54**:405–410.
10. Zeng, G., S. Ye, and T. J. Larson. 1996. Repressor for the sn-glycerol 3-phosphate regulon of *Escherichia coli* K-12: primary structure and identification of the DNA-binding domain. *J. Bacteriol.* **178**:7080–7089.
11. Zwaig, N., W. S. Kistler, and E. C. C. Lin. 1970. Glycerol kinase, the pace-maker for the dissimilation of glycerol in *Escherichia coli*. *J. Bacteriol.* **102**:753–759.