



Demographic and clinical characteristics of patients with α_1 -antitrypsin deficiency genotypes PI*ZZ and PI*SZ in the Spanish registry of EARCO

María Torres-Durán¹, José Luis López-Campos ^{2,3}, Juan Luis Rodríguez-Hermosa^{4,5}, Cristina Esquinas^{2,6}, Cristina Martínez-González ⁷, José María Hernández-Pérez ⁸, Carlota Rodríguez⁹, Ana Bustamante ¹⁰, Francisco Casas-Maldonado¹¹, Miriam Barrecheguren^{2,6}, Cruz González ¹² and Marc Miravittles ^{2,6}

¹Servicio de Neumología, Hospital Álvaro Cunqueiro, NeumoVigo I+i Research Group, IIS Galicia Sur, Vigo, Spain. ²Centro de Investigación Biomédica en Red de Enfermedades Respiratorias (CIBERES), Instituto de Salud Carlos III, Madrid, Spain. ³Unidad Médico-Quirúrgica de Enfermedades Respiratorias, Instituto de Biomedicina de Sevilla (IBiS), Hospital Universitario Virgen del Rocío/Universidad de Sevilla, Sevilla, Spain. ⁴Servicio de Neumología, Hospital Clínico de San Carlos, Departamento de Medicina, Facultad de Medicina, Universidad Complutense de Madrid, Madrid, Spain. ⁵Research Institute of Hospital Clínico San Carlos (IdISSC), Madrid, Spain. ⁶Pneumology Dept, Hospital Universitari Vall d'Hebron; Vall d'Hebron Institut de Recerca (VHIR), Vall d'Hebron Barcelona Hospital Campus, Barcelona, Spain. ⁷Pneumology Dept, Hospital Universitario Central de Asturias, Instituto de Investigación Sanitaria del Principado de Asturias, Oviedo, Spain. ⁸Pneumology Dept, Hospital Universitario Nuestra Señora de La Candelaria, Santa Cruz de Tenerife, Spain. ⁹Servicio de Neumología, Complejo Hospitalario Clínico-Universitario de Santiago, Santiago de Compostela, Spain. ¹⁰Pneumology Section, Hospital Sierrallana-TresMares, Cantabria, Spain. ¹¹Servicio de Neumología, Hospital Clínico Universitario San Cecilio, Departamento de Medicina, Facultad de Medicina, Universidad de Granada, Granada, Spain. ¹²Servicio de Neumología, Hospital Clínico Universitario, Instituto de Investigación INCLIVA, Valencia, Spain.

Corresponding author: Marc Miravittles (marcm@separ.es)



Shareable abstract (@ERSpublications)

Patients with the PI*ZZ genotype have more severe pulmonary disease than those with the PI*SZ genotype. However, asthma is as frequent in PI*SZ as in PI*ZZ. <https://bit.ly/3m10MNN>

Cite this article as: Torres-Durán M, López-Campos JL, Rodríguez-Hermosa JL, *et al.* Demographic and clinical characteristics of patients with α_1 -antitrypsin deficiency genotypes PI*ZZ and PI*SZ in the Spanish registry of EARCO. *ERJ Open Res* 2022; 8: 00213-2022 [DOI: 10.1183/23120541.00213-2022].

Copyright ©The authors 2022

This version is distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0. For commercial reproduction rights and permissions contact permissions@ersnet.org

Received: 2 May 2022
Accepted: 26 May 2022

Abstract

Background The Spanish registry of α_1 -antitrypsin deficiency (AATD) integrated in the European Alpha-1 Research Collaboration (EARCO) provides information about the characteristics of patients, in particular those with the PI*SZ genotype, which is frequent in Spain.

Method Individuals with severe AATD defined as proteinase inhibitor (PI) genotypes PI*ZZ, PI*SZ and other rare deficient variants were included from February 1, 2020, to February 1, 2022. The analysis focused on a comparison of the characteristics of PI*ZZ and PI*SZ patients.

Results 409 patients were included (53.8% men) with a mean \pm SD age of 53.5 \pm 15.9 years. Genotypes were PI*ZZ in 181 (44.7%), PI*SZ in 163 (40.2%), PI*SS in 29 (7.2%) and other in 32 (7.9%). 271 (67.4%) had lung disease: 175 chronic obstructive pulmonary disease (43.5%), 163 emphysema (40.5%) and 83 bronchiectasis (20.6%). Patients with the PI*SZ genotype were younger, more frequently non-index cases and had a lower frequency of respiratory diseases except asthma compared with PI*ZZ patients. Among patients with respiratory diseases, PI*SZ individuals were significantly older both at onset of symptoms and at diagnosis; only asthma was more frequent in PI*SZ than in PI*ZZ individuals. Twelve PI*SZ patients (15.4%) received augmentation therapy compared with 94 PI*ZZ patients (66.2%; $p < 0.001$).

Conclusions There is a high prevalence of PI*SZ in Spain. Patients with the PI*SZ genotype were older at symptom onset and diagnosis and had less severe lung disease compared with PI*ZZ patients. The prevalence of asthma was higher in PI*SZ, and up to 15% of PI*SZ patients received augmentation therapy.

Introduction

α_1 -antitrypsin deficiency (AATD) is considered the most frequent potentially severe rare disease affecting adults, and its prevalence is around two in 10 000 in Europe [1]. AATD may predispose to liver disease in children and adults and pulmonary emphysema in adults [2].



The majority of patients with severe AATD are homozygous carriers of the Z mutation of the gene coding for the α_1 -antitrypsin (AAT) protein (proteinase inhibitor (PI*ZZ)), but hundreds of variants have been identified, some of which are associated with low serum AAT levels [3, 4]. Among these variants, the most frequent is the S variant, which is particularly frequent in Western Spain [5]. The risk of lung disease associated with the heterozygous PI*SZ genotype is controversial. Some carriers have serum levels below those considered protective and may be candidates for augmentation therapy with intravenous AAT, if emphysema develops [6, 7], while some studies suggest that the risk of lung disease associated with PI*SZ is significantly less than that associated with PI*ZZ [8].

The low prevalence of AATD and the extreme variability of its respiratory clinical manifestations make it likely that a single centre will never see enough patients to acquire experience in the management of this condition. Therefore, the European Council [9] and the European Respiratory Society (ERS) [1] recommend organising the care of AATD patients in reference centres and developing prospective registries to better understand the natural history of the disease and the potential impact of different treatments [10]. The Spanish registry of AATD was founded in 1992 [11] and in 2020 it was merged into the new European Alpha-1 Research Collaboration (EARCO) International Registry [12]. The EARCO registry is a prospective international registry developed by the EARCO Clinical Research Collaboration of the ERS [13], and the Spanish registry consists of the Spanish investigators, centres and participants included in the EARCO registry [14].

The current study presents the demographic and clinical characteristics of the participants in the Spanish registry in EARCO and, because Spain is one of the countries with the highest prevalence of the S allele, it provides a unique opportunity to investigate the differential characteristics of individuals with the PI*SZ genotype [15] and compare them with those of PI*ZZ individuals.

Method

Structure of the Spanish registry of AATD

The Spanish registry of patients with AATD is constituted by the Spanish investigators and patients registered in the EARCO International Registry [14]. The registry is open to every clinician who manages AATD patients in Spain. The registry is observational and, therefore, patients are treated according to the attending physicians' criteria. The Spanish registry has one national coordinator (MT-D) designated by the Spanish Society of Pneumology and Thoracic Surgery (SEPAR), and a steering committee formed by clinicians and researchers with expertise in the disease. According to the data processor agreement signed between the national coordinator, representing the SEPAR and the Spanish registry, and the data controller of EARCO (Vall d'Hebron Research Institute, Barcelona, Spain), it is free to access the fully anonymised data of the Spanish patients in the Spanish registry for analysis and evaluation.

The study protocol received central ethics approval by the research ethics committee of the Vall d'Hebron University Hospital of Barcelona, Spain (PR(AG)480/2018) and was subsequently approved by all the participating centres. All participants provided written informed consent. The EARCO registry protocol has been registered at www.clinicaltrials.gov (ID: NCT04180319) and is hosted by www.earco.org. The personal data of the patients are kept under strict confidentiality in compliance with the provisions of the Spanish Organic Law 3/2018 of December 5, 2018, Protection of Personal Data and Guarantee of Digital Rights (LOPDGDD) and its development regulations, and in accordance with the provisions of the General Data Protection Regulation (GDPR) 2016/679 of the European Parliament and of the European Council of April 27, 2016, regarding the protection of personal data.

Objectives of the Spanish AATD registry

The objectives of the Spanish AATD registry are linked to those of EARCO and have been described in detail in a previous publication [16]. Briefly, the main objectives are 1) to generate longitudinal long-term, high-quality clinical data of individuals with AATD; 2) to understand the natural history and prognosis of AATD; 3) to investigate the effect of AAT augmentation and other therapies on the progression of lung disease; and 4) to learn more about the course of the disease in patients with severe AATD with genotypes other than PI*ZZ.

In the current publication, we analyse the baseline data of patients included in the Spanish registry in the first 2 years of EARCO, from February 1, 2020, to February 1, 2022.

Population and measurements

The protocol of the study has been published previously [16]. The inclusion criteria are 1) individuals with diagnosed severe AATD; 2) deficiency defined as AAT serum levels $<11 \mu\text{M}$ ($50 \text{ mg}\cdot\text{dL}^{-1}$) and/or

proteinase inhibitor genotypes PI*ZZ, PI*SZ and compound heterozygotes or homozygotes of other rare deficient variants. The only exclusion criterion is lack of patient consent.

The data collected include the following domains: demographics, proteinase inhibitor genotype and other laboratory analyses, comorbidities, lung function, respiratory symptoms, ultrasound-based elastography of the liver, exacerbations of respiratory disease, quality of life, physical activity, chest computed tomography (as applicable) and treatment.

Data are entered into a secure database through an electronic case report form hosted by the EARCO website (www.earco.org). Data are centrally monitored, and queries are sent for missing or invalid data.

Statistical analysis

Qualitative variables are described with absolute frequencies and percentages. Quantitative variables are described as mean \pm SD or median and quartiles. The Kolmogorov–Smirnov test was used to assess the normality of distributions. The sociodemographic and clinical characteristics were compared according to the genotypes PI*ZZ and PI*SZ. In the case of quantitative variables, a t-test was carried out (Mann–Whitney U-test if normality was not assumed). A Chi-squared test (Fisher test for frequencies <5) was used to compare categorical variables. Cigarette smoking pack-years were transformed into an ordinal variable (0, <10, 10–20, 20–30 and >30) to compare the per cent predicted forced expiratory volume in 1 s (FEV₁ % pred) between genotypes according to the degree of smoking consumption. For all the tests, $p < 0.05$ was considered statistically significant. The statistical package RStudio v2.5.1 (www.rstudio.com) was used for the analyses.

Results

Population included

The Spanish registry included 409 individuals with severe AATD from 17 centres distributed throughout the country; 218 participants (53.8%) were men, mean \pm SD age was 53.5 \pm 15.9 years and only 33 (8%) were active smokers. The distribution of genotypes was PI*ZZ in 181 patients (44.7%), PI*SZ in 163 patients (40.2%), PI*SS in 29 patients (7.2%) and others in 32 patients (7.9%). Up to 70.6% were index cases and the mean age at diagnosis was 46.4 \pm 17.3 years.

Characteristics of lung disease in participants in the Spanish registry

Lung disease was present in 271 participants (67.4%); the most frequent pulmonary disease was chronic obstructive pulmonary disease (COPD) in 175 patients (43.5%), followed by emphysema in 163 (40.5%) and bronchiectasis in 83 (20.6%). The most frequent respiratory symptoms at presentation were dyspnoea in 217 (53%), cough in 96 (23.5%) and sputum production in 68 (16.6%). Up to 77 patients (19.3%) had a history of pneumonia and spirometry showed a mean \pm SD FEV₁ % pred of 81.7 \pm 31.3% and a mean \pm SD per cent predicted transfer coefficient of the lung for carbon monoxide (K_{CO} % pred) of 76.8 \pm 23.4%.

Comparison of demographic and clinical characteristics between PI*ZZ and PI*SZ patients

Patients with the PI*SZ genotype were younger, more frequently non-index cases and had a lower frequency of most respiratory diseases (COPD 25.9%, emphysema 24.7% and bronchiectasis 10% versus 60.3%, 59.8% and 32.4% in the PI*ZZ patients; $p < 0.001$ for all comparisons). However, there was no difference in the prevalence of asthma (13.6% versus 13.4%, $p = 0.96$). Among PI*SZ patients there were more active smokers (23.2% versus 2.8%, $p < 0.001$), but they had better lung function and a lower frequency of exacerbations and previous pneumonia (table 1). Never-smokers showed no differences in FEV₁ % pred, but significant differences in FEV₁ % pred were observed with increased accumulated tobacco consumption in both genotypes (p for trend < 0.001 for PI*ZZ and PI*SZ), with smoking having a higher impact at lower cumulative doses in PI*ZZ patients (figure 1a). Significant differences between both genotypes were also observed for per cent predicted forced vital capacity (FVC % pred) but not for FEV₁/FVC (figure 1b, c).

Comparison of characteristics and severity of lung disease between PI*ZZ and PI*SZ patients

The comparison of the 82 PI*SZ and 146 PI*ZZ patients with respiratory disease is shown in table 2. The distributions of age and sex were no different between the two genotypes, but PI*SZ patients were more frequently active smokers. The proportions of index cases, patients identified by family screening and patients with cardiovascular comorbidity did not differ between the two groups; however, PI*SZ patients had a significantly older age than PI*ZZ patients both at symptom onset (49.9 \pm 18.1 years versus 45.6 \pm 15 years, $p = 0.008$) and diagnosis (52.5 \pm 18 years versus 47.4 \pm 14.9 years, $p = 0.003$). The distribution of respiratory diseases was different between the two genotypes, with more emphysema, COPD and bronchiectasis in PI*ZZ patients and more frequent asthma in PI*SZ patients (table 2 and figure 2a, b). Both FEV₁ % pred

TABLE 1 Demographic and clinical characteristics of the participants in the Spanish registry of the European Alpha-1 Research Collaboration and comparison between Pi*ZZ and Pi*SZ genotypes

Variables	Global	ZZ	SZ	p-value
Subjects (n)	405	181	163	
Age (years)	53.5±15.9	55.4±14.7	51.0±17.3	0.015
Sex, male	218 (53.8)	95 (52.5)	87 (53.4)	0.869
BMI (kg·m ⁻²)	26.4±4.8	26.2±4.9	26.3±4.9	0.730
Smokers	33 (8)	3 (2.8)	22 (23.2)	<0.001
Ex-smokers	222 (54.4)	111 (61.3)	3 (44.9)	
Never-smokers	149 (36.8)	66 (36.7)	68 (41)	
Tobacco exposure (pack-years)	24.5±21.9	20.3±13.8	25.2±25.1	0.953
Chronic respiratory disease	267 (66.4)	146 (81.6)	82 (50.6)	<0.001
COPD	175 (43.5)	108 (60.3)	42 (25.9)	<0.001
Emphysema	163 (40.5)	107 (59.8)	40 (24.7)	<0.001
Bronchiectasis	83 (20.6)	58 (32.4)	16 (10)	<0.001
Asthma	55 (13.7)	24 (13.4)	22 (13.6)	0.963
Liver disease	42 (10.6)	24 (13.4)	16 (10)	0.322
Age at onset of symptoms (years)	46.1±17.2	44.2±16.3	47.9±17.5	0.063
Age at diagnosis of AATD (years)	46.4±17.4	44.5±17.2	46.8±18	0.354
Index case	286 (70.6)	135 (74.6)	106 (65)	0.053
Family screening	98 (24)	39 (21.5)	45 (27.6)	0.121
Charlson index (age corrected)	3.1±2.1	3.2±2	2.9±2	0.347
Cardiovascular disease [#]	89 (22.5)	47 (26.4)	25 (15.7)	0.017
FVC % pred	99.4±22.6	96.1±26.2	103±17.4	0.011
FEV ₁ % pred	81.7±31.4	69.8±32.1	93.3±26	<0.001
FEV ₁ /FVC % pred	65±19	57±19	72±18	<0.001
K _{CO} % pred	76.8±23.4	67.6±22.5	84.7±20.8	<0.001
Augmentation therapy	114 (28.9)	95 (54)	13 (8.2)	0.001
AAT (mg·dL ⁻¹)	44.5±24.4	24.9±17.6	56.7±11.8	<0.001
Number of exacerbations previous year	0.44±1	0.66±1.2	0.28±0.8	<0.001
History of pneumonia	77 (19.3)	44 (24.6)	21 (13.1)	0.007
CAT	7.4±7.2	9.4±7.8	5.6±6.2	<0.001
BODEx	1.2±1.8	1.8±2	0.6 (1.3)	<0.001

Data presented as mean±SD or n (%), unless otherwise indicated. BMI: body mass index; COPD: chronic obstructive pulmonary disease; AATD: α_1 -antitrypsin deficiency; FVC: forced vital capacity; FEV₁: forced expiratory volume in 1 s; K_{CO}: transfer coefficient of the lung for carbon monoxide; AAT: α_1 -antitrypsin; CAT: COPD Assessment Test; BODEx: BMI, Obstruction, Dyspnoea, Exacerbations index. [#]: cardiovascular disease includes hypertension, ischaemic heart disease, congestive heart failure and peripheral vascular disease.

and K_{CO} % pred were significantly lower in Pi*ZZ than Pi*SZ patients (FEV₁ % pred: 61.1±28.2% versus 78.7±28.5%; K_{CO} % pred: 62.9±20.6% versus 84.7±20.8%; p<0.001 for both comparisons). The COPD Assessment Test score was also higher (worse) for Pi*ZZ compared to Pi*SZ patients (10.9±7.9 versus 8.6±7, p<0.036). A total of 12 Pi*SZ patients (15.4%) received augmentation therapy compared with 94 Pi*ZZ patients (66.2%; p<0.001).

Characteristics of lung diseases in patients with AATD

Patients with AATD and asthma had different characteristics from those with emphysema or bronchiectasis. Patients with asthma were younger and more frequently female, with a higher proportion of never-smokers. They also had better lung function and a lower prevalence of comorbidities but a similar frequency of previous pneumonia compared to patients with emphysema. The demographic and clinical characteristics of patients with the most frequent chronic respiratory diseases are shown in table 3. No statistical comparisons were performed because some patients could be included in more than one category due to the coexistence of respiratory diseases in some individuals.

Discussion

The implementation of the EARCO International Registry has been an opportunity for the Spanish AATD registry to join an international initiative and upgrade the database and take advantage of the structure and monitoring system of EARCO to improve the quality of data collection [14]. During the first 2 years of recruitment, and despite the COVID-19 pandemic, a total of 17 centres included more than 400 patients in the EARCO registry, which constitutes the new Spanish AATD registry.




FIGURE 1 Comparison of per cent predicted forced expiratory volume in 1 s (FEV₁ % pred) (a), forced vital capacity (FVC) (b) and FEV₁/FVC (c) between participants in the Spanish registry with the PI*ZZ and PI*SZ genotypes. In each box plot, the median value is indicated by the centre horizontal line, and the 25th and 75th percentiles are indicated by the lower and upper box horizontal lines. Circles indicate the outliers. p-values determined using a t-test. There was a statistically significant difference between pack-years of smoking groups for FEV₁ % pred and FEV₁/FVC for both the PI*ZZ genotype (p-value for trend <0.001) and PI*SZ genotype (p-value for trend <0.001). For FVC % pred, the p-value for trend was p=0.004 for PI*ZZ and p=0.29 for PI*SZ.

The characteristics of the patients included have changed from the previous reports of the Spanish registry. The first report made in 1998 included 223 patients: 73% male, mean age 46 years and mean FEV₁ % pred of 53% [11]. The second report in 2007 included 462 individuals: 63% male, mean age 51 years and mean FEV₁ % pred of 53% [17]. Interestingly the percentage of males is now reduced to 54%, while the mean age has increased to 53 years and mean FEV₁ to 81% predicted. This increase in the mean FEV₁ % pred may, in part, be due to the improvement in population and family screening, earlier detection of carriers and early diagnosis of lung diseases in individuals affected [18–21]. The inclusion of genotypes other than PI*ZZ may also explain the better lung function in the current analysis, although the influence is very limited because the mean FEV₁ % pred in PI*ZZ patients was 69.8%, which is still much better than in previous reports [11, 17].

Several registries have evaluated the characteristics of individuals with the most common deficient PI*ZZ genotype [22–25], but there is limited information about the characteristics of patients with the PI*SZ genotype. Because the S allele is especially frequent in Spain [5, 15], the Spanish registry provides a unique opportunity to investigate the clinical manifestations of PI*SZ individuals and to compare with PI*ZZ carriers. In general, patients with the PI*SZ genotype were younger, more frequently non-index cases and had a lower frequency of most respiratory diseases, despite a higher prevalence of active smokers, but there were no differences in the prevalence of asthma. The possible influence of the S alleles in the prevalence and severity of asthma has been previously reported [26] and justifies the investigation of AATD in all adult asthmatic patients with non-fully reversible airflow obstruction, as indicated in guidelines. It is of note that we did not observe significant differences in lung function between PI*SZ and PI*ZZ never-smokers. However, with increasing intensity of smoking, patients with PI*ZZ presented more severely impaired lung function, while PI*SZ patients only showed deteriorated lung function at smoking consumptions of >20 pack-years, highlighting again the increased susceptibility of homozygous Z patients to the deleterious effect of tobacco smoking [27–29].

TABLE 2 Comparison of demographic and clinical characteristics of the participants in the Spanish registry of European Alpha-1 Research Collaboration with chronic lung disease and PI*ZZ and PI*SZ genotypes

Variables	ZZ	SZ	p-value
Subjects (n)	146	82	
Age (years)	58.6±12.3	57.6±17.2	0.692
Sex, male	75 (51.4)	47 (57.8)	0.386
BMI (kg·m ⁻²)	26.4±4.8	27.6±5.3	0.067
Smokers	2 (1.4)	14 (17.1)	<0.001
Ex-smokers	105 (71.9)	48 (58.5)	
Never-smokers	39 (26.7)	20 (24.4)	
Tobacco exposure (pack-years)	18.8±9.7	20.7±13.5	0.460
COPD	108 (74)	42 (51.2)	0.001
Emphysema	107 (73.3)	40 (48.8)	<0.001
Bronchiectasis	58 (39.7)	16 (19.5)	0.002
Asthma	24 (16.4)	22 (26.8)	0.061
Liver disease	17 (11.8)	8 (9.8)	0.637
Age at onset of symptoms (years)	45.6±15	49.9±18.1	0.008
Age at diagnosis of AATD (years)	47.4±14.9	52.5±18	0.003
Index case	25 (17.1)	15 (18.3)	0.824
Family screening	21 (14.4)	12 (14.6)	0.959
Charlson index (age corrected)	3.5±1.9	4±2.2	0.105
Cardiovascular comorbidity [#]	43 (30.1)	22 (28.2)	0.771
FVC % pred	93.4±26.6	100.4±20.7	0.046
FEV ₁ % pred	61.1±28.2	78.7±28.5	<0.001
FEV ₁ /FVC % pred	51±17	62±19	<0.001
K _{CO} % pred	62.9±20.6	84.7±20.8	<0.001
Augmentation therapy	94 (66.2)	12 (15.4)	<0.001
AAT (mg·dL ⁻¹)	25.8±19.4	57.5±12.2	<0.001
Number of exacerbations previous year	0.80±1.3	0.43±1	0.018
History of pneumonia	39 (26.9)	10 (12.7)	0.014
CAT	10.9±7.9	8.6±7	0.036
BODEx	2.2±2.1	1.1±1.7	<0.001

Data presented as mean±SD or n (%), unless otherwise indicated. BMI: body mass index; COPD: chronic obstructive pulmonary disease; AATD: α_1 -antitrypsin deficiency; FVC: forced vital capacity; FEV₁: forced expiratory volume in 1 s; K_{CO}: transfer coefficient of the lung for carbon monoxide; AAT: α_1 -antitrypsin; CAT: COPD Assessment Test; BODEx: BMI, Obstruction, Dyspnoea, Exacerbations index. [#]: cardiovascular disease includes hypertension, ischaemic heart disease, congestive heart failure and peripheral vascular disease.




FIGURE 2 Distribution of lung diseases in participants of the Spanish registry with the PI*ZZ (a) and PI*SZ (b) genotypes. The size of each circle is proportional to the number of individuals in each group.

TABLE 3 Demographic and clinical characteristics of the participants in the Spanish registry of European Alpha-1 Research Collaboration with different pulmonary diseases

Variables	Emphysema	Bronchiectasis	Asthma
Subjects (n)	163	83	55
Age (years)	61.2±11.7	62.5±11.5	45.8±16.4
Sex, male	96 (58.9)	41 (49.4)	24 (43.6)
BMI (kg·m ⁻²)	26.7±5.1	26.8±4.4	26.4±5.3
Smokers	11 (6.7)	4 (4.8)	6 (10.9)
Ex-smokers	129 (79.1)	52 (62.7)	25 (45.5)
Never-smokers	23 (14.1)	27 (32.5)	24 (43.6)
Tobacco exposure (pack-years)	29.4±22.5	25.5±22	18.7±19
COPD	128 (78.5)	53 (63.9)	14 (25.5)
Emphysema	163 (100)	53 (63.9)	13 (23.6)
Bronchiectasis	53 (32.5)	83 (100)	8 (14.5)
Asthma	13 (8)	8 (9.6)	55 (100)
Liver disease	18 (11.2)	8 (9.6)	5 (9.1)
Age at onset of symptoms (years)	48.7±15.4	48.7±14.9	31.4±17.3
Age at diagnosis of AATD (years)	51.7±14.1	52.9±12.9	38.8±19
Index case	139 (85.3)	75 (90.4)	41 (74.5)
Family screening	21 (12.9)	8 (9.6)	10 (18.2)
Charlson index (age corrected)	3.9±1.7	3.9±1.6	2.3±2.1
Cardiovascular comorbidity [#]	58 (36.7)	28 (34.1)	9 (17)
FVC % pred	94.3±27.7	90.7±25.5	101±4
FEV ₁ % pred	59.6±27.5	67.7±28.7	86.3±27.9
FEV ₁ /FVC % pred	49±15	58±16	69±17
K _{CO} % pred	60.9±19.9	70.4±21.7	81.8±20.9
Augmentation therapy	95 (59.7)	38 (47.5)	8 (16)
AAT (mg·dL ⁻¹)	38.2±26.9	37.4±22.3	45.8±21.9
Number of exacerbations previous year	0.7±1.2	0.7±1.3	0.5±1.2
History of pneumonia	34 (21)	27 (32.5)	12 (23.1)
CAT	11.5 (7.7)	11 (7.8)	7 (6.4)
BODEx	2.3 (2)	1.7 (2)	1 (1.5)

Data presented as mean±SD or n (%), unless otherwise indicated. The different subgroups are not mutually exclusive. BMI: body mass index; COPD: chronic obstructive pulmonary disease; AATD: α_1 -antitrypsin deficiency; FVC: forced vital capacity; FEV₁: forced expiratory volume in 1 s; K_{CO}: transfer coefficient of the lung for carbon monoxide; AAT: α_1 -antitrypsin; CAT: COPD Assessment Test; BODEx: BMI, Obstruction, Dyspnoea, Exacerbations index. #: cardiovascular disease includes hypertension, ischaemic heart disease, congestive heart failure and peripheral vascular disease.

Considering only the individuals with diagnosed lung disease, we observed that PI*SZ carriers were older, both at onset of symptoms and at diagnosis, than PI*ZZ patients and the severity of lung disease was milder in terms of FEV₁ and K_{CO} % pred.

Augmentation therapy was administered to two thirds of patients with the PI*ZZ genotype and lung disease. It is not possible to ascertain why the remaining third of patients was not treated. Two patients were active smokers and treatment was therefore not indicated, but other reasons, including the patient refusing treatment, could explain the lack of augmentation for PI*ZZ patients with lung disease. Although the indications for augmentation are clear, the opinions and attitudes about augmentation may vary greatly among specialists [30], and some may not indicate augmentation for patients with very mild or very severe disease and others may not initiate augmentation if patients do not show clinical, functional or radiological deterioration during follow-up [30, 31].

By contrast, 15.4% of PI*SZ patients with lung disease received augmentation. Treatment for PI*SZ patients is included in the indications described in the summary of product characteristics of both brands of augmentation therapy available in Spain [32, 33]; nevertheless, there is no consensus among specialists about the efficacy of augmentation in PI*SZ patients with lung disease. While some studies demonstrated that PI*SZ patients have a significantly increased risk of lung disease and augmentation would help to prevent its progression [7, 29, 34], others suggest that the risk of lung disease associated with the PI*SZ genotype is similar to that associated with PI*MZ and much less than the homozygous form of PI*ZZ [8, 35].

An interesting observation derived from the data is the different distribution of lung diseases in PI*SZ and PI*ZZ patients. Emphysema and bronchiectasis were more frequent in PI*ZZ, but asthma was relatively more frequent in PI*SZ. Among the PI*ZZ patients, only a minority had lung disease without emphysema, with 16 cases of bronchiectasis (11%), 13 of asthma (9%) and two cases of both diseases combined (1.3%). In contrast, 23.2% of PI*SZ patients had asthma and no evidence of emphysema. We can compare our results with those of a similar cohort of 424 unrelated PI*ZZ patients in the UK. In this study, Wood *et al.* [36] reported a higher prevalence of emphysema (65.8% *versus* 59.8% in our series) and lower prevalence of bronchiectasis (19.5% *versus* 32.4% in Spain). It is of note that 10% of their patients with bronchiectasis had no emphysema, very similar to the 12% observed in our patients. Data from Wood *et al.* [36] were published 16 years ago and it is not clear whether differences observed are due to real differences in phenotypes or due to a higher implementation of AATD screening and more frequent use of high-resolution computed tomography scans during the last years.

The demographic and clinical profiles of patients with emphysema or bronchiectasis in our group were quite similar, but those of asthmatic patients were clearly different. Patients with asthma were younger and diagnosed at a younger age, there was a lower prevalence of smokers, more women and a higher frequency of patients diagnosed by family screening, and patients had better lung function, lower symptom burden and higher serum AAT concentrations. History of pneumonia was also more frequent in PI*ZZ than PI*SZ patients, both in the global population and in individuals with lung disease. These differences could in part be explained by the higher severity of lung disease in PI*ZZ patients, while other risk factors for pneumonia, such as age, were no different between the genotypes.

Spain is only one of the 16 countries currently participating in the EARCO International Registry. Some of the participating countries have joined EARCO through a unique national reference centre, but others, such as Portugal [37] and the UK, have followed the same structure as the Spanish registry with multiple centres contributing directly to the database and designating a national coordinator who has direct access to the anonymised national data. The EARCO International Registry is the main research project of the EARCO initiative, but there are other research projects under development derived from the survey of unmet needs developed by academics, patients and caregivers' representatives. The follow-up of patients included in the registry will provide relevant information about the natural history of the disease [38].

Conclusions

The Spanish registry has shown the evolution of the baseline characteristics of the patients included, who now have an older age and more preserved lung function, suggesting an improvement in early diagnosis and better management of pulmonary disease. In general, patients with the PI*SZ genotype had milder respiratory disease and were less susceptible to tobacco smoking but had a similar prevalence of asthma. Longitudinal follow-up is necessary to better understand the natural history of the AATD.

Acknowledgements: The Spanish registry would like to acknowledge the support of the EARCO Steering committee: Christian Clarenbach and Marc Miravittles (co-chairs), Robert Bals, Jan Stolk, Joanna Chorostowska-Wynimko, Karen O'Hara, Marion Wilkens, José Luis López-Campos, Alice M. Turner, Ilaria Ferrarotti, Gerry McElvaney and Robert A. Stockley.

The following investigators have contributed to the Spanish registry of AATD: Maria Torres-Durán (national coordinator), José Luis López-Campos, Juan Luis Rodríguez-Hermosa, Myriam Calle, Cristina Esquinas (data manager), Cristina Martínez, José María Hernández-Pérez, Carlota Rodríguez, Ana Bustamante, Francisco Casas-Maldonado, Miriam Barrecheguren, Cruz González, Marc Miravittles, Eva Taberner, Mar Fernández-Nieto, Francisco Javier Michel, Virginia Almadana, Lourdes Lázaro, Carlos Martínez-Rivera and María Isabel Parra.

We wish to acknowledge Elise Heuvelin from the European Respiratory Society (Laussane, Switzerland) for her support in the management of EARCO, and Gemma Vilagut and Christina Founti (Bioclever, Barcelona, Spain) for their support in EARCO data monitoring. We also acknowledge the participation of Beatriz Lara (Coventry, UK) and Eduardo Loeb (Barcelona, Spain) in the development of the database and the monitoring of data.

Provenance: Submitted article, peer reviewed.

Availability of data and materials: Information on the EARCO data sharing commitments and requesting access to anonymised aggregated participant data and associated documents can be found at www.earco.org.

Authors' contributions: Study concept and design by M. Torres-Durán, C. Esquinas and M. Miravittles. Data acquisition by M. Torres-Durán, J.L. López-Campos, J.L. Rodríguez-Hermosa, C. Martínez-González, J.M. Hernández-Pérez, C. Rodríguez, A. Bustamante, F. Casas-Maldonado, M. Barrecheguren, C. González and

M. Miravittles. Statistical analysis by C. Esquinas. Drafting of the manuscript by M. Miravittles, C. Esquinas and M. Torres-Durán. Critical revision and approval for submission by M. Torres-Durán, J.L. López-Campos, J.L. Rodríguez-Hermosa, C. Esquinas, C. Martínez-González, J.M. Hernández-Pérez, C. Rodríguez, A. Bustamante, F. Casas-Maldonado, M. Barrecheguren, C. González and M. Miravittles. All authors have read and approved the final manuscript.

Competing interests: M. Torres-Durán has received speaker fees from Chiesi, CSL Behring, Grifols and Resmed, and consulting fees from CSL Behring and Grifols. J.L. López-Campos has received honoraria during the last 3 years for lecturing, scientific advice, participation in clinical studies or writing for publications for AstraZeneca, Boehringer Ingelheim, Chiesi, CSL Behring, Esteve, Ferrer, Gebro, GlaxoSmithKline, Grifols, Menarini, Novartis, Rovi and Teva. J.L. Rodríguez-Hermosa has received speaker fees from Zambon, Bial, Gebro Pharma, GlaxoSmithKline, Chiesi, Boehringer Ingelheim, CSL Behring and Grifols. J.M. Hernández-Pérez has received speaker fees from Grifols, CSL Behring, AstraZeneca, GlaxoSmithKline, Bial laboratory and Teva laboratory, support for attending meetings from Grifols and CSL Behring, and consulting fees from CSL Behring. C. Rodríguez has received speaker fees from GlaxoSmithKline, AstraZeneca, Grifols, Chiesi, Ferrer, Menarini and Boehringer Ingelheim, has provided expert testimony for Chiesi, and has received support for attending meetings from FAES. A. Bustamante has received speaker fees from Boehringer Ingelheim, AstraZeneca, GlaxoSmithKline, Novartis and Ferrer, and funding for travelling or attending meetings from CSL Behring, AstraZeneca and Chiesi. F. Casas-Maldonado has received speaker fees from AstraZeneca, Bial, Boehringer Ingelheim, Chiesi, Gebro Pharma, GlaxoSmithKline, Laboratorios Esteve, Laboratorios Ferrer, Menarini, Novartis, Rovi, Teva, Vertex, Zambon, CSL Behring and Grifols, and consulting fees from AstraZeneca, Chiesi, GlaxoSmithKline, CSL Behring and Grifols. M. Barrecheguren has received speaker fees from Grifols, Menarini, CSL Behring, GlaxoSmithKline and Boehringer Ingelheim, and consulting fees from GlaxoSmithKline, Novartis, CSL Behring and Boehringer Ingelheim. C. González has received speaker fees from Menarini, GlaxoSmithKline, Novartis, Boehringer Ingelheim and Chiesi. M. Miravittles has received speaker fees from AstraZeneca, Boehringer Ingelheim, Chiesi, Cipla, Menarini, Rovi, Bial, Sandoz, Zambon, CSL Behring, Grifols and Novartis, consulting fees from AstraZeneca, Boehringer Ingelheim, Chiesi, GlaxoSmithKline, Bial, Gebro Pharma, Kamada, CSL Behring, Laboratorios Esteve, Ferrer, Mereo Biopharma, Verona Pharma, Teva, Spin Therapeutics, pH Pharma, Novartis, Sanofi and Grifols, and research grants from GlaxoSmithKline and Grifols. The remaining authors report no conflicts of interest.

Support statement: The Spanish registry is funded by an unrestricted grant from Grifols. The EARCO International Registry is funded by unrestricted grants of Grifols, CSL Behring, Kamada, pH Pharma and Takeda to the European Respiratory Society. Funding information for this article has been deposited with the Crossref Funder Registry.

References

- 1 Blanco I, Bueno P, Diego I, *et al.* α_1 -antitrypsin Pi*Z gene frequency and Pi*ZZ genotype numbers worldwide: an update. *Int J Chron Obstruct Pulmon Dis* 2017; 12: 561–569.
- 2 Miravittles M, Dirksen A, Ferrarotti I, *et al.* European Respiratory Society statement: diagnosis and treatment of pulmonary disease in α_1 -antitrypsin deficiency. *Eur Respir J* 2017; 50: 1700610.
- 3 Rodríguez-Frias F, Miravittles M, Vidal R, *et al.* Rare α_1 -antitrypsin variants: are they really so rare? *Thorax* 2012; 6: 79–85.
- 4 López-Campos JL, Casas-Maldonado F, Torres-Duran M, *et al.* Results of a diagnostic procedure based on multiplex technology on dried blood spots and buccal swabs for patients with suspected α_1 -antitrypsin deficiency. *Arch Bronconeumol* 2021; 57: 42–50.
- 5 Blanco I, de Serres FJ, Fernandez-Bustillo E, *et al.* Estimated numbers and prevalence of PI*S and PI*Z alleles of α_1 -antitrypsin deficiency in European countries. *Eur Respir J* 2006; 27: 77–84.
- 6 Alvarez-Granda L, Cabero-Perez MJ, Bustamante-Ruiz A, *et al.* PI SZ phenotype in chronic obstructive pulmonary disease. *Thorax* 1997; 52: 659–661.
- 7 Seersholm N, Kok-Jensen A. Intermediate α_1 -antitrypsin deficiency PiSZ: a risk factor for pulmonary emphysema? *Respir Med* 1998; 92: 241–245.
- 8 McElvaney GN, Sandhaus RA, Miravittles M, *et al.* Clinical considerations in individuals with α_1 -antitrypsin PI*SZ genotype. *Eur Respir J* 2020; 55: 1902410.
- 9 Rare Diseases Task Force. Centres of Reference for Rare Diseases in Europe: State-of-the-art in 2006 and Recommendations of the Rare Diseases Task Force. www.ec.europa.eu/health/archive/ph_threats/non_com/docs/contribution_policy.pdf. Date last accessed: April 19, 2022.
- 10 Miravittles M, Nuñez A, Torres-Durán M, *et al.* the importance of reference centers and registries for rare diseases: the example of α_1 -antitrypsin deficiency. *COPD* 2020; 17: 346–354.
- 11 Miravittles M, Vidal R, Barros-Tizón JC, *et al.* Usefulness of a national registry of α_1 -antitrypsin deficiency. The Spanish experience. *Respir Med* 1998; 92: 1181–1187.

- 12 Miravittles M, Chorostowska-Wynimko J, Ferrarotti I, et al. The European Alpha-1 Research Collaboration (EARCO): a new ERS Clinical Research Collaboration to promote research in α_1 -antitrypsin deficiency. *Eur Respir J* 2019; 53: 1900138.
- 13 Brightling C, Genton C, Bill W, et al. ERS Clinical Research Collaborations: underpinning research excellence. *Eur Respir J* 2018; 52: 1801534.
- 14 Barrecheguren M, Torres-Duran M, Casas-Maldonado F, et al. Spanish implementation of the new international α_1 -antitrypsin deficiency international registry: The European Alpha-1 Research Collaboration (EARCO). *Arch Bronconeumol (Engl Ed)* 2021; 57: 81–82.
- 15 Blanco I, Bueno P, Diego I, et al. α_1 -antitrypsin Pi*SZ genotype: estimated prevalence and number of SZ patients worldwide. *Int J Chron Obstruct Pulmon Dis* 2017; 12: 1683–1694.
- 16 Greulich T, Altraja A, Barrecheguren M, et al. Protocol for the EARCO Registry: a pan-European observational study in patients with α_1 -antitrypsin deficiency. *ERJ Open Res* 2020; 6: 00181-2019.
- 17 Lara B, Blanco I, Martínez MT, et al. Spanish registry of patients with α_1 -antitrypsin deficiency: database evaluation and population analysis. *Arch Bronconeumol* 2017; 53: 13–18.
- 18 Torres-Durán M, Lopez-Campos JL, Barrecheguren M, et al. α_1 -antitrypsin deficiency: outstanding questions and future directions. *Orphanet J Rare Dis* 2018; 13: 114.
- 19 Barrecheguren M, Monteagudo M, Simonet P, et al. Diagnosis of α_1 -antitrypsin deficiency: a population-based study. *Int J Chron Obstruct Pulmon Dis* 2016; 11: 999–1004.
- 20 Esquinas C, Barrecheguren M, Sucena M, et al. Practice and knowledge about diagnosis and treatment of α_1 -antitrypsin deficiency in Spain and Portugal. *BMC Pulm Med* 2016; 16: 64.
- 21 López-Campos JL, Carrasco Hernandez L, Marquez-Martín E, et al. Diagnostic performance of a lateral flow assay for the detection of α_1 -antitrypsin deficiency. *Arch Bronconeumol* 2020; 56: 124–126.
- 22 Survival and FEV₁ decline in individuals with severe deficiency of α_1 -antitrypsin. The Alpha-1 Antitrypsin Deficiency Registry Study Group. *Am J Respir Crit Care Med* 1998; 158: 49–59.
- 23 Piitulainen E, Tanash HA. The clinical profile of subjects included in the Swedish National Register on individuals with severe α_1 -antitrypsin deficiency. *COPD* 2015; 12: Suppl. 1, 36–41.
- 24 Stockley RA. Antitrypsin deficiency assessment and programme for treatment (ADAPT): The United Kingdom registry. *COPD* 2015; 12: Suppl. 1, 63–68.
- 25 Chorostowska-Wynimko J, Struniawski R, Sliwinski P, et al. The national α_1 -antitrypsin deficiency registry in Poland. *COPD* 2015; 12: Suppl. 1, 22–26.
- 26 Miravittles M, Vilà S, Torrella M, et al. Influence of deficient α_1 -antitrypsin phenotypes on clinical characteristics and severity of asthma in adults. *Respir Med* 2002; 96: 186–192.
- 27 Piras B, Ferrarotti I, Lara B, et al. Clinical phenotypes of Italian and Spanish patients with α_1 -antitrypsin deficiency. *Eur Respir J* 2013; 42: 54–64.
- 28 Choate R, Mannino DM, Holm KE, et al. Comparing patients with ZZ versus SZ α_1 -antitrypsin deficiency: findings from AlphaNet's disease management program. *Chronic Obstr Pulm Dis* 2019; 6: 29–39.
- 29 Green CE, Vayalapa S, Hampson JA, et al. PiSZ α_1 -antitrypsin deficiency (AATD): pulmonary phenotype and prognosis relative to PiZZ AATD and PiMM COPD. *Thorax* 2015; 70: 939–945.
- 30 Greulich T, Albert A, Cassel W, et al. Opinions and attitudes of pulmonologists about augmentation therapy in patients with α_1 -antitrypsin deficiency. A survey of the EARCO Group. *Int J Chron Obst Pulm Dis* 2022; 17: 53–64.
- 31 Stockley RA, Miravittles M, Vogelmeier C. Augmentation therapy for α_1 -antitrypsin deficiency: towards a personalised approach. *Orphanet J Rare Dis* 2013; 8: 149.
- 32 Respreeza. Summary of Product Characteristics (SmPC). www.ec.europa.eu/health/documents/community-register/2015/20150820132594/dec_132594_es.pdf. Date last accessed: April 11, 2022.
- 33 Prolastin. Summary of Product Characteristics (SmPC). www.grifols.com/documents/10192/27754341/ft-prolastin-es-en/786cce6a-efb-41b6-867f-59d7a34ca086. Date last accessed: April 11, 2022.
- 34 Dahl M, Hersh CP, Ly NP, et al. The protease inhibitor Pi*S allele and COPD: a meta-analysis. *Eur Respir J* 2005; 26: 67–76.
- 35 Franciosi AN, Carroll TP, McElvaney NG. SZ α_1 -antitrypsin deficiency and pulmonary disease: more like MZ, not like ZZ. *Thorax* 2021; 76: 298–301.
- 36 Wood AM, Simmonds MJ, Bayley DL, et al. The TNF- α gene relates to clinical phenotype in α_1 -antitrypsin deficiency. *Respir Res* 2008; 9: 52.
- 37 Sucena M, Gomes J, Guimarães C, et al. Implementation of European Alpha-1 Research Collaboration (EARCO) in Portugal: the future starts now. *Pulmonology* 2020; 26: 181–183.
- 38 Bilton D, Caine N, Cunningham S, et al. Use of a rare disease patient registry in long-term post-authorisation drug studies: a model for collaboration with industry. *Lancet Respir Med* 2018; 6: 495–496.