



Published in final edited form as:

*Neurochem Res.* 2022 January ; 47(1): 148–162. doi:10.1007/s11064-021-03247-8.

## Pre-steady state kinetics and reverse transport in rat glutamate transporter EAAC1 with an immobilized transport domain

Jiali Wang,

Laura Zielewicz,

Yang Dong,

Christof Grewer

Department of Chemistry, Binghamton University, 4400 Vestal Parkway East, Binghamton, NY 13902

### Abstract

Plasma membrane glutamate transporters move glutamate across the cell membrane in a process that is thought to involve elevator-like movement of the transport domain relative to the static trimerization domain. Conformational changes associated with this elevator-like movement have been blocked by covalent crosslinking of cysteine pairs inserted strategically in several positions in the transporter structure, resulting in inhibition of steady-state transport activity. However, it is not known how these crosslinking restraints affect other partial reactions of the transporter that were identified based on pre-steady-state kinetic analysis. Here, we re-examine two different introduced cysteine pairs in the rat glutamate transporter EAAC1 recombinantly expressed in HEK293 cells, W440C/K268C and K64C/V419C, with respect to the molecular mechanism of their impairment of transporter function. Pre-steady-state kinetic studies of glutamate-induced partial reactions were performed using laser photolysis of caged glutamate to achieve sub-millisecond time resolution. Crosslinking of both cysteine pairs abolished steady-state transport current, as well as the majority of pre-steady-state glutamate-induced charge movements, in both forward and reverse transport mode, suggesting that it is not only the elevator-like movement associated with translocation, but also other transporter partial reactions that are inhibited. In contrast, sodium binding to the empty transporter, and glutamate-induced anion conductance were still intact after the W440C/K268C crosslink. Our results add to the previous mechanistic view of how covalent restraints of the transporter affect function and structural changes linked to individual steps in the transport cycle.

### Keywords

Glutamate transporter; crosslink; electrophysiology; kinetics; laser-photolysis

### Introduction

Plasma membrane glutamate transporters are responsible for transporting the major excitatory neurotransmitter, glutamate, across cell membranes [1–3]. Glutamate transporters take up glutamate into the cell against the glutamate concentration gradient, by co-transporting three sodium ions and one proton, and counter-transporting a potassium ion [1, 3–7]. The stoichiometry of transport is 3:1:1:1 ( $\text{Na}^+:\text{Glu}:\text{H}^+:\text{K}^+$ ), resulting in the net

movement of two positive charges into the cell for each transported glutamate molecule [1, 4, 8, 9]. During the translocation process, glutamate transporters also catalyze uncoupled anion currents that are proportional to glutamate transport activity [4, 10–14].

The structure of glutamate transporters was first studied using the archaeal homologue Glt<sub>ph</sub>, which shares 37% sequence identity and 55% similarity with human transporter subtype hEAAT2, excitatory amino acid transporter 2 [15]. The transporter is assembled as a homotrimer with three identical subunits working independently of each other [16–18]. Two major conformations (outward-facing and inward-facing states, OFS/IFS) of the glutamate transporter were discovered for Glt<sub>ph</sub> and resolved using x-ray crystallography [19, 20]. The inward-facing state was also identified by the Kanner group on the basis of structural modeling of inverted sequence repeats [24]. The generally-accepted molecular mechanism of glutamate transport is predicted to be based on successive steps, including extracellular substrate binding, translocation, and release to the cytosol [21, 22]. The translocation steps are associated with conformational changes of the transport domain, which functions in an elevator-based alternating access mechanism [23]. The trimerization domain, on the other hand, is thought to function as a static scaffold for the dynamic transport domain.

Transporter crosslinking studies, to restrict the transporter chemically to certain conformations, have been performed by several research groups [20, 24, 25]. By crosslinking the trimerization and transport domain between transmembrane domains TM5 and TM8, glutamate uptake was inhibited, likely by blocking the substrate translocation process. In addition, crosslinking was performed to stabilize the inward-facing conformation of Glt<sub>ph</sub> for structure determination. For the present crosslinking study, we chose to introduce disulfide bridges in two major conformations, inward-facing (IFS) and outward-facing (OFS) states. The inward-facing conformation was proposed to be stabilized by the crosslinked K55C/V364C double-cysteine mutant transporter, which crosslinks Glt<sub>ph</sub> between TM2 and HP2 [20, 24]. The outward-facing state was studied using the W440C/K268C pair, located in TM5 and TM8, following the research published by the Kanner group [25].

Here we use the laser-pulse photolysis method to perform rapid kinetic studies on glutamate transporters with the crosslinks described above. The results from these kinetic studies were compared with results from rapid solution exchange experiments at steady state. Laser-pulse photolysis experiments were performed with the glutamate transporter previously [26, 27], in which caged-glutamate was applied to cells, followed by photolysis of caged-glutamate initiated with a nanosecond laser pulse. Rapid transient current signals were recorded on a millisecond time scale, providing information about early steps in the transport cycle, such as glutamate and sodium binding. Here we applied the laser-photolysis method to glutamate transporters with known crosslinks to the transport domain, to analyze their kinetic behavior on a short time scale. Notably, the crosslinks abolish pre-steady state currents previously assigned to glutamate translocation and other pre-steady-state current components closely linked to it. However, Na<sup>+</sup> binding to the empty transporter does not appear to be affected by the crosslinks, suggesting that it does not require large-scale conformational changes of the transport domain.

## Results

### Positions of paired cysteine mutagenesis in EAAC1

Paired cysteine mutagenesis experiments were conducted using the rat glutamate transporter EAAC1 (excitatory amino acid carrier 1), in order to restrict movement of transporter domains relative to each other through covalent crosslinking, for example the transport domain relative to the trimerization domain, followed by rapid kinetic studies of transporter function. Positions were selected based on the  $\text{Glt}_{\text{ph}}$  structure and inspired by recent publications performing crosslinking in the human EAAC1 analog, EAAT3, and  $\text{Glt}_{\text{ph}}$  [19, 20, 24, 25] (Fig. 1 A to C). We constructed two double-cysteine mutants to positions in the transport domain and trimerization domain, W440C/K268C, in TM5 and TM8, and K64C/V419C, in TM2 and HP2. The K64C/V419C mutations were based on conserved residues located in  $\text{Glt}_{\text{ph}}$ , which, through cadmium complexation or the formation of a disulfide bond, crosslink the transporter into the inward-facing conformation [20] (Fig. 1C). The two introduced cysteine residues, located within TM2 and HP2 domains, are predicted to be 3.76 Å apart in the inward-facing conformation (Fig. 1C), while the distance is 21.6 Å in the outward-facing conformation (Fig. 1A), and 14.7 Å in the intermediate state (Fig. 1B). Another double-cysteine mutant, W440C/K268C, was chosen following the research published by the Kanner group [25]. After crosslinking, the W440C/K268C glutamate transporter is predicted to be restricted to sampling the outward-facing state and intermediate state, with distances ranging from 3.5 Å to 7.3 Å between the two residues (Fig. 1 A, B). However, the cysteine pair's distance increased to 14.9 Å when the transport domain was in the inward-facing conformation. From previous results, these two double-cysteine mutants were predicted to be close enough to form disulfide bonds under oxidizing conditions. In previous results, preincubation of HELA cells expressing W440C/K268C double mutant with 1 μM CuPh or CdCl<sub>2</sub> resulted in an inhibition of D-[<sup>3</sup>H]aspartate transport by 60–70%, and over 95% inhibition when using 8μM crosslinking reagent [25].

### Crosslinking results in inhibition of steady-state anion currents

We first tested the activity of the EAAC1<sub>cysless</sub> transporter and the two double-cysteine mutant transporters described in the last paragraph, before and after crosslinking. These experiments were performed in the presence of extracellular Na<sup>+</sup>/glutamate and intracellular K<sup>+</sup>, i.e. the forward transport mode, in which the amino acid substrate is transported across the membrane to the intracellular side. Anion current was observed upon application of glutamate to EAAC1<sub>cysless</sub>, as shown in Fig. 2A. These currents were inwardly directed due to the outflow of internal SCN<sup>-</sup>, caused by the population of an anion conducting state located along the translocation pathway [4, 11, 26, 28]. The glutamate dose response relationship for this current is shown in Fig. 2B with an apparent affinity of  $K_m = 13.4 \pm 2.5 \mu\text{M}$ , a value that is only slightly larger than that of the wild type transporter (Fig. 2C). These results show that the cysless transporter is functional with kinetic properties close to the wild type transporter. The two double-cysteine-mutant transporters were also active (dose response curves shown in Figs. 2C and D), although with slightly smaller glutamate-induced anion currents compared to EAAC1<sub>cysless</sub> (Fig. 2F). However, this reduction in current was not statistically significant. The apparent  $K_m$  values for glutamate of the two mutant transporters are shown in Fig. 2C, with only a minor difference in apparent

glutamate affinity between EAAC1<sub>WT</sub> ( $8.1 \pm 1.5\mu\text{M}$ ), EAAC1<sub>Cysless</sub> ( $13.4 \pm 2.5\mu\text{M}$ ), EAAC1<sub>K64C/V419C</sub> ( $16.3 \pm 3.0\mu\text{M}$ ), and EAAC1<sub>K268C/W440C</sub> ( $25.6 \pm 4.0\mu\text{M}$ ), summarized in Fig. 2E.

As demonstrated in previous studies, crosslinking of two cysteine residues in glutamate transporters can be achieved by either  $\text{Cd}^{2+}$  or copper phenanthroline (CuPh) application [25]. In order to observe the full inhibition effect, we applied  $400\ \mu\text{M}$  CuPh, a higher value compared to the previous studies [25], with or without  $400\ \mu\text{M}$   $\text{CdCl}_2$  in the pipette (intracellular side application) with similar results. An incubation time of 15 minutes with external  $400/400\ \mu\text{M}$  CuPh/ $\text{CdCl}_2$  resulted in an inhibition of glutamate-induced anion current activity by 74% for EAAC1<sub>K268C/W440C</sub> and over 95% for EAAC1<sub>K64C/V419C</sub> (Fig. 2D), respectively. Therefore, EAAC1<sub>K64C/V419C</sub> showed a slightly more pronounced crosslinking effect on the current. Moreover, cells expressing EAAC1<sub>Cysless</sub> were also tested as a control using the same concentration of crosslinking reagent and showed no inhibition effect (Fig. 2D). We also tested the effect of crosslinking on the leak anion conductance. DL-TBOA (DL-*threo*-beta-benzyloxyaspartate), a high affinity glutamate transporter inhibitor was used to block the leak anion conductance. The results shown in Fig. 2D demonstrate that  $\text{Na}^+$ -induced leak anion currents were inhibited by crosslinking of both double mutant transporters, as evidenced by the lack of outward current upon application of TBOA. In contrast, TBOA induced outward current in EAAC1<sub>Cysless</sub> due to outflow of  $\text{SCN}^-$  being blocked by TBOA (Fig. 2D). While these data do not fully exclude the possibility that TBOA does not bind to the transporters after crosslinking, this explanation is less likely because glutamate can still bind to the crosslinked transporter, and TBOA can block  $\text{Na}^+$ -induced transient currents in voltage jump experiments (see below).

We also determined the voltage dependence of the currents to test the crosslink effect. Under ionic condition favoring the forward transport mode, and with  $\text{SCN}^-$  in the pipette, currents before crosslinking were increased by negative transmembrane potential, as expected for glutamate induced anion current caused by  $\text{SCN}^-$  outflow (Figs. 3A and C). Current-voltage relationships (I-V curves) are shown in Figs. 3B and D. Currents were strongly inhibited after crosslinking for both double mutant transporters, although the effect was more complete for EAAC1<sub>K64C/V419C</sub>. Background currents were subtracted by application of  $400\ \mu\text{M}$  TBOA. We attempted to reverse the crosslinking effect by using DTT after current recording in the crosslinked state, with limited success. The reason is that it is more difficult to do long term experiments in patch-clamped cells, as compared to other experimental methods, due to the cells having a limited lifetime when attached to the pipette in the whole cell recording mode.

### **Pre-steady state currents in the forward transport mode are inhibited in the crosslinked state**

EAAC1 pre-steady state currents, in response to rapid application of glutamate, have been described previously [26, 27, 29]. These currents provide important information about early steps in the glutamate transport cycle. Therefore, we measured glutamate-induced pre-steady state currents in the crosslinked transporters, to test which reaction steps are affected by the covalent modification. The time course of pre-steady-state kinetics of EAAC1 currents was

recorded using ionic conditions that favor observation of the transport current, i.e. inside  $K^+$ , outside  $Na^+$  and replacement of  $Cl^-$  by the non-permeant methanesulfonate ion [26, 29]. In the cysless mutant transporter, a glutamate concentration jump induced by laser-pulse photolysis of caged glutamate resulted in a large, inwardly-directed transient current (Fig. 4A), which decayed to a steady state with two-exponential behavior, as described previously for the EAAC1 wild-type transporter [26, 29], and for other EAATs using fast solution exchange methods [28, 30–32]. The transient current fully decayed to the steady state within 25 milliseconds, in analogy to previous results with wild-type EAAC1, indicating that the pre-steady state kinetics are not dramatically affected by introducing the five Cys to Ser mutations.

After introducing the double mutants in the cysless background, both pre-steady state and steady-state currents under forward transport conditions were reduced in K64C/V419C and K268C/W440C mutant transporters before crosslinking, with the K268C/W440C transporters showing a more significant reduction in currents (Figs. 4B and C). This reduction could be caused by reduced expression levels of the double mutant transporters, slowed kinetics and turnover rate, or disulfide bridge formation before CuPh application. In fact, the latter was shown to be the case for the partial reduction of the current in the K268C/W440C mutant transporter [25]. To test for expression levels, we performed immunostaining (Suppl. Fig. 1). Immunofluorescence of the double mutant transporters did not appear to be significantly altered compared to the cysless transporter. After crosslinking with CuPh, both pre-steady-state currents (Figs. 4D and E) and steady-state currents (Fig. 4F) were further reduced, as expected by the crosslink immobilizing the transport domain.

While the pre-steady-state currents in EAAC1<sub>K268C/W440C</sub> could not be quantitatively evaluated before and after crosslinking, due to the small amplitudes, the time constants in EAAC1<sub>K64C/V419C</sub> of the transient current decay, as well as the amplitude ratio between the two decaying phases were virtually unchanged after crosslinking. This indicates that the crosslink was not 100% complete, and some transporter (<30%) remained in the active configuration. Despite the inability to fully block transient current by the crosslink in EAAC1<sub>K64C/V419C</sub>, it is clear that all phases of the transient current are severely affected by the immobilization of the transport domain. This is expected for locking the transporter in the inward-facing conformation, in which external glutamate cannot induce the conformational changes triggered by  $Na^+$  binding to the glutamate-bound transporter, and translocation, which were proposed to be responsible for the pre-steady state charge movement in wild-type EAAC1.

### **Crosslinks eliminate EAAC1 reverse transport current and glutamate release**

$K^+$ -induced relocation of the glutamate-free transporter is an important step in the glutamate transport cycle. Using forward transport conditions, as in the previous paragraphs, intracellular  $K^+$  catalyzes this relocation reaction, to re-expose glutamate binding sites to the extracellular side after translocation is complete. Next, we reversed the ionic gradients across the membrane and used application of extracellular  $K^+$  in the reverse transport mode (in the presence of intracellular  $Na^+$  and glutamate) to test the properties of mutations together with crosslinking on the relocation steps in the reverse transport direction. In

Fig. 5A, a typical steady-state reverse transport current recorded from cells expressing EAAC1<sub>cysless</sub> is shown. As expected [6, 22, 27, 30, 32, 33], this current is outwardly directed due to the coupled outward movement of two positive charges for each K<sup>+</sup> ion taken up. The outward currents were dependent on the K<sup>+</sup> concentration, as reported previously [30]. The K<sup>+</sup> apparent affinity of EAAC1<sub>cysless</sub> was determined as  $5.1 \pm 1.2$  mM, which was in same range as that for EAAC1<sub>WT</sub> ( $3.4 \pm 0.6$  mM) [30].

Next, we performed laser-photolysis experiments, using caged-glutamate inside the recording pipet, to study the rapid kinetics of the reverse translocation process. Here, the extracellular solution contained 140 mM K<sup>+</sup>, and the intracellular solution 140 mM Na<sup>+</sup> and 5 mM caged-glutamate, which diffused into the cytosol within 5 mins after forming the whole cell mode [27]. As shown previously for the wild-type transporter, outwardly-directed currents were observed upon photolysis in EAAC1<sub>cysless</sub>, as illustrated in Fig. 5C. These currents demonstrated a rapidly decaying outward transient component  $\tau = 4.49 \pm 0.84$  ms, which settled to the steady state with a damped oscillation. In contrast, the EAAC1<sub>K64C/V419C</sub> currents showed a larger initial peak relative to the steady-state current, with slower decay kinetics  $\tau = 9.01 \pm 2.34$  ms and no oscillation (Fig. 5D). This indicates that the steady-state turnover rate in the reverse direction is already slowed by the double-mutation before crosslinking, see also Fig. 5H for a summary of steady-state currents. After crosslinking, both steady-state and transient current components were virtually eliminated (Fig. 5F). It was previously proposed that the transient current in the reverse transport direction is caused by the electrogenic reverse translocation process. This assignment is consistent with the data shown in Fig. 5F. In addition, the data indicate that charge movement from partial reaction steps other than translocation may also be impaired by the crosslinking reaction.

For EAAC1<sub>K268C/W440C</sub> pre-steady state currents in the reverse transport mode were very small, before and after crosslinking (Figs. 5E and G). This could be caused by the transporter being significantly crosslinked already before CuPh application, as described above. Overall, the data suggest that this double mutation severely impairs any early reactions occurring in the reverse transport reaction after glutamate application, possibly even glutamate binding.

We also studied K<sup>+</sup>-driven glutamate efflux to test the mutant transporters' function and crosslink effects using iGluSnFR extracellular glutamate binding sensor [34] (Fig.6). The fluorescent sensor was co-expressed with rat EAAC1 wild type or mutant DNA in HEK293 cells. We applied varying external K<sup>+</sup> concentrations and measured resulting glutamate release via fluorescence intensity changes. An increase in fluorescence indicates a change from non-bound state to bound state of glutamate to iGluSnFR sensor. First, 10 mM DTT in 140 mM NaMes was used to wash solutions and the original glutamate efflux was measured upon application of 140 mM KMes, by measuring the increase in fluorescence intensity (Fig. 6A). Cells expressing wild type EAAC1 showed a relative fluorescence increase of  $1.70 \pm 0.06$  -fold, and cells expressing the cysless transporter DNA responded similarly ( $1.55 \pm 0.34$ -fold increase). Both cysteine double-mutant transporters showed much less glutamate efflux under the same conditions (K64C/V419C,  $0.39 \pm 0.05$ ; K268C/W440C,  $0.27 \pm 0.08$ ). This data confirms the results from current recordings, namely that the

cysteine double-mutant transporters may be less effective at transporting glutamate even before crosslinking. Fig. 6B shows the result for the relative change in glutamate efflux for each transporter after crosslinking reagent was applied. The change in glutamate efflux resulting from crosslinking was calculated relative to the original efflux data. Even though the cysteine double-mutants originally showed much less glutamate efflux, they also had a great reduction in efflux as a result of crosslinking, especially when CuPh was used as the crosslinking reagent. EAAC1<sub>K64C/V419C</sub> and EAAC1<sub>K268C/W440C</sub> showed only  $17.9 \pm 8.5\%$  and  $17.4 \pm 11.0\%$  of the original glutamate efflux, respectively, after crosslinking with CuPh. Under the same conditions, the wild type and cysless transporters showed about the same glutamate efflux as they had originally (wild type =  $102 \pm 1\%$ ; cysless =  $103 \pm 1\%$  efflux maintained). The effect of CdCl<sub>2</sub> crosslinking reagent, however, was not as pronounced. The double-cysteine mutants maintained  $68 \pm 5\%$  (K64C/V419C) and  $89 \pm 2\%$  (K268C/W440C) of the original efflux after crosslinking with CdCl<sub>2</sub> (wild type =  $90 \pm 1\%$ ; cysless =  $104 \pm 6\%$ ). Example images of cells expressing EAAC1<sub>K64C/V419C</sub> and iGluSnFR before and after crosslinking with CuPh are shown in Fig. 6C. Before crosslinking, the cells showed a fluorescence increase of 43% with induction by K<sup>+</sup>. After crosslinking, the same cells showed only a fluorescence increase of 3%, indicating that crosslinking reduced the glutamate efflux of these cells by about 94%.

### No effect of crosslinking to Na<sup>+</sup> binding to the empty transporter

It was previously shown that voltage jumps to the glutamate-free transporter result in Na<sup>+</sup>-dependent transient charge movement [35, 36]. When applying step changes to the membrane potential for a period of 40 ms (voltage protocol shown in Fig.7A, top), transient currents were observed, which were sensitive to 400 μM TBOA (used to correct for non-specific transient currents). In the absence of CuPh, these charge movements are shown in (Figs.7A to C). Results after crosslinking are shown in Figs. 7D to F. voltage jump-induced charge movements were preserved in both K64C/V419C and K268C/W440C mutant transporters after crosslinking. We next integrated the transient current signal and plotted charge-voltage relationships, shown in Figs. 7C and F. The Q-V relationships were virtually unchanged by the crosslinks, suggesting that Na<sup>+</sup> binding and release steps in the glutamate-free transporter still function in the immobilized K64C/V419C and K268C/W440C mutant transporters. Control experiments in non-transfected cells are shown in Fig. 7G, demonstrating no charge movement.

## Discussion

In this work, we extended crosslinking studies previously performed on glutamate transporters, to identify the effect of restricting conformational freedom of the transport domain on rapid kinetics and early reactions associated with the substrate transport process. We focused on crosslinking of two disulfide pairs, K268C/W440C, which was first generated by the Kanner group [25], and K64C/V419C, which was shown to stabilize the inward-facing conformation in Glt<sub>Ph</sub> [20, 37]. Other cysteine pairs were also published to show successful crosslinking, based on EAAT1 and GLT-1. In these studies, EAAT1 [38–40] TM2-HP1, TM4-TM7 and TM4-TM8 were covalently linked, in addition to GLT-1 [41, 42] TM5-TM8, TM2-TM4 interactions. In these positions, double cysteine mutants

showed significant crosslinking effects, with oxidizing conditions largely inhibiting the radiolabeled substrate uptake. In particular, a pair of cysteines highly sensitive to oxidation were identified in EAAT1 between TM4 and TM8, with low concentrations of CuPh at 600 nM leading to a complete inhibition effect [40].

In extension of previous studies, we applied the laser-photolysis method with caged glutamate, allowing the observation of early intermediates in the transport cycle by generating glutamate concentration jumps with a sub-millisecond time resolution. In wild-type and cysless glutamate transporters, charge movement in response to glutamate concentration jumps manifests itself in a pre-steady-state component that allows the differentiation of three phases, which can be separated on the time scale (Fig. 4A) [29]. These three phases were previously interpreted to be tightly associated with  $\text{Na}^+/\text{Glu}^-$  binding and translocation steps [29] (i.e. elevator-like movement of the transport domain relative to the membrane normal). The major result of the present study is that it is not only the steady-state transport that is blocked by the application of CuPh, consistent with previous publications, but that the pre-steady state charge movement in the forward transport direction is also largely inhibited (Fig. 4). This result may be expected for the K64C/V419C crosslink, which was proposed to lock the transporter in the inward-facing state [20]. Thus, forward translocation is unlikely to be possible after external application of glutamate. However, the K268C/W440C crosslink also prevented transient charge movement. While EAAC1<sub>K268C/W440C</sub> was much less active than the cysless transporter even before crosslinking, we attribute this effect to intrinsic crosslinking before CuPh application, as has been proposed in [25]. In any case, these results suggest that it is not only the translocation reaction itself, but also a conformational change associated glutamate-induced  $\text{Na}^+$  binding, potentially the formation of the intermediate outward-facing state, that are blocked by the covalent restriction of TM8 relative to TM5.

In the reverse transport mode, both crosslinks virtually eliminated steady-state transport current, and significantly reduced glutamate release through reverse transport (Fig. 6). This result is expected because the immobilization of the transport domain most likely prevents the conformational changes necessary for glutamate release. In addition, very little pre-steady-state charge movement was observed in response to glutamate concentration jumps to the intracellular side of the membrane. For the K268C/W440C crosslink, this lack of charge movement is most likely caused by the inability of intracellular glutamate to bind to the transporter locked in the outward-facing conformation. However, for the K64C/V419C crosslink, the results are more interesting. It was previously shown by Boudker and colleagues that aspartate and  $\text{Na}^+$  can bind to  $\text{Glt}_{\text{ph}}$  with the analogous crosslink [20], which locks the transporter in the inward-facing configuration. This is consistent with the ability of glutamate transporter substrates to bind to brain membranes at low temperatures, when translocation is inhibited (i.e. an immobilized transport domain) [43]. If this is the case, the absence of glutamate-induced transient currents indicate that any of these binding reactions are almost electroneutral, or only weakly electrogenic. Our group had previously proposed that intracellular  $\text{Na}^+$  binding to the empty transporter is electrogenic [27]. This intracellular  $\text{Na}^+$  binding process could be responsible for the small, outwardly-directed charge movement that remains in the K64C/V419C crosslinked transporter (Fig. 5F). Overall, these reverse transport results suggest a strong inhibition of most early reaction



steps in the reverse transport process in response to glutamate binding, including the reverse translocation step.

Interestingly, the glutamate-induced anion current was not fully abolished by oxidative crosslinking in the K268C/W440C mutant transporter (Fig. 2D). This result is consistent with a previous report by the Kanner group, suggesting that the crosslink still allows the anion-conducting state to be populated [25]. One possibility, which was already proposed by Kanner and colleagues [25] could be that the anion conductance is closely linked to the outward-facing intermediate state, which has been previously identified in structural studies [17]. This idea is consistent with proposals by the Fahlke group [11] that the anion conducting state is structurally similar, and in rapid equilibrium with the outward-facing intermediate state. This suggestion would also support previous rapid kinetic studies, in which the anion conducting state was placed along the translocation pathway [29]. While the distance between the two cysteine residues is larger in the intermediate state than in the outward-facing conformation (Fig. 1), it may still allow transient formation of this state, and, thus, anion conductance to function. In contrast, TBOA was unable to significantly block the leak anion conducting state after crosslinking the K268C/W440C mutant transporter (Fig. 2D). This suggests that the leak anion conducting state adopts a different structure than the glutamate-dependent anion conducting state. For the K64C/V419C crosslink, neither glutamate-induced anion conductance, nor TBOA-blockable leak anion conductance was observed (Fig. 2D), consistent with the idea that extracellularly-applied substrates or inhibitors cannot interact with the inward-facing configuration.

Finally, we tested for transient currents in response to voltage jumps in the empty, glutamate-free transporter (Fig.7). It was previously proposed, by us and others [29, 44, 45], that these transient currents are caused by voltage-dependent  $\text{Na}^+$  binding to the extracellular binding site. These transient currents can be isolated from non-specific current components by block with specific, competitive EAAT inhibitors, for example TBOA and kainate [44]. TBOA-sensitive transient currents were observed in crosslinked transporters with both K268C/W440C and K64C/V419C cysteine pairs (Fig.7). For EAAC1<sub>K268C/W440C</sub>, this results suggest that the  $\text{Na}^+$  binding events to the empty transporter are still functional after crosslinking TM5 and TM8. Most likely, the electrogenic  $\text{Na}^+$  binding reaction is not associated with major conformational changes of the transporter, or with structural changes that do not involve movement of TM5 relative to TM8. This conclusion is consistent with structural results from the *apo* and  $\text{Na}^+$  bound Glt<sub>TK</sub> homologue. Here, major structural changes upon  $\text{Na}^+$  binding were only observed in the unwound and highly-conserved NMDGT region of TM7, but not in the relative positioning between transport and trimerization domain [46]. For the transporter with the K64C/V419C crosslink, the results are more difficult to explain. If this crosslink results in an inward-facing state, it should be insensitive to extracellularly-applied TBOA. However, the TBOA-sensitive transient currents are substantial in size. One possibility is that, because of the high TBOA concentration used (about 1000 times  $K_m$ ), the transporter inward-facing-outward-facing equilibrium can be pulled to the outward facing state, despite the crosslink, although this would seemingly require a large distortion of the transporter structure (Fig. 1).

## Conclusions

In this study, we tested K268C/W440C (between TM5 and TM8) and K64C/V419C (between TM2 and HP2) double-cysteine mutant transporters with respect to the effect of crosslinking on partial reactions in the transport cycle. From analysis of pre-steady-state currents in response to rapid glutamate concentration jumps in both forward and reverse transport mode, it was concluded that the crosslinks impair most pre-steady-state processes previously assigned to the glutamate translocation process, as well as Na<sup>+</sup>-binding to the glutamate-bound form of the transporter, and conformational changes associated with it. Therefore, the crosslinks seem to severely restrain conformational flexibility of the transporter, which is necessary for these early reactions in the transport cycle to occur. In contrast, Na<sup>+</sup> binding to the glutamate-free form of the transporter, as well as to a minor extent the glutamate-induced anion conductance, were still operational, at least in the K268C/W440C mutant transporter. Overall, our results add to the previous mechanistic view of how covalent restraints of the transporter affect function and individual reaction steps in the transport cycle.

## Experimental Section

### Cell Culture and Transfection

HEK293 cells (American Type Culture Collection No. CRL 1573) were cultured as described previously[23]. Cell cultures were transiently transfected with wild-type or mutant EAAC1 cDNAs, inserted into a modified pBK-CMV-expression plasmid, using Jetprime transfection reagent according to the protocol supplied by the manufacturer (Polyplus). The cells were used for electrophysiological measurements between 24 and 36 h after transfection. The cysless EAAC1 variant contained Cys to Ser substitutions in positions 9, 158, 218, 255 and 342 and was generated using the QuikChange mutagenesis procedure (Agilent), followed by sequencing. K268C/W440C and K64C/V419C double-mutant transporter DNAs were generated within the EAAC1<sub>cysless</sub> background. The K268C/W440C transporter cDNA was a generous gift provided by Dr. Baruch Kanner. K64C/V419C mutations were generated by Gene Universal Co. Ltd.

### Crosslinking procedures

The solutions of CuPh (Copper 1,10-Phenanthroline) and CdCl<sub>2</sub> were prepared as 10 mM stock solutions in NMG-MES (NMG: N-Methylglucaminegluconate, MES: Methanesulfonate) solution and diluted to 400 μM concentration with external buffer solution (see below) prior to each experiment. DTT (Dithiothreitol) was prepared as a 20 mM stock solutions in NMG-MES solution. Cells were incubated in crosslink or protective solutions at room temperature for 15 mins before electrophysiology experiments were started.

### Solution exchange and laser photolysis

Rapid solution exchange (time resolution 100–200 ms) was performed by means of a quartz tube (opening diameter 350 μm) positioned at a distance of 0.5 mm to the cell. The linear flow rate of the solutions emerging from the opening of the tube was 5–10 cm/s. Laser-pulse

photolysis experiments were performed as described in detail previously [47]. In the reverse transport experiments, MNI-caged glutamate were introduced into the cell through the glass recording electrode. After whole-cell mode was established, a period of 5 mins was allowed for diffusive equilibration with the cell interior. MNI-caged glutamate [48, 49], at concentrations of 1 mM (140 mM and 49 mM Na<sup>+</sup>) to 4 mM (18 mM and 10 mM Na<sup>+</sup>), or free glutamate were applied to the cells and photolysis of the caged glutamate was initiated with a light flash (355nm, 8ns, Minlite II, Continuum). The light was coupled into a quartz fiber (diameter 365 μm) that was positioned in front of the cell in a distance of 300 μm. With maximum light intensities of 500–840 mJ/cm<sup>2</sup> saturating glutamate concentrations could be released, which was tested by comparison of the steady-state current with that generated by rapid perfusion of the same cell with 1 mM caged-glutamate with 140 mM Na<sup>+</sup>. Data were recorded using the pClamp6 software (Axon Instruments), digitized with a sampling rate of 1 kHz (solution exchange) or 25 kHz (laser photolysis and voltage jump) and low-pass filtered at 250 Hz or 3–10 kHz, respectively.

### Electrophysiology

Currents associated with glutamate transporters were measured in the whole-cell current recording configuration. Whole-cell currents were recorded with an EPC7 patch-clamp Amplifier (ALA Scientific, Westbury, NY) under voltage-clamp conditions. The resistance of the recording electrode was 3–6 MΩ. Series resistance was not compensated because of the relatively small whole-cell currents carried by EAAC1, in both anion and transport modes. The composition of the solutions for measuring transport currents or laser-photolysis experiments was: 140 mM NaMes, 2 mM MgGluconate<sub>2</sub>, 2 mM CaMes<sub>2</sub>, 10 mM 4-(2-Hydroxyethyl)piperazine-1-ethanesulfonic acid (HEPES), 10 mM glutamate, pH 7.3 (extracellular), 130 mM KMes (Mes = methanesulfonate), 2 mM MgGluconate<sub>2</sub>, 5 mM Ethylene glycol-bis(2-aminoethylether)-*N,N,N',N'*-tetraacetic acid (EGTA), 10 mM HEPES, pH 7.3 (intracellular). For anion current recordings, intracellular Mes was replaced by SCN<sup>-</sup>. For reverse transport current recordings, KMes are using as external solution, 130 mM NaMes and 5 mM Glutamate as internal solution, as published previously [23, 30, 32].

### Voltage jumps

Voltage jumps (–100 to +60mV) were applied to perturb the electrogenic glutamate translocation equilibrium. To determine EAAC1-specific currents, control currents were recorded in the presence of 400 μM extracellular TBOA (DL-threo-β-benzyloxyaspartic acid) and subtracted from the glutamate-induced currents. Capacitive transient compensation and series resistance compensation of up to 80% was employed using the EPC-7 amplifier. Non-specific transient currents were subtracted in Clampfit software (Molecular Devices).

### Cellular glutamate efflux measured by extracellular glutamate-binding sensor

EAAC1 transporter was co-expressed with iGluSnFR [34, 50] extracellular glutamate-binding sensor in HEK293 cells as a system for measuring glutamate efflux via fluorescence intensity changes. The sensor is bound to the outside of the cell membrane and contains a glutamate-binding component that responds rapidly to extracellular glutamate concentration changes. The external glutamate concentration is reported via fluorescent intensity change.

This method specifically isolates the transport function of EAAC1 in response to various external conditions and eliminates the variable of anion conductance.

Imaging experiments were performed as described previously [44]. All imaging experiments utilized a live-cell flow-through imaging chamber (Warner Instruments, Series 20) together with an inverted fluorescence microscope (Zeiss Axiovert 25). The fluorescence filter set used was FITC, which was obtained from Omega Filters. Bath buffer solution contained NaMes (Mes = methanesulfonate) and KMes equaling a combined concentration of 140 mM, 2 mM CaGluconate<sub>2</sub>, 2 mM MgGluconate<sub>2</sub>, 10 mM 4-(2-Hydroxyethyl) piperazine-1-ethanesulfonic acid (HEPES), pH 7.3 using NaOH. 140 mM NaMes solution was used to initially wash the remaining culturing medium from the cell surface and also in between each test solution to act as a benchmark for changes in fluorescence. Solutions were passed through the imaging chamber for 30–45 seconds before recording an image. For each experiment, the exposure and other imaging features were set as constant and for an appropriate fluorescence intensity below saturation. Images were recorded after each solution exchange and then analyzed using ImageJ software [51]. ImageJ was used to quantify the fluorescence intensity of 5–10 cells per image. The loci were held constant when comparing images in an experiment. The relative fluorescence change ( $\Delta F/F$ ) was calculated as follows:

$$\frac{\Delta F}{F} = \frac{F_{\text{final}} - F_{\text{initial}}}{F_{\text{initial}}} \quad (\text{eq. 1})$$

Where  $F_{\text{initial}}$  was the fluorescence intensity of the image taken after passing the initial solution directly prior to the test solution, and  $F_{\text{final}}$  was the fluorescence intensity of the same point in the image taken following the test solution.

### Data Analysis

All data are shown as mean  $\pm$  SD, collected from recordings of 8 to 10 cells. For statistical analysis, paired two-tailed t tests were used in Microsoft Excel software. To determine  $K_m$  values, non-linear curve fitting was used with a Michaelis-Menten-like equation,  $y = I_{\text{max}} \cdot [\text{Glu}] / (K_m + [\text{Glu}])$  built in the Origin software least-squares-fitting package. Mean values were compared by using two tailed student's t-test results. These results were calculated using Microsoft Excel.

Nonlinear regression fits of laser-pulse photolysis experimental results were performed in Clampfit software (Axon Instruments) by the use of the following equations. The pre-steady-state transport currents and (in the absence of SCN<sup>-</sup>) were fitted with a sum of three exponential functions and a stationary current component:  $I = I_1 \cdot \exp(-t/\tau_{\text{rise}}) + I_2 \cdot \exp(-t/\tau_{\text{decay1}}) + I_3 \cdot \exp(-t/\tau_{\text{decay2}}) + I_{\text{SS}}$ . Here,  $I$  is the current amplitude,  $\tau$  the time constant, and  $t$  the time.

### Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgements


This work was supported by a National Institutes of Health (NIH), Grant 1 R15 GM135843–01 awarded to CG.

## References

- Zerangue N, Kavanaugh MP (1996) Flux coupling in a neuronal glutamate transporter. *Nature* 383:634–637 [PubMed: 8857541]
- Tanaka K, Watase K, Manabe T, Yamada K, Watanabe M, Takahashi K, Iwama H, Nishikawa T, Ichihara N, Kikuchi T, Okuyama S, Kawashima N, Hori S, Takimoto M, Wada K (1997) Epilepsy and exacerbation of brain injury in mice lacking the glutamate transporter GLT-1. *Science* 276:1699–1702 [PubMed: 9180080]
- Pines G, Kanner BI (1990) Counterflow of L-glutamate in plasma membrane vesicles and reconstituted preparations from rat brain. *Biochemistry* 29:11209–11214 [PubMed: 1980217]
- Wadiche JI, Amara SG, Kavanaugh MP (1995) Ion fluxes associated with excitatory amino acid transport. *Neuron* 15:721–728 [PubMed: 7546750]
- Kanner BI, Sharon I (1978) Active transport of L-glutamate by membrane vesicles isolated from rat brain. *Biochemistry* 17:3949–3953 [PubMed: 708689]
- Kavanaugh MP, Bendahan A, Zerangue N, Zhang Y, Kanner BI (1997) Mutation of an amino acid residue influencing potassium coupling in the glutamate transporter GLT-1 induces obligate exchange. *J Biol Chem* 272:1703–1708 [PubMed: 8999849]
- Zhang Y, Bendahan A, Zarbiv R, Kavanaugh MP, Kanner BI (1998) Molecular determinant of ion selectivity of a (Na<sup>+</sup> + K<sup>+</sup>)-coupled rat brain glutamate transporter. *Proc Natl Acad Sci U S A* 95:751–755 [PubMed: 9435265]
- Zerangue N, Kavanaugh MP (1996) Interaction of L-cysteine with a human excitatory amino acid transporter. *J Physiol* 493 ( Pt 2):419–423 [PubMed: 8782106]
- Levy LM, Warr O, Attwell D (1998) Stoichiometry of the glial glutamate transporter GLT-1 expressed inducibly in a Chinese hamster ovary cell line selected for low endogenous Na<sup>+</sup>-dependent glutamate uptake. *J Neurosci* 18:9620–9628 [PubMed: 9822723]
- Bergles DE, Tzingounis AV, Jahr CE (2002) Comparison of coupled and uncoupled currents during glutamate uptake by GLT-1 transporters. *J Neurosci* 22:10153–10162 [PubMed: 12451116]
- Machtens JP, Kortzak D, Lansche C, Leinenweber A, Kilian P, Begemann B, Zachariae U, Ewers D, de Groot BL, Briones R, Fahlke C (2015) Mechanisms of anion conduction by coupled glutamate transporters. *Cell* 160:542–553 [PubMed: 25635461]
- Fairman WA, Vandenberg RJ, Arriza JL, Kavanaugh MP, Amara SG (1995) An excitatory amino-acid transporter with properties of a ligand-gated chloride channel. *Nature* 375:599–603 [PubMed: 7791878]
- Arriza JL, Eliasof S, Kavanaugh MP, Amara SG (1997) Excitatory amino acid transporter 5, a retinal glutamate transporter coupled to a chloride conductance. *Proc Natl Acad Sci U S A* 94:4155–4160 [PubMed: 9108121]
- Leary GP, Stone EF, Holley DC, Kavanaugh MP (2007) The glutamate and chloride permeation pathways are colocalized in individual neuronal glutamate transporter subunits. *J Neurosci* 27:2938–2942 [PubMed: 17360916]
- Yernool D, Boudker O, Jin Y, Gouaux E (2004) Structure of a glutamate transporter homologue from *Pyrococcus horikoshii*. *Nature* 431:811–818 [PubMed: 15483603]
- Grewer C, Balani P, Weidenfeller C, Bartusel T, Tao Z, Rauen T (2005) Individual subunits of the glutamate transporter EAAC1 homotrimer function independently of each other. *Biochemistry* 44:11913–11923 [PubMed: 16128593]
- Verdon G, Boudker O (2012) Crystal structure of an asymmetric trimer of a bacterial glutamate transporter homolog. *Nat Struct Mol Biol* 19:355–357 [PubMed: 22343718]
- Koch HP, Brown RL, Larsson HP (2007) The glutamate-activated anion conductance in excitatory amino acid transporters is gated independently by the individual subunits. *J Neurosci* 27:2943–2947 [PubMed: 17360917]

19. Boudker O, Ryan RM, Yernool D, Shimamoto K, Gouaux E (2007) Coupling substrate and ion binding to extracellular gate of a sodium-dependent aspartate transporter. *Nature* 445:387–393 [PubMed: 17230192]
20. Reyes N, Ginter C, Boudker O (2009) Transport mechanism of a bacterial homologue of glutamate transporters. *Nature* 462:880–885 [PubMed: 19924125]
21. Bendahan A, Armon A, Madani N, Kavanaugh MP, Kanner BI (2000) Arginine 447 plays a pivotal role in substrate interactions in a neuronal glutamate transporter. *J Biol Chem* 275:37436–37442 [PubMed: 10978338]
22. Kanner BI, Bendahan A (1982) Binding order of substrates to the sodium and potassium ion coupled L-glutamic acid transporter from rat brain. *Biochemistry* 21:6327–6330 [PubMed: 6129891]
23. Wang J, Albers T, Grewer C (2018) Energy Landscape of the Substrate Translocation Equilibrium of Plasma-Membrane Glutamate Transporters. *J Phys Chem B* 122:28–39 [PubMed: 29218993]
24. Crisman TJ, Qu S, Kanner BI, Forrest LR (2009) Inward-facing conformation of glutamate transporters as revealed by their inverted-topology structural repeats. *Proc Natl Acad Sci U S A* 106:20752–20757 [PubMed: 19926849]
25. Shabaneh M, Rosental N, Kanner BI (2014) Disulfide cross-linking of transport and trimerization domains of a neuronal glutamate transporter restricts the role of the substrate to the gating of the anion conductance. *J Biol Chem* 289:11175–11182 [PubMed: 24584931]
26. Grewer C, Watzke N, Wiessner M, Rauen T (2000) Glutamate translocation of the neuronal glutamate transporter EAAC1 occurs within milliseconds. *Proc Natl Acad Sci U S A* 97:9706–9711 [PubMed: 10931942]
27. Zhang Z, Tao Z, Gameiro A, Barcelona S, Braams S, Rauen T, Grewer C (2007) Transport direction determines the kinetics of substrate transport by the glutamate transporter EAAC1. *Proc Natl Acad Sci U S A* 104:18025–18030 [PubMed: 17991780]
28. Wadiche JI, Kavanaugh MP (1998) Macroscopic and microscopic properties of a cloned glutamate transporter/chloride channel. *J Neurosci* 18:7650–7661 [PubMed: 9742136]
29. Watzke N, Bamberg E, Grewer C (2001) Early intermediates in the transport cycle of the neuronal excitatory amino acid carrier EAAC1. *J Gen Physiol* 117:547–562 [PubMed: 11382805]
30. Wang J, Zielewicz L, Grewer C (2019) A K(+)/Na(+) co-binding state: Simultaneous versus competitive binding of K(+) and Na(+) to glutamate transporters. *J Biol Chem*
31. Otis TS, Jahr CE (1998) Anion currents and predicted glutamate flux through a neuronal glutamate transporter. *J Neurosci* 18:7099–7110 [PubMed: 9736633]
32. Wang J, Zhang K, Goyal P, Grewer C (2020) Mechanism and potential sites of potassium interaction with glutamate transporters. *J Gen Physiol* 152
33. Szatkowski M, Barbour B, Attwell D (1990) Non-vesicular release of glutamate from glial cells by reversed electrogenic glutamate uptake. *Nature* 348:443–446 [PubMed: 2247147]
34. Marvin JS, Borghuis BG, Tian L, Cichon J, Harnett MT, Akerboom J, Gordus A, Renninger SL, Chen TW, Bargmann CI, Orger MB, Schreiter ER, Demb JB, Gan WB, Hires SA, Looger LL (2013) An optimized fluorescent probe for visualizing glutamate neurotransmission. *Nat Methods* 10:162–170 [PubMed: 23314171]
35. Grewer C, Zhang Z, Mwaura J, Albers T, Schwartz A, Gameiro A (2012) Charge compensation mechanism of a Na<sup>+</sup>-coupled, secondary active glutamate transporter. *J Biol Chem* 287:26921–26931 [PubMed: 22707712]
36. Wadiche JI, Arriza JL, Amara SG, Kavanaugh MP (1995) Kinetics of a human glutamate transporter. *Neuron* 14:1019–1027 [PubMed: 7748550]
37. Ryan RM, Mitrovic AD, Vandenberg RJ (2004) The chloride permeation pathway of a glutamate transporter and its proximity to the glutamate translocation pathway. *J Biol Chem* 279:20742–20751 [PubMed: 14982939]
38. Zhang W, Zhang X, Qu S (2019) Substrate-Induced Motion between TM4 and TM7 of the Glutamate Transporter EAAT1 Revealed by Paired Cysteine Mutagenesis. *Mol Pharmacol* 95:33–42 [PubMed: 30348896]


39. Zhang Y, Zhang X, Qu S (2014) Cysteine mutagenesis reveals alternate proximity between transmembrane domain 2 and hairpin loop 1 of the glutamate transporter EAAT1. *Amino Acids* 46:1697–1705 [PubMed: 24692063]
40. Qu S, Zhang W, He S, Zhang X (2019) Paired-Cysteine Scanning Reveals Conformationally Sensitive Proximity between the TM4b-4c Loop and TM8 of the Glutamate Transporter EAAT1. *ACS Chem Neurosci* 10:2541–2550 [PubMed: 30802031]
41. Rong X, Tan F, Wu X, Zhang X, Lu L, Zou X, Qu S (2016) TM4 of the glutamate transporter GLT-1 experiences substrate-induced motion during the transport cycle. *Sci Rep* 6:34522 [PubMed: 27698371]
42. Zhang X, Qu S (2011) Proximity of transmembrane segments 5 and 8 of the glutamate transporter GLT-1 inferred from paired cysteine mutagenesis. *PLoS One* 6:e21288 [PubMed: 21698173]
43. Aprico K, Beart PM, Crawford D, O'Shea RD (2004) Binding and transport of [3H](2S,4R)-4-methylglutamate, a new ligand for glutamate transporters, demonstrate labeling of EAAT1 in cultured murine astrocytes. *J Neurosci Res* 75:751–759 [PubMed: 14994336]
44. Zielewicz L, Wang J, Ndaru E, Grewer CT (2019) Transient Kinetics Reveal Mechanism and Voltage Dependence of Inhibitor and Substrate Binding to Glutamate Transporters. *ACS Chem Biol*
45. Mennerick S, Shen W, Xu W, Benz A, Tanaka K, Shimamoto K, Isenberg KE, Krause JE, Zorumski CF (1999) Substrate turnover by transporters curtails synaptic glutamate transients. *J Neurosci* 19:9242–9251 [PubMed: 10531428]
46. Guskov A, Jensen S, Faustino I, Marrink SJ, Slotboom DJ (2016) Coupled binding mechanism of three sodium ions and aspartate in the glutamate transporter homologue GltTk. *Nat Commun* 7:13420 [PubMed: 27830699]
47. Niu L, Wieboldt R, Ramesh D, Carpenter BK, Hess GP (1996) Synthesis and characterization of a caged receptor ligand suitable for chemical kinetic investigations of the glycine receptor in the 3-microseconds time domain. *Biochemistry* 35:8136–8142 [PubMed: 8679565]
48. Canepari M, Nelson L, Papageorgiou G, Corrie JE, Ogden D (2001) Photochemical and pharmacological evaluation of 7-nitroindolyl- and 4-methoxy-7-nitroindolyl-amino acids as novel, fast caged neurotransmitters. *J Neurosci Methods* 112:29–42 [PubMed: 11640955]
49. Wieboldt R, Gee KR, Niu L, Ramesh D, Carpenter BK, Hess GP (1994) Photolabile precursors of glutamate: synthesis, photochemical properties, and activation of glutamate receptors on a microsecond time scale. *Proc Natl Acad Sci U S A* 91:8752–8756 [PubMed: 8090718]
50. Zielewicz L, Grewer C (2019) Genetically Encoded Halide Sensor-Based Fluorescent Assay for Rapid Screening of Glutamate Transport and Inhibition. *ACS Sens* 4:2358–2366 [PubMed: 31393114]
51. Arena ET, Rueden CT, Hiner MC, Wang S, Yuan M, Eliceiri KW (2017) Quantitating the cell: turning images into numbers with ImageJ. *Wiley Interdiscip Rev Dev Biol* 6




**Figure 1: Predicted locations of the introduced cysteine pairs in homologue structure *Glt<sub>ph</sub>* in three different conformations.**

The upper panel shows cartoon representation of *Glt<sub>ph</sub>* structures (PDB IDs:2nwx,3v8g,3kbc) in the outward-facing state (A), the outward-facing intermediate state (B), and the inward facing state (C). Three identical subunits are colored as blue, dark green, and light green. The blue-colored subunit (left), illustrates the conformational change. The lower panels illustrate the transport domain in pink and the trimerization domain in yellow. The mutated residues are highlighted and labeled. Distances were calculated based on atom K64(CE), V419(CA), K268(CB), W440(CA) and are labeled in Ångstroms. All images were generated using PyMOL software. The amino acids in parentheses represent EAAT3 numbering.






**Figure 2. Double-cysteine mutations block the anion conductance under oxidizing conditions.** Whole-cell current recording experiments with EAAC1<sup>Cysless</sup> were performed under ionic conditions favoring the observation of the anion conductance, using the permeant anion SCN<sup>-</sup>: 130 mM KSCN in the pipette solution, 140 mM NaMes in the extracellular solution. Extracellular glutamate was applied at the time indicated by the grey bar at V = 0 mV. (A) Typical anion current recordings induced by 10/50/200 μM glutamate. (B, C, D) Dose response curve to determine the EAAC1<sup>Cysless</sup>, EAAC1<sup>K64C/V419C</sup> and EAAC1<sup>K268C/W440C</sup> glutamate apparent affinities. The solid line is a fit to a Michaelis-Menten-like equation. (E) Glutamate K<sub>m</sub> values for wild-type and mutant transporters before crosslinking: EAAC1<sub>wt</sub> K<sub>m</sub> = 8.1 ± 1.5 μM (n=18), EAAC1<sup>Cysless</sup> K<sub>m</sub> = 13.4 ± 2.5 μM (n=17), EAAC1<sup>K64C/V419C</sup> K<sub>m</sub> = 16.3 ± 3.0 μM (n=9) and EAAC1<sup>K268C/W440C</sup> K<sub>m</sub> = 25.6 ± 4 μM (n=12). (F) Maximum anion currents (at saturating glutamate concentration) under forward transport conditions before and after crosslinking. Crosslink reagents were used at the concentration of 400 μM/400 μM CuPh and CdCl<sub>2</sub> from extracellular and intracellular side, respectively. Student's t-test analysis indicated the crosslink effects were significant for K268C/W440C, *p* < 0.00001, and for K64C/V419C, *p* < 0.0000001. For all experiments in the presence of TBOA, the inhibitor was added after crosslinking.




**Figure 3: Glutamate-induced anion current is inhibited by crosslinking of double-mutant transporters.**

(A and C) Voltage-dependent EAAC1<sub>K64C/V419C</sub> and EAAC1<sub>K268C/W440C</sub> anion currents were activated by extracellular application of 1 mM glutamate in the presence of the intracellular permeant anion SCN<sup>-</sup>. The extracellular solution contained 140 mM NaMES, the intracellular solution contained 130 mM KSCN, forward transport conditions. The Voltage jump protocol is show as inset (see *Materials and Methods*). The membrane potential range was -100 mV to +60 mV. (B and D) EAAC1<sub>K64C/V419C</sub> and EAAC1<sub>K268C/W440C</sub> anion current-voltage relationship at steady-state before (black, n=7 and 11) and after crosslinking (red, n=17 and 15). Background currents were obtained by application of 200  $\mu$ M TBOA and subtraction.



**Figure 4: Crosslinking reduces glutamate-induced transient charge movement under forward-transport conditions.**

(A) Current recording after photolytic liberation of glutamate from 1 mM MNI-caged glutamate at the time indicated by the flash symbol in EAAC1<sub>Cysless</sub>-expressing HEK293 cells. Photolysis was initiated with a 355 nm laser flash (400 mJ/cm<sup>2</sup>). Pre-steady-state currents were measured under forward-transport conditions in the absence of permeant anion, with 140 mM NaMES in the extracellular solution, and 130 mM KMes in the intracellular solution. 1mM MNI-caged-glutamate were pre-equilibrated with the cell for 0.5s prior to the laser flash. (B and C) Similar experiment with EAAC1<sub>K64C/V419C</sub> and EAAC1<sub>K268C/W440C</sub> before crosslinking. (D and E) Laser photolysis experiment with EAAC1<sub>K64C/V419C</sub> and EAAC1<sub>K268C/W440C</sub> after crosslinking. (F) Summary of steady-state current amplitude in response to laser-photolysis of 1 mM caged glutamate. The membrane potential was 0 mV in all experiments. Student's t-test analysis indicated the crosslink effect was significant for K268C/W440C,  $p=0.0002$ , and for K64C/V419C,  $p<0.0001$ .



**Figure 5: Inhibition of reverse transport and charge movements by crosslinking.** (A) Typical reverse transport currents of EAAC1<sub>cysless</sub> (in the absence of permeant anion) after application of extracellular K<sup>+</sup> (indicated by grey bar). The intracellular solution contained 130 mM NaMES and 5 mM glutamate. (B) Determination of K<sup>+</sup> apparent affinity from the dose response curve fitted with a Michaelis-Menten-like equation (red line). EAAC1<sub>cysless</sub> K<sub>m</sub> for K<sup>+</sup> was 5.07 ± 1.2mM. (C) Reverse transport laser photolysis experiment with EAAC1<sub>cysless</sub>. The time of the laser pulse is indicated with the flash symbol. The extracellular solution contained 140 mM K<sup>+</sup>, the intracellular solution contained 130 mM NaMES and 5 mM MNI-caged glutamate. (D and E) Similar experiment with EAAC1<sub>K64C/V419C</sub> (n=35) and EAAC1<sub>K268C/W440C</sub> (n=20) before crosslinking. (F and G) Laser photolysis experiment with EAAC1<sub>K64C/V419C</sub> (n=12) and EAAC1<sub>K268C/W440C</sub>


(n=26) after crosslinking. **(H)** Comparison of steady-state reverse transport currents before and after crosslinking. The membrane potential was 0 mV in all experiments. Student's t-test results indicated the crosslink effect was significant for K268C/W440C,  $p=0.01$ , and for K64C/V419C,  $p<0.000000001$

Author Manuscript

Author Manuscript


Author Manuscript

Author Manuscript



**Figure 6: K64C/V419C, K268C/W440C mutant transporters show significant decrease in glutamate efflux after crosslinking.**

iGluSnFr glutamate fluorescent sensor and EAAC1 transporter DNA (wild type, cysless, K64C/V419C or K268C/W440C) were co-expressed in HEK293 cells. **(A)** Relative glutamate efflux from cells (as measured by fluorescent sensor) induced by K<sup>+</sup> without the presence of crosslinking reagents (wild type= 1.70±0.06 (n=4); cysless= 1.55±0.34 (n=2); K64C/V419C= 0.39±0.05 (n=7); K268C/W440C= 0.27±0.08 (n=7)). Student's t-test analysis indicate significant difference compared to wild type for K64C/V419C  $p < 0.00000001$  and K268C/W440C  $p < 0.00000001$ . **(B)** Percent of original fluorescence intensity maintained after crosslinking with CuPh (wild type= 102 ± 1 % (n=3); cysless= 103 ± 1 % (n=2); K64C/V419C= 18 ± 9 % (n=5); K268C/W440C= 17 ± 11 % (n=5)) and CdCl<sub>2</sub> (wild type= 90 ± 1 % (n=3); cysless= 104 ± 6 % (n=2); K64C/V419C= 68 ± 5 % (n=4); K268C/W440C= 89 ± 2 % (n=4)). T-test results indicate significant difference compared to wild type for K64C/V419C  $p < 0.001$  and K268C/W440C  $p < 0.001$  for CuPh and K64C/V419C  $p < 0.05$  for CdCl<sub>2</sub>. **(C)** Example images for the determination of glutamate efflux in EAAC1<sub>K64C/V419C</sub>. The top panel shows one example of original efflux data for a selection of cells whereby the fluorescence intensity increased by 43% upon application of external K<sup>+</sup>. After crosslinking with 400 μM CuPh (bottom panel), the efflux of these cells decreased to only a 2.6% change with application of K<sup>+</sup>, which is an overall 94% decrease in efflux after crosslinking.



**Figure 7: Crosslinked transporters show transient current after voltage jumps in the absence of glutamate.**

A voltage-jump protocol shown in the inset of (A) was used to isolate  $\text{Na}^+$ -induced transient currents in the glutamate-free transporter in the absence of permeant anion. (A and B) Original transient current traces before crosslinking in K64C/V419C ( $n=7$ ) and K268C/W441C ( $n=8$ ) mutant transporters. Unspecific currents were subtracted using  $400\mu\text{M}$  TBOA. (D and E) Similar experiments as in (A and B) but after crosslinking ( $n=6$  and 5). (C and F) Charge movement as a function of the membrane potential obtained by integrating the transient current signals. The intracellular solution contained 130 mM KMes, the extracellular solution contained 140 mM NaMes. (G) Result of the same voltage jump experiment performed in non-transfected cells.