

CORRECTIONS

Corrigendum to Performance and meat quality traits of slow-growing chickens stimulated in ovo with galactooligosaccharides and exposed to heat stress [Poultry Science Volume 101, Issue 8, August 2022, 101972]

S. Tavaniello,^{*} A. Slawinska,[†] F. Sirri,[‡] M. Wu,^{*} D. De Marzo,[§] M. Siwek,[#] and G. Maiorano^{*,1}

^{*}Department of Agricultural, Environmental and Food Sciences, University of Molise, 86100 Campobasso, Italy;

[†]Faculty of Biological and Veterinary Sciences, Nicolaus Copernicus University in Toruń, 87-100 Toruń, Poland;

[‡]Department of Agricultural and Food Sciences, Alma Mater Studiorum - University of Bologna, 40064 Ozzano dell'Emilia, Italy; [§]Department of DETO, Section of Veterinary Science and Animal Production, University of Bari Aldo Moro, 70010 Valenzano, Italy; and [#]Department of Animal Biotechnology and Genetics, Politechnika Bydgoska, 85-084 Bydgoszcz, Poland

2022 Poultry Science 101:102080

<https://doi.org/10.1016/j.psj.2022.102080>

The authors regret that there was an error in Table 1 diet composition. The corrected Table 1 is shown here. The authors would like to apologize for any inconvenience caused.

Table 1. Composition of the diet supplied to the birds of all the experimental groups.

	Period		
	Starter (0–14 d)	Grower (15–36 d)	Finisher (37–50 d)
<i>Dietary components (Kg)</i>			
Corn	42.17	34.96	12.73
White corn	0.00	0.00	15.00
Wheat	10.00	20.00	25.01
Sorghum	0.00	0.00	5.00
Soybean meal	23.11	20.63	17.60
Expanded soybean	10.00	10.00	13.00
Sunflower	3.00	3.00	3.00
Corn gluten	4.00	3.00	0.00
Soybean oil	3.08	4.43	5.48
Dicalcium phosphate	1.52	1.20	0.57
Calcium carbonate	0.91	0.65	0.52
Sodium bicarbonate	0.15	0.10	0.15
Salt	0.27	0.27	0.25
Coline chloride	0.10	0.10	0.10
Lysine sulfate	0.59	0.55	0.46
DL-methionine	0.27	0.29	0.30
Threonine	0.15	0.14	0.14
Enzyme-roxazyme G2g	0.08	0.08	0.08
Phytase 0.1%	0.10	0.10	0.10
Vitamin-mineral premix ¹	0.50	0.50	0.50
<i>Calculated nutrient content</i>			
DM, %	88.57	88.65	88.64
CP, %	22.70	21.49	19.74
Lipid, %	7.06	8.24	9.74
Fiber, %	3.08	3.04	3.07
Ash, %	5.85	5.17	4.49
Lysine, %	1.38	1.29	1.21
Methionine, %	0.67	0.62	0.59
Methionine+cysteine, %	1.03	0.97	0.91
Calcium, %	0.91	0.80	0.59
Phosphate, %	0.63	0.57	0.46
ME, Kcal/Kg	3.076	3.168	3.264

¹Provided the following per kg of diet: vitamin A (retinyl acetate), 13,000 IU; vitamin D3 (cholecalciferol), 4,000 IU; vitamin E (DL- α -tocopheryl acetate), 80 IU; vitamin K (menadione sodium bisulfite), 3mg; riboflavin, 6 mg; panthothenic acid, 6 mg; niacin, 20 mg; pyridoxine, 2 mg; folic acid, 0.5 mg; biotin, 0.10 mg; thiamine, 2.5 mg; vitamin B12 20 μ g; Mn, 100 mg; Zn, 85 mg; Fe, 30 mg; Cu, 10 mg; I, 1.5 mg; Se, 0.2 mg; ethoxyquin, 100 mg.

© 2022 The Authors. Published by Elsevier Inc. on behalf of Poultry Science Association Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

DOI of original article: <http://dx.doi.org/10.1016/j.psj.2022.101972>

¹Corresponding author: maior@unimol.it