



New epidemiological trends of respiratory syncytial virus bronchiolitis during COVID-19 pandemic

Emanuele Castagno¹ · Irene Raffaldi¹ · Francesco Del Monte² · Silvia Garazzino³ · Claudia Bondone¹

Received: 27 May 2022 / Accepted: 15 September 2022 / Published online: 26 September 2022
© Children's Hospital, Zhejiang University School of Medicine 2022

Restrictive measures adopted worldwide against COVID-19 (e.g., wearing masks, frequent hand washing, and social distancing) produced not only a decrease in SARS-CoV-2 diffusion but also a drastic reduction in respiratory diseases caused by other pathogens, as reported by many authors [1, 2]. The spread of respiratory viruses, particularly respiratory syncytial virus (RSV), has been completely changed during the winter of 2020–2021 [3, 4], but an unusual resurgence of RSV infection has been recently reported in different countries [3–7]. A drastic reduction in bronchiolitis hospitalizations also occurred during the winter of 2020–2021, with RSV being the main etiologic agent of bronchiolitis [3, 6].

We described the unusual RSV bronchiolitis outbreak in young infants in autumn 2021 and the difference in RSV bronchiolitis epidemiology during the COVID-19 pandemic season (March 2020–September 2021) compared to previous seasons.

We retrospectively analyzed all infants aged <6 months with acute bronchiolitis admitted to the Pediatric Emergency Department (PED) of our tertiary teaching children's hospital from October to November 2021 compared to those admitted during the same period from 2017 to 2020. Data about demographics, triage priority codes, oxygen supplementation, and outcomes were collected. The diagnosis of RSV infection was made by RNA detection on a nasal swab. The study was approved by our local ethics

committee. Informed consent to participate in the study was obtained from participants' parents or legal guardians. Statistical analysis was performed using R software (version 4.0.2). The results were reported as medians and interquartile ranges (IQRs) or frequencies and percentages, as appropriate. All tests were two-sided; P value < 0.05 was considered significant.

Comparing October–November 2021 to October–November 2017–2020, we observed an impressive outbreak of acute bronchiolitis in the first six months of life, especially due to RSV (159 vs. 1; $P < 0.001$) but also a lower hospital admission rate ($P = 0.001$) and shorter length of hospitalization ($P = 0.016$). No significant difference was observed for the need of oxygen supplementation ($P = 0.120$) or admission rate to Pediatric Intensive Care Unit (PICU) ($P = 0.999$), suggesting similar severity (Table 1). To address the great number of patients coming to the PED, a temporary short observation unit with six beds was opened in November 2021, and 9.2% of patients were moved to other hospitals due to a lack of beds. We did not observe any outbreaks of RSV bronchiolitis during spring and summer 2021 (Fig. 1) or a higher percentage of prematurity and PICU admissions, as otherwise reported by other authors [6].

Our analysis confirmed previous observations of different new epidemiological trends of RSV bronchiolitis worldwide during the COVID-19 pandemic [1, 5, 8]. In New Zealand, the incidence rate during the bronchiolitis peak in 2021 was higher than the average of peaks in 2015–2019 but without more severe cases [8]. Hatter et al. hypothesized that in countries with a higher COVID-19 burden in the Northern Hemisphere, the RSV epidemic might have been more severe during the winter season of 2021 than in the prepandemic years due to the immunity debt secondary to long periods of low exposure to RSV [8]. This prediction has come true in our area in autumn 2021, when school, kindergarten, and work social activities fully reprised. On the other hand, we have not observed any summer peak, as reported in Japan and Spain

✉ Emanuele Castagno
ecastagno@cittadellasalute.to.it

¹ Present Address: Department of Pediatric Emergency, Regina Margherita Children's Hospital - A.O.U. Città Della Salute e della Scienza di Torino, Piazza Polonia 94, 10126 Turin, Italy


² Department of Public Health and Pediatrics, University of Turin, Turin, Italy

³ Pediatric Infectious Disease Unit, Regina Margherita Children's Hospital - A.O.U. Città della Salute e della Scienza di Torino, Piazza Polonia 94, 10126 Turin, Italy

Table 1 Epidemiological and clinical features of infants with acute bronchiolitis

Variables	October–November 2021	October–November 2017–2020	<i>P</i> value
Total admissions for acute bronchiolitis, <i>n</i> (%)	174 (100.0)	18 (100.0)	
RSV positive, <i>n</i> (%)	159 (91.3)	1 (5.5)	<0.001
Males, <i>n</i> (%)	107 (61.5)	9 (5.2)	0.448
Prematurity, <i>n</i> (%)	21 (12.1)	5 (2.9)	0.075
Median age, d, <i>n</i> (IQR)	58 (34.0–95.5)	69.5 (32.7–116.2)	0.212
Triage priority code, <i>n</i> (%)			
Red (emergency)	30 (17.2)	5 (27.8)	0.332
Yellow (high priority)	128 (73.6)	13 (72.2)	0.999
Green (low priority)	16 (9.2)	0 (0.0)	0.999
White (very low priority)	0 (0.0)	0 (0.0)	0.999
Need for respiratory support, <i>n</i> (%)	111 (63.8)	15 (83.3)	0.120
Low-flow nasal cannulae, <i>n</i> (%)	109 (62.6)	7 (4.0)	
High-flow nasal cannulae, <i>n</i> (%)	55 (31.6)	10 (5.7)	
NIV/cPAP, <i>n</i> (%)	7 (4.0)	1 (0.6)	
Mechanical ventilation, <i>n</i> (%)	3 (1.7)	0 (0.0)	
Admission to hospital, <i>n</i> (%)	107 (61.5)	18 (100.0)	0.001
Admission to PICU, <i>n</i> (%)	12 (6.8)	1 (5.5)	0.999
Median length of hospitalization, d, <i>n</i> (IQR)	5 (1.0–9.0)	8 (6.0–11.5)	0.016
Moved to other Hospitals, <i>n</i> (%)	16 (9.2)	0 (0.0)	

RSV respiratory syncytial virus, IQR interquartile range, NIV noninvasive ventilation, cPAP continuous positive air pressure, PICU pediatric intensive care unit

**Fig. 1** Seasonal trends of respiratory syncytial virus bronchiolitis between 2017 and 2022

[5, 6]. We fully agree with Bermúdez Barrezueta et al. about the necessity of potentiating the epidemiological surveillance systems that control RSV circulation for a better management of health resources, including the better timing of palivizumab prophylaxis [6].

In conclusion, we observed an unusual early RSV outbreak in autumn 2021, suggesting a possible change in the epidemiology of the virus during the COVID-19 pandemic. Despite the noticeable increase, no more severe cases were recorded compared to 2017–2020.

Author contributions IR and CE contributed to conceptualization, formal analysis, and writing –original draft. MF was involved in data curation and writing–original draft. BC and GS contributed to writing–original draft, and writing–review and editing.

Funding No funding was received for this paper.

Data availability The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Declarations

Conflict of interest No financial or non-financial benefits have been received or will be received from any party related directly or indirectly to the subject of this article.

Ethical approval The study was approved by our local Ethical Committee (approval number: 0049894, May 3, 2022). Informed consent to participate in the study has been obtained from participants' parents or legal guardians.

References

1. Agca H, Akalin H, Saglik I, Hacimustafaoglu M, Celebi S, Ener B. Changing epidemiology of influenza and other respiratory viruses in the first year of COVID-19 pandemic. *J Infect Public Health*. 2021;14:1186–90.
2. Di Mattia G, Nenna R, Mancino E, Rizzo V, Pierangeli A, Villani A, et al. During the COVID-19 pandemic where has respiratory syncytial virus gone? *Pediatr Pulmonol*. 2021;56:3106–9.
3. Van Brusselen D, De Troeyer K, ter Haar E, Vander Auwera A, Poschet K, Van Nuijs S, et al. Bronchiolitis in COVID-19 times: a nearly absent disease? *Eur J Pediatr*. 2021;180:1969–73.
4. Vittucci AC, Piccioni L, Coltella L, Ciarlito C, Antilici L, Bozzola E, et al. The disappearance of respiratory viruses in children during the COVID-19 pandemic. *Int J Environ Res Public Health*. 2021;18:9550.
5. Ujiie M, Tsuzuki S, Nakamoto T, Iwamoto N. Resurgence of respiratory syncytial virus infections during COVID-19 pandemic, Tokyo. *Japan Emerg Infect Dis*. 2021;27:2969–70.
6. Bermúdez Barreza L, Matías Del Pozo V, López-Casillas P, Brezmes Raposo M, Gutiérrez Zamorano M, Pino Vázquez MA. Variation in the seasonality of the respiratory syncytial virus during the COVID-19 pandemic. *Infection*. 2022;22:1–5.
7. Cason C, Zamagni G, Cozzi G, Tonegutto D, Ronfani L, Oretti C, et al. Spread of respiratory pathogens during the COVID-19 pandemic among children in the northeast of Italy. *Front Microbiol*. 2022;24:804700.
8. Hatter L, Eathorne A, Hills T, Bruce P, Beasley R. Respiratory syncytial virus: paying the immunity debt with interest. *Lancet Child Adolesc Health*. 2021;5:e44–5.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.