

Review



Beneficial Role of Vitamin D on Endothelial Progenitor Cells (EPCs) in Cardiovascular Diseases

Atanu Sen ,¹ Vinnyfred Vincent ,² Himani Thakkar ,² Ransi Abraham ,¹ Lakshmy Ramakrishnan ¹

¹Department of Cardiac Biochemistry, All India Institute of Medical Sciences, New Delhi, India

²Department of Biochemistry, All India Institute of Medical Sciences, New Delhi, India

OPEN ACCESS

Received: Feb 4, 2022

Revised: Mar 23, 2022

Accepted: Apr 4, 2022

Published online: Apr 29, 2022

Correspondence to

Lakshmy Ramakrishnan

Department of Cardiac Biochemistry, All India Institute of Medical Sciences, Ansari Nagar, New Delhi, Delhi 110029, India.
Email: lakshmy_ram@yahoo.com

Copyright © 2022 The Korean Society of Lipid and Atherosclerosis.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Atanu Sen
<https://orcid.org/0000-0002-5023-6260>

Vinnyfred Vincent
<https://orcid.org/0000-0002-4268-0951>

Himani Thakkar
<https://orcid.org/0000-0002-2825-5190>

Ransi Abraham
<https://orcid.org/0000-0003-1954-3112>

Lakshmy Ramakrishnan
<https://orcid.org/0000-0002-8582-3705>

Funding

None.

ABSTRACT

Cardiovascular diseases (CVDs) are the leading cause of death in the world. Endothelial progenitor cells (EPCs) are currently being explored in the context of CVD risk. EPCs are bone marrow derived progenitor cells involved in postnatal endothelial repair and neovascularization. A large body of evidence from clinical, animal, and *in vitro* studies have shown that EPC numbers in circulation and their functionality reflect endogenous vascular regenerative capacity. Traditionally vitamin D is known to be beneficial for bone health and calcium metabolism and in the last two decades, its role in influencing CVD and cancer risk has generated significant interest. Observational studies have shown that low vitamin D levels are associated with an adverse cardiovascular risk profile. Still, Mendelian randomization studies and randomized control trials (RCTs) have not shown significant effects of vitamin D on cardiovascular events. The criticism regarding the RCTs on vitamin D and CVD is that they were not designed to investigate cardiovascular outcomes in vitamin D-deficient individuals. Overall, the association between vitamin D and CVD remains inconclusive. Recent clinical and experimental studies have demonstrated the beneficial role of vitamin D in increasing the circulatory level of EPC as well as their functionality. In this review we present evidence supporting the beneficial role of vitamin D in CVD through its modulation of EPC homeostasis.

Keywords: Endothelial progenitor cell; Cardiovascular diseases; Vitamin D

INTRODUCTION

Differentiation of cells from mesodermal origin to angioblast and subsequently to endothelial cells (ECs) was thought to exclusively take place during embryonic development. But this dogma was overturned in 1997 when Asahara et al.¹ reported the existence of endothelial progenitor cells (EPCs) as putative ECs in adult human peripheral blood and their role in neovascularization in postnatal life. This cell type has become of significant interest in regenerative medicine due to their potential role in repair of endothelium in cardiovascular disease (CVD).

Conflict of Interest

The authors have no conflicts of interest to declare.

Author Contributions

Conceptualization: Sen A; Data curation: Sen A, Vincent V, Thakkar H; Formal analysis: Sen A, Vincent V, Thakkar H; Resources: Ramakrishnan L; Supervision: Ramakrishnan L; Validation: Sen A; Visualization: Vincent V, Thakkar H; Writing - original draft: Sen A; Writing - review & editing: Sen A, Vincent V, Thakkar H, Abraham R, Ramakrishnan L.

Deficiency of vitamin D is a common public health problem and unfortunately most often remain unrecognized and untreated. Exposure of ultraviolet radiation of sunlight to skin tissues is considered as one of the major source of vitamin D, while deficiency of this vitamin is associated with indoor lifestyle, sun avoidance strategies, darker skin, distance from equator and winter season.²⁻⁴ Vitamin D is well known for its role in mineral homeostasis and bone health. Since last 2 decades clinical and epidemiological studies have shed light on circulatory vitamin D deficiency and its association with the increased risk of CVDs and its risk factors including diabetes, hypertension etc.⁵ Recent clinical, *in vitro* and *in vivo* studies have also shown that the vitamin D plays a pivotal role in vascular health by promoting cell proliferation, cell-cell adhesions, barrier integrity and mobilization of EPCs.⁶⁻⁹ Since EPCs are known to be important in postnatal neovasculogenesis and repair of the damaged endothelium with significant role in prevention of cardiovascular disorders, individuals with reduced EPCs level or impaired functions would benefit from vitamin D based interventions.¹⁰

In this review we will discuss the prevailing knowledge about the role of vitamin D on EPCs in context to CVD.

EPCs IN CVD

1. EPC

Asahara et al.¹ reported for the first time in 1997 the existence of bone marrow derived EPCs in peripheral circulation. In the past, regeneration of damaged endothelium was attributed to the migration and proliferation of adjacent mature ECs but it is now recognized that local and circulating EPCs have a crucial role in repair and regeneration of the damaged endothelium and in neovascularization.¹ In response to various stimulatory factors including angiogenic growth factors, ischemia, vascular trauma, and various cytokines, EPCs can be mobilized from the bone marrow to the sites of injury to play a role in re-endothelization and neovascularization.¹¹

EPCs are characterized by the surface expression of vascular endothelial growth factor receptor-2 (VEGFR-2); also known as kinase insert domain receptor (KDR), and progenitor cell surface markers (CD34, CD133, and CXCR4).¹² However, the definition and characterization of the true EPC population on the basis of cellular markers are still debated, as EPCs are a heterogenous population of cells originating from precursors within the bone marrow and are present in different stages of endothelial differentiation in the peripheral blood.¹³ It was believed that the differentiation of mesodermal cells to angioblasts and subsequent endothelial differentiation occurs exclusively during embryonic development, but now we know that CD34⁺ progenitor cell population from adults can also give rise to the endothelial phenotype. Homing of EPCs from the bone marrow to sites of cardiovascular injury occurs in response to variety of cytokines including stromal derived factor (SDF)-1 α that attracts binding of progenitors expressing the CXCR4 receptor, granulocyte colony stimulating factor (GS-CSF), granulocyte macrophage colony stimulating factor (GM-CSF), vascular endothelial growth factor (VEGF) and nitric oxide (NO). Downregulation of VEGF and other cytokines and reduced bioavailability of NO contribute to impaired migratory capacity of EPC.^{14,18}

Emerging evidence suggests that EPCs show two different morphologies during *in vitro* culture, the spindle shaped early EPCs and late EPCs with cobble stone morphology. Similar




Fig. 1. Bright field microscopic images of early (A) and late (B) EPCs with spindle shape and typical cobble stone morphology. Both the images are at 10× magnification with 100 µm scale bar.

morphology of early and late EPCs in *in vitro* culture (**Fig. 1**) was observed in our studies also. The former originates from monocytes with limited proliferation potential and express CD14, a monocytic marker. The late EPCs, also known as outgrowth endothelial cells (OECs) or endothelial colony forming cells (ECFCs) exhibit a typical cobblestone morphology, originate in the bone marrow and possess a greater proliferation and survival potential. The late EPCs are negative for CD14 and pan leukocyte marker CD45. Late EPC colonies express endothelial (KDR, vWF, and CD105) as well as progenitor cell markers (CD34 and CD133). These two subtypes are differentiated based on their morphology and functional activity in terms of rate of proliferation, *in vitro* angiogenesis, and rate of survival in culture. OECs or ECFCs show higher proliferation rate than early EPCs and only OECs/ECFCs has been shown to have direct involvement in *in vitro* angiogenesis.¹⁹⁻²¹

2. EPCs and coronary artery disease (CAD)

The role of EPCs as a marker of endothelial dysfunction and cardiovascular injury has usefulness in assessment of CAD progression.¹ Reduced circulating progenitor cells defined as CD34⁺ mononuclear cells and/or cells with expression of CD133 was shown to be associated with the risk of death in patients with CAD which links impaired endogenous regeneration capacity with increased mortality.²² Schmidt-Lucke et al.²³ reported that lower levels of circulating EPC counts were associated with a higher incidence of cardiovascular events in patients with stable CAD or acute coronary syndrome (ACS). Similarly, an inverse correlation was found by Werner et al.,²⁴ between the level of circulating EPCs and the risk of cardiovascular events among patients with angiographically confirmed CAD. Other studies indicate that progression of CAD is associated with not only the reduced circulatory level of EPCs, but also their impaired functional activity. Impaired migratory response of EPCs was inversely correlated with disease severity in CAD patients.²⁵

Reduced circulating EPC counts and their impaired functional activity, assessed with cell migration assays *in vitro*, is observed in CAD.²⁶ The early cellular senescence and lower EPC counts in patients with premature CAD correlates with shorter telomere length and reduced telomerase activity.²⁷ Likewise, Hammadah et al.²⁸ reported a positive association between leukocyte telomere length (LTL) and circulating progenitor cells (CPCs) characterized as CD34⁺ and subsets as CD34⁺/CD133⁺, CD34⁺/CXCR4⁺ and CD34⁺/CXCR4⁺/CD133⁺ in CAD patients. Moreover, both LTL and CPC levels were independent and additive predictors of adverse cardiovascular outcomes in CAD patients.²⁸ A recent prospective cohort study demonstrated that, CAD patients with renal insufficiency had lower CPC (CD34⁺, CD34⁺/

CD133⁺, CD34⁺/CXCR4⁺ and CD34⁺/VEGFR2⁺) counts among older participants (≥70 years of age) and low CPC count, a marker of impaired indigenous regenerative capacity, an independent predictor of adverse cardiovascular outcomes in patients with renal insufficiency. However, among patients with renal insufficiency, those with higher CPC count had similar cardiovascular (CV) outcomes as those without renal insufficiency, which cumulatively suggest that CV outcomes in patients with CAD and renal insufficiency is modulated by indigenous regenerative capacity contributed by CPCs.²⁹ In another cohort study, higher CPC (CD34⁺, CD34⁺/CD133⁺, and CD34⁺/CXCR4⁺) count was shown to be one of the reasons behind the obesity paradox in CAD (among individuals with CAD, obesity appears to provide a favorable outcome). The lower risk of CV outcomes was limited to obese individuals with high CPC count which preserve the endogenous regenerative capacity.³⁰

3. EPCs and heart failure (HF)

Endothelial dysfunction also occurs in patients with congestive heart failure (CHF).³¹⁻³³ However, limited evidence is available in support of EPCs and CD34⁺ cell mobilization during HF. Valgimigli et al.³⁴ reported a higher circulating level of EPCs in patients with CHF compared to control. An inverse correlation was observed between the severity of HF and circulating EPC count with significantly higher CD34⁺ counts in patients with mild HF compared to those with severe HF.³⁵ In acute CHF, stimulation of hematopoietic progenitor cells occurs, whereas in chronic stable stages of disease, depletion of the progenitor cell pool is observed, giving rise to a biphasic pattern in HF.³⁵ Circulating EPC numbers and colony forming units (CFU) (a functional parameter for the EPCs) are independent predictors of adverse outcomes in CHF.³⁶

4. EPCs and CVD risk factors

As far as the risk factors of CVD including hypertension, diabetes, metabolic syndrome, dyslipidemia, and smoking are concerned, Oliveras et al.³⁷ found a reduced EPC level in peripheral blood and the level was consistent even after *in vitro* culture in patients with refractory hypertension. Further this finding suggest that the remarkable decline of EPC level is independent of other risk factors which in turn further reinforces a likely role of EPCs in the development of atherosclerosis. Fadini et al.³⁸ demonstrated lower circulating levels of CD34⁺ and KDR⁺ EPC populations in subjects with greater carotid intima-media thickness (c-IMT). Reduced circulating EPC counts and impaired functionality was observed in metabolic syndrome,³⁹ diabetes,⁴⁰ and those with peripheral vascular disease.⁴¹ Hill et al.⁴² demonstrated a significant association of CFU-EC level with endothelial function as assessed by flow mediated brachial artery reactivity in individuals with various degree of cardiovascular risk factors but no history of CVDs. Moreover, this study speculate circulating EPC level as better predictor of vascular health than the presence or absence of conventional risk factors.

Diet and exercise also influence the circulating levels of EPC and their functional activity. A significant positive effect of 4 weeks exercise training on EPC number as well as cellular functionality was seen in young patients with chronic HF and this effect was consistent even in older patients with chronic HF which implies the potential benefit of rehabilitation intervention for patients with chronic HF.⁴³ In a randomized controlled trial (RCT), the Mediterranean diet was positively associated with higher circulating levels of CD34⁺KDR⁺ and CD34⁺KDR⁺CD133⁺ cells compared to a low-fat diet in the subjects with newly diagnosed type 2 diabetes.⁴⁴ In an another prospective randomized single-blind control trial, patients with coronary heart disease (CHD) showed improved endothelial functions as shown by higher flow mediated dilatation and increased circulatory EPCs level after consumption

of Mediterranean diet when compared with low fat diet. In addition, reduced intracellular oxidative stress, cell apoptosis and cellular senescence of ECs as well as higher cell proliferation and angiogenesis were seen in patients who followed the Mediterranean diet.⁴⁵

VITAMIN D AND CV HEALTH

Vitamin D is a secosteroid molecule traditionally associated with bone and calcium metabolism. Typically Vitamin D exists in 2 forms; D2 (ergocalciferol) and D3 (cholecalciferol).⁴⁶ The main sources of vitamin D3 in humans are either dietary such as fish oil or by endogenous synthesis from 7-dehydrocholesterol in the malpighian layer of the epidermis after exposure to sunlight, hence the name sunshine vitamin.⁴⁷ Hydroxylation of cholecalciferol by hepatic and microsomal enzymes occurs in the liver to form 25-hydroxy cholecalciferol (25-HCC) which gets further hydroxylated in the kidney by renal 1 α hydroxylase to form the active hormonal form of vitamin D, 1,25-dihydroxy cholecalciferol (1,25-DHCC), also known as calcitriol.⁴⁸ This active form of vitamin D binds to nuclear vitamin D receptors (VDR) to regulate expression of more than 200 genes that modulate a wide range of biological functions including calcium absorption from the intestine, calcium deposition in bone, stimulation of insulin and inhibition of renin production, stimulation of macrophage cathelicidin production etc.^{47,49} The vitamin D/VDR complex in conjunction with other nuclear hormone receptors, particularly the family of retinoid X receptor, bind to the vitamin D response element (VDRE), a hormone response element on DNA that can influence gene expression.⁵⁰ In the cardiovascular system, VDRs are present on vascular smooth muscle cells, endothelium and cardiomyocytes.⁵¹

1. Vitamin D and CVD: observational and interventional studies

Vitamin D status assumes greater importance in view of recent evidence from studies which suggest that in addition to its known effects on the musculoskeletal system, vitamin D may also influence cardiovascular health. The earliest observational study suggesting a protective role for vitamin D in ischemic heart disease came from the UK where Grimes et al.⁵² observed that mortality in ischemic heart disease was inversely proportional to hours of sunlight exposure. The NHANES conducted between 1988 and 1994 on 16,603 men and women aged >18 years reported greater frequency of 25-HCC deficiency in subjects with ischemic heart disease and stroke than in the general population.⁵³ Other studies have observed that severe vitamin D deficiency is associated with a higher risk of developing adverse cardiovascular events,⁵⁴ including myocardial infarction (MI)⁵⁵⁻⁵⁷ and sudden cardiac death/HF.⁵⁸ Vitamin D insufficiency (<30 ng/mL) was found to be directly associated with slow epicardial coronary flow, endothelial dysfunction as well as subclinical atherosclerosis in patients with normal or near normal coronary arteries³³ and decreased levels of vitamin D binding proteins correlates with number of affected coronary arteries in younger survivors of MI.³⁴ A large meta-analysis of cohort studies reported that vitamin D concentrations below 37 nmol/L was associated with sharp non-linear increase of CVD risk and mortality.⁵⁹ In contrast, another study found no association between vitamin D levels and severity of coronary lesions in patients with ST-segment elevation MI.⁵⁷ Apart from the direct effect of vitamin D on cardiomyocytes, vitamin D indirectly modifies CVD risk through its association with cardiovascular risk factors including diabetes, hypertension, systemic inflammation, and oxidative stress.⁶⁰

Lower vitamin D levels are associated with insulin resistance in observational and case control studies,^{61,62} and with increased risk of diabetes.^{63,64} Vitamin D supplementation was

associated with a slower rise in glucose levels in subjects with impaired fasting glucose.⁶⁵ The European Community Concerted Action on the Epidemiology and Prevention of Diabetes (EURODIAB) study reported a 33% lower risk of developing type 1 diabetes in children receiving vitamin D supplementation.⁶⁶

Vitamin D levels are inversely correlated with systolic blood pressure levels, a major risk factor of CAD.^{67,68} In the Health Professionals' follow-up study and the Nurses' health study, subjects with vitamin D <15 ng/mL were at 3- to 6-fold higher risk of developing incident of hypertension during a 4-year follow-up, compared to those with optimal vitamin D status.⁶⁹ Supplementation of vitamin D lowers systolic blood pressure,^{70,71} possibly by suppression of renin activity and sensitization of vascular smooth muscle cells.⁷² *In vivo* study reported sustained elevation of renin expression in vitamin D receptor-null mice resulting in increased production of angiotensin-II (Ang-II), a strong vasoconstrictor that promotes development of hypertension and left ventricular hypertrophy.⁷³ Moreover, vitamin D attenuates expression of the angiotensin-1 receptor in ECs of renal arteries from hypertensive patients, leading to improvement of endothelial function and decreased production of reactive oxygen species (ROS).⁷⁴

Vitamin D can also exert its cardioprotective effects through an anti-inflammatory action, by inhibition of smooth muscle cell proliferation, suppression of proatherogenic T lymphocytes, preservation of endothelial function,⁷⁴⁻⁸⁰ inhibition of foam cell formation, cholesterol uptake by macrophages, enabling high-density lipoprotein (HDL) transport⁸¹ and by protection against advanced glycation product formation.⁸² Its anti-inflammatory actions are mediated through a variety of pathways including the cyclooxygenase pathway, by upregulation of anti-inflammatory cytokines, reduction of cytokine-induced expression of cell adhesion molecules, reduction of matrix metalloproteinase (MMP)-9, inhibition of prostaglandins and downregulation of renin-angiotensin-aldosterone-system (RAAS).⁸³⁻⁸⁵ Supplementation of vitamin D in patients with CHF is associated with significant increase in anti-inflammatory cytokine interleukin (IL)-10.⁸⁶ Experimental studies have provided evidence that vitamin D reduces inflammation by lowering the release of tumor necrosis factor (TNF)- α and upregulation of anti-inflammatory cytokine IL-10 as well as expression of IL-10 receptor.^{87,88} Furthermore, in the Framingham offspring study, the plasma 25-hydroxyvitamin D (25[OH] D) was inversely associated with urinary isoprostane, an indicator of oxidative stress.⁸⁹

2. Vitamin D and CVD: RCT

Though the evidence from experimental and observational studies supports beneficial effect of vitamin D on cardiovascular health, the outcomes of interventional studies investigating the role of vitamin D supplementation on CVD is equivocal and conflict the results of observational and experimental studies. Randomized placebo-controlled trial by Hin et al.⁹⁰ reported no changes in cardiovascular risk factors including arterial stiffness in healthy elderly people (≥ 65 years of age) consuming daily supplementation of 4,000 IU vitamin D. A prospective RCT with MI patients demonstrated that the acute effect of vitamin D supplementation (4,000 IU) for 5 days was able to attenuate the inflammatory cytokines, such as C-reactive protein (CRP) and IL-6, but cell adhesion molecules intercellular adhesion molecule-1 (ICAM-1), E-selectin and other factors including vascular endothelial growth factor, TNF- α remains unchanged. However, the limitations of this study were small sample size and short duration of intervention.⁹¹ In another double blind RCT, weekly ergocalciferol (50,000 IU) administration for 12 weeks to CAD patients did not result in significant improvement of vascular and endothelial functions.⁹² Likewise, supplementation of vitamin D bolus (100,000 IU) to 5,110 community resident adults aged 50–84 years with

baseline 25(OH)D <50 nmol/L for a mean duration of 3.34 years showed no significant effect on the incidence of CVD.⁹³ In the same line, a US based nationwide randomized placebo-controlled trial conducted on 25,871 individuals found no association between vitamin D supplementation and risk of cardiovascular events and cancer after a median duration of 5.3 years.⁹⁴ In contrast, a randomized placebo-controlled double blind study of stable CAD patients receiving 0.5 µg vitamin D3/day for 6 months, reported a significant decrement of SYNTAX scores for CAD severity as well as significant decrease in high-sensitivity CRP level and renin-angiotensin system activity.⁹⁵

ROLE OF VITAMIN D ON EPCs AND ECs

1. *In vitro* studies

Vitamin D exposure at supra-physiological (10^{-9} mol/L) concentrations to cultured early EPCs improved cell proliferation and colony formation in type-2 diabetic patients. However, there was no effect of calcitriol on EPCs angiogenic marker expression and transcription factor Kruppel-like factor 10 (KLF10) which is known to participate in various aspects of cellular growth, development, and differentiation.⁷ A dose-dependent increase in production of NO via activation of endothelial nitric oxide synthase (eNOS) was observed in human umbilical vein endothelial cells (HUVECs) when treated with vitamin D. Increased NO production resulted in significant increase in phosphorylation of eNOS and intracellular kinases including p38, AKT and ERK which are known to be involved in intracellular signaling pathway associated with NO synthesis. Concomitant administration of ZK191784 (VDR agonist) and vitamin D to HUVECs resulted in greater production of number as compared to vitamin D or VDR agonist treatment alone, implying involvement of the VDR in production of NO.⁹⁶ In a recent study, *in vitro* vitamin D treatment in a dose dependent manner on early EPCs from systemic lupus erythematosus (SLE) patients shows improvement in EPC number and functionality in terms of migratory and proliferative potential as well as upregulation of EPC's NO biosynthesis. In addition to that, intracellular EPC's NO⁻ level was positively correlated with EPC number and functions which further may predicts the role of vitamin D in the improvement of EPC number and functions via NO⁻ signaling pathway.⁹⁷

In another study, vitamin D treatment on ECFCs derived from cord blood sample of healthy pregnant women during delivery exhibited a significantly increased functional activity reflected by higher formation of entire length of tubular structure on matrigel matrix and an increased cell proliferation compared to vehicle treated controls. In addition, an increased VEGF mRNA expression and elevated activity of pro-MMP-2 was demonstrated in ECFCs when treated with vitamin D. However, these positive effects of vitamin D were reversed when VDR and VEGF cascade pathways were specifically blocked, suggesting that the effects on ECFCs are secondary to up regulation of VEGF expression. Since angiogenesis is crucially associated with pathophysiology of preeclampsia, this study speculated the beneficial effect of vitamin D supplementation in early pregnancy particularly during placental development in reducing the risk of developing preeclampsia.⁹⁸

Several studies have shown markedly reduced migratory capacity and angiogenesis of cord blood-derived ECFCs when exposed to serum from preeclamptic compared to normal women.^{99,100} Vitamin D restored the functional properties and abolished the negative effect of preeclampsia on ECFCs. This positive effect of vitamin D appeared to be mediated through VEGF signaling pathway as Flk-1/KDR (VEGF) receptor tyrosine kinase inhibitor SU5416

was able to suppress the ECFC's tubule formation, an effect similar to VDR blockade by pyridoxal phosphate. Unlike the findings of the previous study by Grundmann et al.,⁹⁸ the positive effect of vitamin D on ECFC's angiogenesis in this study was independent of the FBS concentration in treated media.¹⁰⁰ In addition, attenuation of ECFCs migration was seen in response to conditioned media obtained from placenta under aberrant oxygen conditions; 2% O₂ (hypoxia) or 21% hyperoxia compared to normoxic (8% O₂). Vitamin D at a concentration of 10 nM restored the ECFC's functionality. The reduced tube formation and migration from VDR blockade was partially reversed by vitamin D, but VEGF pathway inhibition could not be restored by vitamin D. These findings suggest that vitamin D can overcome some of negative effects of preeclamptic condition on ECFC, an effect that is mediated via the VDR.⁹⁹

Adverse prenatal exposure is one of the leading causes of developmental origin of chronic diseases such as cardiovascular diseases in adulthood.¹⁰¹⁻¹⁰³ Based on this paradigm prenatal exposure of preeclampsia (PE) is well known as an independent risk factor for long-term cardiovascular morbidity of the offspring and studies suggest that vitamin D deficiency is one of the factors to be responsible for the increased incidence of PE.¹⁰⁴ A study suggested the role of human fetal EPCs in maternal endothelial function during pregnancy as evidenced by fetal EPC's transmigration to the maternal blood stream and homing to locations of maternal uterine vasculogenesis.¹⁰⁵ In an *in vitro* model, Brodowski et al.¹⁰⁶ explained that fetal blood derived ECFCs when exposed to fetal serum from PE pregnancy, become impaired in integration and invasion into the monolayer of mature HUVECs, which may explain the role of fetal circulating factors in disruption of endothelial functions and the mechanism by which angiogenesis and endothelial homeostasis are disrupted in PE. Exogenous vitamin D administration at physiological concentrations led to a substantial reversal of fetal serum mediated inhibitory effect on ECFCs.¹⁰⁶

In another *in vitro* study, Zhong et al.¹⁰⁷ reported stimulation of VDR expression by vitamin D in a time and dose dependent manner on ECs. Vitamin D also up regulated the expression of VEGF and its receptor as well as antioxidant CuZn-superoxide dismutase expression in ECs. Hypoxia mediated oxidative stress by CoCl₂, a hypoxia mimetic agent can lead to downregulation of VDR and CuZn superoxide dismutase expression and this was prevented by administration of vitamin D in *in vitro* culture.¹⁰⁷

Uberti et al.¹¹⁰ reported that vitamin D alone or in combination with VDR ligand ZK191784 prevent human EC death from oxidative stress via modulation of the interplay between apoptosis and autophagy. Vitamin D negated the effects by attenuating apoptosis related gene expression involved in both intrinsic and extrinsic pathways and augmenting activation of pro-autophagic beclin 1 as well as phosphorylation of ERK1/2 and Akt.¹⁰⁸

An increase in expression of vascular cell adhesion molecule-1 (VCAM-1) and MT1-MMP is seen in human ECs when stimulated with lipopolysaccharide (LPS) and incubated with platelets. This effect was significantly attenuated by pre-treatment of cells with vitamin D. Moreover, a reduced CD62P expression in platelets in direct contact with pretreated ECs has been found compared to platelets incubated with untreated ECs. Vitamin D was able to suppress the platelet activation and expression of VCAM-1, MT1-MMP in human ECs.¹⁰⁹

Schröder-Heurich et al.⁶ reported that the vasoprotective effect of vitamin D could be through its impact on ECFC homeostasis. The effect included maintenance of stability of ECFC monolayers by enhancement of endothelial interconnection through vascular endothelial

cadherin (VE-cadherin) junctions. In addition, this study also showed the positive effect of vitamin D on directional ECFC mobilization mediated through enhanced expression of adhesion proteins and change in F-actin formation.⁶

The formation of advanced glycation end products (AGE) and their tissue accumulation is common phenomena in diabetes, chronic renal failure and aging,¹¹⁰ and is associated with development of multiple complications, particularly vascular atherosclerosis.^{111,112} In an *in vitro* study by Talmor et al.,¹¹³ a beneficial effect of vitamin D on human ECs affected by AGE was seen. Cells induced with AGE-human serum albumin (HSA) showed reduced eNOS gene expression and enzyme activity, but administration of calcitriol to AGE-HSA treated cells ameliorated the effect of AGE by blunting the AGEs receptor mRNA expression and proteins. In addition, calcitriol administration in ECs effectively reduced AGEs-induced elevated levels of IL-6 mRNA and nuclear factor (NF)- κ B-p65 DNA binding activity, a phenomenon associated with elevated expression of I κ B α .¹¹³

MicroRNA (MiRNA), the non-coding RNA, play a role in regulating expression of a number of genes by binding at the 3' untranslated region region of target mRNA of protein coding genes and induce the translational repression and/or mRNA cleavage.^{114,115} MiRNAs miR-659 and miR-510 are up regulated in human ECs pretreated with AGE-HSA whereas miR-181c, miR-411, miR-126, miR-15a and miR-20b are down regulated. Administration of calcitriol reverted this pattern of miR expression.¹¹⁶ Paricalcitol, a vitamin D analog and calcitriol downregulated markers involved in the inflammatory response of ECs exposed to AGE-HSA.¹¹⁷

Gestational diabetes mellitus (GDM) is a common cause of manifestation of CVD in later life of the mother and the offspring.¹¹⁸ The fetal ECFCs from gestational diabetes mellitus pregnancies exhibited a significant impairment in functional activity as well as the number of colonies *in vitro* when compared with fetal ECFCs from uncomplicated pregnancies. In the same study, fetal ECFCs obtained from uncomplicated pregnancies showed a significant functional deficit in terms of cell migration and tubule formation upon exposure to mildly hyperglycemic conditions. But a significant reduction in the inhibitory effect of GDM and mild hyperglycemic on fetal ECFCs functionality was seen after *in vitro* treatment of vitamin D.¹¹⁹

Endothelial dysfunction has been seen in type-2 diabetes mellitus patients with vitamin D insufficiency or deficiency and supplementation of vitamin D can improve the EC function.¹²⁰ Bioinformatics analysis explains the vitamin D induced molecular mechanism of differentially expressed miRNA, mRNA, circular RNA (circRNA) and long noncoding RNA (lncRNA) in bone marrow derived EPCs pretreated with high level of glucose. Following vitamin D administration, competent endogenous RNA (ceRNA) which comprises circRNA, lncRNA has been found to attenuate the effects of miRNAs on the expression of mRNAs in EPCs and may suggest the role of vitamin D in alleviation of EPCs dysfunctions through the associated ceRNA regulatory network in diabetes.¹²¹ The mechanism of action of vitamin D on endothelial and or endothelial progenitor cells has been summarized in **Fig. 2**.

Inflammation to endothelium is the hallmark of endothelial activation and development of atherosclerosis.¹²² This cellular inflammatory effect results in stimulation of excessive cytokines and adhesion molecule expression which further exacerbate endothelial dysfunction.^{123,124} Angiotensin-II is known to upregulate NF- κ B and which further induce expression of TNF- α , IL-6, cell adhesion molecules ICAM-1, VCAM-1 and E-selectin, resulting vascular injury.¹²⁴ However, vitamin D has a role in inhibition of inflammatory reaction




Fig. 2. Schematic representation of effect of vitamin D3 on EPCs and/or endothelial cells. Binding of vitamin D to nuclear VDR in combination with RXR on VDRE triggers the expression of number of genes, such as eNOS, VEGF, CuZn dismutase, and pro-autophagic bacilin-1. Vitamin D3 mediates activation of eNOS by phosphorylation resulting in the augmentation of NO synthesis. Intracellular Ca²⁺ influx is caused by vitamin D and subsequent calcium calmodulin complex stabilizes the catalytic state of eNOS which catalyzes the synthesis of NO from L-Arginine and molecular oxygen. In association with VEGF, SDF-1 and MMP-9, NO leads to the migration of EPC from the permissive vascular zone of bone marrow into the circulation. The upregulation of CuZn dismutase by vitamin D3 via VDR activation attenuates the cellular oxidative stress, while upregulation of pro-autophagic bacilin-1 by VDR mediated vitamin D3 reduces the cellular apoptosis. Increased VEGF mRNA expression and protein synthesis mediates EPC migration, proliferation, and *in vitro* angiogenesis. Upregulation of microRNA (miR-659, miR-510) and downregulation of microRNA (miR-181c, miR-411, miR-126, miR-15a, miR-20b) seen in hyperglycemic conditions like diabetes are reverted by vitamin D3 via VDR signaling pathway. The adhesion of leukocyte and platelets to the endothelium is suppressed by vitamin D3 through downregulation of VCAM-1 and ICAM-1 expression as well as proinflammatory cytokine IL-6. This illustration was created using BioRender.com.
EPC, endothelial progenitor cell; VDR, vitamin D receptor; RXR, retinoid X receptor; VDRE, vitamin D response element; eNOS, endothelial nitric oxide synthase; VEGF, vascular endothelial growth factor; NO, nitric oxide; SDF, stromal derived factor; MMP, matrix metalloproteinase; VCAM-1, vascular cell adhesion molecule-1; ICAM-1, intercellular adhesion molecule-1; IL, interleukin.

and thus protect vascular damage.⁹⁵ Recent study demonstrate that, impaired cell viability, migration ability and angiogenesis as well as cell apoptosis has been noticed in EPCs when treated with Ang-II and vitamin D administration ameliorate the effect of Ang-II via inhibition of cellular oxidative stress, NF-κB activation and inflammatory cytokines.¹²⁵

2. In vivo studies

A recent *in vivo* study demonstrated amelioration of hyperlipidemia as well as progression of atherosclerosis by intravenous transplantation of genetically modified late outgrowth EPCs with VDR over-expression in ApoE-knockout mice. Additionally, VDR over-expressed EPCs (ovVDR-EPCs) transplanted group had higher HDL-cholesterol level, lower levels of total cholesterol, low-density lipoprotein, apoB and Lp(a), elevated serum NO levels, increased expression of serum and vessel wall eNOS, reduced expression and activity of

MMPs and decreased expression and activity of their tissue inhibitor (tissue inhibitors of metalloproteinase, TIMPs) compared to controls. These findings demonstrate the beneficial role of EPCs over-expressing VDR in protecting against atherosclerotic risk and suggest that transfusion of ovVDR-EPCs might be used in angiogenic therapy.¹²⁶

SLE, an autoimmune disease is associated with increased risk of CVD and endothelial dysfunction.^{127,128} Premature CVD in patients with SLE is precipitated by failure of endothelial repair.¹²⁸ Vitamin D deficiency introduced by dietary restriction in lupus prone MRL/lpr mice caused endothelial dysfunction. Improved differentiation of EPCs was observed after vitamin D3 supplementation. Vitamin D deficiency was associated with upregulation of interferon-stimulated gene expression in both MRL/lpr mice and patients with SLE.¹²⁹

The initiation and progression of atherosclerotic plaque formation results from EC activation followed by leukocyte recruitment and adhesion to the activated endothelium. The knockdown of VDR expression in *in vitro* human ECs induces a significant reduction of peripheral blood mononuclear cells (PBMC) rolling velocity over the endothelial monolayer and enhanced rolling flux and adhesion of PBMCs to the endothelium. The enhanced rolling velocity and adhesion of PBMCs was seen to be mediated by upregulation of VCAM-1 and ICAM-1 in VDR knockdown ECs along with significant elevation of IL-6 secretion. Down-regulation of VDR in ECs exhibited reduced level of I κ B α as well as aggregation of p65 transcription factor (a member of NF- κ B family) in the nucleus. VDR deletion in apoE^{-/-} mice exhibited elevated aortic expression of VCAM-1, ICAM-1 and IL-6. *In vivo* deletion of VDR is associated with larger aortic arch and aortic root lesions along with higher macrophage content in apoE^{-/-} mice. The lack of VDR signaling in ECs may lead to increased leukocyte-EC interactions causing atherosclerotic plaque development in apoE^{-/-}VDR^{-/-} mice.⁹

3. Clinical studies

The receptor for vitamin D is widely distributed in almost all tissues and hence the use of vitamin D as a therapeutic agent has been shown to be beneficial in many conditions including CVDs and cancers.¹³⁰ At least 60 genes are regulated by vitamin D, which acts as a steroid hormone upon binding to the hormone response element of the genes.^{131,132} A positive correlation between serum levels of vitamin D and circulating EPCs as well as cultured angiogenic cells suggests a possible role of vitamin D as a developmental hormone for stem cells that differentiate into the endothelial phenotype.^{133,134} Study done by Mikirova et al.,¹³⁵ indicated a positive correlation between serum level of vitamin D (25[OH]D) and circulating EPC as well as cultured angiogenic cells suggesting a possible role of vitamin D3 as a developmental hormone for stem cells which differentiates into endothelial phenotype.

Reduced VDR expression on circulating EPCs has been reported in CAD patients, particularly among those with elevated HbA1c.⁸ VDR mRNA expression was significantly decreased when EPCs were treated with high glucose (22 mmol/L) for 24 and 48 hours. Similarly, VDR expression on EPCs among hemodialysis patients was lower than that in control subjects, while upregulation of VDR expression on EPCs had been observed among those treated with oral/IV vitamin D.¹³⁶ Depletion of circulatory proangiogenic progenitor mononuclear cells as CD14⁺CD309⁺Tie-2⁺ cells were closely associated with the vitamin D status especially vitamin D3 in patients with metabolic syndrome (MetS) without known CVD. This study suggests a link between vitamin D deficiency and impaired endogenous endothelial repair system which in turn might lead to vascular complications in patients with MetS.¹³⁷ Reduced circulating progenitor cells (CD34⁺) in patients with rheumatoid arthritis (RA) was positively

Beneficial Role of Vitamin D on EPCs in CVDs

Table 1. Association between serum vitamin D and circulatory EPC levels

References	No. of subjects	Subject characteristics	Serum vitamin D concentration	Levels of EPCs in circulation	Associations observed
Mikirova et al. ¹³⁵	n=41	Healthy adults	<40 ng/mL >40 ng/mL <30 ng/mL >30 ng/mL	Mean % 0.045 0.068 0.046 0.060	Positive correlation between serum VitD concentration and circulatory EPC level.
Yiu et al. ¹³⁴	n=96, VitD deficient n=115, VitD insufficient n=69, VitD sufficient	Diabetic	<20 ng/mL <30 ng/mL ≥30 ng/mL	CD34 ⁺ /KDR ⁺ (%): 0.746±0.465 CD133 ⁺ /KDR ⁺ (%): 0.219±0.189 CD34 ⁺ /KDR ⁺ (%): 0.628±0.354 CD133 ⁺ /KDR ⁺ (%): 0.283±0.197 CD34 ⁺ /KDR ⁺ (%): 0.747±0.412 CD133 ⁺ /KDR ⁺ (%): 0.364±0.172	Circulatory level of CD133 ⁺ /KDR ⁺ EPC found to be positively correlated with VitD deficiency and insufficiency group.
Chan et al. ¹³⁹	n=297	CVD (coronary artery disease, ischemic stroke)	<20 ng/mL 20–29 ng/mL 30–39 ng/mL ≥40 ng/mL	log, unit (×10 ⁻³ /mL) 0.97±0.31 1.05±0.42 1.08±0.41 1.15±0.39	Positive association between serum 25(OH)D and CD34 ⁺ KDR ⁺ , CD133 ⁺ KDR ⁺ EPC.
Gurses et al. ¹⁴⁰	n=27, control group n=31, VitD deficient group	Premenopausal women with VitD deficient	34.40±10.30 ng/mL (for control group) 10.60±4.70 ng/mL (for VitD deficient group)	CD34 ⁺ /KDR ⁺ EPC (/ μ L): 64.50±17.10 CD133 ⁺ /KDR ⁺ EPC (/ μ L): 49.40±23.10 CD34 ⁺ /KDR ⁺ EPC (/ μ L): 26.20±20.50 CD133 ⁺ /KDR ⁺ EPC (/ μ L): 27.40±18.50	Positive correlation between baseline VitD level and circulatory level of CD34 ⁺ /KDR ⁺ , CD133 ⁺ /KDR ⁺ cells.
Berezin et al. ¹³⁷	n= 10, 25(OH)D3 n=12, 25(OH)D3 n= 14, 25(OH)D3 level n=11, 25(OH)D3	MetS without known CVD	>100 nmol/L 50–100 nmol/L 30–50 nmol/L <30 nmol/L	CD14 ⁺ CD309 ⁺ Tie2 ⁺ cells/ μ L 0.039 (0.032–0.047) 0.035 (0.028–0.044) 0.030 (0.022–0.041) 0.028 (0.016–0.033)	Positive association VitD deficiency significant predictor for reduced circulatory proangiogenic progenitor cells as CD14 ⁺ CD309 ⁺ Tie2 ⁺ .

VitD, vitamin D; EPC, endothelial progenitor cell; CD, cluster differentiation; KDR, kinase insert domain receptor; Tie2, tyrosine kinase receptor 2; 25(OH)D, 25-hydroxyvitamin D; MetS, metabolic syndrome.

correlated with lower vitamin D levels as well as with endothelial functions including pulse wave velocity and carotid intima-media thickness which may predict the contribution of vitamin D in endothelial homeostasis in patients with RA.¹³⁸ Chan et al.¹³⁹ have reported that statin therapy significantly raises serum vitamin D levels and vitamin D level was found to associate independently with higher circulating levels of CD34⁺KDR⁺, CD133⁺KDR⁺ EPC and a reduction in burden of carotid atherosclerosis in patients with cardiovascular disease.

Premenopausal women with vitamin D deficiency (<20 ng/mL) had impaired endothelial function as evaluated by flow mediated dilatation (FMD), and lower CD34⁺/KDR⁺ (26.20±20.50 / μ L) and CD133⁺/KDR⁺ (27.40±18.50 / μ L) EPC counts, together with lower levels of IL-10 and elevated level of IL-17. Six months (50,000 IU/week) vitamin D supplementation significantly increased vitamin D levels and ameliorated endothelial function as evidenced by higher FMD, CD34⁺/KDR⁺ and CD133⁺/KDR⁺ EPC counts as well as a shift of cytokine profile towards more anti-inflammatory cytokines.¹⁴⁰ In contrast, a 12-week randomized control trial of oral vitamin D supplementation of 5,000 IU/day in patients with suboptimal serum 25(OH)D levels (<30 ng/mL) did not show any significant improvement in FMD, pulse wave velocity and circulating levels of EPC in type 2 diabetic patients.¹⁴¹ With reference to RCTs, the beneficial effect of vitamin D supplementation on CVDs is a topic which has not been explored adequately and hence concrete evidence does not yet exist. **Table 1** summarizes the clinical studies on vitamin D status and its effect on circulatory EPCs.

CONCLUSION AND PERSPECTIVES

An optimum serum level of vitamin D is important not only to maintain the body's normal calcium homeostasis but also to reduce the risk of cardiovascular events, diabetes and

hypertension. Vitamin D exerts its beneficial role in several ways including antithrombotic, antihypertensive, anti-inflammatory, anti-fibrotic and antidiabetic effects. However, despite of showing the beneficial effect of vitamin D in the prevention of CVD, there are conflicting results from experimental and clinical studies on the role of vitamin D in the pathogenesis of CVD. This notion is contradicted by the data from the RCTs, RCTs have not shown beneficial effects of vitamin D on CVD, these trials were not designed to investigate cardiovascular outcomes in vitamin D-deficient individuals.¹⁴² There could be a subset of individuals with CAD who could benefit from vitamin D supplementation.

Recent research suggest that vitamin D also plays an important role in vascular remodeling and healing of damaged endothelium by increasing the circulatory level and functionality of EPCs. Since 1997, when Asahara et al.¹ reported the existence of EPCs and their role in postnatal neovascularization, a large number of studies have shown the importance of EPCs in the process of repair of damaged endothelium during vascular trauma and ischemia, and recent *in vitro* and *in vivo* studies have revealed some important signaling pathways through which vitamin D exerts its beneficial effects on EPCs. In the field of regenerative medicine, EPCs have been recognized as one of the novel therapeutic agents in vascular disease and several pilot studies on animals have already shown encouraging results. Further studies focused on investigating the mechanism of action of vitamin D on EPC number and cell viability will likely yield metabolic and molecular pathways that more precisely elucidate the role of vitamin D in EPCs. Determining the mechanistic pathways of beneficial functions of vitamin D on EPCs will be critical for the translation of vitamin D as therapeutic agent in a subset of individuals with defective EPC functions or vitamin D deficiency.

REFERENCES

1. Asahara T, Murohara T, Sullivan A, Silver M, van der Zee R, Li T, et al. Isolation of putative progenitor endothelial cells for angiogenesis. *Science* 1997;275:964-967.
[PUBMED](#) | [CROSSREF](#)
2. Naeem Z. Vitamin D deficiency- an ignored epidemic. *Int J Health Sci (Qassim)* 2010;4:V-VI.
[PUBMED](#)
3. Holick MF. Sunlight and vitamin D for bone health and prevention of autoimmune diseases, cancers, and cardiovascular disease. *Am J Clin Nutr* 2004;80:1678S-1688S.
[PUBMED](#) | [CROSSREF](#)
4. Grimes DS, Hindle E, Dyer T. Sunlight, cholesterol and coronary heart disease. *QJM* 1996;89:579-589.
[PUBMED](#) | [CROSSREF](#)
5. Danik JS, Manson JE. Vitamin D and cardiovascular disease. *Curr Treat Options Cardiovasc Med* 2012;14:414-424.
[PUBMED](#) | [CROSSREF](#)
6. Schröder-Heurich B, von Hardenberg S, Brodowski L, Kipke B, Meyer N, Borns K, et al. Vitamin D improves endothelial barrier integrity and counteracts inflammatory effects on endothelial progenitor cells. *FASEB J* 2019;33:9142-9153.
[PUBMED](#) | [CROSSREF](#)
7. Hammer Y, Soudry A, Levi A, Talmor-Barkan Y, Leshem-Lev D, Singer J, et al. Effect of vitamin D on endothelial progenitor cells function. *PLoS One* 2017;12:e0178057.
[PUBMED](#) | [CROSSREF](#)
8. Ai S, He Z, Ding R, Wu F, Huang Z, Wang J, et al. Reduced vitamin D receptor on circulating endothelial progenitor cells: a new risk factor of coronary artery diseases. *J Atheroscler Thromb* 2018;25:410-421.
[PUBMED](#) | [CROSSREF](#)
9. Bozic M, Álvarez Á, de Pablo C, Sanchez-Niño MD, Ortiz A, Dolcet X, et al. Impaired vitamin D signaling in endothelial cell leads to an enhanced leukocyte-endothelium interplay: implications for atherosclerosis development. *PLoS One* 2015;10:e0136863.
[PUBMED](#) | [CROSSREF](#)

10. Aragona CO, Imbalzano E, Mamone F, Cairo V, Lo Gullo A, D'Ascola A, et al. Endothelial progenitor cells for diagnosis and prognosis in cardiovascular disease. *Stem Cells Int* 2016;2016:8043792.
[CROSSREF](#)
11. Peplow PV. Growth factor- and cytokine-stimulated endothelial progenitor cells in post-ischemic cerebral neovascularization. *Neural Regen Res* 2014;9:1425-1429.
[PUBMED](#) | [CROSSREF](#)
12. Fadini GP, Mehta A, Dhindsa DS, Bonora BM, Sreejit G, Nagareddy P, et al. Circulating stem cells and cardiovascular outcomes: from basic science to the clinic. *Eur Heart J* 2020;41:4271-4282.
[PUBMED](#) | [CROSSREF](#)
13. Lee PS, Poh KK. Endothelial progenitor cells in cardiovascular diseases. *World J Stem Cells* 2014;6:355-366.
[PUBMED](#) | [CROSSREF](#)
14. Zheng H, Fu G, Dai T, Huang H. Migration of endothelial progenitor cells mediated by stromal cell-derived factor-1alpha/CXCR4 via PI3K/Akt/eNOS signal transduction pathway. *J Cardiovasc Pharmacol* 2007;50:274-280.
[PUBMED](#) | [CROSSREF](#)
15. Yamaguchi J, Kusano KF, Masuo O, Kawamoto A, Silver M, Murasawa S, et al. Stromal cell-derived factor-1 effects on *ex vivo* expanded endothelial progenitor cell recruitment for ischemic neovascularization. *Circulation* 2003;107:1322-1328.
[PUBMED](#) | [CROSSREF](#)
16. Natori T, Sata M, Washida M, Hirata Y, Nagai R, Makuuchi M. G-CSF stimulates angiogenesis and promotes tumor growth: potential contribution of bone marrow-derived endothelial progenitor cells. *Biochem Biophys Res Commun* 2002;297:1058-1061.
[PUBMED](#) | [CROSSREF](#)
17. Kalka C, Masuda H, Takahashi T, Gordon R, Tepper O, Gravereaux E, et al. Vascular endothelial growth factor(165) gene transfer augments circulating endothelial progenitor cells in human subjects. *Circ Res* 2000;86:1198-1202.
[PUBMED](#) | [CROSSREF](#)
18. Thum T, Fraccarollo D, Schultheiss M, Froese S, Galuppo P, Widder JD, et al. Endothelial nitric oxide synthase uncoupling impairs endothelial progenitor cell mobilization and function in diabetes. *Diabetes* 2007;56:666-674.
[PUBMED](#) | [CROSSREF](#)
19. Medina RJ, O'Neill CL, Sweeney M, Guduric-Fuchs J, Gardiner TA, Simpson DA, et al. Molecular analysis of endothelial progenitor cell (EPC) subtypes reveals two distinct cell populations with different identities. *BMC Med Genomics* 2010;3:18.
[PUBMED](#) | [CROSSREF](#)
20. Gulati R, Jevremovic D, Peterson TE, Chatterjee S, Shah V, Vile RG, et al. Diverse origin and function of cells with endothelial phenotype obtained from adult human blood. *Circ Res* 2003;93:1023-1025.
[PUBMED](#) | [CROSSREF](#)
21. Mukai N, Akahori T, Komaki M, Li Q, Kanayasu-Toyoda T, Ishii-Watabe A, et al. A comparison of the tube forming potentials of early and late endothelial progenitor cells. *Exp Cell Res* 2008;314:430-440.
[PUBMED](#) | [CROSSREF](#)
22. Patel RS, Li Q, Ghasemzadeh N, Eapen DJ, Moss LD, Janjua AU, et al. Circulating CD34⁺ progenitor cells and risk of mortality in a population with coronary artery disease. *Circ Res* 2015;116:289-297.
[PUBMED](#) | [CROSSREF](#)
23. Schmidt-Lucke C, Rössig L, Fichtlscherer S, Vasa M, Britten M, Kämper U, et al. Reduced number of circulating endothelial progenitor cells predicts future cardiovascular events: proof of concept for the clinical importance of endogenous vascular repair. *Circulation* 2005;111:2981-2987.
[PUBMED](#) | [CROSSREF](#)
24. Werner N, Wassmann S, Ahlers P, Schiegl T, Kosiol S, Link A, et al. Endothelial progenitor cells correlate with endothelial function in patients with coronary artery disease. *Basic Res Cardiol* 2007;102:565-571.
[PUBMED](#) | [CROSSREF](#)
25. Vasa M, Fichtlscherer S, Aicher A, Adler K, Urbich C, Martin H, et al. Number and migratory activity of circulating endothelial progenitor cells inversely correlate with risk factors for coronary artery disease. *Circ Res* 2001;89:E1-E7.
[PUBMED](#) | [CROSSREF](#)
26. Huang C, Zhang L, Wang Z, Pan H, Zhu J. Endothelial progenitor cells are associated with plasma homocysteine in coronary artery disease. *Acta Cardiol* 2011;66:773-777.
[PUBMED](#) | [CROSSREF](#)

27. Vemparala K, Roy A, Bahl VK, Prabhakaran D, Nath N, Sinha S, et al. Early accelerated senescence of circulating endothelial progenitor cells in premature coronary artery disease patients in a developing country - a case control study. *BMC Cardiovasc Disord* 2013;13:104.
[PUBMED](#) | [CROSSREF](#)
28. Hammadah M, Al Mheid I, Wilmot K, Ramadan R, Abdelhadi N, Alkhoder A, et al. Telomere shortening, regenerative capacity, and cardiovascular outcomes. *Circ Res* 2017;120:1130-1138.
[PUBMED](#) | [CROSSREF](#)
29. Mehta A, Tahhan AS, Liu C, Dhindsa DS, Nayak A, Hooda A, et al. Circulating progenitor cells in patients with coronary artery disease and renal insufficiency. *JACC Basic Transl Sci* 2020;5:770-782.
[PUBMED](#) | [CROSSREF](#)
30. Mehta A, Meng Q, Li X, Desai SR, D'Souza MS, Ho AH, et al. Vascular regenerative capacity and the obesity paradox in coronary artery disease. *Arterioscler Thromb Vasc Biol* 2021;41:2097-2108.
[PUBMED](#) | [CROSSREF](#)
31. Chong AY, Blann AD, Patel J, Freestone B, Hughes E, Lip GY. Endothelial dysfunction and damage in congestive heart failure: relation of flow-mediated dilation to circulating endothelial cells, plasma indexes of endothelial damage, and brain natriuretic peptide. *Circulation* 2004;110:1794-1798.
[PUBMED](#) | [CROSSREF](#)
32. Fischer D, Rossa S, Landmesser U, Spiekermann S, Engberding N, Hornig B, et al. Endothelial dysfunction in patients with chronic heart failure is independently associated with increased incidence of hospitalization, cardiac transplantation, or death. *Eur Heart J* 2005;26:65-69.
[PUBMED](#) | [CROSSREF](#)
33. Marti CN, Gheorghiane M, Kalogeropoulos AP, Georgiopoulou VV, Quyyumi AA, Butler J. Endothelial dysfunction, arterial stiffness, and heart failure. *J Am Coll Cardiol* 2012;60:1455-1469.
[PUBMED](#) | [CROSSREF](#)
34. Valgimigli M, Rigolin GM, Fucili A, Porta MD, Soukhomovskaia O, Malagutti P, et al. CD34⁺ and endothelial progenitor cells in patients with various degrees of congestive heart failure. *Circulation* 2004;110:1209-1212.
[PUBMED](#) | [CROSSREF](#)
35. Nonaka-Sarukawa M, Yamamoto K, Aoki H, Nishimura Y, Tomizawa H, Ichida M, et al. Circulating endothelial progenitor cells in congestive heart failure. *Int J Cardiol* 2007;119:344-348.
[PUBMED](#) | [CROSSREF](#)
36. Michowitz Y, Goldstein E, Wexler D, Sheps D, Keren G, George J. Circulating endothelial progenitor cells and clinical outcome in patients with congestive heart failure. *Heart* 2007;93:1046-1050.
[PUBMED](#) | [CROSSREF](#)
37. Oliveras A, Soler MJ, Martínez-Estrada OM, Vázquez S, Marco-Feliu D, Vila JS, et al. Endothelial progenitor cells are reduced in refractory hypertension. *J Hum Hypertens* 2008;22:183-190.
[PUBMED](#) | [CROSSREF](#)
38. Fadini GP, Coracina A, Baesso I, Agostini C, Tiengo A, Avogaro A, et al. Peripheral blood CD34⁺KDR⁺ endothelial progenitor cells are determinants of subclinical atherosclerosis in a middle-aged general population. *Stroke* 2006;37:2277-2282.
[PUBMED](#) | [CROSSREF](#)
39. Jialal I, Devaraj S, Singh U, Huet BA. Decreased number and impaired functionality of endothelial progenitor cells in subjects with metabolic syndrome: implications for increased cardiovascular risk. *Atherosclerosis* 2010;211:297-302.
[PUBMED](#) | [CROSSREF](#)
40. Egan CG, Lavery R, Caporali F, Fondelli C, Laghi-Pasini F, Dotta F, et al. Generalised reduction of putative endothelial progenitors and CXCR4-positive peripheral blood cells in type 2 diabetes. *Diabetologia* 2008;51:1296-1305.
[PUBMED](#) | [CROSSREF](#)
41. Fadini GP, Miorin M, Facco M, Bonamico S, Baesso I, Grego F, et al. Circulating endothelial progenitor cells are reduced in peripheral vascular complications of type 2 diabetes mellitus. *J Am Coll Cardiol* 2005;45:1449-1457.
[PUBMED](#) | [CROSSREF](#)
42. Hill JM, Zalos G, Halcox JP, Schenke WH, Waclawiw MA, Quyyumi AA, et al. Circulating endothelial progenitor cells, vascular function, and cardiovascular risk. *N Engl J Med* 2003;348:593-600.
[PUBMED](#) | [CROSSREF](#)
43. Sandri M, Viehmann M, Adams V, Rabald K, Mangner N, Höllriegel R, et al. Chronic heart failure and aging - effects of exercise training on endothelial function and mechanisms of endothelial regeneration: results from the Leipzig Exercise Intervention in Chronic heart failure and Aging (LEICA) study. *Eur J Prev Cardiol* 2016;23:349-358.
[PUBMED](#) | [CROSSREF](#)

44. Maiorino MI, Bellastella G, Petrizzo M, Gicchino M, Caputo M, Giugliano D, et al. Effect of a Mediterranean diet on endothelial progenitor cells and carotid intima-media thickness in type 2 diabetes: follow-up of a randomized trial. *Eur J Prev Cardiol* 2017;24:399-408.
[PUBMED](#) | [CROSSREF](#)
45. Yubero-Serrano EM, Fernandez-Gandara C, Garcia-Rios A, Rangel-Zuñiga OA, Gutierrez-Mariscal FM, Torres-Peña JD, et al. Mediterranean diet and endothelial function in patients with coronary heart disease: an analysis of the CORDIOPREV randomized controlled trial. *PLoS Med* 2020;17:e1003282.
[PUBMED](#) | [CROSSREF](#)
46. Laird E, Ward M, McSorley E, Strain JJ, Wallace J. Vitamin D and bone health; potential mechanisms. *Nutrients* 2010;2:693-724.
[PUBMED](#) | [CROSSREF](#)
47. Nair R, Maseeh A. Vitamin D: the “sunshine” vitamin. *J Pharmacol Pharmacother* 2012;3:118-126.
[PUBMED](#) | [CROSSREF](#)
48. Al Mheid I, Quyyumi AA. Vitamin D and cardiovascular disease: controversy unresolved. *J Am Coll Cardiol* 2017;70:89-100.
[PUBMED](#) | [CROSSREF](#)
49. Nitsa A, Toutouza M, Machairas N, Mariolis A, Philippou A, Koutsilieris M. Vitamin D in cardiovascular disease. *In Vivo* 2018;32:977-981.
[PUBMED](#) | [CROSSREF](#)
50. Bikle DD. Vitamin D: production, metabolism, and mechanisms of action. In: Feingold KR, Anawalt B, Boyce A, Chrousos G, Dungan K, Grossman A, et al., editors. *Endotext* [Internet]. South Dartmouth (MA): MDText.com, Inc.; 2000 [cited 2020 May 9]. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK278935/>.
51. Chen S, Law CS, Grigsby CL, Olsen K, Hong TT, Zhang Y, et al. Cardiomyocyte-specific deletion of the vitamin D receptor gene results in cardiac hypertrophy. *Circulation* 2011;124:1838-1847.
[PUBMED](#) | [CROSSREF](#)
52. Grimes DS, Hindle E, Dyer T. Sunlight, cholesterol and coronary heart disease. *QJM* 1996;89:579-589.
[PUBMED](#) | [CROSSREF](#)
53. Kendrick J, Targher G, Smits G, Chonchol M. 25-hydroxyvitamin D deficiency is independently associated with cardiovascular disease in the Third National Health and Nutrition Examination Survey. *Atherosclerosis* 2009;205:255-260.
[PUBMED](#) | [CROSSREF](#)
54. Wang TJ, Pencina MJ, Booth SL, Jacques PF, Ingelsson E, Lanier K, et al. Vitamin D deficiency and risk of cardiovascular disease. *Circulation* 2008;117:503-511.
[PUBMED](#) | [CROSSREF](#)
55. Giovannucci E, Liu Y, Hollis BW, Rimm EB. 25-hydroxyvitamin D and risk of myocardial infarction in men: a prospective study. *Arch Intern Med* 2008;168:1174-1180.
[PUBMED](#) | [CROSSREF](#)
56. Hlaing SM, Garcia LA, Contreras JR, Norris KC, Ferrini MG, Artaza JN. 1,25-vitamin D3 promotes cardiac differentiation through modulation of the WNT signaling pathway. *J Mol Endocrinol* 2014;53:303-317.
[PUBMED](#) | [CROSSREF](#)
57. Goleniewska B, Kacprzak M, Zielińska M. Vitamin D level and extent of coronary stenotic lesions in patients with first acute myocardial infarction. *Cardiol J* 2014;21:18-23.
[PUBMED](#) | [CROSSREF](#)
58. Pilz S, März W, Wellnitz B, Seelhorst U, Fahrleitner-Pammer A, Dimai HP, et al. Association of vitamin D deficiency with heart failure and sudden cardiac death in a large cross-sectional study of patients referred for coronary angiography. *J Clin Endocrinol Metab* 2008;93:3927-3935.
[PUBMED](#) | [CROSSREF](#)
59. Zhang R, Li B, Gao X, Tian R, Pan Y, Jiang Y, et al. Serum 25-hydroxyvitamin D and the risk of cardiovascular disease: dose-response meta-analysis of prospective studies. *Am J Clin Nutr* 2017;105:810-819.
[PUBMED](#) | [CROSSREF](#)
60. Goliasch G, Blessberger H, Azar D, Heinze G, Wojta J, Bieglmayer C, et al. Markers of bone metabolism in premature myocardial infarction (≤ 40 years of age). *Bone* 2011;48:622-626.
[PUBMED](#) | [CROSSREF](#)
61. Mathieu C, Gysemans C, Giulietti A, Bouillon R. Vitamin D and diabetes. *Diabetologia* 2005;48:1247-1257.
[PUBMED](#) | [CROSSREF](#)
62. Liu E, Meigs JB, Pittas AG, McKeown NM, Economos CD, Booth SL, et al. Plasma 25-hydroxyvitamin D is associated with markers of the insulin resistant phenotype in nondiabetic adults. *J Nutr* 2009;139:329-334.
[PUBMED](#) | [CROSSREF](#)

63. Scragg R, Sowers M, Bell C; Third National Health and Nutrition Examination Survey. Serum 25-hydroxyvitamin D, diabetes, and ethnicity in the Third National Health and Nutrition Examination Survey. *Diabetes Care* 2004;27:2813-2818.
[PUBMED](#) | [CROSSREF](#)
64. Chonchol M, Scragg R. 25-Hydroxyvitamin D, insulin resistance, and kidney function in the Third National Health and Nutrition Examination Survey. *Kidney Int* 2007;71:134-139.
[PUBMED](#) | [CROSSREF](#)
65. Pittas AG, Lau J, Hu FB, Dawson-Hughes B. The role of vitamin D and calcium in type 2 diabetes. A systematic review and meta-analysis. *J Clin Endocrinol Metab* 2007;92:2017-2029.
[PUBMED](#) | [CROSSREF](#)
66. Vitamin D supplement in early childhood and risk for Type I (insulin-dependent) diabetes mellitus. The EURODIAB Substudy 2 Study Group. *Diabetologia* 1999;42:51-54.
[PUBMED](#) | [CROSSREF](#)
67. Duprez D, de Buyzere M, de Backer T, Clement D. Relationship between vitamin D3 and the peripheral circulation in moderate arterial primary hypertension. *Blood Press* 1994;3:389-393.
[PUBMED](#) | [CROSSREF](#)
68. Kristal-Boneh E, Froom P, Harari G, Ribak J. Association of calcitriol and blood pressure in normotensive men. *Hypertension* 1997;30:1289-1294.
[PUBMED](#) | [CROSSREF](#)
69. Forman JP, Giovannucci E, Holmes MD, Bischoff-Ferrari HA, Tworoger SS, Willett WC, et al. Plasma 25-hydroxyvitamin D levels and risk of incident hypertension. *Hypertension* 2007;49:1063-1069.
[PUBMED](#) | [CROSSREF](#)
70. Sugden JA, Davies JI, Witham MD, Morris AD, Struthers AD. Vitamin D improves endothelial function in patients with type 2 diabetes mellitus and low vitamin D levels. *Diabet Med* 2008;25:320-325.
[PUBMED](#) | [CROSSREF](#)
71. Pfeifer M, Begerow B, Minne HW, Nachtigall D, Hansen C. Effects of a short-term vitamin D3 and calcium supplementation on blood pressure and parathyroid hormone levels in elderly women. *J Clin Endocrinol Metab* 2001;86:1633-1637.
[PUBMED](#) | [CROSSREF](#)
72. Burgess ED, Hawkins RG, Watanabe M. Interaction of 1,25-dihydroxyvitamin D and plasma renin activity in high renin essential hypertension. *Am J Hypertens* 1990;3:903-905.
[PUBMED](#) | [CROSSREF](#)
73. Li YC, Kong J, Wei M, Chen ZF, Liu SQ, Cao LP. 1,25-dihydroxyvitamin D3 is a negative endocrine regulator of the renin-angiotensin system. *J Clin Invest* 2002;110:229-238.
[PUBMED](#) | [CROSSREF](#)
74. Dong J, Wong SL, Lau CW, Lee HK, Ng CF, Zhang L, et al. Calcitriol protects renovascular function in hypertension by down-regulating angiotensin II type 1 receptors and reducing oxidative stress. *Eur Heart J* 2012;33:2980-2990.
[PUBMED](#) | [CROSSREF](#)
75. Rigby WF, Denome S, Fanger MW. Regulation of lymphokine production and human T lymphocyte activation by 1,25-dihydroxyvitamin D3. Specific inhibition at the level of messenger RNA. *J Clin Invest* 1987;79:1659-1664.
[PUBMED](#) | [CROSSREF](#)
76. Canning MO, Grotenhuis K, de Wit H, Ruwhof C, Drexhage HA. 1-alpha,25-Dihydroxyvitamin D3 (1,25(OH)(2)D(3)) hampers the maturation of fully active immature dendritic cells from monocytes. *Eur J Endocrinol* 2001;145:351-357.
[PUBMED](#) | [CROSSREF](#)
77. Aihara K, Azuma H, Akaike M, Ikeda Y, Yamashita M, Sudo T, et al. Disruption of nuclear vitamin D receptor gene causes enhanced thrombogenicity in mice. *J Biol Chem* 2004;279:35798-35802.
[PUBMED](#) | [CROSSREF](#)
78. Guillot X, Semerano L, Saidenberg-Kermanac'h N, Falgarone G, Boissier MC. Vitamin D and inflammation. *Joint Bone Spine* 2010;77:552-557.
[PUBMED](#) | [CROSSREF](#)
79. Takeda M, Yamashita T, Sasaki N, Nakajima K, Kita T, Shinohara M, et al. Oral administration of an active form of vitamin D3 (calcitriol) decreases atherosclerosis in mice by inducing regulatory T cells and immature dendritic cells with tolerogenic functions. *Arterioscler Thromb Vasc Biol* 2010;30:2495-2503.
[PUBMED](#) | [CROSSREF](#)

80. Manson JE, Bassuk SS, Lee IM, Cook NR, Albert MA, Gordon D, et al. The VITamin D and Omega-3 Trial (VITAL): rationale and design of a large randomized controlled trial of vitamin D and marine omega-3 fatty acid supplements for the primary prevention of cancer and cardiovascular disease. *Contemp Clin Trials* 2012;33:159-171.
[PUBMED](#) | [CROSSREF](#)
81. Oh J, Weng S, Felton SK, Bhandare S, Riek A, Butler B, et al. 1,25(OH)₂ vitamin D inhibits foam cell formation and suppresses macrophage cholesterol uptake in patients with type 2 diabetes mellitus. *Circulation* 2009;120:687-698.
[PUBMED](#) | [CROSSREF](#)
82. Al Mheid I, Patel RS, Tangpricha V, Quyyumi AA. Vitamin D and cardiovascular disease: is the evidence solid? *Eur Heart J* 2013;34:3691-3698.
[PUBMED](#) | [CROSSREF](#)
83. Kunadian V, Ford GA, Bawamia B, Qiu W, Manson JE. Vitamin D deficiency and coronary artery disease: a review of the evidence. *Am Heart J* 2014;167:283-291.
[PUBMED](#) | [CROSSREF](#)
84. Lee JH, O'Keefe JH, Bell D, Hensrud DD, Holick MF. Vitamin D deficiency an important, common, and easily treatable cardiovascular risk factor? *J Am Coll Cardiol* 2008;52:1949-1956.
[PUBMED](#) | [CROSSREF](#)
85. Ferder M, Inserra F, Manucha W, Ferder L. The world pandemic of vitamin D deficiency could possibly be explained by cellular inflammatory response activity induced by the renin-angiotensin system. *Am J Physiol Cell Physiol* 2013;304:C1027-C1039.
[PUBMED](#) | [CROSSREF](#)
86. Schleithoff SS, Zittermann A, Tenderich G, Berthold HK, Stehle P, Koerfer R. Vitamin D supplementation improves cytokine profiles in patients with congestive heart failure: a double-blind, randomized, placebo-controlled trial. *Am J Clin Nutr* 2006;83:754-759.
[PUBMED](#) | [CROSSREF](#)
87. Zhu Y, Mahon BD, Froicu M, Cantorna MT. Calcium and 1 alpha,25-dihydroxyvitamin D₃ target the TNF-alpha pathway to suppress experimental inflammatory bowel disease. *Eur J Immunol* 2005;35:217-224.
[PUBMED](#) | [CROSSREF](#)
88. Canning MO, Grotenhuis K, de Wit H, Ruwhof C, Drexhage HA. 1-alpha,25-dihydroxyvitamin D₃ (1,25(OH)₂D₃) hampers the maturation of fully active immature dendritic cells from monocytes. *Eur J Endocrinol* 2001;145:351-357.
[PUBMED](#) | [CROSSREF](#)
89. Shea MK, Booth SL, Massaro JM, Jacques PF, D'Agostino RB Sr, Dawson-Hughes B, et al. Vitamin K and vitamin D status: associations with inflammatory markers in the Framingham Offspring Study. *Am J Epidemiol* 2008;167:313-320.
[PUBMED](#) | [CROSSREF](#)
90. Hin H, Tomson J, Newman C, Kurien R, Lay M, Cox J, et al. Optimum dose of vitamin D for disease prevention in older people: BEST-D trial of vitamin D in primary care. *Osteoporos Int* 2017;28:841-851.
[PUBMED](#) | [CROSSREF](#)
91. Arnsen Y, Itzhaky D, Mosseri M, Barak V, Tzur B, Agmon-Levin N, et al. Vitamin D inflammatory cytokines and coronary events: a comprehensive review. *Clin Rev Allergy Immunol* 2013;45:236-247.
[PUBMED](#) | [CROSSREF](#)
92. Sokol SI, Srinivas V, Crandall JP, Kim M, Tellides G, Lebastchi AH, et al. The effects of vitamin D repletion on endothelial function and inflammation in patients with coronary artery disease. *Vasc Med* 2012;17:394-404.
[PUBMED](#) | [CROSSREF](#)
93. Scragg R, Stewart AW, Waayer D, Lawes CM, Toop L, Sluyter J, et al. Effect of monthly high-dose vitamin D supplementation on cardiovascular disease in the vitamin D assessment study: a randomized clinical trial. *JAMA Cardiol* 2017;2:608-616.
[PUBMED](#) | [CROSSREF](#)
94. Manson JE, Cook NR, Lee IM, Christen W, Bassuk SS, Mora S, et al. Vitamin D supplements and prevention of cancer and cardiovascular disease. *N Engl J Med* 2019;380:33-44.
[PUBMED](#) | [CROSSREF](#)
95. Wu Z, Wang T, Zhu S, Li L. Effects of vitamin D supplementation as an adjuvant therapy in coronary artery disease patients. *Scand Cardiovasc J* 2016;50:9-16.
[PUBMED](#) | [CROSSREF](#)
96. Molinari C, Uberti F, Grossini E, Vacca G, Carda S, Invernizzi M, et al. 1 α ,25-dihydroxycholecalciferol induces nitric oxide production in cultured endothelial cells. *Cell Physiol Biochem* 2011;27:661-668.
[PUBMED](#) | [CROSSREF](#)

97. Huang Z, Liu L, Huang S, Li J, Feng S, Huang N, et al. Vitamin D (1,25-(OH)₂D₃) improves endothelial progenitor cells function via enhanced NO secretion in systemic lupus erythematosus. *Cardiol Res Pract* 2020;2020:6802562.
[PUBMED](#) | [CROSSREF](#)
98. Grundmann M, Haidar M, Placzko S, Niendorf R, Darashchonak N, Hubel CA, et al. Vitamin D improves the angiogenic properties of endothelial progenitor cells. *Am J Physiol Cell Physiol* 2012;303:C954-C962.
[PUBMED](#) | [CROSSREF](#)
99. Brodowski L, Burlakov J, Myerski AC, von Kaisenberg CS, Grundmann M, Hubel CA, et al. Vitamin D prevents endothelial progenitor cell dysfunction induced by sera from women with preeclampsia or conditioned media from hypoxic placenta. *PLoS One* 2014;9:e98527.
[PUBMED](#) | [CROSSREF](#)
100. von Versen-Höyneck F, Brodowski L, Dechend R, Myerski AC, Hubel CA. Vitamin D antagonizes negative effects of preeclampsia on fetal endothelial colony forming cell number and function. *PLoS One* 2014;9:e98990.
[PUBMED](#) | [CROSSREF](#)
101. Barker DJ. Maternal and fetal origins of coronary heart disease. *J R Coll Physicians Lond* 1994;28:544-551.
[PUBMED](#)
102. Gluckman PD, Hanson MA, Cooper C, Thornburg KL. Effect of *in utero* and early-life conditions on adult health and disease. *N Engl J Med* 2008;359:61-73.
[PUBMED](#) | [CROSSREF](#)
103. Bodnar LM, Catov JM, Simhan HN, Holick MF, Powers RW, Roberts JM. Maternal vitamin D deficiency increases the risk of preeclampsia. *J Clin Endocrinol Metab* 2007;92:3517-3522.
[PUBMED](#) | [CROSSREF](#)
104. Robinson CJ, Wagner CL, Hollis BW, Baatz JE, Johnson DD. Maternal vitamin D and fetal growth in early-onset severe preeclampsia. *Am J Obstet Gynecol* 2011;204:556.e1-556.e4.
[PUBMED](#) | [CROSSREF](#)
105. Sipos PI, Rens W, Schlecht H, Fan X, Wareing M, Hayward C, et al. Uterine vasculature remodeling in human pregnancy involves functional macrochimerism by endothelial colony forming cells of fetal origin. *Stem Cells* 2013;31:1363-1370.
[PUBMED](#) | [CROSSREF](#)
106. Brodowski L, Schröder-Heurich B, Hubel CA, Vu TH, von Kaisenberg CS, von Versen-Höyneck F. Role of vitamin D in cell-cell interaction of fetal endothelial progenitor cells and umbilical cord endothelial cells in a preeclampsia-like model. *Am J Physiol Cell Physiol* 2019;317:C348-C357.
[PUBMED](#) | [CROSSREF](#)
107. Zhong W, Gu B, Gu Y, Groome LJ, Sun J, Wang Y. Activation of vitamin D receptor promotes VEGF and CuZn-SOD expression in endothelial cells. *J Steroid Biochem Mol Biol* 2014;140:56-62.
[PUBMED](#) | [CROSSREF](#)
108. Uberti F, Lattuada D, Morsanuto V, Nava U, Bolis G, Vacca G, et al. Vitamin D protects human endothelial cells from oxidative stress through the autophagic and survival pathways. *J Clin Endocrinol Metab* 2014;99:1367-1374.
[PUBMED](#) | [CROSSREF](#)
109. Stach K, Kälsch AI, Nguyen XD, Elmas E, Kralev S, Lang S, et al. 1 α ,25-dihydroxyvitamin D₃ attenuates platelet activation and the expression of VCAM-1 and MT1-MMP in human endothelial cells. *Cardiology* 2011;118:107-115.
[PUBMED](#) | [CROSSREF](#)
110. Brownlee M, Cerami A, Vlassara H. Advanced glycosylation end products in tissue and the biochemical basis of diabetic complications. *N Engl J Med* 1988;318:1315-1321.
[PUBMED](#) | [CROSSREF](#)
111. Makita Z, Radoff S, Rayfield EJ, Yang Z, Skolnik E, Delaney V, et al. Advanced glycosylation end products in patients with diabetic nephropathy. *N Engl J Med* 1991;325:836-842.
[PUBMED](#) | [CROSSREF](#)
112. Vlassara H, Fuh H, Donnelly T, Cybulsky M. Advanced glycation endproducts promote adhesion molecule (VCAM-1, ICAM-1) expression and atheroma formation in normal rabbits. *Mol Med* 1995;1:447-456.
[PUBMED](#) | [CROSSREF](#)
113. Talmor Y, Golan E, Benchetrit S, Bernheim J, Klein O, Green J, et al. Calcitriol blunts the deleterious impact of advanced glycation end products on endothelial cells. *Am J Physiol Renal Physiol* 2008;294:F1059-F1064.
[PUBMED](#) | [CROSSREF](#)
114. Ambros V. The functions of animal microRNAs. *Nature* 2004;431:350-355.
[PUBMED](#) | [CROSSREF](#)

115. Farh KK, Grimson A, Jan C, Lewis BP, Johnston WK, Lim LP, et al. The widespread impact of mammalian microRNAs on mRNA repression and evolution. *Science* 2005;310:1817-1821.
[PUBMED](#) | [CROSSREF](#)
116. Zitman-Gal T, Green J, Pasmanik-Chor M, Golan E, Bernheim J, Benchetrit S. Vitamin D manipulates miR-181c, miR-20b and miR-15a in human umbilical vein endothelial cells exposed to a diabetic-like environment. *Cardiovasc Diabetol* 2014;13:8.
[PUBMED](#) | [CROSSREF](#)
117. Zitman-Gal T, Golan E, Green J, Bernheim J, Benchetrit S. Vitamin D receptor activation in a diabetic-like environment: potential role in the activity of the endothelial pro-inflammatory and thioredoxin pathways. *J Steroid Biochem Mol Biol* 2012;132:1-7.
[PUBMED](#) | [CROSSREF](#)
118. Marco LJ, McCloskey K, Vuillermin PJ, Burgner D, Said J, Ponsonby AL. Cardiovascular disease risk in the offspring of diabetic women: the impact of the intrauterine environment. *Exp Diabetes Res* 2012;2012:565160.
[PUBMED](#) | [CROSSREF](#)
119. Gui J, Rohrbach A, Borns K, Hillemanns P, Feng L, Hubel CA, et al. Vitamin D rescues dysfunction of fetal endothelial colony forming cells from individuals with gestational diabetes. *Placenta* 2015;36:410-418.
[PUBMED](#) | [CROSSREF](#)
120. Sugden JA, Davies JI, Witham MD, Morris AD, Struthers AD. Vitamin D improves endothelial function in patients with type 2 diabetes mellitus and low vitamin D levels. *Diabet Med* 2008;25:320-325.
[PUBMED](#) | [CROSSREF](#)
121. Yu P, Song H, Gao J, Li B, Liu Y, Wang Y. Vitamin D (1,25-(OH)₂D₃) regulates the gene expression through competing endogenous RNAs networks in high glucose-treated endothelial progenitor cells. *J Steroid Biochem Mol Biol* 2019;193:105425.
[PUBMED](#) | [CROSSREF](#)
122. Jiang S, Li Y, Lin T, Yuan L, Li Y, Wu S, et al. IL-35 inhibits angiogenesis through VEGF/Ang2/Tie2 pathway in rheumatoid arthritis. *Cell Physiol Biochem* 2016;40:1105-1116.
[PUBMED](#) | [CROSSREF](#)
123. Chen X, Xiu M, Xing J, Yu S, Min D, Guo F. Lanthanum chloride inhibits LPS mediated expressions of pro-inflammatory cytokines and adhesion molecules in HUVECs: involvement of NF-κB-Jmjd3 signaling. *Cell Physiol Biochem* 2017;42:1713-1724.
[PUBMED](#) | [CROSSREF](#)
124. Liang B, Wang X, Zhang N, Yang H, Bai R, Liu M, et al. Angiotensin-(1-7) attenuates angiotensin II-induced ICAM-1, VCAM-1, and MCP-1 expression via the MAS receptor through suppression of P38 and NF-κB pathways in HUVECs. *Cell Physiol Biochem* 2015;35:2472-2482.
[PUBMED](#) | [CROSSREF](#)
125. Xu W, Hu X, Qi X, Zhu R, Li C, Zhu Y, et al. Vitamin D ameliorates angiotensin II-induced human endothelial progenitor cell injury via the PPAR-γ/HO-1 pathway. *J Vasc Res* 2019;56:17-27.
[PUBMED](#) | [CROSSREF](#)
126. Xiang W, Hu ZL, He XJ, Dang XQ. Intravenous transfusion of endothelial progenitor cells that overexpress vitamin D receptor inhibits atherosclerosis in ApoE-deficient mice. *Biomed Pharmacother* 2016;84:1233-1242.
[PUBMED](#) | [CROSSREF](#)
127. El-Magadmi M, Bodill H, Ahmad Y, Durrington PN, Mackness M, Walker M, et al. Systemic lupus erythematosus: an independent risk factor for endothelial dysfunction in women. *Circulation* 2004;110:399-404.
[PUBMED](#) | [CROSSREF](#)
128. Zeller CB, Appenzeller S. Cardiovascular disease in systemic lupus erythematosus: the role of traditional and lupus related risk factors. *Curr Cardiol Rev* 2008;4:116-122.
[PUBMED](#) | [CROSSREF](#)
129. Reynolds JA, Rosenberg AZ, Smith CK, Sergeant JC, Rice GI, Briggs TA, et al. Brief report: vitamin D deficiency is associated with endothelial dysfunction and increases type I interferon gene expression in a murine model of systemic lupus erythematosus. *Arthritis Rheumatol* 2016;68:2929-2935.
[PUBMED](#) | [CROSSREF](#)
130. Heaney RP. Vitamin D in health and disease. *Clin J Am Soc Nephrol* 2008;3:1535-1541.
[PUBMED](#) | [CROSSREF](#)
131. Omdahl JL, Morris HA, May BK. Hydroxylase enzymes of the vitamin D pathway: expression, function, and regulation. *Annu Rev Nutr* 2002;22:139-166.
[PUBMED](#) | [CROSSREF](#)

132. Rachez C, Freedman LP. Mechanisms of gene regulation by vitamin D(3) receptor: a network of coactivator interactions. *Gene* 2000;246:9-21.
[PUBMED](#) | [CROSSREF](#)
133. Swain LD, Schwartz Z, Boyan BD. 1,25-(OH)2D3 and 24,25-(OH)2D3 regulation of arachidonic acid turnover in chondrocyte cultures is cell maturation-specific and may involve direct effects on phospholipase A2. *Biochim Biophys Acta* 1992;1136:45-51.
[PUBMED](#) | [CROSSREF](#)
134. Yiu YF, Chan YH, Yiu KH, Siu CW, Li SW, Wong LY, et al. Vitamin D deficiency is associated with depletion of circulating endothelial progenitor cells and endothelial dysfunction in patients with type 2 diabetes. *J Clin Endocrinol Metab* 2011;96:E830-E835.
[PUBMED](#) | [CROSSREF](#)
135. Mikirova NA, Belcaro G, Jackson JA, Riordan NH. Vitamin D concentrations, endothelial progenitor cells, and cardiovascular risk factors. *Panminerva Med* 2010;52:81-87.
[PUBMED](#)
136. Cianciolo G, La Manna G, Cappuccilli ML, Lanci N, Della Bella E, Cuna V, et al. VDR expression on circulating endothelial progenitor cells in dialysis patients is modulated by 25(OH)D serum levels and calcitriol therapy. *Blood Purif* 2011;32:161-173.
[PUBMED](#) | [CROSSREF](#)
137. Berezin AE, Kremzer AA, Martovitskaya YV, Berezina TA. Non-classical progenitor mononuclears in metabolic syndrome: the role of serum 25-hydroxyvitamin D3. *Clin Med Biochem* 2016;2:1000115.
[CROSSREF](#)
138. Lo Gullo A, Mandraffino G, Bagnato G, Aragona CO, Imbalzano E, D'Ascola A, et al. Vitamin D status in rheumatoid arthritis: inflammation, arterial stiffness and circulating progenitor cell number. *PLoS One* 2015;10:e0134602.
[PUBMED](#) | [CROSSREF](#)
139. Chan YH, Lau KK, Yiu KH, Li SW, Tam S, Lam TH, et al. Vascular protective effects of statin-related increase in serum 25-hydroxyvitamin D among high-risk cardiac patients. *J Cardiovasc Med (Hagerstown)* 2015;16:51-58.
[PUBMED](#) | [CROSSREF](#)
140. Gurses KM, Tokgozoglu L, Yalcin MU, Kocyigit D, Dural M, Canpinar H, et al. Markers of subclinical atherosclerosis in premenopausal women with vitamin D deficiency and effect of vitamin D replacement. *Atherosclerosis* 2014;237:784-789.
[PUBMED](#) | [CROSSREF](#)
141. Yiu YF, Yiu KH, Siu CW, Chan YH, Li SW, Wong LY, et al. Randomized controlled trial of vitamin D supplement on endothelial function in patients with type 2 diabetes. *Atherosclerosis* 2013;227:140-146.
[PUBMED](#) | [CROSSREF](#)
142. Pilz S, Verheyen N, Gröbler MR, Tomaschitz A, März W. Vitamin D and cardiovascular disease prevention. *Nat Rev Cardiol* 2016;13:404-417.
[PUBMED](#) | [CROSSREF](#)