

Original Research

Exploring the Effects of a Neck Strengthening Program on Purposeful Soccer Heading Biomechanics and Neurocognition

Katelyn M. Waring, Edward R. Smith, Gary P. Austin, Thomas G. Bowman^a

Keywords: CNS vital signs, concussion prevention, cervical strength

<https://doi.org/10.26603/001c.38327>

International Journal of Sports Physical Therapy

Vol. 17, Issue 6, 2022

Background

Cervical (neck) strengthening has been proposed as an important factor in concussion prevention. The purpose of the study was to determine if a six-week cervical strengthening program affected neurocognition and purposeful soccer heading biomechanics. The hypothesis was that the neck strengthening program would improve strength, maintain neurocognition, and alter purposeful soccer heading biomechanics.

Study Design

Randomized controlled trial.

Methods

Twenty collegiate soccer athletes (8 males, 12 females, age=20.15±1.35 years, height=171.67±9.01 cm, mass=70.56±11.03 kg) volunteered to participate. Time (pre, post) and group (experimental, control) served as the independent variables. Four composite scores from the CNS Vital Signs computer based neurocognitive test (CNSVS; verbal memory, visual memory, executive function, reaction time) and aspects of heading biomechanics from inertial measurement units (xPatch; peak linear acceleration, peak rotational acceleration, duration, Gadd Severity Index [GSI]) served as the dependent variables. Each athlete completed a baseline measure of neck strength (anterior neck flexors, bilateral anterolateral neck flexors, bilateral cervical rotators) and CNSVS after heading 10 soccer balls at two speeds (11.18 and 17.88 m/s) while wearing the xPatch. The experimental group completed specific cervical neck strengthening exercises twice a week for six weeks using a Shingo Imara™ cervical neck resistance apparatus while the control group did not. After six weeks, the participants completed the same heading protocol followed by measurement of the same outcome variables. The alpha value was set to $p < 0.05$ *a priori*.

Results

The interaction between time and group was significant for visual memory ($F_{1,17}=5.16$, $p=0.04$, $\eta^2=0.23$). Interestingly, post hoc results revealed visual memory decreased for the control group from pretest (46.90±4.46) compared to posttest (43.00±4.03; mean difference=3.90, 95% CI=0.77-7.03, $p=0.02$). Interactions for all other dependent variables were not statistically significant ($p > 0.05$).

^a **Corresponding author:**

Thomas G. Bowman
1501 Lakeside Drive
Lynchburg, VA 24501
Bowman.t@lynchburg.edu
434-544-8869 (phone)
434-544-8365 (fax)

Conclusions

The cervical neck strengthening protocol allowed maintenance of visual memory scores but did not alter other neurocognitive measures or heading biomechanics. The link between cervical neck strengthening and concussion predisposition should continue to be explored.

Level of Evidence

Level 1b

INTRODUCTION

In sports across the globe, some of the most investigated injuries are those related to the head. Most people associate concussions and head-related injuries with American football. However, a study using the National Collegiate Athletic Association Injury Surveillance System found that 8.6% of injuries to women's soccer athletes were concussions as well as 5.8% of injuries in men's soccer athletes.¹ In fact, this number is potentially much larger considering more than 50% of concussions go unreported.²⁻⁶ Therefore, the number of concussions reported during soccer activity is likely a conservative measure of incidence.

In addition to concern over concussion in sport, repetitive head impacts without immediate clinical manifestation may cause long term consequences including negative brain health.⁷⁻¹³ On average, players head the ball six to 12 times in a competitive soccer match where the ball could travel over 22 m/s in amateur games.¹⁴ Since soccer is the world's most popular sport and includes more than 265 million players worldwide, soccer heading is a common cause for repetitive head impacts.¹⁵ However, the relationship between subconcussive heading and brain injury remains poorly understood.¹⁶⁻¹⁸ There have been multiple studies where researchers have found cognitive dysfunction or negative brain alterations after purposeful heading.^{2-6,10,13} However, another study has shown that there is no detrimental relationship between the number of purposeful headers and neurocognitive measures.¹⁸ Therefore, it remains unknown how repetitive head impacts may affect long term brain health.

Since head impact magnitude has been linked to injury predisposition,¹⁹ it is important to understand how to mitigate the energy received when the head is impacted during soccer participation. Purposeful heading includes repetitive low-level impacts and the magnitude transferred to the head can depend on many factors. Of these factors, neck strength is one that has received considerable recent attention in the literature. A correlation has been shown between neck strength and head impact kinematics where individuals with higher neck strength measurements had lower head accelerations upon impact.²⁰⁻²² Weaker mean overall neck strength was significantly associated with concussion and that for every pound of neck strength that athletes gain, their chances of a concussion decreases by five percent.²⁰ Therefore, the current study investigated a way to minimize head accelerations by testing one possible intervention. The purpose of the study was to determine if a six-week cervical strengthening program affected strength, neurocognition, and purposeful soccer heading biomechan-

ics. The following research questions underpinned the purpose: (1) Would a six-week neck strengthening program improve neck strength in soccer players?, (2) What effect would a six-week neck strengthening program have on head impact biomechanics during purposeful soccer ball heading?, and (3) What effect would a six-week neck strengthening program have on neurocognitive outcomes after a bout of purposeful soccer ball heading? The hypothesis was that the strengthening program would increase neck strength which would in turn decrease the impact magnitude the head experiences upon purposeful heading and maintain or improve (learning effect) players' neurocognition after purposeful heading.

METHODS

PARTICIPANTS

A total of 20 varsity collegiate soccer players participated in this study (12 female, 8 males, age = 20.15±1.35 years, height = 171.67±9.01 cm, mass = 70.56±11.03 kg). Each participant was over the age of 18 and signed a consent form approved by the host institution's human subjects review board. The participants also played a variety of positions such as forward (N=4), midfielder (N=8), defender (N=5) and goalie (N=3). A breakdown of demographics between the two groups can be found in [Table 1](#).

PROCEDURES

Prior to beginning data collection, the Institutional Review Board at the University of Lynchburg approved the study. All men's and women's soccer athletes at one institution sponsoring National Collegiate Athletic Association Division III athletics were sent an email that provided information on the study and were asked to respond if they were interested in participating. To begin the study, participants signed the informed consent form to be made aware of all the risks and benefits of the study. Next, the participants' height, mass, neck girth, and neck segment length were measured. Participants' height was measured in centimeters (cm) with a stadiometer (Seca Model 222, Hamburg Germany) and mass was measured in kilograms (kg) using a scale (Tanita BWB-800 Tokyo, Japan). While the participants were sitting straight and looking at an object at eye level, the head-neck segment length and neck girth was measured in cm with a metric tape measure. The participants' head-neck segment lengths were measured in a straight line using a tape measure from the seventh cervical vertebrae to the most superior region of the head ob-

Table 1. Demographic information across experimental and control groups.

	Experimental	Control	Comparison
Age	20.57±1.13	19.44±1.42	p=0.04, Cohen's d=1.03
Height	177.43±7.79	169.00±8.56	p=0.13, Cohen's d=0.79
Mass	75.94±11.44	66.23±9.93	p=0.09, Cohen's d=0.92
Neck Girth	37.50±2.78	34.06±2.79	p=0.03, Cohen's d=1.14
Neck Length	24.71±0.76	23.11±1.97	p=0.14, Cohen's d=0.77

served in the frontal plane.²³ The participants' neck girth was measured just above the thyroid cartilage.²⁴

After anthropometrics were recorded, half of the male and half of the female athletes were randomly selected for the experimental group while the other half served as the control group. A random number generator was used to determine group membership, and the researcher who collected all data was blinded to group membership. Before completing the baseline tests, the athletes participated in a neck warm up to reduce the possibility of injury. The neck warm up consisted of neck rotations (15 seconds (s) clockwise and 15 s counterclockwise) and neck stretching (two repetitions each of 15 s for flexion and extension).²³ Each athlete completed a series of neck strength tests, purposeful heading biomechanics tests, and a battery of cognitive tests to serve as baselines for a comparison later in the study. The experimental group followed a neck strengthening program for six weeks as part of the normal strength and conditioning program, and the control group did not perform neck strengthening exercises. However, both groups participated in the same soccer specific strength and conditioning program as prescribed by the team Certified Strength and Conditioning Specialist. After the six weeks of neck strength training, the participants completed the same series of neck strength tests, purposeful heading biomechanics tests, and battery of cognitive tests to compare to the baseline testing results.

NECK STRENGTH TESTING

One research team member measured neck strength as described by Kendall et al.²⁵ with an isometric dynamometer (MicroFET®2 Digital Handheld Dynamometer, Hogan Scientific, Salt Lake City, UT). Using an isometric dynamometer has been found to be a reliable and valid measure of cervical muscle strength in a seated position.²⁶ However, strength was measured in a supine or prone position as described by Kendall et al.²⁵ Strength measurements of the anterior neck flexors (Figure 1A; in supine the participants attempted to lift the head straight up), bilateral anterolateral neck flexors (Figure 1B; in supine the participants attempted to lift the head with the head turned completely in one direction/the other direction), bilateral cervical rotators (Figure 1C; in supine the participants turned the head to the left/right), and bilateral posterolateral neck extensors (Figure 1D; in prone participants raised their heads with their head completely turned to left/right) were recorded. In order to familiarize the participants with the




Figure 1. Positions for neck strength measurements: anterior neck flexors (A), bilateral anterolateral neck flexors (B), bilateral cervical rotators (C), bilateral posterolateral neck extensor (D)

testing method, they completed two practice trials followed by three recorded trials for each direction of motion. Participants rested for 30 seconds between each trial and each trial lasted for three seconds. The mean for each position was calculated based on the three trials and used for analysis.

PURPOSEFUL HEADING BIOMECHANICS TESTING

A JUGS soccer machine (JUGS Inc., Tualatin, OR) was used to simulate a kick that soccer players would be most likely to head in a game or practice. The speed of the ball coming out of the machine was adjusted to two different speeds to simulate low and high velocities of balls that players head during games or practices. A size five soccer ball pumped to 5,624.55 kgf/m² was dispensed out of the JUGS machine 35 m away from participants at 11.18 and 17.88 m/s as described in a previous study.¹ Each participant received five balls at each speed with a one-minute rest between balls. The participants were assigned a numbered inertial measurement unit (xPatch sensor, X2 Biosystems, Seattle, WA) that they wore while heading the 10 soccer balls. The sensor was applied over the right mastoid process with an adhesive patch. The xPatch sensor was used to measure the magnitude of the impact of the ball on the head in peak linear acceleration (PLA; g) and peak rotational acceleration (PRA; deg/s²), common metrics used to characterize head impact biomechanics as they are thought to be re-

lated to injury risk.¹⁹ The xPatch sensor also provided impact duration (time the ball and head made contact), and Gadd Severity Index (GSI). Higher durations would result in greater force transmission which may have clinical implications leading to the inclusion of duration as a dependent variable. The GSI scores provided information regarding the likelihood of an impact causing catastrophic head injury.²⁷ Although the xPatch has been shown to overestimate head impact magnitude,²⁸ it serves to facilitate frequency and magnitude comparisons between groups when all of the participants wear the sensors²⁹ which was the case in the current study. Head mounted sensors like the xPatch have been found to provide an accurate detection of the peak angular acceleration when compared to other helmet mounted systems.²⁹ Once the purposeful heading biomechanics test was completed, the sensor was removed, placed back on the charging dock, and the data were downloaded using a laptop computer (Apple Inc, Macbook Air, Cupertino, California).

COGNITIVE TESTING

After participants finished the purposeful heading biomechanics test, they completed the CNS Vital Signs (CNSVS) computer based cognitive test. The test was used to establish the participants' neurocognitive performance and symptom load. The CNSVS provided data regarding the participant's verbal memory, visual memory, executive function, reaction time, and symptom severity scores. Higher values indicate better performance for verbal memory, visual memory, and executive function. Lower reaction time scores indicate superior performance and higher symptom severity scores indicate more symptoms. Age adjusted standard scores for verbal memory, visual memory, and executive function were used while reaction time was measured in milliseconds. Symptom severity scores (range=0-144) summed the individual symptom scores on a 0-6 scale (0=none at all, 1-2=mild, 3-4=moderate, 5-6 severe). Scores were reported in the CNSVS detailed test result reports. CNSVS has been found to have reasonable test-retest reliability previously in broad populations.^{30,31}

NECK STRENGTHENING PROTOCOL

After the baseline testing was complete, the participants in the experimental group participated in a six-week neck strengthening program (three times per week) created by a Certified Strength and Conditioning Specialist. The Shingo Imara™ (Shingo Imara, Ann Arbor, Michigan) was used to provide resistance during neck strengthening exercises. The participants completed the number of repetitions and sets represented in [Table 2](#) for each of four exercises, resisted cervical flexion, extension, and lateral flexion on both sides from a seated position ([Figure 2](#)).

A Certified Strength and Conditioning Specialist monitored all exercise prescription, progression, and strength training sessions. The exercises were chosen because they were thought to target the neck muscles most utilized for purposeful heading.

Table 2. Strength training prescription during the six-week protocol

Week	Sets	Repetitions
Week 1	1	10
Week 2	1	12
Week 3	1	15
Week 4	2	10
Week 5	2	12
Week 6	2	15



Figure 2. Example of cervical extension strengthening exercise using the Shingo Imara (Shingo Imara, Ann Arbor, Michigan)

STATISTICAL ANALYSIS

The data were collected and organized in an Excel (2013 version, Microsoft Inc., Redmond, WA) spreadsheet for further analysis. In this study there were two independent variables, group (experimental and control, between factor) and time (pre and post, within factor). The dependent variables were neck strengthening measurements for each direction; PLA, PRA, duration, and GSI for heading biomechanics at each of the two speeds; and verbal memory, visual memory, executive function, reaction time, and symptom severity scores from the cognitive testing. Therefore, a 2x2 mixed model ANOVA was used to analyze the data from each dependent variable separately in SPSS (Version 26, IBM, Inc, Armonk, NY). Partial eta squared (η_p^2) was calculated as an effect size for the interactions and Cohen's d as an effect size for post hoc tests. Results were interpreted with $\eta_p^2=0.01$ indicating a small effect, $\eta_p^2=0.06$ indicating a medium effect and $\eta_p^2=0.14$ indicating a large

Table 3. Mean and standard deviations for strength testing dependent variables

	Strengthening Group		Control Group		Interaction
	Pre	Post	Pre	Post	
Anterior Strength	20.72±5.54	28.29±7.25	16.41±4.50	20.11±4.96	p=0.04, $\eta_p^2=0.22$
Right Anterolateral Strength	19.91±5.49	26.68±5.78	16.45±3.80	18.97±3.95	p=0.04, $\eta_p^2=0.22$
Left Anterolateral Strength	19.58±5.45	26.43±7.04	16.91±4.69	19.18±4.11	p=0.02, $\eta_p^2=0.28$
Right Rotation Strength	19.32±4.98	21.37±5.19	17.20±3.70	16.52±3.08	p=0.16, $\eta_p^2=0.11$
Left Rotation Strength	19.61±4.86	21.20±3.84	18.03±4.22	17.09±3.28	p=0.09, $\eta_p^2=0.16$
Right Posterolateral Strength	27.04±7.40	35.15±7.21	25.94±5.03	29.96±5.50	p=0.10, $\eta_p^2=0.15$
Left Posterolateral Strength	27.88±6.64	36.37±9.78	27.09±5.71	30.30±6.42	p=0.10, $\eta_p^2=0.14$

effect³² while Cohen's *d* was interpreted as *d*=0.2 as small, *d*=0.5 as medium, and *d*=0.8 as large.³³ The alpha value was set to *p*<0.05 *a priori* and Bonferroni post hoc tests were used to determine where significant pairwise differences existed for significant interactions.

RESULTS

NECK STRENGTH

Means and standard deviations for strength testing dependent variables can be found in [Table 3](#). The interaction between time and group was significant for anterior ($F_{1,17}=41.78$, *p*=0.04, $\eta_p^2=0.22$), right anterolateral ($F_{1,17}=4.80$, *p*=0.04, $\eta_p^2=0.22$), and left anterolateral ($F_{1,17}=6.48$, *p*=0.02, $\eta_p^2=0.28$) strength measurements. Post hoc tests showed strength improved pre- to post-intervention measures in the anterior direction for both the strengthening (*p*<0.001) and the control (*p*=0.02) groups, but only for the strengthening group for right anterolateral (*p*<0.001) and left anterolateral (*p*<0.001) directions. The interaction between time and group was not significant for right rotation ($F_{1,17}=2.12$, *p*=0.16, $\eta_p^2=0.11$, *1-β*=0.28), left rotation ($F_{1,17}=3.20$, *p*=0.09, $\eta_p^2=0.16$, *1-β*=0.39), right posterolateral ($F_{1,17}=2.98$, *p*=0.10, $\eta_p^2=0.15$, *1-β*=0.37), or left posterolateral ($F_{1,17}=2.86$, *p*=0.10, $\eta_p^2=0.14$, *1-β*=0.36) strength measurements.

PURPOSEFUL HEADING BIOMECHANICS

Means and standard deviations for heading biomechanics dependent variables can be found in [Table 4](#). There was no interaction present between group and time for PLA at 11.18 m/s ($F_{1,11}=0.66$, *p*=0.43, $\eta_p^2=0.06$, *1-β*=0.12) or PLA at 17.88 m/s ($F_{1,11}=0.98$, *p*=0.34, $\eta_p^2=0.08$, *1-β*=0.15). There was also no interaction present between group and time for PRA at 11.18 m/s ($F_{1,11}=0.003$, *p*=0.96, $\eta_p^2<0.01$, *1-β*=0.05) or for PRA at 17.88 m/s ($F_{1,11}=0.002$, *p*=0.97, $\eta_p^2<0.001$, *1-β*=0.05). There was no interaction present between group and time for head impact duration at 11.18 m/s ($F_{1,11}=0.41$, *p*=0.53, $\eta_p^2=0.04$, *1-β*=0.09) or at 17.88 m/s ($F_{1,11}=0.08$, *p*=0.79, $\eta_p^2=0.01$, *1-β*=0.06). Finally, there was no interaction present between group and time for GSI at 11.18 m/s ($F_{1,11}=1.03$, *p*=0.33, $\eta_p^2=0.09$, *1-β*=0.15) or at 17.88 m/s ($F_{1,10}=0.55$, *p*=0.48, $\eta_p^2=0.05$, *1-β*=0.10).

COGNITIVE TEST

Mean values and standard deviations for cognitive dependent variables can be found in [Table 5](#). When determining the effects on neurocognition and symptoms, the interaction between time and group was significant for visual memory ($F_{1,17}=5.16$, *p*=0.04, $\eta_p^2=0.23$). Interestingly, post hoc results revealed visual memory decreased for the control group from pretest (46.90±4.46) compared to posttest (43.00±4.03; mean difference=3.90, 95% CI=0.77-7.03, *p*=0.02, Cohen's *d*=0.92). The interaction between time and group was not significant for verbal memory ($F_{1,17}=0.01$, *p*=0.91, $\eta_p^2<0.001$, *1-β*=0.05), executive function ($F_{1,17}=0.71$, *p*=0.41, $\eta_p^2=0.04$, *1-β*=0.13), reaction time ($F_{1,17}=1.05$, *p*=0.32, $\eta_p^2=0.06$, *1-β*=0.16), or symptom severity score ($F_{1,17}=2.40$, *p*=0.14, $\eta_p^2=0.12$, *1-β*=0.31).

DISCUSSION

The purpose of the study was to determine if a six-week cervical strengthening program affected strength, neurocognition, and purposeful soccer heading biomechanics. The neck strengthening program only significantly improved strength for the muscles in the anterior and anterolateral (right and left) directions. In the second part of the study, there were no significant findings for the heading biomechanics portion of the experiment. Finally, the single significant finding for the neurocognitive component was a decrease in visual memory from pretest to posttest only in the control group.

NECK STRENGTH

The neck strengthening protocol significantly increased anterior and anterolateral neck strength. Rotational strength and posterolateral strength failed to improve despite strong effect sizes that indicate clinically meaningful strength improvement. The exercises selected may be the reason for the finding. Rotation, anterolateral, and posterolateral movements were not specifically targeted because the movements were not considered as important during purposeful heading. However, the protocol exactly as described by Kendall et al²⁵ was used. Despite the lack of significant improvement in rotational and posterolateral strength, the

Table 4. Mean and standard deviations for heading biomechanics dependent variables

	Strengthening Group		Control Group		Interaction
	Pre	Post	Pre	Post	
PLA at 11.18 m/s	22.70±2.11	20.01±1.59	22.10±2.72	25.21±5.44	p=0.43, $\eta_p^2=0.06$
PRA at 11.18 m/s	4456.20±916.54	3547.10±835.10	5027.84±793.66	4046.62±508.83	p=0.96, $\eta_p^2<0.01$
Duration at 11.18 m/s	10.35±0.99	8.78±1.28	11.50±3.22	15.13±5.95	p=0.53, $\eta_p^2=0.04$
GSI at 11.18 m/s	24.78±5.08	16.88±3.23	26.30±8.59	43.72±23.59	p=0.33, $\eta_p^2=0.09$
PLA at 17.88 m/s	30.84±4.45	22.83±3.29	25.24±1.45	20.89±3.56	p=0.34, $\eta_p^2=0.08$
PRA at 17.88 m/s	4279.75±1126.39	3383.56±946.99	4838.79±219.14	3843.23±614.17	p=0.97, $\eta_p^2<0.001$
Duration at 17.88 m/s	11.34±1.38	8.91±2.05	12.02±2.56	8.59±1.51	p=0.79, $\eta_p^2=0.01$
GSI at 17.88 m/s	41.98±12.13	22.30±7.04	31.70±4.11	19.51±5.39	p=0.48, $\eta_p^2=0.05$

PLA=peak linear acceleration, PRA=peak rotational acceleration, GSI=Gadd Severity Index

Table 5. Mean values and standard deviations for cognitive dependent variables

	Strengthening Group		Control Group		Interaction
	Pre	Post	Pre	Post	
Visual Memory	46.67±4.77	47.67±4.90	46.89±4.73	42.78±4.21	p=0.04, $\eta_p^2=0.23$
Verbal Memory	51.00±4.69	49.00±5.41	51.56±4.64	49.11±5.65	p=0.91, $\eta_p^2<0.001$
Executive Function	53.78±8.06	48.44±10.92	51.67±8.76	49.00±9.07	p=0.41, $\eta_p^2=0.04$
Reaction Time	598.67±48.61	638.56±45.20	630.00±73.40	638.00±109.01	p=0.32, $\eta_p^2=0.06$
Symptom Severity Score	9.22±12.03	4.56±6.23	7.44±7.32	11.67±14.82	p=0.14, $\eta_p^2=0.12$

Shingo Imara™ device was successful in improving neck strength with targeted exercise prescription and supervision by a Certified Strength and Conditioning Specialist. Improving neck strength may be important in athletic performance enhancement, even if the positive effect on head injury prevention remains not fully understood. Details regarding the findings related to the three purposes are provided in the sections below.

PURPOSEFUL HEADING BIOMECHANICS

Similar to the current findings, Mihalik et al.³⁴ also found that increasing neck strength failed to lead to lower linear and rotational accelerations of the head during impact. We also reported small to medium effect sizes suggesting limited clinical meaningfulness. It remains unknown why the increases in cervical strength observed failed to alter head impact biomechanics. Although neck musculature activity while performing the purposeful heading trials was not measured, the head impacts were anticipated by the participants as they knew the ball was coming toward them in both groups (experimental and control). Perhaps deep neck stability exercises would provide different results.

Others have shown that there is a correlation between neck strength and head impact kinematics where individuals with higher neck strength measurements had lower head accelerations upon impact.^{20–22} Weaker mean overall neck strength has been significantly associated with concussion and for every pound of neck strength that athletes gain, their chances of a concussion decreases by five percent.²⁰ Lamond et al.³⁵ found that anticipated headers, such as the ones in the current study, had lower linear and rotational head accelerations than unanticipated deflections or hits. Despite whether they were anticipated or not, head impacts to the front of the head, the maneuver used in the current study, have been found to create linear and rotational accelerations that were well below those associated with traumatic brain injury.³⁵ The magnitudes in the current study align similarly with the literature for similar impact types.³⁵

COGNITIVE TEST

Kaminski et al.¹⁷ used a test similar to CNSVS called Automated Neuropsychological Assessment Metrics (ANAM) to evaluate youth soccer players' neurocognition after purposeful soccer heading over a season and found that there was little to no relationship between heading and measures of neurocognitive performance. In another study by Kaminski et al.,¹⁸ collegiate and varsity high school women's soccer athletes had their cognitive function and balance evaluated before and after the course of a season while heading frequency was recorded. The study used the two-part Wechsler Digit Span test to determine cognitive function which included two of the same neurocognitive sections as the testing used in the current study, visual memory and verbal memory.¹⁸ Their results showed that there were no statistically significant changes in cognitive function or in balance. The researchers in these two studies did not determine if neck strength played a role in neurocognitive performance,

but findings did show that repetitive heading showed no change in neurocognitive performance. However, one study found cognitive and vestibular impairment immediately following a bout of 20 purposeful headers within three minutes.³⁶ The current findings suggest maintenance of visual memory (statistically significant differences and large effect size) for the neck strengthening group but no effect of group on verbal memory, executive function, or reaction time (no statistically significant differences and small effect sizes) immediately after 10 purposeful headers with longer rest periods between headers. On average, participants' visual memory scores decreased by 3.90 (95% CI=0.77-7.03) points in the control group. However, these decreases are within one standard deviation of normative data and would be unlikely to flag a patient as impaired.³⁰ Symptom severity score also did not change despite a medium to large effect size. The symptom severity score was included although it is not a measure of cognition because it is often used as part of concussion examination, either as part of the SCAT5 or computer-based neurocognitive testing.³⁷

LIMITATIONS AND FUTURE DIRECTIONS

The results could be explained by many different factors, but there are some that could be addressed and modified in future research. The goal was to observe the effect of strengthening exercises for the neck on heading biomechanics and neurocognition from pre- to posttest. However, the protocol to strengthen the participants' necks only strengthened the subjects' necks in the anterior and left and right anterolateral directions. In the future, researchers may obtain different results if the strengthening exercises are selected to attempt to strengthen all directions of the neck. Neck strength was measured in either a supine or prone position as described previously.²⁵ However, reliability and a validity of strength measures has only been measured in a seated position,²⁶ and cannot be assumed for measurements taken in supine or prone positions. Also Lamond et al.³⁵ found that unanticipated heading resulted in larger linear and rotational accelerations of the head than anticipated heading. It would be interesting to study the effects of neck strengthening during unanticipated head impacts as they are likely more concerning since they deliver higher accelerations to the head. Finally, neck muscle activation timing was not tested during purposeful heading impacts. Future research should determine if the timing of muscle activation is an important consideration with regards to reducing impact magnitude during purposeful heading.

CONCLUSION

The strengthening program the participants completed resulted in statistically significant improvements in anterior and anterolateral neck strength, but not in rotation or posterolateral strength. The neck strengthening protocol did not affect heading biomechanics of the collegiate soccer athletes in this study. Improving strength in the two anterior directions allowed maintenance of visual memory

scores but did not alter other neurocognitive measures following repetitive soccer heading. More research should be completed to determine the role neck strength plays in reducing head injury risk.

.....

CONFLICT OF INTEREST

The authors have no conflicts to disclose.

ACKNOWLEDGMENTS

Funded by the University of Lynchburg through the Percy Wootton Student-Faculty Research Award and the Schewel Student-Faculty Research Fund.

Submitted: March 29, 2021 CDT, Accepted: November 17, 2021 CDT



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-NC-4.0). View this license's legal deed at <https://creativecommons.org/licenses/by-nc/4.0> and legal code at <https://creativecommons.org/licenses/by-nc/4.0/legalcode> for more information.

REFERENCES

1. Bretzin AC, Mansell JL, Tierney RT, McDevitt JK. Sex differences in anthropometrics and heading kinematics among division i soccer athletes: a pilot study. *Sports Health*. 2017;9(2):168-173. doi:10.1177/1941738116678615
2. Baugh CM, Meehan WP III, Kroshus E, McGuire TG, Hatfield LA. College football players less likely to report concussions and other injuries with increased injury accumulation. *J Neurotrauma*. 2019;36(13):2065-2072. doi:10.1089/neu.2018.6161
3. Chrisman SP, Quitiquit C, Rivara FP. Qualitative study of barriers to concussive symptom reporting in high school athletics. *J Adolesc Health*. 2013;52(3):330-335.e3. doi:10.1016/j.jadohealth.2012.10.271
4. McCrea M, Hammeke T, Olsen G, Leo P, Guskiewicz K. Unreported concussion in high school football players: implications for prevention. *Clin J Sport Med*. 2004;14(1):13-17. doi:10.1097/00042752-200401000-00003
5. Register-Mihalik JK, Guskiewicz KM, McLeod TCV, Linnan LA, Mueller FO, Marshall SW. Knowledge, attitude, and concussion-reporting behaviors among high school athletes: a preliminary study. *J Athl Train*. 2013;48(5):645-653. doi:10.4085/1062-6050-48.3.20
6. Rivara FP, Schiff MA, Chrisman SP, Chung SK, Ellenbogen RG, Herring SA. The effect of coach education on reporting of concussions among high school athletes after passage of a concussion law. *Am J Sports Med*. 2014;42(5):1197-1203. doi:10.1177/0363546514521774
7. Koerte IK, Ertl-Wagner B, Reiser M, Zafonte R, Shenton ME. White matter integrity in the brains of professional soccer players without a symptomatic concussion. *JAMA*. 2012;308(18):1859. doi:10.1001/jama.2012.13735
8. Koerte IK, Lin AP, Muehlmann M, et al. Altered neurochemistry in former professional soccer players without a history of concussion. *J Neurotrauma*. 2015;32(17):1287-1293. doi:10.1089/neu.2014.3715
9. Koerte IK, Mayinger M, Muehlmann M, et al. Cortical thinning in former professional soccer players. *Brain Imaging Behav*. 2016;10(3):792-798. doi:10.1007/s11682-015-9442-0
10. Lipton ML, Kim N, Zimmerman ME, et al. Soccer heading is associated with white matter microstructural and cognitive abnormalities. *Radiology*. 2013;268(3):850-857. doi:10.1148/radiol.13130545
11. Matser EJT. Neuropsychological impairment in amateur soccer players. *JAMA*. 1999;282(10):971. doi:10.1001/jama.282.10.971
12. Svaldi DO, McCuen EC, Joshi C, et al. Cerebrovascular reactivity changes in asymptomatic female athletes attributable to high school soccer participation. *Brain Imaging Behav*. 2017;11(1):98-112. doi:10.1007/s11682-016-9509-6
13. Zhang MR, Red SD, Lin AH, Patel SS, Sereno AB. Evidence of cognitive dysfunction after soccer playing with ball heading using a novel tablet-based approach. *PLOS ONE*. 2013;8(2):e57364. doi:10.1371/journal.pone.0057364
14. Spiotta AM, Bartsch AJ, Benzel EC. Heading in soccer: dangerous play? *Neurosurgery*. 2012;70(1):1-11. doi:10.1227/neu.0b013e31823021b2
15. Levitch CF, Zimmerman ME, Lubin N, et al. Recent and long-term soccer heading exposure is differentially associated with neuropsychological function in amateur players. *J Int Neuropsychol Soc*. 2018;24(2):147-155. doi:10.1017/s1355617717000790
16. Caccese J, Kaminski TW. Neurocognitive changes in men's and women's soccer players across a collegiate career. In: *International Research in Science and Soccer II*. Routledge; 2015.
17. Kaminski TW, Cousino ES, Glutting JJ. Examining the relationship between purposeful heading in soccer and computerized neuropsychological test performance. *Res Q Exerc Sport*. 2008;79(2):235-244. doi:10.1080/02701367.2008.10599486
18. Kaminski TW, Wikstrom AM, Gutierrez GM, Glutting JJ. Purposeful heading during a season does not influence cognitive function or balance in female soccer players. *J Clin Exp Neuropsychol*. 2007;29(7):742-751. doi:10.1080/13825580600976911
19. Rowson S, Duma SM. Brain injury prediction: assessing the combined probability of concussion using linear and rotational head acceleration. *Ann Biomed Eng*. 2013;41(5):873-882. doi:10.1007/s10439-012-0731-0

20. Collins CL, Fletcher EN, Fields SK, et al. Neck strength: a protective factor reducing risk for concussion in high school sports. *J Prim Prev*. 2014;35(5):309-319. [doi:10.1007/s10935-014-0355-2](https://doi.org/10.1007/s10935-014-0355-2)
21. Eckner JT, Oh YK, Joshi MS, Richardson JK, Ashton-Miller JA. Effect of neck muscle strength and anticipatory cervical muscle activation on the kinematic response of the head to impulsive loads. *Am J Sports Med*. 2014;42(3):566-576. [doi:10.1177/0363546513517869](https://doi.org/10.1177/0363546513517869)
22. Streifer M, Brown AM, Porfido T, Anderson EZ, Buckman JF, Esopenko C. The potential role of the cervical spine in sports-related concussion: clinical perspectives and considerations for risk reduction. *J Orthop Sports Phys Ther*. 2019;49(3):202-208. [doi:10.2519/jospt.2019.8582](https://doi.org/10.2519/jospt.2019.8582)
23. Tierney RT, Sitler MR, Swanik CB, Swanik KA, Higgins M, Torg J. Gender differences in head-neck segment dynamic stabilization during head acceleration. *Med Sci Sports Exerc*. 2005;37(2):272-279. [doi:10.1249/01.mss.0000152734.447516.aa](https://doi.org/10.1249/01.mss.0000152734.447516.aa)
24. Plagenhoef S, Evans FG, Abdelnour T. Anatomical data for analyzing human motion. *Res Q Exerc Sport*. 1983;54(2):169-178. [doi:10.1080/02701367.1983.10605290](https://doi.org/10.1080/02701367.1983.10605290)
25. Kendall F, McCreary E, Provance P. *Muscles, Testing and Function*. 4th ed. Lippincott, Williams and Wilkins; 1993.
26. Kubas C, Chen YW, Echeverri S, et al. Reliability and validity of cervical range of motion and muscle strength testing. *J Strength Cond Res*. 2017;31(4):1087-1096. [doi:10.1519/jsc.0000000000001578](https://doi.org/10.1519/jsc.0000000000001578)
27. Gadd CW. *Use of a Weighted-Impulse Criterion for Estimating Injury Hazard*. SAE International; 1966. [doi:10.4271/660793](https://doi.org/10.4271/660793)
28. Wu LC, Nangia V, Bui K, et al. In vivo evaluation of wearable head impact sensors. *Ann Biomed Eng*. 2016;44(4):1234-1245. [doi:10.1007/s10439-015-1423-3](https://doi.org/10.1007/s10439-015-1423-3)
29. Cummiskey B, Schiffmiller D, Talavage TM, et al. Reliability and accuracy of helmet-mounted and head-mounted devices used to measure head accelerations. *Proc Inst Mech Eng Part P J Sports Eng Technol*. 2017;231(2):144-153. [doi:10.1177/1754337116658395](https://doi.org/10.1177/1754337116658395)
30. Gualtieri CT, Johnson LG. Reliability and validity of a computerized neurocognitive test battery, CNS Vital Signs. *Arch Clin Neuropsychol*. 2006;21(7):623-643. [doi:10.1016/j.acn.2006.05.007](https://doi.org/10.1016/j.acn.2006.05.007)
31. Littleton AC, Register-Mihalik JK, Guskiewicz KM. Test-retest reliability of a computerized concussion test: cns vital signs. *Sports Health*. 2015;7(5):443-447. [doi:10.1177/1941738115586997](https://doi.org/10.1177/1941738115586997)
32. Lakens D. Calculating and reporting effect sizes to facilitate cumulative science: a practical primer for t-tests and ANOVAs. *Front Psychol*. 2013;4:863. [doi:10.3389/fpsyg.2013.00863](https://doi.org/10.3389/fpsyg.2013.00863)
33. Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. Routledge Academic; 1988.
34. Mihalik JP, Guskiewicz KM, Marshall SW, Greenwald RM, Blackburn JT, Cantu RC. Does cervical muscle strength in youth ice hockey players affect head impact biomechanics? *Clin J Sport Med*. 2011;21(5):416-421. [doi:10.1097/jsm.0b013e31822c8a5c](https://doi.org/10.1097/jsm.0b013e31822c8a5c)
35. Lamond LC, Caccese JB, Buckley TA, Glutting J, Kaminski TW. Linear acceleration in direct head contact across impact type, player position, and playing scenario in collegiate women's soccer players. *J Athl Train*. 2018;53(2):115-121. [doi:10.4085/1062-6050-90-17](https://doi.org/10.4085/1062-6050-90-17)
36. Ashton J, Coyles G, Malone JJ, Roberts JW. Immediate effects of an acute bout of repeated soccer heading on cognitive performance. *Sci Med Footb*. 2020;0(0):1-7. [doi:10.1080/24733938.2020.1846769](https://doi.org/10.1080/24733938.2020.1846769)
37. McCrory P, Meeuwisse W, Dvorak J, et al. Consensus statement on concussion in sport—the 5th international conference on concussion in sport held in Berlin, October 2016. *Br J Sports Med*. Published online April 26, 2017;bjssports-2017-097699. [doi:10.1136/bjssports-2017-097699](https://doi.org/10.1136/bjssports-2017-097699)