

4. Gangemi S, Minciullo PL, Miroddi M, Chinou I, Calapai G, Schmidt RJ. Contact dermatitis as an adverse reaction to some topically used European herbal medicinal products - part 2: *Echinacea purpurea*-*Lavandula angustifolia*. *Contact Dermatitis*. 2015;72(4):193-205.
5. Farmer PW. The patch test as a means of diagnosis in contact dermatitis. *Med J Aust*. 1941;1:9-11.
6. Woelk H, Burkard G, Grünwald J. Benefits and risks of hypericum extract LI 160: drug monitoring study with 3250 patients. *J Geriatr Psychiatry Neurol*. 1994;7:S34-S38.

**How to cite this article:** O'Hagan T, Bisconti I, Leck C, et al. Delayed-type hypersensitivity reaction to St. John's wort extract confirmed by patch testing. *Contact Dermatitis*. 2022; 87(4):381-383. doi:10.1111/cod.14172

# Hand dermatitis aggravated by contact allergy to methylisothiazolinone in protective nitrile gloves

Raphaël André<sup>1</sup>  | Yassaman Alipour Tehrani<sup>1</sup>  | Aurélie Bugey<sup>2</sup> | Patrick Edder<sup>2</sup> | Pierre Piletta<sup>1</sup>

<sup>1</sup>Dermatology Department, Geneva Hospitals, University of Geneva, Geneva, Switzerland

<sup>2</sup>The Official Food and Veterinary Control Authority of Geneva, Geneva, Switzerland

## Correspondence

Raphaël André, Dermatology Department, Geneva Hospitals, 4 rue Gabrielle-Perret-Gentil, 1211 Geneva, Switzerland.

Email: [raphael.andre@hcuge.ch](mailto:raphael.andre@hcuge.ch)

## Funding information

Universite de Geneve

**KEYWORDS:** case report, contact dermatitis, hand dermatitis, methylisothiazolinone, nitrile gloves, protective gloves

Methylisothiazolinone (MI) is a frequent sensitizer in cosmetics, household detergents and industrial products.<sup>1</sup> In the occupational setting, the use of protective (nitrile) gloves is often recommended to limit exposure to this preservative, and to prevent allergic contact dermatitis. We here describe a case of hand dermatitis aggravated by MI sensitization, and we discuss different exposure sources.

## CASE REPORT

A 57-year-old non atopic man, working as a plumber, suffered from hand eczema since 10 years, which improved when being off work. Although he used protective nitrile gloves (ActivArmr Hycron 27-805; ANSELL), he described worsening of his hand dermatitis 3 days after resuming work. Patch tests were performed with the European baseline series, preservatives and a piece of his (used) gloves (semi-open test). Patch test materials were mounted on IQ Ultra patch test chambers (Chemotechnique Diagnostics) occluded for 2 days, and read on Day 2 (D2) and D4. We observed extremely positive (++++) reactions to MI 0.2% aq., but no reactions to methylchlorisothiazolone/MI 0.01%, benzisothiazolinone (BIT) 0.1% pet. nor to octylisothiazolinone (OIT) 0.1% pet. The test with the (used) glove showed + and ++ reactions on D2 and D4, respectively.

Chemical analysis of a series of used and new gloves were performed in the Official Food and Veterinary Control Authority (Table 1). Briefly, 0.5 g of nitrile was dissolved in 5 ml of methanol, sonicated during 15 min, and centrifuged during 10 min at 3500 RCF. After filtration, the extract was analysed by high pressure liquid chromatography coupled to a diode array detector following a previously published procedure.<sup>2</sup>

Significant concentrations of MI were identified in the patient's used nitrile glove (26.4 ppm). Interestingly, we also observed traces of MI in a new (unused) nitrile glove (4.9 ppm), and BIT in both new and used gloves (7.8 ppm and 6.9 ppm, respectively). OIT was present only in a new glove, again at only a low level (4.6 ppm). Details of protocol are explained in Appendix S1.

## DISCUSSION

Results pertaining to the new gloves, containing MI, as well as BIT and OIT, were interpreted as being likely due to the industrial manufacturing process of the gloves. Allergic contact dermatitis due to a high BIT content (>20 ppm) in (reusable) polyvinyl chloride gloves has previously been described.<sup>3</sup> In our case, however, neither BIT nor OIT showed any reactions on patch testing. Our case is interesting

**TABLE 1** Chemical analyses of new and used gloves

	MI content (mg/kg)	MCI content (mg/kg)	BIT content (mg/kg)	OIT content (mg/kg)	DCOIT content (mg/kg)
New glove (hand)	4.9	ND	7.8	4.6	ND
New glove (wrist)	4.4	ND	7.7	ND	ND
Used glove (hand)	26.4	ND	6.9	ND	ND
Used glove (wrist)	7.7	ND	5.3	ND	ND

Abbreviations: BIT, benzisothiazolinone; DCOIT, dichlorooctylisothiazolinone; MCI, methylchloroisothiazolinone; MI, methylisothiazolinone; ND, not detected; OIT, octylisothiazolinone.

because it demonstrates a combined mechanism of MI hand dermatitis: (i) although not traceable, a probable direct contact with MI present in (a) product(s) of the patient's working environment (canalisations and plumbing at private citizens home), (ii) confirmed contact with nitrile protective gloves, likely contaminated by MI from the work environment, and (iii) additional exposure to (low/residual levels of) MI in new protective gloves. Although in the present case it concerned nitrile gloves, it has been previously described that some rubber gloves (i.e., natural latex rubber) might not always prevent MI penetration<sup>4</sup>; however, this particular feature (penetration of MI through nitrile rubber gloves) was not investigated in our case. Finally, we cannot exclude that MI-dependent contact dermatitis was further maintained due to frequent exposure to rinse-off cosmetics or detergents.<sup>5</sup> In our patient, regular changes of new gloves every 3 days lead to a relative improvement, but no clearance, of his hand dermatitis. In practice, clinicians must be aware that, occasionally, protective nitrile gloves might contain isothiazolinones, inherently or by contamination, which might potentially aggravate existing hand dermatitis.

#### AUTHOR CONTRIBUTIONS

**Raphaël André:** Writing – original draft; investigation; validation.  
**Yassaman Alipour Tehrany:** Writing – review and editing. **Aurélié Bugey:** Conceptualization; investigation; writing – review and editing.  
**Patrick Edder:** Investigation. **Pierre Piletta:** Conceptualization; methodology; validation; writing – review and editing; investigation.

#### ACKNOWLEDGEMENT

Open access funding provided by Universite de Geneve. Open access funding provided by Universite de Geneve.

#### CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

#### ORCID

Raphaël André  <https://orcid.org/0000-0002-1362-7836>

Yassaman Alipour Tehrany  <https://orcid.org/0000-0001-9483-3238>

#### REFERENCES

1. Gonçalo M, Goossens A. Whilst Rome burns: the epidemic of contact allergy to methylisothiazolinone. *Contact Dermatitis*. 2013;68:257-258.
2. Alipour Tehrany Y, Quenan S, Bugey A, Piletta P. Allergic contact dermatitis caused by octylisothiazolinone in a leather sofa. *Contact Dermatitis*. 2018;79:188-189.
3. Aalto-Korte K, Ackermann L, Henriks-Eckerman M-L, et al. 1,2-Benzisothiazolin-3-one in disposable polyvinyl chloride gloves for medical use. *Contact Dermatitis*. 2007;57:365-370.
4. Espasandin-Arias M, Goossens A. Natural rubber gloves might not protect against skin penetration of methylisothiazolinone. *Contact Dermatitis*. 2014;70:249-251.
5. Herman A, Aerts O, Jacobs M-C, et al. Evolution of methylisothiazolinone sensitization: a Belgian multicentric study from 2014 to 2019. *Contact Dermatitis*. 2021;85:643-649.

#### SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

**How to cite this article:** André R, Alipour Tehrany Y, Bugey A, Edder P, Piletta P. Hand dermatitis aggravated by contact allergy to methylisothiazolinone in protective nitrile gloves. *Contact Dermatitis*. 2022;87(4):383-384. doi:[10.1111/cod.14173](https://doi.org/10.1111/cod.14173)