CORRECTION Open Access

Correction: Predicting the growth performance of growing-finishing pigs based on net energy and digestible lysine intake using multiple regression and artificial neural networks models

Li Wang, Qile Hu, Lu Wang, Huangwei Shi, Changhua Lai^{*} and Shuai Zhang^{*}

Correction: J Anim Sci Biotechnol 13, 57 (2022) https://doi.org/10.1186/s40104-022-00707-1

After publication of this article [1], it was brought to our attention that Figs. 2 and 3 were misplaced, the correct Figs. 2 and 3 are shown below:

The original publication has been corrected.

Published online: 09 October 2022

Reference

 Wang L, et al. Predicting the growth performance of growing-finishing pigs based on net energy and digestible lysine intake using multiple regression and artificial neural networks models. J Anim Sci Biotechnol. 2022;13:57. https://doi.org/10.1186/s40104-022-00707-1.

The original article can be found online at https://doi.org/10.1186/s40104-022-00707-1.

*Correspondence: laichanghua999@163.com; zhangshuai16@cau.edu.cn

State Key Laboratory of Animal Nutrition, College of Animal Science and Technology, China Agricultural University, Beijing 100193, People's Republic of China



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/loublicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data




Fig. 2 The response of ADG on different SID Lys intake (**a**) and NE intake (**b**). The curves were generated by the best fitted MR models in training. Only SID Lys intake and SID Lys intake² were considered as input variables in Fig. 2a while other variables were neglected. Only NE intake and NE intake² were considered as input variables in Fig. 2b




Fig. 3 The structure of the best-fitted artificial neural networks in predicting ADG (**a**) and F/G (**b**). H_1 was the value in the 1st node in the hidden layer; I_1 was the 1st input; am was the bias; O_1 was the value of the 1st output variable; H_1 was the value of the 1st node; D_1 was the bias; D_2 was the bias; D_3 was the activation function