

REVIEW

Fiber-enriched botanicals: A therapeutic tool against certain metabolic ailments

Waseem Khalid¹ | Muhammad Sajid Arshad¹  | Ayesha Jabeen¹ |
Faqir Muhammad Anjum^{2,3} | Tahira Batool Qaisrani⁴ | Hafiz Ansar Rasul Suleria⁵

¹Department of Food Science, Faculty of Life Sciences, Government College University, Faisalabad, Pakistan

²University of the Gambia, Banjul, The Gambia

³IFANCA Halal Apex (Pvt.) Limited, Faisalabad, Pakistan

⁴Department of Agricultural Engineering and Technology, Ghazi University, Dera Ghazi Khan, Pakistan

⁵Department of Agriculture and Food Systems, The University of Melbourne, Melbourne, Australia

Correspondence

Muhammad Sajid Arshad, Department of Food Science, Faculty of Life Sciences, Government College University, Faisalabad, Pakistan.

Emails: sajid_ft@yahoo.com, msajidarshad@gcuf.edu.pk

Abstract

Plant-based foods are natural sources including vegetables, fruits, cereals and legumes. These foods consist of various types of nutrients in which carbohydrate is the basic component. However, some plant-based diets contain carbohydrates in the form of fiber. The fiber is usually a nondigestible polysaccharide that is not digested in the human body. It is present in the form of soluble or insoluble in different part of foods like peel, bran, pulp and grain. Pectin, beta-glucan, mucilage, psyllium, resistant starch and inulin are soluble fiber, and cellulose, hemicellulose and lignin are insoluble fiber attained from plant foods. The major function enhances immunity by creating gastrointestinal barrier, mucus production, immune cell activity and IgA level. Previous evidences showed that peoples with strong immunity have fewer chances of viral disease. A recent viral disease named COVID-19 spread in the world and millions of peoples died due to this viral disease. Coronavirus mostly attacks humans that suffer with weak immune system. It is due chronic diseases like diabetes and CVD (cardiovascular disease). The current review shows that fiber-containing plant-based foods boost immunity and aid human against COVID-19. The therapeutic role of fiber in the human body is to control the risk of hypertension and diabetes because a high-fiber diet has the ability to lower cholesterol, blood pressure and blood sugar. Fibers aid in GIT (gastrointestinal tract) and prevent constipation because it absorbs water and adds bulk to stool.

KEYWORDS

COVID-19, diseases, fiber, immune system, plant-based foods

1 | INTRODUCTION

1.1 | Overview of fiber

Fiber, which is also called roughage. It is a component of plant-based foods (grains, nuts, beans, vegetables and fruits) which are not digested by our body (Thilagavathi et al., 2020). This nondigestible

part passed out from the body without any digestion process. As such, it keeps the digestive system healthy and clean, helps with the movement of the bowels and flushes cholesterol and other harmful substances out of the body (Kandasamy & Shanmugapriya, 2015).

Fiber is a carbohydrate that is not digested in the human digestive system (Mudgil & Barak, 2013). Whole fruits, beans, whole grains and vegetables are a great source of fiber (Erkkilä et al., 2005).

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *Food Science & Nutrition* published by Wiley Periodicals LLC.

The plant-based food sources of fiber and their types are presented in Table 1. Most carbohydrates are broken down into small sugar molecules (Meier et al., 2017). But the fiber cannot be broken into simple molecules and passes out of the body in undigested form (Buttriss & Stokes, 2008). Fiber aids in regulating the glycemic index in humans and helps to maintain hunger and blood sugar balance (Brownlee, 2011).

Dietary fiber is a plant-based nutrient that is also called bulk. It is a carbohydrate but unique from other carbohydrates. It is nondigestible, but some of the components are partially digestible. Therefore, it passes through the intestinal tract without any change (Soliman, 2019).

Dietary fiber is classified by origin, chemical composition and physicochemical properties. Along with that, it is subcategorized by the degree of polymerization, like chain length. Predominantly, all the properties of dietary fiber can impact fermentation by microbes (Mei et al., 2010). Plant-based fibers can be derived from fruits, vegetables, cereals, grains, legumes and nuts. The fiber originated from plants has varied chemical and physiochemical properties (McRorie & Fahey, 2013). The banana is a fruit that contains inulin which is a fructan and resistant starch, while the apple is rich in pectin. Diet having plant-based food is rich in different types of dietary fiber and provides the most diverse composition of the microbiota (Jha et al., 2019; Judprasong et al., 2011). Physiochemical characteristics, including solubility, viscosity and fermentability, influence the therapeutic effects of consumption along with fermentation (Schieber et al., 2001).

The lower retention of glucose and bile corrosive coupling is the physiological advantage of psyllium, gelatin and beta-glucan, which aid the growth of microbiota in the gastrointestinal tract. Polydextrose and soluble corn fiber are soluble fibers that are poorly digested by human alimentary enzymes (Holscher et al., 2015). Fructan is an inulin type-fiber which is present in artichokes,

bananas, garlic, agave, asparagus, leeks, wheat and onions (Holscher et al., 2014).

The level of polymerization of fructans are affected the maturation profile in people. Proof of physiological advantages of fructans in clinical investigations is restricted. The investigation of rodents has exhibited that the utilization of inulin-type strands differentially reduces body weight, convergence of blood glucose and blood cholesterol (Rendón-Huerta et al., 2012).

According to several studies, higher consumption of dietary fiber is beneficial for a variety of health outcomes, while many results showed that there has been no attempt to systematically capture the breadth of outcomes associated with intake of dietary fiber or assess the quality and strength of evidence on the association of dietary fiber intake, different medical conditions and health outcomes (Veronese et al., 2018). Generally, the fiber in food (fermented fiber) is the basis of a healthy diet. While prebiotics are specialized food ingredients that affect the final fermentation of products. It is beneficial for the host's health and specific bacteria (Platta, 2020). The study of rodents has demonstrated that the consumption of inulin-type fibers differentially reduces body weight and the concentration of blood glucose and blood cholesterol (Rendón-Huerta et al., 2012).

Holscher (2017) demonstrated the effect of prebiotics and fiber utilization on the metabolic capacity and composition of human gastrointestinal microbiota. The impacts of physiochemical properties of complex sugar can improved the health of human microbiota.

1.1.1 | Soluble fiber in plant-based foods

Soluble fiber is present in barley, nuts, beans, seeds, peas, lentils, fruits, vegetables and oat bran. This is also present in psyllium, which is a fiber supplement. This fiber is generally derived from the inner

TABLE 1 Types and plant-based food sources of fiber

Fiber	Type (soluble or insoluble)	Plant-based food sources	References
Pectin	Soluble	Pears, apples, guavas, quince, plums, gooseberries, oranges and other citrus fruits contain large amounts of pectin, while soft fruits, like cherries, grapes and strawberries, contain small amounts of pectin	(Dranca & Oroian, 2018)
Mucilage	Soluble	Flaxseeds, chia seeds, <i>Aloe vera</i> , okra and figs	(Muñoz et al., 2012)
Beta-glucan	Soluble	Barley, oats and whole grains	(Kaur et al., 2020)
Psyllium	Soluble	Psyllium husk	(Khalid et al., 2021; Qaisrani et al. 2014)
Resistant Starch	soluble	Oats, rice, grains, beans, legumes, potato, green bananas and maize	(Fuentes-Zaragoza et al., 2010)
Inulin	soluble	Onions, wheat, garlic, Jerusalem artichoke, chicory, root, wheat and barley	(Murphy, 2001; Roberfroid, 1993, 2005, 2007)
Cellulose	Insoluble	Bran, legumes, nuts, peas, roots, cabbage, apple skins, pineapple and banana leaves, sugarcane rind and coconut husks	Reddy & Yang, 2005)
Hemicellulose	Insoluble	Bran, nuts, legumes and whole grains	(Kalyani Nair et al., 2010)
Lignin	Insoluble	Pasta, rice, cereal, citrus fruits, whole grain and oat	(Mongeau & Brooks, 2001)

flesh of foods that are plant-based (Mudgil, 2017). The soluble fiber is mainly associated with the pulp or flesh of foods like oranges and potatoes (Lachman et al., 2005). Fructan is a type of inulin that is found in artichokes, bananas, garlic, agave, asparagus, leeks, wheat and onions (Holscher et al., 2014). The soluble fiber attracts water and gets converted into a gel by digestion. These fibers dissolve in water and form a viscous and gelatinous structure in the human digestive system (Wang et al., 2021).

Soluble fibers can change many metabolic and physiological processes involved in lowering cholesterol levels. These included stomach emptying, carbohydrate and fat digestion and absorption, nutrient intake, postprandial blood response, fermentation in the colon, liver metabolism and fecal excretion (Rideout et al., 2008).

Soluble dietary fibers are present in all food products separately or in combination with insoluble dietary fiber in small amounts, which contribute towards dietary fiber. Soluble dietary fibers are associated with a significant role in human physiological function. The potential benefits of soluble fibers are including a reduction in blood pressure and cholesterol levels, the prevention of gastrointestinal problems, protection against several types of cancer (including prostate, breast and prostate cancer) and increased bioavailability of minerals (Chawla & Patil, 2010).

Pectin is a soluble fiber which is present in jellies and jams, mixed with partially digested foods in the same way as gum, inulin, beta-glucan and mucilage (Yanniotis et al., 2007). The consumption of viscous forming and water-soluble fibers, including pectin and psyllium reduces the cholesterol level in human plasma (Butt et al., 2007).

Psyllium is included in nonfermentable fiber, while its greater viscosity and solubility cause unique therapeutic effects including reduction in blood cholesterol levels and improved glycemic control (Holscher et al., 2015).

The beta-glucan and pectin are highly fermentable along with high viscosity and solubility (Schieber et al., 2001). These fibers are found in whole grains including barley and oats (beta-glucan) and fruits like apples (pectin) (McRorie & Fahey, 2013). Reduced absorption of glucose and bile acid-binding are the physiological benefits of psyllium, pectin and beta-glucan which strongly impact the microbiota of the gastrointestinal tract. The soluble and nonviscous fibers are fermented by gastrointestinal microbiota like resistant maltodextrins, resistant starch, inulin, soluble corn fiber and ply-dextrose (Holscher et al., 2015).

The soluble fiber improves a healthy heart by modulating the cholesterol level in the body and lowering blood pressure. For example, pectin helps to reduce fat absorption in the human body from different foods, while beta-glucan is responsible for lowering the content of bad cholesterol. The soluble fiber lowers and regulates blood glucose levels in type 2 diabetic patients. Healthy blood glucose level leads to reduced inulin requirements in diabetics (Post et al., 2012).

1.1.2 | Insoluble fiber in plant-based foods

The insoluble fiber cannot dissolve in water and remains intact when food moves through the gastrointestinal tract. Fiber is generally

a part of plant-based foods which is not digested and absorbed in the human body (McRorie & McKeown, 2017). Wheat brans, beans, nuts, whole wheat flour and vegetables including kale, potatoes, cauliflower and green beans are good sources of insoluble fiber.

Insoluble dietary fiber is abundant, inexpensive and rich in dietary fiber as a by-product (Daou & Zhang, 2014). Advanced dietary fiber has been recently reported to promote heavy metal adsorption capacity by using chemical, physical and enzymatic methods (Feng et al., 2009; Park et al., 2013). The insoluble fiber is present in vegetables, whole grains and wheat bran. The insoluble fiber helps to cure constipation and complications like helping stool to pass out quickly through the intestine and bulking it up. Insoluble fiber helps to reduce the risk of colorectal cancer by speeding up the process of removal of waste from the digestive tract. The removal of harmful substances from the bloodstream of the human body depends upon the removal of water from the intestines along with stool. Due to lower loss of water from the body it is absorbed into the bloodstream again through intestinal walls (Huang et al., 2014).

The insoluble fiber aids in the passage of food from the body because it adds a bulky appearance to the stool and helps it pass quickly through the stomach and intestines, by increasing fecal bulk and helping the stool pass out easily (Forootan et al., 2018).

After entering the digestive system, insoluble fiber attracts water and sweeps waste materials from the large intestine. It is made up of cellulose and lignin, derived from the tough and outer skin of the plant. Generally, insoluble fiber is present in the skin of fruits and vegetables including pears, potatoes and apples. Foods packed with insoluble fiber generally have a chewy and tough texture. Insoluble fiber includes the skins of fruits and vegetables, the outer layer of cereal grains and wheat bran (Huang et al., 2014).

Cellulose is a dietary fiber that is poorly fermented by microbes present in the gut (Trompette et al., 2014). Insoluble fibers like cellulose are poorly fermented by microbes present in the gut. Their presence in the diet increases the transit rate of the gut and reduces the amount of time required for colonic bacterial fermentation of nondigested foodstuffs (McRorie, 2015).

2 | FIBER-RICH PLANT-BASED FOODS

2.1 | Fruits and their parts which are rich sources of fiber

The fruit is usually a sweet and fleshy product that contains pulp, skin and seeds. It is highly consumed in the world as food (Kumoro et al., 2020). These two types are fresh fruit and dried fruit, which are composed of different nutrients and many chemicals (Idah et al., 2007). Apples, pears, oranges, grapefruit, apricot, peach, plum, banana, mango, watermelon, melon, strawberries, raspberries, blueberries, kiwifruit, avocados, cherries, lemon, pineapple, pomegranate, papaya and guavas all contain both types of fiber.

Fruits are a rich source of soluble and insoluble fiber (Mcintosh & Miller, 2001). The fiber is present in the skin and pulp of fruits.

The pectin, gum, mucilage, beta-glucan and alginates are soluble fibers, and lignin or cellulose is an insoluble fiber attained from fruits (Bunzel et al., 2005).

The fruit ripening after harvesting causes three-dimensional hydrated cell wall fiber components pectin that completely dissembled cellulose and hemicellulose. It allow access to microbes and enhance fermentation susceptibility (Low et al., 2015). The fruit is eaten and its digestion by the gastrointestinal tract breaks it into smaller particles and damages the cell walls of the fruit, which empowers superior enzymatic fermentation and breakdown of colonic bacteria. Some starchy fruits like plantain and banana also contained fiber from both cell wall and resistant starch stored within the cell (Garcia-Mantrana et al., 2018). Basically, it is anticipated that the soluble fibers are completely fermented and equated quickly with the insoluble fibers, but freshly ripped fruits containing cell walls having insoluble fibers in a hybrid hydrated and moderately (Williams et al., 2017).

As per previous investigations, 100 g of dry apple contained about 46 g of insoluble fiber, 14 g of soluble fiber, 8–12 g of gelatin and fructooligosaccharide and 4.9% of complete dietary fiber correspondingly (Nawirska & Kwaśniewska, 2005). Similarly, banana contains around 43–49 g of dietary fiber, 1 g of inulin, 10–20 g of gelatin and 6 g of fructooligosaccharide in 100 g of dry matter (Emaga et al., 2007; Mohapatra et al., 2010). Indisputably, 100g of the husk of yellow enthusiasm organic product contains 20 g of gelatin and 60 g of complete insoluble dietary fiber (Yapo & Koffi, 2008). The level of polymerization of homogalacturonan in apple and banana strips and yellow enthusiasm natural product skin was depicted to be 78%, 67% and 93% individually (Yapo & Koffi, 2008).

The consumption of dietary fiber in the form of whole fruit has beneficial health effects including lowering LDL cholesterol, improving gut health, reducing cardiovascular diseases, reducing the risk of excessive weight gain, obesity, chronic heart diseases and type 2 diabetes, improving cancers and stroke (Veronese et al., 2018).

2.2 | Different types of vegetables and their constituents rich sources of fiber

Vegetables are also like fruits, have low-calorie fiber and can be consumed along with the skin of some fruits. Humans and animals consumed vegetables as food (Pennington & Fisher, 2009). The edible parts of plants are fruits, flowers, leaves, stems, seeds and roots. Brussels sprouts, artichokes, broccoli, collard greens, carrots, turnips, beets, celery root, Swiss chard, parsnips and kale are different types of vegetables (Khalid et al., 2021; Verkerk et al., 2009). Generally, the darker color of vegetables shows that they are rich in fiber content. Vegetables like beets, broccoli and carrots are rich in fiber. Artichokes contain the highest amount of fiber (a medium-sized artichoke contains 10 g of fiber) (Villanueva-Suárez et al., 2019).

Worldwide, potatoes are a staple vegetable and consumed largely. Potatoes are rich in starch and provide protein of high biological value which is not present in leafy vegetables (Camire et al.,

2009). Whole potatoes are rich in potassium, vitamin C and dietary fiber (Khalid et al., 2020). According to several prospective studies, there is a relationship between potato consumption and health (Halton et al., 2006).

2.3 | Different types of cereals and their constituents rich sources of fiber

Food products made with whole grain are rich in fiber. Cereals having wheat bran or oats are rich in fiber. On the other hand, cereals having rice or corn have less fiber. The whole grain cereals without excessive preservatives, flavoring, sugar and coloring showed more fiber than the highly processed cereals (Barrett et al., 2020). Oat bran is rich in soluble fiber which is beneficial for lowering the cholesterol level in the blood. Rice bran, wheat and corn are rich in insoluble fiber which is responsible for preventing constipation (Bernstein et al., 2013).

Oats and wheat are rich in fiber. Barley is a cereal that is often used in soups; 1 bowl of barley contains 6 g of fiber which is consumed in breakfast generally. Barley is rich in zinc which is a mineral that helps our body to build, repair and heal. Rice bran contains a low amount of dietary fiber up to 1.8–2.7 g per 100 g while oat bran contains 5.3–8.4 g per 100 g (Marlett et al., 1994).

Rice, wheat and oat have less hypolipidemic impact than some other grains because of less substance of dissolvable fiber. While rice-wheat has 12%–23% of oil, a higher rate than other grain sources, oil has unsaponifiable issue fixation of up to 4.2% (Gerhardt et al., 1998). This γ -oryzanol, β -sitosterol, tocotrienols and unsaturated fats contribute to the decrease in cholesterol. These all contribute to the decrease in cholesterol (Berger et al., 2005).

Diet contain water-dissolvable fiber that is played a basic part in decreasing the cholesterol level (Anderson et al., 1994). Oat gum, guar gum and gelatin are solvent strands which introduce hypocholesterolemic properties into humans (Cui & Roberts, 2009). As per an investigation, guar gum, oat wheat, gelatin, psyllium and insect bean gum reduce around 3%–20% of the cholesterol level (Anderson et al., 1994). The water-dissolvable filaments do not have any effect on cholesterol levels (AL-Rawi, 2007).

The capacity of different types of fiber to bind bile acids or bile salts is quite variable, and a high capacity seems to correspond with a hypocholesterolemic effect (Andersson & Hellstrand, 2012). This multiplies the union of bile salts, and catabolism of low-thickness lipoprotein is improved and constrained by the low-thickness lipoprotein receptors (Wood et al., 2007).

In vitro study suggested that bile acid-binding by rice, oat, wheat, and corn brans is determined using a mixture of bile acids normally secreted in human bile. The result showed that cholesterol-lowering by rice bran appears to be related to bile acid-binding. Bile acid-binding by oat bran (soluble fiber) may not be cholesterol-lowering. Bile acid binding by wheat bran can prevent cancer and provide healthful properties (Kahlon & Chow, 2000).

2.4 | Different types of legumes and their constituents rich sources of fiber

Legumes belong to a class of vegetables which also include peas, lentils and beans. These are among the most nutritious and versatile foods available globally (Graham & Vance, 2003). Peas, beans, lentils, peanuts, soybeans and chickpeas are some edible legumes. The types of legumes are differentiated on the basis of appearance, taste, use and nutrition. Legumes are generally used to describe the seeds of plants (Sánchez-Chino et al., 2015). Legumes generally have no cholesterol, low fat, high folate, iron, magnesium and potassium (Messina, 2014).

Legumes contain soluble and insoluble fiber and beneficial fats. The legumes have insoluble fiber which slows down the digestion process and releases carbohydrates gradually, which tends to help keep the blood sugar level at its optimum level (Trinidad et al., 2010). Generally, seeds of legumes comprise additional dietary fiber than cereals and are a virtuous source of metabolically active soluble dietary fiber (Hughes & Swanson, 1989).

Navy beans are one of the richest sources of fiber. Black beans contain a large amount of magnesium and iron. Pinto beans are consumed as a staple food in the United States. The total dietary fiber content fluctuates from lentils 11.5% to guar 33.2%. Guar is an annoying source of soluble dietary fiber. The ratio of water-soluble to insoluble fiber in legumes revealed that there is a higher percentage of insoluble fiber than soluble fiber. The foremost percentage of fiber in legumes comprises cellulose and hemicellulose, while a small proportion is of lignin and pectin. Guar seeds contain the highest amounts of lignin 2.0%, pectin 3.0% and cellulose 12.5%. Chickpeas are the best source of hemicellulose. The results of studies showed that chickpeas, field bean and guar contain more proportion of soluble fiber than other species of legumes, which are more nutritious and should be added to the diet (Khan et al., 2007).

3 | IMMUNITY-BOOSTING ROLE OF FIBER

Immunity is a complex biological system enriched in the ability to recognize and tolerate whatever belongs to the self and to perceive and dismiss. Immunity is the ability of multicellular living beings to oppose harmful microorganisms. Immunity has two types including innate and adaptive. Innate provides the first line of defense against infection whereas, adaptive provides a specific immune response directed at an invading pathogen. A plant-based eating routine reinforces your resistant immune system to secure you against germs and microorganisms. A sound safe immune system is fundamental for diminishing the danger of pathogens since it can perceive transformations in cells (Klasing, 2007).

Plant-based food sources expanded the intestinal valuable microbes which are useful and makeup 85% of the immune system (Arshad et al., 2020). The utilization of a lot of water, minerals like magnesium and zinc, micronutrients, spices, food plentiful in vitamins (C, D and E) and a better way of life can boost immunity against

different diseases. The study suggested that glutathione is the most abundant molecular weight antioxidant. It is played a defensive role against oxidative damage to cells (Polonikov, 2020). The plant-based nourishments play a fundamental part in improving the resistance of individuals to controlling COVID-19 (Arshad et al., 2020).

A high fiber plant-rich eating routine with a lot of natural products, vegetables and entire grains seems to help the development and support of valuable microorganisms. Certain accommodating organisms separate filaments into short-chain unsaturated fats, which have been appeared to invigorate invulnerable cell movement (Randhawa et al., 2015).

Fiber assumes a significant part in stomach-related wellbeing. Fiber is the fuel the colon cells use to keep them solid. Fiber likewise assists with keeping the stomach-related plot streaming, by keeping your solid discharges delicate and normal. An eating regimen wealthy in fiber adds to the upkeep of a solid gut microbiota related to expanded variety and capacities (Hecht, 2004).

Soluble and insoluble fibers are found in various food sources, for example, vegetables, nuts, seeds, natural products and oats. Nonetheless, not a wide range of fibers are available in similar food classes. Fiber is found in dull food sources, for example, cereals, vegetables and tubers, and nondeveloped natural products like green bananas. While gelatins are more advanced in foods grown from the ground vegetables, though beta-glucans and arabinoxylans are available in oats (Lovegrove et al., 2017). Although fibers are available in a wide range of plant-based food sources but utilization is low in Western nations. The stronghold of nourishments with extricated or orchestrated nonedible starches or their utilization in dietary enhancements accordingly establishes a procedure to expand fiber admission. A wide range of these carb polymers and oligosaccharides are industrially accessible (Deehan & Walter, 2016). A portion of these mixtures are considered "prebiotics" for the reason that they provide medical advantages by specifically prompting advantageous bacterial populaces in the gut (Bindels et al., 2015). The task of just certain fibers as prebiotics dependent on these models is by one way or another subjective (Deehan et al., 2018). Basically, all filaments will initiate explicit moves in microbiota arrangement because of serious connections and compositional movements add to medical advantages (Bindels et al., 2015). Interestingly, the systems that have been shown to add to medical advantages don't depend on a specific usage of the starch but instead on metabolic mixtures (Koh et al., 2016), physiological changes (pH bringing down) or security of the bodily fluid layer (Schroeder et al., 2018; Zou et al., 2018). It could merit considering a move in the focal point of the prebiotic idea away from the particular impact of explicit dietary segments on gut microbial networks and rather toward environmental and useful outcomes of fiber maturation (Bindels et al., 2015).

Fiber helps in digestion by enhancing bowel movement, supporting the immune system and providing food for the gut bacteria (Anderson et al., 2009). Bacteria present in our body also need food like living organisms to survive and fiber present in our daily diet is the food for these bacteria. The food for good bacteria is called

prebiotics, which are present in fiber-rich foods like garlic, onions and jackfruit. When the gut bacteria get fiber-rich foods called prebiotics, then these bacteria enhance the immune system, digestive health and any other health benefits. The eating of fiber on daily base that can enhance gut microbe and perform their function properly to improve the immune system and gut health. The gastrointestinal tract is the principal immune organ in humans (Slavin, 2013). The lymphoid tissues accompanying with gut comprise almost 60% of lymphocytes present in the human body including Peyer's patches, nonaggregated and intracellular lymphocytes. Figure 1 represents the mechanism of boosting immunity by fiber-rich plant foods.

The working of the immune system of the gut is influenced by the concentration of prebiotics in the diet which encourages the growth of health-promoting bacteria in the colon (Roberfroid, 2007; Vos et al., 2007). Most probiotics are nondigestible carbohydrates that are fermented in the colon (Roberfroid, 2005). Inulin and other oligofructoses are the most studied dietary fiber. It can help to promote the growth of *bifidobacteria* in the colon. *Bifidobacteria* and *Lactobacilli* are the health-promoting bacteria that generate short-chain fatty acids and also stimulate the immune system (Vos et al., 2007). Many animal studies have shown that inulin has a favorable effect on the immune system and human studies also supported this hypothesis (Roberfroid, 2007). Oligofructose and inulin are not assimilated by pancreatic and brush border enzymes due to their movement inside the colon deprived of any restraint.

Inulin is fermented absolutely by the microbacteria in the colon which stimulates the progression of bifidobacteria. The fermentation results in the production of short-chain fatty acids. The other fibers are also fermented in the gut to generate short-chain fatty acids. The effects of bifidobacteria on non-oligofructose fiber are not well defined yet. The health benefits of bifidobacteria are including depressing of intestinal pH to form acids after acclimatization of carbohydrates, protection from an intestinal infection, reduction of harmful bacteria,

manufacturing of vitamins and minerals, bulking activity to inhibit constipation, inspiration of immune response, reduction of risks of colorectal cancer, assistance with indigestion and fascination with minerals exclusively calcium (Roberfroid, 2007; Vos et al., 2007).

The study was conducted in which indicate that the supplementation of inulin increases the bacterial content in feces and has an important effect on circulating lymphocytes (Watzl et al., 2005). Oligofructoses and inulin have been studied completely and have the beneficial effect of fermentable soluble fibers on gum arabic, oat beta-glucan and others (Schley & Field, 2002).

The prebiotic fiber has a major role in the health and nutrition of infants which is generating great interest. The supplemented mixture of prebiotic fiber has various health benefits including a reduction in respiratory infections and atopic dermatitis, enhancing bowel function and promoting postnatal immune development (Veereman, 2007).

The potential of prebiotic fibers is to cure inflammatory bowel diseases. According to different studies on animals, it has been proved that prebiotic fiber reduces gut inflammation (Ewaschuk & Dieleman, 2006; Guarner, 2005). Recent study has indicated that prebiotic fibers reduce the risk of liver infection after having a liver transplant (Guarner, 2007). The previous human studies showed favorable responses of fibers against Cohn's disease, ulcerative colitis and pouchitis (Ewaschuk & Dieleman, 2006; Guarner, 2005). Pectin is a dietary fiber that has different structural characteristics. The structure of specific pectin influences the gastrointestinal immune barrier by interacting with immune cells or impacting intestinal microbiota directly. The pectin impact depends upon the structural characteristics of pectin including degree of esterification of methyl, acetylation, rhamnogalacturonan I and rhamnogalacturonan II neutral side chains (Prado et al., 2020).

The previous study reviewed that there is an interaction of the pectin structures with the immune barriers of gastrointestinal. The




FIGURE 1 Plants derived fiber (soluble or insoluble) enter into GIT (gastrointestinal tract) and performed some functions. The major function enhances immunity by creating gastrointestinal barrier, mucus production, immune cell activity and IgA level

effects of pectin include enhancing epithelial integrity, strengthening the mucus layer, inhibiting dendritic cell and macrophage response. The interaction between the immune barrier of the gastrointestinal and pectin is governed by recognition receptors included, Galectin 3, Toll-like receptors 2 and 4. Additionally, some of the pectin can stimulate the abundance and diversity of beneficial microbial communities. The immune barriers of gastrointestinal enhance the number of short-chain fatty acids. Pectin enhances the intestinal immune barrier by inhibiting the adhesion of commensal bacteria and pathogens to the epithelial cells. Different studies illustrated that pectin is a powerful dietary fiber which prevents inflammatory conditions (Beukema et al., 2020).

4 | FAVORABLE AGAINST DIFFERENT DISEASES

Fiber is naturally present in vegetables, fruits, cereals and legumes. There are several benefits to human health including heart health, stomach regulation, small or large intestine movement and preventing rectum cancer. Figure 2 shows the dietary sources of fiber and potential health benefits.

4.1 | COVID-19

Coronavirus disease (COVID-19) is an emerging respiratory infection caused by the severe acute respiratory syndrome coronavirus (SARS-CoV-2), which has in recent times become a pandemic (Lai et al., 2020). Most COVID-19 patients show mild to moderate symptoms, but about 15% of patients progress to severe pneumonia, and about 5% of patients eventually develop acute respiratory distress syndrome (ARDS), septic shock and multiple organ failure (Huang et al., 2020; Xu et al., 2020). Flu-like symptoms of COVID-19 usually appear 5–6 days after infection, and in some cases include cough, sore throat, fever, muscle and body aches and even loss of smell or taste (Galanakis, 2020).

In humans, coronaviruses are included in the spectrum of viruses that cause the common cold and the recent severe acute respiratory syndrome. Emerging infectious diseases such as SARS-CoV-2

pose a major threat to public health. The new coronavirus has spread speedily to many countries and has been declared a pandemic by the World Health Organization. It usually causes the virus infection of COVID-19. It is very likely that people with low immune response are infected (Arshad et al., 2020). The different entry routes of COVID-19 into the body as shown in Figure 3.

The host immune response to SARS-CoV-2 seems to play a key role in the pathogenesis and clinical manifestations of the disease. Severe acute respiratory syndrome coronavirus not only activates the antiviral immune response, but also causes an unrestrained inflammatory response characterized by the pronounced release of pro-inflammatory cytokines in patients with severe COVID-19, resulting in lymphopenia and lymphocytes. However, it is the abnormalities of granulocytes and monocytes. These immune abnormalities caused by SARS-CoV-2 may lead to microbial infections, septic shock and severe multiple organ dysfunction (Zhu et al., 2020). For that reason, it is necessary to clarify the underlying mechanism of immune abnormality in COVID-19 patients to guide the clinical management of the disease. In addition, reasonable treatment of the immune response of SARS-CoV-2 (including enhancing antiviral immunity while suppressing systemic inflammation) may be the key to successful treatment (Lin et al., 2020).

Patients with COVID-19 and hypertension or diabetes are more probable to develop serious diseases. Delineating the mechanisms by which these chronic diseases lead to disease deterioration and a better understanding of the escape mechanism of SARS-CoV-2 may provide clues for the clinical treatment of severe cases (Sanders et al., 2020).

In the new era, the food industry and food supply chain should solve four important issues. First, as consumers want to protect themselves and their immune systems by adopting healthier diets, the availability of bioactive ingredients in foods and functional foods become critical as the demand for these products may increase. Secondly, in order to prevent the virus from spreading among producers, retailers and consumers, food safety is an important issue. Third, food security problems have arisen because one billion people are locked in houses. Last but not least, the sustainability of the food system during the pandemic is another issue that the sector should address in order to limit related crises in the future (Galanakis, 2020).




FIGURE 2 Major dietary sources of fiber and potential health benefits




FIGURE 3 There are different entry routes of COVID-19 in to the body. Corona viruses enter into the body through the nose, mouth and eyes. Viruses enter into brain and lungs and attack nerve and bronchiole cells. However, the virus damages the cells and creates a health problem

The previous study indicates that the immune system is directly related to viral diseases patients. The decrease in peripheral T cell subsets is an exceptional characteristic in patients suffering with severe acute respiratory syndrome (SARS-COV-2) (Li et al., 2004). In patients recovered from this disease, a prompt reinstatement of peripheral T cell subsets is identified. Thus, peripheral T cell number can assist as a precise diagnostic tool for SARS-COV-2 (Li et al., 2004). The comparable phenomenon was also described in another study, in which the immune system was proved to be diminished during SARS-COV-2 (Cui et al., 2003). With the extrication of the association between immune responses and COVID-19, immune characteristics are now being documented as potential biomarkers for disease progression as well as potential therapeutic targets for COVID-19 (Perico et al., 2020).

The foods that originated from plants are intended to be used to improve the immunity of groups of people belonging to all ages against COVID-19 (Arshad et al., 2020). It withdraws perceptions about the properties of bioactive ingredients present in foods and herbs for the provision of the immune system of humans against infections before deliberating the possibility of transmission of COVID-19 through the food chain. It also highlights the food security issues globally arising from the fact that one-third of the world's population is on lockdown. Finally, it emphasizes the prominence of sustainability in the food chain in order to avoid or reduce the incidence of food and health crises in the future (Galanakis, 2020).

4.2 | Fiber reduces the risk of constipation

Constipation is a condition when fecal material moves through the colon very slowly (Sinclair, 2011). The fluid in the stool is absorbed back into our body, stool becomes dry and hard, which makes stool difficult to pass through the colon. Inadequate sleep, anxiety, poor nutrition, emotional stress, age and limited exercise are the possible causes of constipation. Various other diseases are also responsible for constipation, associated with sudden weight loss, fatigue, pain, bloody stool and changes in bowel habits (Costilla & Foxx-Orenstein, 2014).

Fiber is a plant-based food which is not digested in the human digestive system (García Ochoa et al., 2008). The two types of the

fiber are soluble and insoluble fibers (Wang et al., 2002). Soluble fiber helps to enhance the bulk form of the stool. Apples, bananas, beans, oats and barley are good sources of soluble fiber. Insoluble fiber helps to enhance the transit of food through the digestive system which prevents constipation. Whole grains, wheat bran, legumes and vegetables are good sources of insoluble fiber. Some foods contain both soluble and insoluble fibers (Fernández-López et al., 2009; Vitaglione et al., 2008).

The soluble fiber makes the stool bulky (McRorie & McKeown, 2017). Apples, oats, beans, barley and bananas are included in the foods which are rich in soluble fiber. The insoluble fiber helps to enhance the resistance of food in the digestive system and helps to prevent constipation. The dietary fiber present in food increases the weight and size of the stool and softens it. The bulky formed stool passes easily and lessens the risks of constipation. In the case of loose and watery stool, fiber is consumed to solidify the stool because the fiber absorbs water and maintains the bulky form of the stool (Foorootan et al., 2018). Fiber helps to maintain bowel health. Fibers present in cereal have a cell wall that resists digestion and retains water in the cellular structures of fiber. Wheat bran acts as an effective natural laxative (McDougall et al., 1996).

Fresh fruits, vegetables and legumes including lentils and beans should be consumed. The fiber present in legumes and citrus fruits helps to stimulate the growth of colonic flora, increase the weight of the stool and enhance the number of bacteria in the stool. The growth of bacteria present in the colon enhances the health of the intestines. Adults should consume water-rich foods like vegetables and fruits which are rich in fiber and help against constipation. They should be consumed according to the daily recommendation. Some studies have shown that there is a relationship between lifestyle, nutritional status and body mass index (BMI) (Karabudak et al., 2019).

Total dietary fiber has physiological effects which increase the movement in the intestine. The fiber in food accelerates the functions of the intestine by reducing the transit time of stool and its softening (Weber et al., 2014). During clinical practice, the initial management of primary constipation is done with the increased intake of fiber (Barichella et al., 2016). According to Western-style societies, in less developed countries the main cause of constipation is the low intake of fiber. Many societies like Western Africa have made a guideline that average dietary fiber intake should include

35g of insoluble fiber, to ease the bowel movement (Sura & Christie, 2014).

An adequate amount of fiber intake softens the stool and then decreases the risk of constipation. According to previous data, the rate of constipation was higher in people whose diet was deficient in fiber (Beleli et al., 2015; Bunn et al., 2015). An adequate amount of fluid should be taken to normalize the functions of the bowels. Additionally, the increased intake of fluid can increase the efficacy of fiber-rich foods (Sura & Christie, 2014). The nutritional status and body weight affect constipation, which is a common problem in obese patients (Pashankar & Loening-Baucke, 2005).

According to previous literature, there is a direct relationship between an individual's constipation level and intake of dietary fiber (Kranz et al., 2012). While, according to limited studies, there is also a relationship between nutritional status, fluid intake, body mass index (BMI) and lifestyle (Misra et al., 2006).

The study was conducted to investigate the relationship between lifestyle, body weight and constipation in young healthy adults. The health of the digestive system plays an important role in our active everyday life like being free from diarrhea, bowel syndrome, bloating, abdominal pain, inflammatory bowel disease, flatulence and constipation. Dietary fibers are used to maintain the proper health of the digestive system, but there are not seen any comprehensive solutions for the health maintenance of the digestive system (Forootan et al., 2018).

Another previous review focused on summarizing the health effects of guar fiber on the digestive system. While guar fiber has been emerged to be credible for such solutions (Noack et al., 2013). According to studies, a regular intake of 5–10 g/d of guar fiber is beneficial for treating the diseases associated with the digestive system. Guar fiber is a natural fiber. It is beneficial for the protection and promotion of the health of the digestive system either separately or in combination with probiotics (Rao & Quartarone, 2019).

In children, constipation depends upon several symptoms which are likely to persist till adulthood. Whole grains and dietary fiber are used as the treatment of constipation during childhood. According to the recommendations, the fiber intake by children varies generally among the organizations. The increased intake of dietary fiber and whole grains can relieve constipation during childhood, while some additional trials are necessary for the normal recommendations (Stewart & Schroeder, 2013).

4.3 | Colorectal cancer and fibers

Dietary fibers belong to a group of heterogeneous compounds which may vary in their biological effects. High-fiber diet intake is responsible for reduction in the risk of colorectal cancer. The researchers assay that the fiber from whole grains and cereals can reduce the risk of colorectal cancer. Fiber can reduce the risks of colon cancer due to lessen transit time of the gastrointestinal and

increasing the stool bulk and fermentation of volatile fatty acids (Hullings et al., 2020).

Fiber-rich food lowers the risk of hemorrhoids and small pouches in the human colon. Different studies have shown that fiber-rich diet lessens the risk of colorectal cancer. Some of the fibers are fermented in the colon (Alonso-Coello et al., 2006).

4.3.1 | Normalizes bowel movements

The dietary fiber present in the diet increases the weight and size of the stool and softens it. Bulky stool can be passed easily and decreases the chances of constipation. In the case of loose and watery stool, fiber-rich food should be consumed which solidify the stool by absorbing water and adding bulk to the stool (Nagarajan et al., 2015).

4.3.2 | Helps maintain bowel health

Diet rich in fiber can reduce the chances of developing hemorrhoids and small pouches in human colon. According to different studies, high-fiber diet lowers the risk of colorectal cancer. Some of the fiber is fermented in the colon (Alonso-Coello et al., 2006).

Dietary fiber plays a major role in the regulation of normal functions of intestines in the maintenance of healthy intestinal mucosa (Montagne et al., 2003). Some evidence has shown that the diet deficient in fiber predisposes to colon carcinogenesis (Zhu et al., 2011). The complication is that there is not any recommended level of fiber intake till now. The physicians advice their patients to consume moderate diet containing fruits, vegetables and whole grains (Demark-Wahnefried et al., 2008). This provided a varied source of fiber-rich foods, and their consumption optimizes the transit of the intestines depending upon the needs of every individual. Fiber supplements have been shown to be effective against colon (Klurfeld et al., 2018). Study suggested that diet rich in fiber like whole grain foods instead of fat-rich foods can reduce the colon cancer. The results showed that diet rich in fiber like whole grain foods should be consumed instead of fat-rich foods (Larsson et al., 2005).

Studies have shown that there is a reduced risk of colon cancer when people eat diet high in total fiber and certain whole grain foods instead of high fat foods (Kyrø et al., 2013). The animal models have shown that the dietary fiber has inhibitory effects on neoplasms depending on the nature and source of fiber (Young et al., 2005). Wheat bran inhibits the colon tumor more consistently than other types of fibers like oat and corn bran due to the presence of phytic acid which inhibits colon carcinogenesis. Human studies have revealed that the dietary fiber has modifying effect on bacterial enzymes which are involved in the production of colon tumors depending upon the fiber consumption (Holscher, 2017). Wheat bran significantly decreases the level of tumor promoters present in the colon which is not done by oat or corn bran (Zhu et al., 2011).

4.4 | Reduce the blood sugar (diabetes) and cholesterol level

The soluble fiber dissolves in the human stomach and forms a jelly and goopy fluid, and the insoluble fiber stays as such in the human digestive system. Soluble and insoluble fibers both help with digestion, while soluble fiber has some other advantages also. Soluble fiber-rich foods are slowly digested than processed and refined foods like cakes, cookies and white bread. Dietary fiber affects the activity of enzymes present in the intestine and then regulates the metabolic activity of the body. It lessens the speed of digestion which relieves the release of glucose and control the blood sugar up to an optimal level. Sugar is absorbed from food by the human body very slightly which increases blood sugar level slowly and prevents the human body from a sudden increase in blood sugar level (Brennan & Tudorica, 2008).

According to many studies, high carbohydrates in the diet have adverse effects but can be minimized by increasing the content of carbohydrates and fiber simultaneously in diabetic patients' diet (Czyzewska-Majchrzak et al., 2014). Particularly, many studies demonstrated that a diet rich in carbohydrates and fiber can enhance the glucose control of blood and lessen the cholesterol level of plasma by comparing the diet low in carbohydrates and fibers for diabetic patients. Additionally, a diet rich in fiber and carbohydrates cannot increase insulin and triglyceride concentration in plasma (Reynolds et al., 2019).

The ability of dietary fibers to retard nutrient absorption and food digestion has a very important impact on the metabolism of carbohydrates and lipids. The beneficial effects of fiber-rich foods are also exerted by foods not rich in fiber. The fiber content and physical form of food influence the accessibility of nutrients by the activity of enzymes, which delay the digestion and absorption processes. Identification of foods with low glycemic response helps to enlarge the number of foods mainly for diabetic patients. A diet deficient in saturated fat and cholesterol is mainly recommended to diabetic patients to control cardiovascular diseases. A diet rich in saturated fat and cholesterol should be replaced by a diet rich in fiber and unsaturated fat and consumed by diabetic patients. High intake of dietary fiber, mainly soluble fiber can improve glycemic control, lower the lipid concentrations of plasma and lessen hyperinsulinemia in diabetic patients of type 2 (Guo et al., 2018).

Fiber can reduce glucose levels in the blood. Fiber inhibits the glucose passage from intestinal walls to bloodstream and controls the blood glucose levels. According to previous studies, there is a significant relationship between fiber intake and blood glucose level in diabetic patients. Study suggested that there is a link between fiber intake and glucose level of the blood. The results showed that low fiber intake increases the glucose level of blood (Bertalina & Rajjani, 2018).

Psyllium is a dietary fiber used to lower blood cholesterol and blood sugar (Ziai et al., 2005). Soluble fiber lessens the absorption of cholesterol into the bloodstream. The intake of soluble fiber lessens

the concentration of LDL cholesterol (Jesch & Carr, 2017). Psyllium is a fiber which reduces the concentration of LDL cholesterol in the bloodstream. Psyllium is a type of fiber which is found in psyllium husk (Slavin, 2013).

The previous study focused on the use of seaweed to prepare the dietary fiber by microbial transformation of *Lactobacillus rhamnosus* and *Lactobacillus plantarum* (Gupta et al., 2011). Studies have shown that the concentration of soluble dietary fiber increases after the action of mixed lactic acid bacteria and the rate of incrimination was 21.1%. After fermentation, the inhibition rate of α -amylase and α -glucosidase increased by 25.9% and 16.6% respectively. While, after fermentation, seaweed increases the concentration of soluble dietary fiber and increases the inhibition of enzymes responsible for digestion. Dietary fibers present in seaweed are responsible for hypoglycemic effects (Zhou et al., 2019).

4.4.1 | Fiber lowers cholesterol

Soluble fiber is also present in brussels sprouts, pears, apples and kidney beans. These soluble fibers reduce the absorption of cholesterol in the bloodstream (Gunness & Gidley, 2010). The soluble fiber present in oatmeal reduces the concentration of low-density lipoprotein cholesterol which is also called bad cholesterol (Othman et al., 2011).

The soluble fiber lowers the cholesterol level by binding it to the small intestine. During passing through the small intestine, the fiber attaches with the cholesterol particles and prevents them from entering the bloodstream and then excreting them from the body through feces (Gunness & Gidley, 2010).

The main biological mechanisms of soluble dietary fibers used to lower the cholesterol are excretion of bile salts by inhibiting their absorption in the small intestine, lowering of stimulation of insulin for hepatic cholesterol synthesis which is the result of reduced glycemic response and effects of soluble dietary fiber. The studies suggest that the micelles of bile salts bind with soluble dietary fibers which inhibit their reabsorption. While different studies have been conducted to research the glycemic effect of soluble dietary fibers, which suggested that the micelles of bile salts can interact with soluble dietary fibers which inhibit their absorption (Tosh & Bordenave, 2020). The three physiochemical mechanisms are proposed together with the suggestion of in vitro experiments to test them (Gunness & Gidley, 2010).

Previous study shown that there is a positive relationship between diet having soluble dietary fibers (guar gum, psyllium, pectin and β -glucan) and lowering of cardiovascular diseases and cholesterol in blood serum (Theuwissen & Mensink, 2008).

4.5 | Hypertension

Fibers consist of indigestible parts of plants which are not digested in our digestive system. Fibers are mainly carbohydrates, and their

role is to keep the digestive system healthy (Mudgil & Barak, 2013). The fiber-rich diet is associated with significant reduction of both systolic and diastolic blood pressure in the people with high blood pressure (Keenan et al., 2002). Studies have shown that fiber-rich foods have many other health benefits like reduction in blood pressure and inflammation (Papanikolaou & Fulgoni, 2008). Soluble fiber can slow down the absorption of sugar and improve the blood glucose level in diabetic patients (Abutair et al., 2016; Aleixandre & Miguel, 2016).

The main factors associated with hypertension and blood pressure are increased population trends of obesity, overweight and aging. The cases of hypertension can be controlled by changing the lifestyle factors like alcohol consumption, poor dietary patterns, sedentary lifestyles, smoking and excess energy intake. Studies have shown that increased intake of fiber up to 7–15 g/d above the normal level is associated with reduction of hypertension and blood pressure. Meta-analysis trials have shown that increased intake of fiber is effective for lowering the blood pressure in elders and obese and elevated hypertension and blood pressure in younger, leaner and normotensives. Soluble fiber including beta-glucan is effective in lowering the blood pressure more than the insoluble fiber obtained from wheat bran. The mechanism of fiber to lower the blood pressure included the reduction of risks of weight gain and obesity, improving the vascular health by lowering the LDL cholesterol and lowering the inflammation via colonic microbiota (Dreher, 2018a, 2018b).

The role of fiber in preventing heart disease is related to its ability to lower cholesterol and blood pressure. Soluble fiber present in oat grain including beta-glucan is responsible for the reduction of cholesterol of plasma. According to many studies, dietary fiber may improve insulin resistance, weight loss and glycemia. Study was conducted on hypertensive rats in which dietary fiber affects the arterial blood pressure. Besides that, some types of fiber improve the arterial blood pressure. Besides that, some types of fiber improve the arterial blood pressure, and it has been proved by the studies on hypertensive and prehypertensive subjects. This study reviewed the researches done to analyze the antihypertensive effects of dietary fiber and their effects on the cardiovascular diseases (Aleixandre & Miguel, 2016).

5 | CONCLUSION

It is concluded that plant-based foods composed of rich nutritive components. These nutritive components play a separate role in human health. From all of these components, fiber is a nondigestible polysaccharide available in plant food like vegetables, fruits, legumes and cereals. Pectin, beta-glucan, mucilage, psyllium, resistant starch and inulin are present in soluble form, whereas cellulose, hemicellulose and lignin are in insoluble form in different varieties of plant food. The fiber vital function is to boost immunity that aids human from viral diseases. The COVID-19 is viral disease spread overall the world. However, the attack of coronavirus are detected lower in those peoples who have strong immune systems.

Fiber played a medicinal role to protect the human body from acute or chronic diseases. Fiber can help to reduce the chances of hypertension and hyperglycemia because a high-fiber diet is the ability to lower cholesterol, blood pressure and blood sugar. It boosts the gut microbes that protect GIT and prevent constipation because it absorbs water and adds bulk to stool.

ACKNOWLEDGEMENT

The authors are highly thankful to Government College University Faisalabad, Pakistan for providing the facilities.

DATA AVAILABILITY STATEMENT

No data available because it is a review article.

ORCID

Muhammad Sajid Arshad  <https://orcid.org/0000-0001-9564-886X>

REFERENCES

- Abutair, A. S., Naser, I. A., & Hamed, A. T. (2016). Soluble fibers from psyllium improve glycemic response and body weight among diabetes type 2 patients (randomized control trial). *Nutrition Journal*, 15(1), 1–7. <https://doi.org/10.1186/s12937-016-0207-4>
- Aleixandre, A., & Miguel, M. (2016). Dietary fiber and blood pressure control. *Food & Function*, 7(4), 1864–1871.
- Alonso-Coello, P., Mills, E. D., Heels-Ansdell, D., Lopez-Yarto, M., Zhou, Q. I., Johanson, J. F., & Guyatt, G. (2006). Fiber for the treatment of hemorrhoids complications: A systematic review and meta-analysis. *American Journal of Gastroenterology*, 101(1), 181–188.
- AL-Rawi, M. M. (2007). Efficacy of oat bran (*Avena sativa* L.) in comparison with atorvastatin in treatment of hypercholesterolemia in albino rat liver. *The Egyptian Journal of Hospital Medicine*, 29(1), 511–521.
- Anderson, J. W., Baird, P., Davis Jr, R. H., Ferreri, S., Knudtson, M., Koraym, A., Waters, V., & Williams, C. L. (2009). Health benefits of dietary fiber. *Nutrition Reviews*, 67(4), 188–205. <https://doi.org/10.1111/j.1753-4887.2009.00189.x>
- Anderson, J. W., Jones, A. E., & Riddell-Mason, S. (1994). Ten different dietary fibers have significantly different effects on serum and liver lipids of cholesterol-fed rats. *The Journal of Nutrition*, 124(1), 78–83. <https://doi.org/10.1093/jn/124.1.78>
- Andersson, K. E., & Hellstrand, P. (2012). Dietary oats and modulation of atherogenic pathways. *Molecular Nutrition & Food Research*, 56(7), 1003–1013.
- Arshad, M. S., Khan, U., Sadiq, A., Khalid, W., Hussain, M., Yasmeen, A., Asghar, Z., & Rehana, H. (2020). Coronavirus disease (COVID-19) and immunity booster green foods: A mini review. *Food Science & Nutrition*, 8(8), 3971–3976. <https://doi.org/10.1002/fsn3.1719>
- Barichella, M., Pacchetti, C., Bolliri, C., Cassani, E., Iorio, L., Pusani, C., Pinelli, G., Privitera, G., Cesari, I., Faierman, S. A., Caccialanza, R., Pezzoli, G., & Cereda, E. (2016). Probiotics and prebiotic fiber for constipation associated with Parkinson disease: An RCT. *Neurology*, 87(12), 1274–1280. <https://doi.org/10.1212/WNL.0000000000003127>
- Barrett, E. M., Foster, S. I., & Beck, E. J. (2020). Whole grain and high-fibre grain foods: How do knowledge, perceptions and attitudes affect food choice? *Appetite*, 149, 104630. <https://doi.org/10.1016/j.appet.2020.104630>
- Beleli, C. A. V., Antonio, M. A. R. G. M., dos Santos, R., Pastore, G. M., & Lomazi, E. A. (2015). Effect of 4' galactooligosaccharide on

- constipation symptoms. *Jornal de Pediatria*, 91(6), 567–573. <https://doi.org/10.1016/j.jpmed.2015.01.010>
- Berger, A., Rein, D., Schäfer, A., Monnard, I., Gremaud, G., Lambelet, P., & Bertoli, C. (2005). Similar cholesterol-lowering properties of rice bran oil, with varied γ -oryzanol, in mildly hypercholesterolemic men. *European Journal of Nutrition*, 44(3), 163–173.
- Bernstein, A. M., Titgemeier, B., Kirkpatrick, K., Golubic, M., & Roizen, M. F. (2013). Major cereal grain fibers and psyllium in relation to cardiovascular health. *Nutrients*, 5(5), 1471–1487. <https://doi.org/10.3390/nu5051471>
- Bertalina, A. H., & Rajiani, I. (2018). Analysis of factors associated with blood sugar levels in type 2 diabetes mellitus patients. *Indian Journal of Public Health Research & Development*, 9(12), 383.
- Beukema, M., Faas, M. M., & de Vos, P. (2020). The effects of different dietary fiber pectin structures on the gastrointestinal immune barrier: Impact via gut microbiota and direct effects on immune cells. *Experimental & Molecular Medicine*, 52(9), 1364–1376.
- Bindels, L. B., Delzenne, N. M., Cani, P. D., & Walter, J. (2015). Towards a more comprehensive concept for prebiotics. *Nature Reviews Gastroenterology & Hepatology*, 12(5), 303. <https://doi.org/10.1038/nrgastro.2015.47>
- Brennan, C. S., & Tudorica, C. M. (2008). Evaluation of potential mechanisms by which dietary fibre additions reduce the predicted glycaemic index of fresh pastas. *International Journal of Food Science & Technology*, 43(12), 2151–2162. <https://doi.org/10.1111/j.1365-2621.2008.01831.x>
- Brownlee, I. A. (2011). The physiological roles of dietary fibre. *Food Hydrocolloids*, 25(2), 238–250. <https://doi.org/10.1016/j.foodhyd.2009.11.013>
- Bunn, D., Jimoh, F., Wilsher, S. H., & Hooper, L. (2015). Increasing fluid intake and reducing dehydration risk in older people living in long-term care: A systematic review. *Journal of the American Medical Directors Association*, 16(2), 101–113. <https://doi.org/10.1016/j.jamda.2014.10.016>
- Bunzel, M., Seiler, A., & Steinhart, H. (2005). Characterization of dietary fiber lignins from fruits and vegetables using the DFRC method. *Journal of Agricultural and Food Chemistry*, 53(24), 9553–9559. <https://doi.org/10.1021/jf0520037>
- Butt, M. S., Shahzadi, N., Sharif, M. K., & Nasir, M. (2007). Guar gum: A miracle therapy for hypercholesterolemia, hyperglycemia and obesity. *Critical Reviews in Food Science and Nutrition*, 47(4), 389–396. <https://doi.org/10.1080/10408390600846267>
- Buttriss, J. L., & Stokes, C. S. (2008). Dietary fibre and health: An overview. *Nutrition Bulletin*, 33(3), 186–200. <https://doi.org/10.1111/j.1467-3010.2008.00705.x>
- Camire, M. E., Kubow, S., & Donnelly, D. J. (2009). Potatoes and human health. *Critical Reviews in Food Science and Nutrition*, 49(10), 823–840. <https://doi.org/10.1080/10408390903041996>
- Chawla, R. P. G. R., & Patil, G. R. (2010). Soluble dietary fiber. *Comprehensive Reviews in Food Science and Food Safety*, 9(2), 178–196. <https://doi.org/10.1111/j.1541-4337.2009.00099.x>
- Costilla, V. C., & Foxx-Orenstein, A. E. (2014). Constipation: Understanding mechanisms and management. *Clinics in Geriatric Medicine*, 30(1), 107–115. <https://doi.org/10.1016/j.cger.2013.10.001>
- Cui, S. W., & Roberts, K. T. (2009). Dietary fiber: Fulfilling the promise of added-value formulations. In S. Kapis, I. T. Norton, & J. B. Ubbink (Eds), *Modern biopolymer science* (pp. 399–448). Academic Press. <https://doi.org/10.1016/B978-0-12-374195-0.00013-6>
- Cui, W., Fan, Y., Wu, W., Zhang, F., Wang, J. Y., & Ni, A. P. (2003). Expression of lymphocytes and lymphocyte subsets in patients with severe acute respiratory syndrome. *Clinical Infectious Diseases*, 37(6), 857–859. <https://doi.org/10.1086/378587>
- Czyzewska-Majchrzak, L., Grzelak, T., Kramkowska, M., Czyzewska, K., & Witmanowski, H. (2014). The use of low-carbohydrate diet in type 2 diabetes-benefits and risks. *Annals of Agricultural and Environmental Medicine*, 21(2), 320–326.
- Daou, C., & Zhang, H. (2014). Functional and physiological properties of total, soluble, and insoluble dietary fibres derived from defatted rice bran. *Journal of Food Science and Technology*, 51(12), 3878–3885.
- Deehan, E. C., Duar, R. M., Armet, A. M., Perez-Munoz, M. E., Jin, M., & Walter, J. (2018). Modulation of the gastrointestinal microbiome with nondigestible fermentable carbohydrates to improve human health. *Bugs as Drugs: Therapeutic Microbes for the Prevention and Treatment of Disease*, 453–483. <https://doi.org/10.1128/978155819705.ch20>
- Deehan, E. C., & Walter, J. (2016). The fiber gap and the disappearing gut microbiome: Implications for human nutrition. *Trends in Endocrinology & Metabolism*, 27(5), 239–242. <https://doi.org/10.1016/j.tem.2016.03.001>
- Demark-Wahnefried, W., Rock, C. L., Patrick, K., & Byers, T. (2008). Lifestyle interventions to reduce cancer risk and improve outcomes. *American Family Physician*, 77(11), 1573–1578.
- Dranca, F., & Oroian, M. (2018). Extraction, purification and characterization of pectin from alternative sources with potential technological applications. *Food Research International*, 113, 327–350. <https://doi.org/10.1016/j.foodres.2018.06.065>
- Dreher, M. L. (2018a). Fiber and hypertension. In M. L. Dreher (Ed), *Dietary fiber in health and disease* (pp. 291–303). Humana Press. https://doi.org/10.1007/978-3-319-50557-2_14
- Dreher, M. L. (2018b). Whole fruits and fruit fiber emerging health effects. *Nutrients*, 10(12), 1833. <https://doi.org/10.3390/nu10121833>
- Emaga, T. H., Andrianaivo, R. H., Wathélet, B., Tchango, J. T., & Paquot, M. (2007). Effects of the stage of maturation and varieties on the chemical composition of banana and plantain peels. *Food Chemistry*, 103(2), 590–600. <https://doi.org/10.1016/j.foodchem.2006.09.006>
- Erkkilä, A. T., Herrington, D. M., Mozaffarian, D., & Lichtenstein, A. H. (2005). Cereal fiber and whole-grain intake are associated with reduced progression of coronary-artery atherosclerosis in postmenopausal women with coronary artery disease. *American Heart Journal*, 150(1), 94–101. <https://doi.org/10.1016/j.ahj.2004.08.013>
- Ewaschuk, J. B., & Dieleman, L. A. (2006). Probiotics and prebiotics in chronic inflammatory bowel diseases. *World Journal of Gastroenterology*, 12(37), 5941. <https://doi.org/10.3748/wjg.v12.i37.5941>
- Feng, N., Guo, X., & Liang, S. (2009). Adsorption study of copper (II) by chemically modified orange peel. *Journal of Hazardous Materials*, 164(2–3), 1286–1292. <https://doi.org/10.1016/j.jhazmat.2008.09.096>
- Fernández-López, J., Sendra-Nadal, E., Navarro, C., Sayas, E., Viuda-Martos, M., & Alvarez, J. A. P. (2009). Storage stability of a high dietary fibre powder from orange by-products. *International Journal of Food Science & Technology*, 44(4), 748–756. <https://doi.org/10.1111/j.1365-2621.2008.01892.x>
- Forootan, M., Bagheri, N., & Darvishi, M. (2018). Chronic constipation: A review of literature. *Medicine*, 97(20), e10631. <https://doi.org/10.1097/MD.0000000000010631>
- Fuentes-Zaragoza, E., Riquelme-Navarrete, M. J., Sánchez-Zapata, E., & Pérez-Álvarez, J. A. (2010). Resistant starch as functional ingredient: A review. *Food Research International*, 43(4), 931–942. <https://doi.org/10.1016/j.foodres.2010.02.004>
- Galanakis, C. M. (2020). The food systems in the era of the coronavirus (COVID-19) pandemic crisis. *Foods*, 9(4), 523. <https://doi.org/10.3390/foods9040523>
- García Ochoa, O. E., Infante, R. B., & Rivera, C. J. (2008). Hacia una definición de fibra alimentaria. In *Anales Venezolanos de Nutrición* (Vol. 21, No. 1, pp. 25–30). Fundación Bengoa.
- García-Mantrana, I., Selma-Royo, M., Alcantara, C., & Collado, M. C. (2018). Shifts on gut microbiota associated to Mediterranean diet adherence and specific dietary intakes on

- general adult population. *Frontiers in Microbiology*, 9, 890. <https://doi.org/10.3389/fmicb.2018.00890>
- Gerhardt, A. L., & Gallo, N. B. (1998). Full-fat rice bran and oat bran similarly reduce hypercholesterolemia in humans. *The Journal of Nutrition*, 128(5), 865–869. <https://doi.org/10.1093/jn/128.5.865>
- Graham, P. H., & Vance, C. P. (2003). Legumes: importance and constraints to greater use. *Plant Physiology*, 131(3), 872–877. <https://doi.org/10.1104/pp.017004>
- Guarner, F. (2005). Inulin and oligofructose: Impact on intestinal diseases and disorders. *British Journal of Nutrition*, 93(S1), S61–S65. <https://doi.org/10.1079/BJN20041345>
- Guarner, F. (2007). Prebiotics in inflammatory bowel diseases. *British Journal of Nutrition*, 98(S1), S85–S89. <https://doi.org/10.1017/S0007114507832958>
- Gunness, P., & Gidley, M. J. (2010). Mechanisms underlying the cholesterol-lowering properties of soluble dietary fibre polysaccharides. *Food & Function*, 1(2), 149–155.
- Guo, Q., Kang, J., Bai, Y., & Xu, F. (2018). Dietary Fiber: Chemistry, structure, and properties. *Journal of Chemistry*, 2018, 1–2. <https://doi.org/10.1155/2018/1328797>
- Gupta, S., Abu-Ghannam, N., & Scannell, A. G. (2011). Growth and kinetics of *Lactobacillus plantarum* in the fermentation of edible Irish brown seaweeds. *Food and Bioprocess Technology*, 89(4), 346–355. <https://doi.org/10.1016/j.fbp.2010.10.001>
- Halton, T. L., Willett, W. C., Liu, S., Manson, J. E., Stampfer, M. J., & Hu, F. B. (2006). Potato and french fry consumption and risk of type 2 diabetes in women. *The American Journal of Clinical Nutrition*, 83(2), 284–290. <https://doi.org/10.1093/ajcn/83.2.284>
- Hecht, J. (2004). *City of light: The story of fiber optics*. Oxford University Press on Demand.
- Holscher, H. D. (2017). Dietary fiber and prebiotics and the gastrointestinal microbiota. *Gut Microbes*, 8(2), 172–184. <https://doi.org/10.1080/19490976.2017.1290756>
- Holscher, H. D., Caporaso, J. G., Hooda, S., Brulc, J. M., Fahey, G. C. Jr, & Swanson, K. S. (2015). Fiber supplementation influences phylogenetic structure and functional capacity of the human intestinal microbiome: Follow-up of a randomized controlled trial. *The American Journal of Clinical Nutrition*, 101(1), 55–64. <https://doi.org/10.3945/ajcn.114.092064>
- Holscher, H. D., Doligale, J. L., Bauer, L. L., Gourineni, V., Pelkman, C. L., Fahey, G. C., & Swanson, K. S. (2014). Gastrointestinal tolerance and utilization of agave inulin by healthy adults. *Food & Function*, 5(6), 1142–1149. <https://doi.org/10.1002/hep.1840200612>
- Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J., Hu, Y. I., Zhang, L. I., Fan, G., Xu, J., Gu, X., Cheng, Z., Yu, T., Xia, J., Wei, Y., Wu, W., Xie, X., Yin, W., Li, H., Liu, M., ... Cao, B. (2020). Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The Lancet*, 395(10223), 497–506. [https://doi.org/10.1016/S0140-6736\(20\)30183-5](https://doi.org/10.1016/S0140-6736(20)30183-5)
- Huang, Y. L., Tsai, Y. H., & Chow, C. J. (2014). Water-insoluble fiber-rich fraction from pineapple peel improves intestinal function in hamsters: Evidence from cecal and fecal indicators. *Nutrition Research*, 34(4), 346–354. <https://doi.org/10.1016/j.nutres.2014.03.001>
- Hughes, J. S., & Swanson, B. G. (1989). Soluble and insoluble dietary fiber in cooked common bean (*Phaseolus vulgaris*) seeds. *Food Structure*, 8(1), 4. <https://digitalcommons.usu.edu/foodmicrostructure/vol8/iss1/4>
- Hullings, A. G., Sinha, R., Liao, L. M., Freedman, N. D., Graubard, B. I., & Loftfield, E. (2020). Whole grain and dietary fiber intake and risk of colorectal cancer in the NIH-AARP Diet and Health Study cohort. *The American Journal of Clinical Nutrition*, 112(3), 603–612. <https://doi.org/10.1093/ajcn/nqaa161>
- Idah, P. A., Ajisegiri, E. S. A., & Yisa, M. G. (2007). Fruits and vegetables handling and transportation in Nigeria. *AU Journal of Technology*, 10(3), 175–183.
- Jesch, E. D., & Carr, T. P. (2017). Food ingredients that inhibit cholesterol absorption. *Preventive Nutrition and Food Science*, 22(2), 67. <https://doi.org/10.3746/pnf.2017.22.2.67>
- Jha, R., Fohse, J. M., Tiwari, U. P., Li, L., & Willing, B. P. (2019). Dietary fiber and intestinal health of monogastric animals. *Frontiers in Veterinary Science*, 6, 48. <https://doi.org/10.3389/fvets.2019.00048>
- Judprasong, K., Tanjor, S., Puwastien, P., & Sungpuag, P. (2011). Investigation of Thai plants for potential sources of inulin-type fructans. *Journal of Food Composition and Analysis*, 24(4–5), 642–649. <https://doi.org/10.1016/j.jfca.2010.12.001>
- Kahlon, T. S., & Chow, F. I. (2000). In vitro binding of bile acids by rice bran, oat bran, wheat bran, and corn bran. *Cereal Chemistry*, 77(4), 518–521. <https://doi.org/10.1094/CHEM.2000.77.4.518>
- Kalyani Nair, K., Kharb, S., & Thompkinson, D. K. (2010). Inulin dietary fiber with functional and health attributes—A review. *Food Reviews International*, 26(2), 189–203. <https://doi.org/10.1080/87559121003590664>
- Kandasamy, P., & Shanmugapriya, C. (2015). Medicinal and nutritional characteristics of fruits in human health. *Journal of Medicinal Plants Studies*, 4(4), 124–131.
- Karabudak, E., Koksul, E., & Macit, M. S. (2019). The relationship between body weight, fiber and fluid intake status and functional constipation in young adults. *Nutrition & Food Science*, 49(1), 129–140.
- Kaur, R., Sharma, M., Ji, D., Xu, M., & Agyei, D. (2020). Structural features, modification, and functionalities of beta-glucan. *Fibers*, 8(1), 1. <https://doi.org/10.3390/fib8010001>
- Keenan, J. M., Pins, J. J., Frazel, C., Moran, A., & Turnquist, L. (2002). Oat ingestion reduces systolic and diastolic blood pressure in patients with mild or borderline hypertension: A pilot trial. *Journal of Family Practice*, 51(4), 369.
- Khalid, W., Arshad, M. S., Yasin, M., Imran, A., & Ahmad, M. H. (2021). Quality characteristics of gamma irradiation and kale leaf powder treated ostrich and chicken meat during storage. *International Journal of Food Properties*, 24(1), 1335–1348. <https://doi.org/10.1080/10942912.2021.1963274>
- Khalid, W., Khalid, M. Z., Aziz, A., Tariq, A., Ikram, A., Rehan, M., Younas, S., Bashir, A., & Fatima, A. (2020). Nutritional composition and health benefits of potato: A review. *Advance Food & Nutrition Science*, 5, 7–16. <https://doi.org/10.21065/25631640.5.7>
- Khan, A. R., Alam, S., Ali, S., Bibi, S., & Khalil, I. A. (2007). Dietary fiber profile of food legumes. *Sarhad Journal of Agriculture*, 23(3), 763.
- Klasing, K. C. (2007). Nutrition and the immune system. *British Poultry Science*, 48(5), 525–537. <https://doi.org/10.1080/00071660701671336>
- Klurfeld, D. M., Davis, C. D., Karp, R. W., Allen-Vercoe, E., Chang, E. B., Chassaing, B., Fahey, G. C., Hamaker, B. R., Holscher, H. D., Lampe, J. W., Marette, A., Martens, E., O'Keefe, S. J., Rose, D. J., Saarela, M., Schneeman, B. O., Slavin, J. L., Sonnenburg, J. L., Swanson, K. S., ... Lynch, C. J. (2018). Considerations for best practices in studies of fiber or other dietary components and the intestinal microbiome. *American Journal of Physiology-Endocrinology and Metabolism*, 315(6), E1087–E1097. <https://doi.org/10.1152/ajpendo.00058.2018>
- Koh, A., De Vadder, F., Kovatcheva-Datchary, P., & Bäckhed, F. (2016). From dietary fiber to host physiology: Short-chain fatty acids as key bacterial metabolites. *Cell*, 165(6), 1332–1345. <https://doi.org/10.1016/j.cell.2016.05.041>
- Kranz, S., Brauchla, M., Slavin, J. L., & Miller, K. B. (2012). What do we know about dietary fiber intake in children and health? The effects of fiber intake on constipation, obesity, and diabetes in children. *Advances in Nutrition*, 3(1), 47–53. <https://doi.org/10.3945/an.111.001362>
- Kumoro, A. C., Alhanif, M., & Wardhani, D. H. (2020). A critical review on tropical fruits seeds as prospective sources of nutritional and bioactive compounds for functional foods development: A case of Indonesian exotic fruits. *International Journal of Food Science*, 2020, 1–15. <https://doi.org/10.1155/2020/4051475>

- Kyrø, C., Skeie, G., Loft, S., Landberg, R., Christensen, J., Lund, E., Nilsson, L. M., Palmqvist, R., Tjønneland, A., & Olsen, A. (2013). Intake of whole grains from different cereal and food sources and incidence of colorectal cancer in the Scandinavian HELGA cohort. *Cancer Causes & Control*, 24(7), 1363–1374.
- Lachman, J. A. R. O. M. È. R., Hamouz, K. A. R. E. L., & Orsák, M. (2005). Red and purple potatoes—a significant antioxidant source in human nutrition. *Chemické Listy*, 99(7), 474–482.
- Lai, C. C., Shih, T. P., Ko, W. C., Tang, H. J., & Hsueh, P. R. (2020). Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and coronavirus disease-2019 (COVID-19): The epidemic and the challenges. *International Journal of Antimicrobial Agents*, 55(3), 105924. <https://doi.org/10.1016/j.ijantimicag.2020.105924>
- Larsson, S. C., Giovannucci, E., Bergkvist, L., & Wolk, A. (2005). Whole grain consumption and risk of colorectal cancer: A population-based cohort of 60 000 women. *British Journal of Cancer*, 92(9), 1803–1807.
- Li, T., Qiu, Z., Zhang, L., Han, Y., He, W., Liu, Z., & Wang, A. (2004). Significant changes of peripheral T lymphocyte subsets in patients with severe acute respiratory syndrome. *The Journal of Infectious Diseases*, 189(4), 648–651. <https://doi.org/10.1086/381535>
- Lin, L., Lu, L., Cao, W., & Li, T. (2020). Hypothesis for potential pathogenesis of SARS-CoV-2 infection—a review of immune changes in patients with viral pneumonia. *Emerging Microbes & Infections*, 9(1), 727–732. <https://doi.org/10.1080/22221751.2020.1746199>
- Lovegrove, A., Edwards, C. H., De Noni, I., Patel, H., El, S. N., Grassby, T., Zielke, C., Ulmius, M., Nilsson, L., Butterworth, P. J., Ellis, P. R., & Shewry, P. R. (2017). Role of polysaccharides in food, digestion, and health. *Critical Reviews in Food Science and Nutrition*, 57(2), 237–253. <https://doi.org/10.1080/10408398.2014.939263>
- Low, D. Y., Williams, B. A., D'Arcy, B. R., Flanagan, B. M., & Gidley, M. J. (2015). In vitro fermentation of chewed mango and banana: Particle size, starch and vascular fibre effects. *Food & Function*, 6(8), 2464–2474.
- Marlett, J. A., Hosig, K. B., Vollendorf, N. W., Shinnick, F. L., Haack, V. S., & Story, J. A. (1994). Mechanism of serum cholesterol reduction by oat bran. *Hepatology*, 20(6), 1450–1457.
- McDougall, G. J., Morrison, I. M., Stewart, D., & Hillman, J. R. (1996). Plant cell walls as dietary fibre: Range, structure, processing and function. *Journal of the Science of Food and Agriculture*, 70(2), 133–150. [https://doi.org/10.1002/\(SICI\)1097-0010\(199602\)70:2<133:AID-JSFA495>3.0.CO;2-4](https://doi.org/10.1002/(SICI)1097-0010(199602)70:2<133:AID-JSFA495>3.0.CO;2-4)
- McIntosh, M., & Miller, C. (2001). A diet containing food rich in soluble and insoluble fiber improves glycemic control and reduces hyperlipidemia among patients with type 2 diabetes mellitus. *Nutrition Reviews*, 59(2), 52–55. <https://doi.org/10.1111/j.1753-4887.2001.tb06976.x>
- McRorie, J. W. Jr (2015). Psyllium is not fermented in the human gut. *Neurogastroenterology and Motility*, 27(11), 1681–1682. <https://doi.org/10.1111/nmo.12649>
- McRorie, J. W., & Fahey, G. C. (2013). A review of gastrointestinal physiology and the mechanisms underlying the health benefits of dietary fiber: Matching an effective fiber with specific patient needs. *Clinical Nursing Studies*, 1(4), 82–92. <https://doi.org/10.5430/cns.v1n4p82>
- McRorie, J. W. Jr, & McKeown, N. M. (2017). Understanding the physics of functional fibers in the gastrointestinal tract: An evidence-based approach to resolving enduring misconceptions about insoluble and soluble fiber. *Journal of the Academy of Nutrition and Dietetics*, 117(2), 251–264. <https://doi.org/10.1016/j.jand.2016.09.021>
- Mei, X., Mu, T. H., & Han, J. J. (2010). Composition and physicochemical properties of dietary fiber extracted from residues of 10 varieties of sweet potato by a sieving method. *Journal of Agricultural and Food Chemistry*, 58(12), 7305–7310. <https://doi.org/10.1021/jf101021s>
- Meier, K. K., Jones, S. M., Kaper, T., Hansson, H., Koetsier, M. J., Karkehabadi, S., & Kelemen, B. (2017). Oxygen activation by Cu LPMOs in recalcitrant carbohydrate polysaccharide conversion to monomer sugars. *Chemical Reviews*, 118(5), 2593–2635. <https://doi.org/10.1021/acs.chemrev.7b00421>
- Messina, V. (2014). Nutritional and health benefits of dried beans. *The American Journal of Clinical Nutrition*, 100(suppl_1), 437S–442S. <https://doi.org/10.3945/ajcn.113.071472>
- Misra, S., Lee, A., & Gensel, K. (2006). Chronic constipation in overweight children. *Journal of Parenteral and Enteral Nutrition*, 30(2), 81–84. <https://doi.org/10.1177/014860710603000281>
- Mohapatra, D., Mishra, S., & Sutar, N. (2010). Banana and its by-product utilization: An overview. *Journal of Scientific and Industrial Research*, 69, 323–329.
- Mongeau, R., & Brooks, S. P. (2001). Chemistry and analysis of lignin. *Food Science and Technology*, 1, 321–374.
- Montagne, L., Pluske, J. R., & Hampson, D. J. (2003). A review of interactions between dietary fibre and the intestinal mucosa, and their consequences on digestive health in young non-ruminant animals. *Animal Feed Science and Technology*, 108(1–4), 95–117. [https://doi.org/10.1016/S0377-8401\(03\)00163-9](https://doi.org/10.1016/S0377-8401(03)00163-9)
- Mudgil, D. (2017). The interaction between insoluble and soluble fiber. In R. A. Samaan (Ed), *Dietary fiber for the prevention of cardiovascular disease* (pp. 35–59). Academic Press. <https://doi.org/10.1016/B978-0-12-805130-6.00003-3>
- Mudgil, D., & Barak, S. (2013). Composition, properties and health benefits of indigestible carbohydrate polymers as dietary fiber: A review. *International Journal of Biological Macromolecules*, 61, 1–6. <https://doi.org/10.1016/j.ijbiomac.2013.06.044>
- Muñoz, L. A., Cobos, A., Diaz, O., & Aguilera, J. M. (2012). Chia seeds: Microstructure, mucilage extraction and hydration. *Journal of Food Engineering*, 108(1), 216–224. <https://doi.org/10.1016/j.jfoodeng.2011.06.037>
- Murphy, O. (2001). Non-polyol low-digestible carbohydrates: Food applications and functional benefits. *British Journal of Nutrition*, 85(S1), S47–S53. <https://doi.org/10.1079/BJN2000261>
- Mwaikambo, L. (2006). Review of the history, properties and application of plant fibres. *African Journal of Science and Technology*, 7(2), 121.
- Nagarajan, N., Morden, A., Bischof, D., King, E. A., Kosztowski, M., Wick, E. C., & Stein, E. M. (2015). The role of fiber supplementation in the treatment of irritable bowel syndrome: A systematic review and meta-analysis. *European Journal of Gastroenterology & Hepatology*, 27(9), 1002–1010. <https://doi.org/10.1097/MEG.0000000000000425>
- Nawirska, A., & Kwaśniewska, M. (2005). Dietary fibre fractions from fruit and vegetable processing waste. *Food Chemistry*, 91(2), 221–225. <https://doi.org/10.1016/j.foodchem.2003.10.005>
- Noack, J., Timm, D., Hospattankar, A., & Slavin, J. (2013). Fermentation profiles of wheat dextrin, inulin and partially hydrolyzed guar gum using an in vitro digestion pretreatment and in vitro batch fermentation system model. *Nutrients*, 5(5), 1500–1510. <https://doi.org/10.3390/nu5051500>
- Othman, R. A., Moghadasian, M. H., & Jones, P. J. (2011). Cholesterol-lowering effects of oat β -glucan. *Nutrition Reviews*, 69(6), 299–309. <https://doi.org/10.1111/j.1753-4887.2011.00401.x>
- Papanikolaou, Y., & Fulgoni, V. L. III (2008). Bean consumption is associated with greater nutrient intake, reduced systolic blood pressure, lower body weight, and a smaller waist circumference in adults: results from the National Health and Nutrition Examination Survey 1999–2002. *Journal of the American College of Nutrition*, 27(5), 569–576. <https://doi.org/10.1080/07315724.2008.10719740>
- Park, K. H., Lee, K. Y., & Lee, H. G. (2013). Chemical composition and physicochemical properties of barley dietary fiber by chemical modification. *International Journal of Biological Macromolecules*, 60, 360–365. <https://doi.org/10.1016/j.ijbiomac.2013.06.024>

- Pashankar, D. S., & Loening-Baucke, V. (2005). Increased prevalence of obesity in children with functional constipation evaluated in an academic medical center. *Pediatrics*, 116(3), e377–e380. <https://doi.org/10.1542/peds.2005-0490>
- Pennington, J. A., & Fisher, R. A. (2009). Classification of fruits and vegetables. *Journal of Food Composition and Analysis*, 22, S23–S31. <https://doi.org/10.1016/j.jfca.2008.11.012>
- Perico, L., Benigni, A., Casiraghi, F., Ng, L. F., Renia, L., & Remuzzi, G. (2020). Immunity, endothelial injury and complement-induced coagulopathy in COVID-19. *Nature Reviews Nephrology*, 17(1), 46–64. <https://doi.org/10.1038/s41581-020-00357-4>
- Platta, A. (2020). Assessment of the consumption of fiber food products by a selected women's group. *Scientific Journal of Gdynia Maritime University*, 114, 17.
- Polonikov, A. (2020). Endogenous deficiency of glutathione as the most likely cause of serious manifestations and death in COVID-19 patients. *ACS Infectious Diseases*, 6(7), 1558–1562. <https://doi.org/10.1021/acsinfectdis.0c00288>
- Post, R. E., Mainous, A. G., King, D. E., & Simpson, K. N. (2012). Dietary fiber for the treatment of type 2 diabetes mellitus: A meta-analysis. *The Journal of the American Board of Family Medicine*, 25(1), 16–23. <https://doi.org/10.3122/jabfm.2012.01.110148>
- Prado, S. B., Beukema, M., Jermendi, E., Schols, H. A., de Vos, P., & Fabi, J. P. (2020). Pectin interaction with immune receptors is modulated by ripening process in papayas. *Scientific Reports*, 10(1), 1–11.
- Qaisrani, T. B., Butt, M. S., Hussain, S., & Ibrahim, M. (2014). Characterization and utilization of psyllium husk for the preparation of dietetic cookies. *International Journal of Modern Agriculture*, 3(3), 81–91.
- Randhawa, M. A., Khan, A. A., Javed, M. S., & Sajid, M. W. (2015). Green leafy vegetables: A health promoting source. In *Handbook of fertility* (pp. 205–220). Academic Press. <https://doi.org/10.1016/B978-0-12-800872-0.00018-4>
- Rao, T. P., & Quartarone, G. (2019). Role of guar fiber in improving digestive health and function. *Nutrition*, 59, 158–169. <https://doi.org/10.1016/j.nut.2018.07.109>
- Reddy, N., & Yang, Y. (2005). Properties and potential applications of natural cellulose fibers from cornhusks. *Green Chemistry*, 7(4), 190–195.
- Rendón-Huerta, J. A., Juárez-Flores, B., Pinos-Rodríguez, J. M., Aguirre-Rivera, J. R., & Delgado-Portales, R. E. (2012). Effects of different sources of fructans on body weight, blood metabolites and fecal bacteria in normal and obese non-diabetic and diabetic rats. *Plant Foods for Human Nutrition*, 67(1), 64–70. <https://doi.org/10.1007/s11130-011-0266-9>
- Reynolds, A., Mann, J., Cummings, J., Winter, N., Mete, E., & Te Morenga, L. (2019). Carbohydrate quality and human health: A series of systematic reviews and meta-analyses. *The Lancet*, 393(10170), 434–445. [https://doi.org/10.1016/S0140-6736\(18\)31809-9](https://doi.org/10.1016/S0140-6736(18)31809-9)
- Rideout, T. C., Harding, S. V., Jones, P. J., & Fan, M. Z. (2008). Guar gum and similar soluble fibers in the regulation of cholesterol metabolism: Current understandings and future research priorities. *Vascular Health and Risk Management*, 4(5), 1023. <https://doi.org/10.2147/vhrm.s3512>
- Roberfroid, M. (1993). Dietary fiber, inulin, and oligofructose: A review comparing their physiological effects. *Critical Reviews in Food Science & Nutrition*, 33(2), 103–148. <https://doi.org/10.1080/10408399309527616>
- Roberfroid, M. B. (2005). Introducing inulin-type fructans. *British Journal of Nutrition*, 93(S1), S13–S25. <https://doi.org/10.1079/BJN20041350>
- Roberfroid, M. B. (2007). Inulin-type fructans: Functional food ingredients. *The Journal of Nutrition*, 137(11), 2493S–2502S. <https://doi.org/10.1093/jn/137.11.2493S>
- Sánchez-Chino, X., Jiménez-Martínez, C., Dávila-Ortiz, G., Álvarez-González, I., & Madrigal-Bujaidar, E. (2015). Nutrient and non-nutrient components of legumes, and its chemopreventive activity: A review. *Nutrition and Cancer*, 67(3), 401–410. <https://doi.org/10.1080/01635581.2015.1004729>
- Sanders, J. M., Monogue, M. L., Jodlowski, T. Z., & Cutrell, J. B. (2020). Pharmacologic treatments for coronavirus disease 2019 (COVID-19): A review. *JAMA*, 323(18), 1824–1836.
- Schieber, A., Stintzing, F. C., & Carle, R. (2001). By-products of plant food processing as a source of functional compounds—recent developments. *Trends in Food Science & Technology*, 12(11), 401–413. [https://doi.org/10.1016/S0924-2244\(02\)00012-2](https://doi.org/10.1016/S0924-2244(02)00012-2)
- Schley, P. D., & Field, C. J. (2002). The immune-enhancing effects of dietary fibres and prebiotics. *British Journal of Nutrition*, 87(S2), S221–S230. <https://doi.org/10.1079/BJN/2002541>
- Schroeder, B. O., Birchenough, G. M., Ståhlman, M., Arike, L., Johansson, M. E., Hansson, G. C., & Bäckhed, F. (2018). Bifidobacteria or fiber protects against diet-induced microbiota-mediated colonic mucus deterioration. *Cell Host & Microbe*, 23(1), 27–40. <https://doi.org/10.1016/j.chom.2017.11.004>
- Sinclair, M. (2011). The use of abdominal massage to treat chronic constipation. *Journal of Bodywork and Movement Therapies*, 15(4), 436–445. <https://doi.org/10.1016/j.jbmt.2010.07.007>
- Slavin, J. (2013). Fiber and prebiotics: Mechanisms and health benefits. *Nutrients*, 5(4), 1417–1435. <https://doi.org/10.3390/nu5041417>
- Soliman, G. A. (2019). Dietary fiber, atherosclerosis, and cardiovascular disease. *Nutrients*, 11(5), 1155. <https://doi.org/10.3390/nu11051155>
- Stewart, M. L., & Schroeder, N. M. (2013). Dietary treatments for childhood constipation: Efficacy of dietary fiber and whole grains. *Nutrition Reviews*, 71(2), 98–109. <https://doi.org/10.1111/nure.12010>
- Sura, S. P., & Christie, J. (2014). Chronic constipation. *Constipation*, 1, 41–65. https://doi.org/10.1007/978-1-4939-0332-0_3
- Theuwissen, E., & Mensink, R. P. (2008). Water-soluble dietary fibers and cardiovascular disease. *Physiology & Behavior*, 94(2), 285–292. <https://doi.org/10.1016/j.physbeh.2008.01.001>
- Thilagavathi, T., Pandiyan, M., Suganyadevi, M., Sivaji, M., Yuvaraj, M., & Sasmitha, R. (2020). Dietary fibre-health benefits. *Biotica Research Today*, 2(6), 519–522.
- Tosh, S. M., & Bordenave, N. (2020). Emerging science on benefits of whole grain oat and barley and their soluble dietary fibers for heart health, glycemic response, and gut microbiota. *Nutrition Reviews*, 78(Suppl. 1), 13–20. <https://doi.org/10.1093/nutrit/nuz085>
- Trinidad, T. P., Mallillin, A. C., Loyola, A. S., Sagum, R. S., & Encabo, R. R. (2010). The potential health benefits of legumes as a good source of dietary fibre. *British Journal of Nutrition*, 103(4), 569–574.
- Trompette, A., Gollwitzer, E. S., Yadava, K., Sichelstiel, A. K., Sprenger, N., Ngom-Bru, C., & Marsland, B. J. (2014). Gut microbiota metabolism of dietary fiber influences allergic airway disease and hematopoiesis. *Nature Medicine*, 20(2), 159–166.
- Veereman, G. (2007). Pediatric applications of inulin and oligofructose. *The Journal of Nutrition*, 137(11), 2585S–2589S. <https://doi.org/10.1093/jn/137.11.2585S>
- Verkerk, R., Schreiner, M., Krumbain, A., Ciska, E., Holst, B., Rowland, I., & Dekker, M. (2009). Glucosinolates in Brassica vegetables: The influence of the food supply chain on intake, bioavailability and human health. *Molecular Nutrition & Food Research*, 53(S2), S219.
- Veronese, N., Solmi, M., Caruso, M. G., Giannelli, G., Osella, A. R., Evangelou, E., Maggi, S., Fontana, L., Stubbs, B., & Tzoulaki, I. (2018). Dietary fiber and health outcomes: An umbrella review of systematic reviews and meta-analyses. *The American Journal of Clinical Nutrition*, 107(3), 436–444. <https://doi.org/10.1093/ajcn/nqx082>
- Villanueva-Suárez, M. J., Mateos-Aparicio, I., Pérez-Cózar, M. L., Yokoyama, W., & Redondo-Cuenca, A. (2019). Hypolipidemic effects of dietary fibre from an artichoke by-product in Syrian hamsters. *Journal of Functional Foods*, 56, 156–162. <https://doi.org/10.1016/j.jff.2019.03.013>

- Vitaglione, P., Napolitano, A., & Fogliano, V. (2008). Cereal dietary fibre: A natural functional ingredient to deliver phenolic compounds into the gut. *Trends in Food Science & Technology*, 19(9), 451–463. <https://doi.org/10.1016/j.tifs.2008.02.005>
- Vos, A. P., M'rabet, L., Stahl, B., Boehm, G., & Garssen, J. (2007). Immunomodulatory effects and potential working mechanisms of orally applied nondigestible carbohydrates. *Critical Reviews™ in Immunology*, 27(2), 97–140. <https://doi.org/10.1615/CritRevImmunol.v27.i2.10>
- Wang, J., Rosell, C. M., & de Barber, C. B. (2002). Effect of the addition of different fibres on wheat dough performance and bread quality. *Food Chemistry*, 79(2), 221–226. [https://doi.org/10.1016/S0308-8146\(02\)00135-8](https://doi.org/10.1016/S0308-8146(02)00135-8)
- Wang, Y.-X., Li, L.-Y., Zhang, T., Wang, J.-Q., Huang, X.-J., Hu, J.-L., Yin, J.-Y., & Nie, S.-P. (2021). Fractionation, physicochemical and structural characterization of polysaccharides from barley water-soluble fiber. *Food Hydrocolloids*, 113, 106539. <https://doi.org/10.1016/j.foodhyd.2020.106539>
- Watzl, B., Girrbaach, S., & Roller, M. (2005). Inulin, oligofructose and immunomodulation. *British Journal of Nutrition*, 93(S1), S49–S55. <https://doi.org/10.1079/BJN20041357>
- Weber, T. K., Toporovski, M. S., Tahan, S., Neufeld, C. B., & de Morais, M. B. (2014). Dietary fiber mixture in pediatric patients with controlled chronic constipation. *Journal of Pediatric Gastroenterology and Nutrition*, 58(3), 297–302. <https://doi.org/10.1097/MPG.0000000000000224>
- Williams, B. A., Grant, L. J., Gidley, M. J., & Mikkelsen, D. (2017). Gut fermentation of dietary fibres: Physico-chemistry of plant cell walls and implications for health. *International Journal of Molecular Sciences*, 18(10), 2203. <https://doi.org/10.3390/ijms18102203>
- Wood, R. J., Fernandez, M. L., Sharman, M. J., Silvestre, R., Greene, C. M., Zern, T. L., & Volek, J. S. (2007). Effects of a carbohydrate-restricted diet with and without supplemental soluble fiber on plasma low-density lipoprotein cholesterol and other clinical markers of cardiovascular risk. *Metabolism*, 56(1), 58–67. <https://doi.org/10.1016/j.metabol.2006.08.021>
- Xu, Z., Shi, L., Wang, Y., Zhang, J., Huang, L., Zhang, C., Liu, S., Zhao, P., Liu, H., Zhu, L. I., Tai, Y., Bai, C., Gao, T., Song, J., Xia, P., Dong, J., Zhao, J., & Wang, F.-S. (2020). Pathological findings of COVID-19 associated with acute respiratory distress syndrome. *The Lancet Respiratory Medicine*, 8(4), 420–422. [https://doi.org/10.1016/S2213-2600\(20\)30076-X](https://doi.org/10.1016/S2213-2600(20)30076-X)
- Yanniotis, S., Petraki, A., & Soumpasi, E. (2007). Effect of pectin and wheat fibers on quality attributes of extruded cornstarch. *Journal of Food Engineering*, 80(2), 594–599. <https://doi.org/10.1016/j.jfoodeng.2006.06.018>
- Yapo, B. M., & Koffi, K. L. (2008). Dietary fiber Components in yellow passion fruit rind—A potential fiber source. *Journal of Agricultural and Food Chemistry*, 56(14), 5880–5883. <https://doi.org/10.1021/jf073247p>
- Young, G. P., Hu, Y., Le Leu, R. K., & Nyskohus, L. (2005). Dietary fibre and colorectal cancer: A model for environment–gene interactions. *Molecular Nutrition & Food Research*, 49(6), 571–584.
- Zhou, L., Zhao, L., Su, Y., & Li, K. (2019). In vitro hypoglycemic potential of dietary fiber from fermented seaweed. *IOP Conference Series: Materials Science and Engineering*, 612(2), 022072.
- Zhu, N., Zhang, D., Wang, W., Li, X., Yang, B., Song, J., & Tan, W. (2020). A novel coronavirus from patients with pneumonia in China, 2019. *New England Journal of Medicine*, 382(8), 727–733. <https://doi.org/10.1056/NEJMoa2001017>
- Zhu, Y., Conklin, D. R., Chen, H., Wang, L., & Sang, S. (2011). 5-Alk (en) ylresorcinols as the major active components in wheat bran inhibit human colon cancer cell growth. *Bioorganic & Medicinal Chemistry*, 19(13), 3973–3982. <https://doi.org/10.1016/j.bmc.2011.05.025>
- Ziai, S. A., Larijani, B., Akhoondzadeh, S., Fakhrzadeh, H., Dastpak, A., Bandarian, F., Rezai, A., Badi, H. N., & Emami, T. (2005). Psyllium decreased serum glucose and glycosylated hemoglobin significantly in diabetic outpatients. *Journal of Ethnopharmacology*, 102(2), 202–207. <https://doi.org/10.1016/j.jep.2005.06.042>
- Zou, J., Chassaing, B., Singh, V., Pellizzon, M., Ricci, M., Fythe, M. D., Kumar, M. V., & Gewirtz, A. T. (2018). Fiber-mediated nourishment of gut microbiota protects against diet-induced obesity by restoring IL-22-mediated colonic health. *Cell Host & Microbe*, 23(1), 41–53. <https://doi.org/10.1016/j.chom.2017.11.003>

How to cite this article: Khalid, W., Arshad, M. S., Jabeen, A., Muhammad Anjum, F., Qaisrani, T. B., & Suleria, H. A. R. (2022). Fiber-enriched botanicals: A therapeutic tool against certain metabolic ailments. *Food Science & Nutrition*, 10, 3203–3218. <https://doi.org/10.1002/fsn3.2920>